

ent-Halimane Diterpenes and a Guaiane Sesquiterpene from *Cladogynos orientalis*

Mayuree Kanlayavattanakul,^{†,‡} Nijisiri Ruangrungsi,^{*,†} Toshiko Watanabe,[‡] Masatoshi Kawahata,[‡] Bruno Therrien,[§] Kentaro Yamaguchi,[§] and Tsutomu Ishikawa[‡]

Department of Pharmacognosy, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand, and Graduate School of Pharmaceutical Sciences and Chemical Analysis Center, Chiba University, Chiba 263-8522, Japan

Received April 6, 2004

Four new *ent*-halimane diterpenes (**1–4**) and one new guaiane sesquiterpene (**5**) were isolated from the CHCl₃ extract of the roots of *Cladogynos orientalis*, together with six known compounds. The structures of compounds **1–5** were established using spectroscopic methods, and the stereochemistry of chettaphanin I (**6**) was confirmed by X-ray crystallography.


Cladogynos orientalis Zipp. ex Span. (syn. *Adenochlaena siamensis* Ridl.) (Euphorbiaceae), commonly known as “Chettaphangki” in Thailand, is the only member of the genus *Cladogynos*.¹ It is distributed sporadically in central and northeast Thailand, and the roots are used as a carminative in Thai folk medicine.² However, very little is known about the chemical constituents of *C. orientalis* and, so far, there have been only two previous phytochemical investigations, which reported the isolation and structure elucidation of two *ent*-halimane diterpenes, chettaphanin I and II,^{3,4} from the roots of this plant. As a part of our chemical studies on Thai medicinal plants, we describe herein the isolation of four new *ent*-halimane diterpenes (**1–4**) and one new guaiane sesquiterpene (**5**), along with six known compounds, chettaphanin I (**6**),^{3,5} chettaphanin II,^{4,5} spathulenol,⁶ cyperenoic acid,^{7,8} taraxerol,^{9,10} and acetoxyaleuritolate,^{11–13} from the CHCl₃ extract of the roots of *C. orientalis*. The absolute stereochemistry of **6** was established by X-ray crystallographic analysis.

Results and Discussion

Among four new diterpenes, the molecular formula of compound **1** was established as C₂₀H₂₄O₄ from its HR-FABMS and NMR data. The IR spectrum showed absorption bands due to a conjugated keto carbonyl (1671 cm⁻¹), a lactone ring (1757, 1276 cm⁻¹), and a furan ring (3122, 1509, 872 cm⁻¹). In the ¹H NMR spectrum, one secondary methyl group at δ_H 0.86 (d, *J* = 7.0 Hz, H₃-17) and two tertiary methyl groups at δ_H 1.07 and 1.32 (each s, H₃-20 and H₃-19) were observed. The signals at δ_H 6.73, 7.41, and 7.95 (1H each, H-14, H-15, and H-16) were characteristic of a β-substituted furan ring. In the ¹³C NMR spectrum, 20 carbon signals appeared including three methyl carbons at δ_C 15.2 (C-17), 16.5 (C-19), and 21.9 (C-20), four furan ring carbons at δ_C 108.7 (C-14), 129.3 (C-13), 144.2 (C-15), and 147.6 (C-16), a lactone carbonyl carbon at δ_C 178.3 (C-18), and a ketone carbon at δ_C 193.6 (C-12). The signal patterns of the ¹³C NMR spectrum were similar to those of methyl 15,16-epoxy-2-ethylenedioxy-12-oxo-5(10),13(16)-14-*ent*-halimatrien-18-oate.⁵ This compound has been obtained as a semisynthetic product from *ent*-halimic acid (a bicyclic diterpene with known absolute configuration). In the case of **1**, there was evidence that a lactone ring was formed between C-2 and C-4, based on HMBC correlations from H₃-19 and H_a-3 to C-18, from H_a-1 and H_b-1 to C-2, and from H-2 to C-4 and C-10. The assignments of each signal were completed by analysis of HMQC, HMBC, and ¹H–¹H COSY NMR correlations (Figure 1). Thus, **1** was assigned as 6-[2-(furan-3-yl)-2-oxoethyl]-1,5,6-trimethyl-10-oxatricyclo[7.2.1.0^{2,7}]dodec-2(7)-en-11-one.

Compound **2** was assigned to have the molecular formula C₂₀H₂₆O₃ from its HRFABMS and NMR data. The IR absorption bands indicated the presence of a lactone ring (1773, 1290 cm⁻¹) and a furan ring (3120, 1459, 873 cm⁻¹). Comparison of the NMR spectra of **1** and **2** (Tables 1 and 2) showed similarities except for the substitution of the keto carbonyl group at C-12 in **1** with a methylene group (δ_C 19.1) in **2**. The examination of 2D NMR data allowed us to deduce that **2** is 6-[2-(furan-3-yl)ethyl]-1,5,6-trimethyl-10-oxatricyclo[7.2.1.0^{2,7}]dodec-2(7)-en-11-one.

Compound **3** was assigned as a reductive cleaved product between the alkoxy bond of the lactone ring in **2** from its HRFABMS and NMR data. The ¹H (Table 1) and ¹³C NMR (Table 2) spectra of **3** were similar to those of **2** except that the C-2 signal was assignable as a methylene group. The IR spectrum showed absorption bands due to a carboxylic acid (3600–2400, 1699 cm⁻¹), and the ¹³C NMR spectrum also supported the presence of a carboxylic acid group (δ_C



* To whom correspondence should be addressed. Tel: +66-2-2188359.

Fax: +66-2-2558227. E-mail: Nijisiri.R@Chula.ac.th.

[†] Chulalongkorn University.

[‡] Graduate School of Pharmaceutical Sciences, Chiba University.

[§] Chemical Analysis Center, Chiba University.




Figure 1. Selected HMBC (curved arrows) and ^1H - ^1H COSY (solid bold lines) correlations for **1**, **4**, and **5**.

Table 1. ^1H NMR Data of Compounds **1**–**5**^a

position	1	2	3	4	5
1	2.33 (dd, 17.9, 2.7, H _a -1) ^b 2.40 (dddd, 17.9, 2.8, 2.5, H _b -1) ^b	2.39–2.45 (m)	1.89–2.02 (m, H _a -1) 2.07–2.17 (m, H _b -1)	5.90 (s)	
2	4.76 (ddd, 6.0, 2.8, 2.7) ^b	4.81 (ddd, 5.5, 2.8, 2.5) ^c	1.74–1.81 (m)		2.10–2.17 (m, H _a -2) 2.43–2.56 (m, H _b -2)
3	1.93 (d, 11.0, H _a -3) ^b 2.13 (dd, 11.0, 6.0, H _b -3) ^b	1.96 (d, 11.0, H _a -3) ^c 2.13 (dd, 11.0, 5.5, H _b -3) ^c	1.64–1.69 (m, H _a -3) 1.89–2.02 (m, H _b -3)	2.38 (d, 16.3, H _a -3) 2.39 (d, 16.3, H _b -3)	1.26–1.32 (m, H _a -3) 1.93–2.01 (m, H _b -3) 2.43–2.56 (m)
4					2.43–2.56 (m)
6	2.10–2.19 (m)	1.99–2.21 (m)	1.34–1.44 (m, H _a -6) 1.89–2.02 (m, H _b -6)	1.89 (dd, 13.3, 4.8, H _a -6) 2.34 (dd, 13.3, 4.8, H _b -6)	1.67–1.77 (m, H _a -6) 2.43–2.56 (m, H _b -6)
7	1.42–1.49 (m, H _a -7) 1.74–1.81 (m, H _b -7)	1.38–1.46 (m, H _a -7) 1.61–1.65 (m, H _b -7)	1.50–1.56 (m)	1.37–1.41 (m, H _a -7) 2.12–2.20 (m, H _b -7)	2.43–2.56 (m)
8	2.01–2.08 (m)	1.67–1.76 (m)	1.74–1.81 (m)	1.92–1.97 (m)	
9					3.97–4.01 (m)
10					1.67–1.77 (m, H _a -9) 1.93–2.01 (m, H _b -9)
11	2.74 (d, 15.5, H _a -11) 2.85 (d, 15.5, H _b -11)	1.58–1.59 (m, H _a -11) 1.67–1.76 (m, H _b -11)	1.64–1.69 (m)	4.80 (s)	
12		1.99–2.21 (m, H _a -12) 2.27–2.35 (m, H _b -12)	2.07–2.17 (m, H _a -12) 2.33–2.40 (m, H _b -12)		4.78 (s, H _a -12) 5.02 (s, H _b -12)
13					1.80 (s)
14	6.73 (dd, 2.0, 1.0)	6.24 (s)	6.26 (dd, 0.8, 0.8)	6.40 (dd, 0.8, 0.8)	0.98 (d, 7.0)
15	7.41 (dd, 2.0, 1.5)	7.33 (dd, 1.5, 1.5)	7.34 (dd, 1.5, 1.5)	7.33 (dd, 1.8, 1.8)	1.06 (d, 7.5)
16	7.95 (dd, 1.5, 0.5)	7.19 (d, 1.0)	7.20 (s)	7.47 (d, 1.0)	
17	0.86 (d, 7.0)	0.88 (d, 7.0)	0.87 (d, 7.0)	0.88 (d, 7.0)	
18					
19	1.32 (s)	1.31 (s)	1.30 (s)	1.42 (s)	
20	1.07 (s)	0.90 (s)	0.86 (s)	1.17 (s)	
OCH ₃				3.54 (s)	
OH					1.57 (s)

^a Chemical shift values are in ppm from CDCl₃ and *J* values (in Hz) are presented in parentheses. The assignments are based on decoupling, HMQC, HMBC, and ^1H - ^1H COSY experiments. ^b Assignment of coupling constants: δ_{H} 1.93 (d, $J_{\text{H}_a-3, \text{H}_b-3} = 11.0$ Hz), 2.13 (dd, $J_{\text{H}_b-3, \text{H}_a-3} = 11.0$ Hz, $J_{\text{H}_b-3, \text{H}-2} = 6.0$ Hz), 2.33 (dd, $J_{\text{H}_a-1, \text{H}_b-1} = 17.9$ Hz, $J_{\text{H}_a-1, \text{H}-2} = 2.7$ Hz), 2.40 (dddd, $J_{\text{H}_b-1, \text{H}_a-1} = 17.9$ Hz, $J_{\text{H}_b-1, \text{H}-2} = 2.8$ Hz, $J_{\text{H}_b-1, \text{H}_b-3} = 2.8$ Hz, $J_{\text{H}_b-1, \text{H}-6} = 2.5$ Hz), 4.76 (ddd, $J_{\text{H}-2, \text{H}_b-3} = 6.0$ Hz, $J_{\text{H}-2, \text{H}_a-1} = 2.7$ Hz, $J_{\text{H}-2, \text{H}_b-1} = 2.8$ Hz). ^c Assignment of coupling constants: δ_{H} 1.96 (d, $J_{\text{H}_a-3, \text{H}_b-3} = 11.0$ Hz), 2.13 (dd, $J_{\text{H}_b-3, \text{H}_a-3} = 11.0$ Hz, $J_{\text{H}_b-3, \text{H}-2} = 5.5$ Hz), 4.81 (ddd, $J_{\text{H}-2, \text{H}_b-3} = 5.5$ Hz, $J_{\text{H}-2, \text{H}_a-1}$ and $\text{H}_b-1 = 2.5$ and 2.8 Hz).

183.1). Thus, **3** was deduced to be 5-[2-(furan-3-yl)ethyl]-1,5,6-trimethyl-1,2,3,4,5,6,7,8-octahydronaphthalene-1-carboxylic acid.

The relative stereochemistry of compounds **1**–**3** could not be completely established by application of NOE experiments. However, it would be reasonable to deduce that the three methyl groups at C-4, C-8, and C-9 in these diterpenes are in a *cis* orientation to one another because of the co-isolation of the structurally related **6** in this study, the absolute stereochemistry of which was determined by X-ray crystallography, as described below.

The molecular formula of compound **4** was analyzed as C₂₁H₂₄O₅ from its HRFABMS and NMR data. The IR spectrum showed absorption bands due to an ester carbonyl group (1763, 1276 cm⁻¹), a conjugated keto carbonyl group (1676 cm⁻¹), and a furan ring (3150, 1458, 920 cm⁻¹). The



^{13}C NMR spectrum was similar to that of chettaphanin I (**6**),⁶ except for the lack of a C-12 keto carbonyl group, instead of the presence of olefinic carbons at δ_{C} 103.2 (C-11) and 146.2 (C-12). Detailed assignments of each signal using 2D NMR data (Figure 1) including NOE correlations (Figure 2) led to a tricyclic system through an enol ether linkage to the structure of **4**. The close proximity of the methyl ester group at C-4 to the H-16 furan ring proton was also indicated by a NOE experiment.

It is reasonable to suppose that the ether bridge between C-5 and C-12 in **4** could be built up by intramolecular hemiacetal formation of the 12-keto group with a *cis*-oriented OH-5 group, as in compound **7**, the C-5 epimer of **6**, followed by dehydration, as shown in Scheme 1. However, the epimer is unknown. The absolute stereochemistry of **6** has been determined by chemical correlation to natural

Table 2. ^{13}C NMR Data of Compounds 1–5^a

carbon	1	2	3	4	5
1	31.6	31.2	25.1	121.8	139.6
2	74.0	74.4	19.5	196.4	33.8
3	41.1	41.2	35.4	45.5	30.8
4	43.6	43.5	47.4	51.4	46.1
5	132.1	133.5	131.0	79.5	140.9
6	22.2	24.5	25.9	31.7	26.0
7	25.5	26.2	26.8	26.5	49.7
8	33.2	32.4	33.3	42.5	68.3
9	40.3	39.9	40.9	41.4	42.0
10	132.4	133.9	136.0	157.7	29.2
11	47.7	37.9	36.5	103.2	148.0
12	193.6	19.1	19.5	146.2	112.3
13	129.3	125.3	125.8	121.3	23.0
14	108.7	110.9	111.0	107.2	20.0
15	144.2	142.7	142.6	143.2	21.6
16	147.6	138.5	138.4	139.5	
17	15.2	15.7	16.0	14.3	
18	178.3	178.8	183.1	173.9	
19	16.5	17.0	22.9	20.1	
20	21.9	21.4	20.8	22.3	
OCH ₃				52.1	

^a Chemical shift values are in ppm from CDCl₃. The assignments are based on decoupling, HMQC, HMBC, and ¹H–¹H COSY experiments.

**Figure 2.** Selected NOE correlations for 4 and 5.**Scheme 1.** Possible Formation of 4 from 7, the C-5 Epimer of 6

ent-halimic acid and by NOE experiments,⁵ in which these functionalities are situated in a *trans* arrangement. Accordingly, the stereochemistry of 6 isolated in this study was re-examined. A single crystal of 6 was prepared carrying CHCl₃ in its molecule by recrystallization from hexane–CHCl₃. The X-ray crystallographic analysis of the CHCl₃-containing crystal (Figure 3) indicated that the reported stereochemistry of 6 including the absolute configuration was correct, in which C-5 is in the *S* configuration.

Compound 5 was analyzed as a sesquiterpene with the molecular formula C₁₅H₂₄O from its HRFABMS and NMR data. The IR spectrum showed an absorption band due to a hydroxyl group (3448 cm⁻¹). The ¹H and ¹³C NMR data of 5 were almost identical to those of the known α -guaiene,¹⁴ except for an additional hydroxyl signal. The downfield shift of the ¹H NMR signal due to H-8 (δ_{H} 3.39) implied 5 to be 8-hydroxy- α -guaiene. The protons and carbons in 5 were completely assigned by analysis of its 2D NMR spectra (Figure 1). The NOE correlation (Figure 2) between

H-8 and H-7 and H-10 indicated that substitutions at those positions were situated in a *cis* orientation to each other. Thus, 5 was identified as (4*S**,7*R**,8*R**,10*S**)-8-hydroxy- α -guaiene.

All isolated products were tested for cytotoxic activity toward a human small cell lung cancer cell line (NCI-H187) and for antituberculosis activity using *M. tuberculosis* H₃₇Ra. However, none of these compounds exhibited significant cytotoxicity (IC₅₀ < 5 $\mu\text{g}/\text{mL}$) or antituberculosis activity (MIC < 12.5 $\mu\text{g}/\text{mL}$).

Experimental Section

General Experimental Procedures. Melting points were determined using a MP-S3 micromelting point hot stage (Yanagimoto). Optical rotations were measured on a JASCO P-1020 polarimeter. UV absorption spectra were measured on a JASCO V-560 UV spectrophotometer. IR spectra were recorded on a JASCO FT/IR-300E spectrometer. ¹H NMR (400, 500 MHz) and ¹³C NMR (100, 125 MHz) spectra were measured on JEOL JNM-ECP400 and JNM-GSX500A NMR spectrometers. Mass spectra were obtained by a JEOL JMS-HX110 (FABMS) spectrometer. TLC was performed using Merck precoated plates (Silica gel 60 F₂₅₄) of 0.25 mm thickness. Silica gel FL100D (Fuji Silysia Chemical Ltd.) was used for column chromatography.

Plant Material. The plant material was collected from the World Biosphere Reserve, Sakaeraj Environmental Research Station, Nakorn-Rachasima Province, Thailand, in October 2002. Authentication was achieved by comparison with a herbarium specimen (BKF No. 28024) at the National Park, Wildlife and Plant Conservation Department, Ministry of Natural Resources and Environment, Bangkok, Thailand. A voucher specimen (NSR 090251) has been deposited at the herbarium of the Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok, Thailand.

Extraction and Isolation. The dried, powdered roots of *C. orientalis* (4.5 kg) were extracted successively with CHCl₃ (5 \times 20 L) and then with MeOH (3 \times 20 L). Removal of the solvent from the extract under reduced pressure gave 208.6 g (CHCl₃) and 227.3 g (MeOH) of dark brown oily residues. The CHCl₃-soluble part (208.6 g) was chromatographed on a silica gel column with hexane–CHCl₃–MeOH (1:0:0 \rightarrow 0:1:0 \rightarrow 0:0:1). The eluted fractions were evaluated by TLC to give eight main fractions. Fraction 2 (18.2 g), eluted with a gradient of hexane–EtOAc (50:1 \rightarrow 0:1), was purified by column chromatography (hexane–EtOAc, 20:1) to furnish compound 5 (50.7 mg). Fraction 3 (6.8 g), eluted with hexane–EtOAc (20:1 \rightarrow 0:1), was rechromatographed on a silica gel column (hexane–CH₂Cl₂, 2:1) to give the following compounds, in order of increasing polarity: spathulenol (62.5 mg), compound 4 (32.7 mg), and compound 3 (3.2 mg). Fraction 4 (16.6 g) was subjected to a silica gel column (hexane–ether, 5:1 \rightarrow 0:1) to give fractions 4-1 to 4-9. Fraction 4-2 (2.1 g) was crystallized from a hexane–CHCl₃ mixture to afford acetoxyleuritolate (162.5 mg). Fraction 4-7 (1.4 g) was crystallized from hexane–CHCl₃ to afford taraxerol (79.0 mg), and the mother liquid was further purified with silica gel column chromatography (hexane–CH₂Cl₂, 1:1) to give chettaphanin II (25.2 mg). Fraction 5 (20.2 g), eluted with hexane–CHCl₃ (1:2 \rightarrow 0:1) and then CHCl₃–EtOAc (1:0 \rightarrow 0:1), was chromatographed over silica gel (CH₂Cl₂–EtOAc, 1:0 \rightarrow 0:1) to give compound 2 (4.1 mg) and compound 1 (33.4 mg). Repeated chromatography of fraction 6 (9.8 g) with hexane–EtOAc (3:1) yielded chettaphanin I (6) (258.0 mg). Fraction 7 (113.9 g) was crystallized from hexane–CHCl₃ to yield cyperenoic acid (295.9 mg).

6-[2-(Furan-3-yl)-2-oxoethyl]-1,5,6-trimethyl-10-oxatricyclo[7.2.1.0^{2,7}]dodec-2(7)-en-11-one (1): pale yellow amorphous solid; mp 103–105 $^{\circ}\text{C}$; [α]_D²³ –151.5 $^{\circ}$ (c 0.017, CHCl₃); UV (MeOH) λ_{max} (log ϵ) 230 (4.13) nm; IR (KBr) ν_{max} 3122, 1757, 1671, 1509, 1276, 872 cm⁻¹; ¹H and ¹³C NMR data, see Tables 1 and 2; HRFABMS *m/z* 329.1727 (calcd for C₂₀H₂₅O₄, [M + H]⁺ 329.1753).




Figure 3. ORTEP drawing of **6**. The chloroform molecule is omitted for clarity.

6-[2-(Furan-3-yl)ethyl]-1,5,6-trimethyl-10-oxatricyclo[7.2.1.0^{2,7}]dodec-2(7)-en-11-one (2): pale yellow oil; $[\alpha]_{\text{D}}^{23}$ -88.6° (c 0.0017, MeOH); UV (MeOH) λ_{max} ($\log \epsilon$) 204 (4.19) nm; IR (film) ν_{max} 3120, 1773, 1459, 1290, 1024, 873 cm^{-1} ; ^1H and ^{13}C NMR data, see Tables 1 and 2; HRFABMS m/z 315.1990 (calcd for $\text{C}_{20}\text{H}_{27}\text{O}_3$, $[\text{M} + \text{H}]^+$ 315.1960).

5-[2-(Furan-3-yl)ethyl]-1,5,6-trimethyl-1,2,3,4,5,6,7,8-octahydronaphthalene-1-carboxylic acid (3): pale yellow oil; $[\alpha]_{\text{D}}^{23}$ -23.2° (c 0.0013, MeOH); UV (MeOH) λ_{max} ($\log \epsilon$) 299 (3.23) nm; IR (film) ν_{max} 3600–2400 (br), 2929, 1699, 1458, 1190, 938 cm^{-1} ; ^1H and ^{13}C NMR data, see Tables 1 and 2; HRFABMS m/z 317.2108 (calcd for $\text{C}_{20}\text{H}_{29}\text{O}_3$, $[\text{M} + \text{H}]^+$ 317.2117).

Methyl 9-(furan-3-yl)-2,7,13-trimethyl-4-oxo-10-oxatricyclo[5.3.3.0^{1,6}]trideca-5,8-diene-2-carboxylate (4): pale yellow oil; $[\alpha]_{\text{D}}^{23}$ $+56.1^\circ$ (c 0.015, MeOH); UV (MeOH) λ_{max} ($\log \epsilon$) 239 (4.19) nm; IR (film) ν_{max} 3150, 1736, 1676, 1458, 1276, 920 cm^{-1} ; ^1H and ^{13}C NMR data, see Tables 1 and 2; HRFABMS m/z 357.1685 (calcd for $\text{C}_{21}\text{H}_{25}\text{O}_5$, $[\text{M} + \text{H}]^+$ 357.1702).

8-Hydroxy- α -guaiene (5): pale yellow oil; $[\alpha]_{\text{D}}^{23}$ -65.1° (c 0.03, MeOH); UV (MeOH) λ_{max} ($\log \epsilon$) 264 (2.99) nm; IR (film) ν_{max} 3448, 3100, 2926, 1457, 1023 cm^{-1} ; ^1H and ^{13}C NMR data, see Tables 1 and 2; HRFABMS m/z 259.1487 (calcd for $\text{C}_{15}\text{H}_{24}\text{O}_2$, $[\text{M} + \text{K}]^+$ 259.1464).

X-ray data for Chettaphanin I (6)·CHCl₃:¹⁵ colorless prisms, mp 157–158 °C by crystallization of **6** from hexane–CHCl₃ and selected for data collection. Crystal data: $\text{C}_{22}\text{H}_{29}\text{Cl}_3\text{O}_6$, $M = 495.80$ g mol⁻¹, orthorhombic, $P2_12_12_1$, $a = 7.338(3)$ Å, $b = 11.777(5)$ Å, $c = 26.354(12)$ Å, $U = 2277.5(18)$ Å³, $T = 173$ K, $Z = 4$, $\mu(\text{Mo K}\alpha) = 0.439$ mm⁻¹, 5438 reflections measured, 2094 were unique ($R_{\text{int}} = 0.1894$) and used in all calculations. The final $wR(F^2)$ was 0.3330 (all data). The data were measured using a Bruker SMART CCD diffractometer, using Mo K α graphite-monochromated radiation ($\lambda = 0.71073$ Å). The structures were solved by direct methods using the program SHELXS-97.¹⁶ The refinement and all further calculations were carried out using SHELXL-97. The H atoms were included in calculated positions and treated as riding atoms using the SHELXL default parameters. The non-H atoms were refined anisotropically, using weighted full-matrix least-squares on F^2 . Figure 3 is drawn with ORTEP.¹⁷

Cytotoxicity Assays.¹⁸ Cytotoxicity was assessed using the sulforhodamine B (SRB) assay using human tumor cell lines of NCI-H187 (small cell lung cancer). The cells were incubated at 37 °C for 72 h, at which time the SRB was added. The results are expressed as an IC₅₀ ($\mu\text{g/mL}$), and the reference substance was ellipticine.

Antituberculosis Assays.¹⁹ Antituberculosis activity was performed by a microplate alamar blue assay. *M. tuberculosis* H₃₇Ra was used as a tested microorganism. The MICs of the tested compounds were measured in $\mu\text{g/mL}$, and the reference substances were isoniazid and kanamycin sulfate.

Acknowledgment. M.K. would like to acknowledge financial support from the Thailand Research Fund (TRF) through the Royal Golden Jubilee Ph.D. Program (RGJ scholarship) and the Association of International Education, Japan (ALEJ). We are grateful to Prof. Dr. M. Nagai (Hoshi University) for the gift of an authentic sample of taraxerol and the Bioassay Research Facility of BIOTEC, NSTDA, Science Park, Thailand, for conducting biological activity tests.

Supporting Information Available: X-ray data for chettaphanin I. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- Smitinand, T. *Thai Plant Names (Botanical Names-Vernacular Names)*; Funny Publishing: Bangkok, 1990; p 7.
- Shaw, H. K. *Kew Bull.* **1972**, *26*, 191–233.
- Sato, A.; Kurabayashi, M.; Nagahori, H.; Ogiso, A. *Tetrahedron Lett.* **1970**, *11*, 1095–1098.
- Sato, A.; Kurabayashi, M.; Ogiso, A.; Mishima, H. *Tetrahedron Lett.* **1971**, *12*, 839–842.
- Marcos, I. S.; Hernandez, F. A.; Sexmero, M. J.; Basabe, D. P.; Pedrero, A. B.; Garcia, N.; Urones, J. G. *Tetrahedron* **2003**, *59*, 685–694.
- Inagaki, F.; Abe, A. *J. Chem. Soc., Perkin Trans. 2* **1985**, 1773–1778.
- Boonyarathanakornkit, L.; Che, C. T.; Fong, H. H. S.; Farnsworth, N. R. *Planta Med.* **1988**, *74*, 61–63.
- Jacobs, H.; Lachmansing, S. S.; Ramdayal, F. *J. Nat. Prod.* **1987**, *50*, 835–842.
- Sakurai, N.; Yaguchi, Y.; Inoue, T. *Phytochemistry* **1987**, *26*, 217–219.
- The ^{13}C NMR chemical shifts of C-10 and C-12 of taraxerol were confirmed as δ_{C} 35.8 and 37.8, because these were due to quaternary and methylene carbons, respectively, in a DEPT experiment. In a previous report,⁹ these assignments were transposed.
- Carpenter, R. C.; Sitheeswaran, S.; Sultanbawa, M. U. S.; Ternai, B. *Org. Magn. Reson.* **1980**, *14*, 462–465.
- Carpenter, R. C.; Sitheeswaran, S.; Sultanbawa, M. U. S.; Balasubramiam, S. *Phytochemistry* **1980**, *19*, 1171–1174.
- The ^{13}C NMR chemical shifts of C-22 and C-29 of acetoxyleuritolate were confirmed as δ_{C} 33.3 and 31.8, because these were due to methylene and methyl carbons, respectively, in a DEPT experiment. In a previous report,¹¹ these assignments were transposed.
- Rakotonirainy, O.; Gaydou, E. M.; Faure, R.; Bombarda, I. *J. Essent. Oil Res.* **1997**, *9*, 321–327.
- The R value (0.12 for $I > 2\sigma(I)$, 0.23 for all data) is larger than usual. This may result from a significant portion of weak data in the data set. The confirmation of the absolute configuration of **6** is, however, identified within experimental error (absolute parameter 0.2(2)). Crystallographic data (excluding structure factors) for **6** have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication number CCDC-232426. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK [fax: +44(0)-1223-336033 or www.ccdc.cam.ac.uk/data_request/cif or e-mail: data_request@ccdc.cam.ac.uk].
- Sheldrick, G. M. *SHELXS-97-Program for Crystal Structure Solution*; University of Göttingen: Göttingen, Germany, 1997.
- Farrugia, L. J. *J. Appl. Crystallogr.* **1997**, *30*, 565.
- Skehan, P.; Storeng, R.; Scudiero, D.; Monks, A.; McMahon, J.; Vistica, D.; Warren, J. T.; Bokesch, H.; Kenney, S.; Boyd, M. R. *J. Natl. Cancer Inst.* **1990**, *82*, 1107–1112.
- Collins, L. A.; Franzblau, S. G. *Antimicrob. Agents Chemother.* **1997**, *41*, 1004–1009.