

## Brominated Cyclodipeptides from the Marine Sponge *Geodia barretti* as Selective 5-HT Ligands

Erik Hedner,<sup>†,‡</sup> Martin Sjögren,<sup>†,‡</sup> Per-Anders Frändberg,<sup>§</sup> Tobias Johansson,<sup>§</sup> Ulf Göransson,<sup>‡</sup> Mia Dahlström,<sup>‡</sup> Per Jonsson,<sup>‡</sup> Fred Nyberg,<sup>§</sup> and Lars Bohlin<sup>\*‡</sup>

Division of Pharmacognosy, Department of Medicinal Chemistry, Biomedical Centre, Uppsala University, PO Box 574, SE-751 23 Uppsala, Sweden, Division of Biological Research on Drug Dependence, Department of Pharmaceutical Biosciences, Biomedical Centre, Uppsala University, PO Box 574, SE-751 23 Uppsala, Sweden, and Department of Marine Ecology, Tjärnö Marine Biological Laboratory, Göteborg University, SE-452 96 Strömstad, Sweden

Received April 20, 2006

The brominated cyclodipeptides baretin (cyclo[(6-bromo-8-tryptophan)arginine]) and 8,9-dihydrobaretin (cyclo[(6-bromotryptophan)arginine]) isolated from the marine sponge *Geodia barretti* have previously been shown to inhibit settlement of barnacle larvae in a dose-dependent manner in concentrations ranging from 0.5 to 25  $\mu\text{M}$ . To further establish the molecular target and mode of action of these compounds, we investigated their affinity to human serotonin receptors. The tryptophan residue in the baretins resembles that of endogenous serotonin [5-hydroxytryptamine]. A selection of human serotonin receptors, including representatives from all subfamilies (1–7), were transfected into HEK-293 cells. Baretin selectively interacted with the serotonin receptors 5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, and 5-HT<sub>4</sub> at concentrations close to that of endogenous serotonin, with the corresponding  $K_i$  values being 1.93, 0.34, and 1.91  $\mu\text{M}$ , respectively. 8,9-Dihydrobaretin interacted exclusively with the 5-HT<sub>2C</sub> receptor with a  $K_i$  value of 4.63  $\mu\text{M}$ ; it failed to show affinity to 5-HT<sub>2A</sub> and 5-HT<sub>4</sub>, indicating that the double bond between the tryptophan and arginine residue plays an important role in the interaction with the receptor proteins.

Interactions among organisms have led to the evolution of bioactive chemical compounds selected for diverse purposes such as deterring fouling and feeding, capturing prey, and combating invasive pests and pathogens. The production of bioactive compounds by plants, animals, and microorganisms has long been exploited in the search for drug candidates or for compounds to serve as leads in drug development.<sup>1,2</sup> Traditionally, such bio-prospecting for drug candidates has focused on terrestrial microorganisms and plants; the equivalent research in marine systems is in its infancy, but the much larger diversity of major lineages in the sea promises a wealth of new molecular structures with as yet unknown functions.

In the ocean, sessile sponges have proved to be a rich source of bioactive compounds, many of which are believed to constitute a chemical defense against predators or foulers aimed at protecting the body surface.<sup>3</sup> We have previously reported on the production of two brominated cyclodipeptides in the marine sponge *Geodia barretti* Bowerbank (family Geodiidae, class Demospongiae, order Astrophorida), compounds that strongly inhibit the settlement of the barnacle larva of *Balanus improvisus* and the blue mussel *Mytilus edulis*.<sup>4,5</sup> These two cyclodipeptides, baretin (87/13 mixture of *Z/E* isomers) and 8,9-dihydrobaretin, have both been isolated, elucidated, and synthesized by our group<sup>5,6</sup> and, in the case of baretin, by others.<sup>7</sup> Baretin and 8,9-dihydrobaretin both inhibit settlement of barnacle larvae of *B. improvisus* in a dose-dependent manner in concentrations ranging from 0.5 to 25  $\mu\text{M}$ . Their respective EC<sub>50</sub> values are 0.9 (baretin) and 7.9  $\mu\text{M}$  (8,9-dihydrobaretin).<sup>5</sup> The dose-dependence of settlement inhibition is indicative of a specific molecular target in the barnacle larva. In order to further establish the mode of action of the baretins, we investigated their affinity to a wide range of human serotonin receptors from all subfamilies (1–7). Serotonin receptors were chosen on the basis of the chemical structures of the baretins; like serotonin, the baretins carry an indole nucleus by way of their tryptophan residue. However, serotonin is hydroxylated in the

5-position, whereas baretin and 8,9-dihydrobaretin are substituted with a bromine atom in the 6-position. This brominated tryptophan residue can also be found as a 6-Br-Trp moiety in the 41-amino-acid peptide  $\sigma$ -conotoxin, a polypeptide that selectively inhibits the 5-HT<sub>3</sub> receptor through competitive antagonism.<sup>8</sup> The structural similarities of baretin and 8,9-dihydrobaretin to ligands with high affinity for 5-HT receptors, such as the drug tegaserod, which was recently approved for treatment of irritable bowel syndrome (IBS),<sup>9</sup> also prompted us to investigate a possible interaction with the serotonergic system. Here, we describe the affinity and selectivity of the marine cyclodipeptides baretin and 8,9-dihydrobaretin to serotonin receptors.

### Results and Discussion

Of the panel of receptors tested, covering all the major 5-HT subfamilies, the HEK-293 cell membranes expressing 5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, and 5-HT<sub>4</sub> receptors showed receptor–ligand affinity with baretin below a concentration of 10  $\mu\text{M}$  (Table 1). Baretin displaced the radioligands ([*N*-methyl-<sup>3</sup>H]LSD for 5-HT<sub>2A</sub>, <sup>2C</sup>; [*N*-methyl-<sup>3</sup>H]GR113808 for 5-HT<sub>4</sub>) with dose-responsive kinetics. The affinity constant  $K_i$  for baretin at the 5-HT<sub>2A</sub> receptor was determined to be 1.93  $\mu\text{M}$  (Figure 2). Baretin does not have the high affinity to 5-HT<sub>2A</sub> as the more selective ligand methysergide displays, but is selective enough to bind nearly as good as endogenous 5-HT (0.69  $\mu\text{M}$ ). In the case of 5-HT<sub>2C</sub>, both baretin and 8,9-dihydrobaretin were able to displace [*N*-methyl-<sup>3</sup>H]LSD, producing  $K_i$  values of 0.34 and 4.63  $\mu\text{M}$ , respectively (Figure 3). Once again methysergide showed a better affinity ( $K_i = 2.5$  nM) and the displacement with 5-HT gained a 10-fold lower  $K_i$  than for baretin. Competing with [*N*-methyl-<sup>3</sup>H]GR113808 for binding to the 5-HT<sub>4</sub> receptor, baretin showed a  $K_i$  of 1.91  $\mu\text{M}$ , where 5-HT had a  $K_i$  of 0.50  $\mu\text{M}$ . 8,9-Dihydrobaretin was not able to displace the radioligand at the 5-HT<sub>4</sub> receptor. Tegaserod was introduced as a selective ligand and produced a  $K_i$  of 0.031  $\mu\text{M}$ , approximately 10 times better than 5-HT, 100 times better than baretin.

Taken together, baretin clearly has its advantages at the 5-HT<sub>2C</sub> receptor, producing a selectivity ratio of 5.68 (5-HT<sub>2A</sub>/5-HT<sub>2C</sub>) between the two 5-HT<sub>2</sub> receptor subtypes included in the study. The small differences between baretin and 8,9-dihydrobaretin

\* To whom correspondence should be addressed. Tel: +46-18-4714492. Fax: +46-18-509101. E-mail: lars.bohlin@fkog.uu.se.

<sup>†</sup> These authors contributed equally.

<sup>‡</sup> Department of Medicinal Chemistry, Uppsala University.


<sup>§</sup> Department of Pharmaceutical Biosciences, Uppsala University.

<sup>‡</sup> Department of Marine Ecology, Göteborg University.

**Table 1.** Affinities of Baretin, 8,9-Dihydrobaretin, 5-Hydroxytryptamine, Methysergide (5-HT<sub>2A,2C</sub>), and Tegaserod (5-HT<sub>4</sub>) to the 5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, and 5-HT<sub>4</sub> Receptors Expressed in HEK-293 Cell Membranes (data presented as mean ± SEM with three independent experiments performed in triplicate)

receptor <sup>a</sup>	<i>K<sub>i</sub></i> [ $\mu$ M]			
	baretin	8,9-dihydrobaretin	5-hydroxytryptamine	selective ligand <sup>b</sup>
5-HT <sub>1A</sub>	> 10	> 10		
5-HT <sub>1D</sub>	> 10	> 10		
5-HT <sub>2A</sub>	1.93 ± 0.59	> 10	0.69 ± 0.1	0.01 ± 0.003
5-HT <sub>2C</sub>	0.34 ± 0.14	4.63 ± 0.13	0.020 ± 0.004	0.0025 ± 0.001
5-HT <sub>3A</sub>	> 10	> 10		
5-HT <sub>4</sub>	1.91 ± 0.12	> 10	0.50 ± 0.07	0.031 ± 0.01
5-HT <sub>5A</sub>	> 10	> 10		
5-HT <sub>6</sub>	> 10	> 10		
5-HT <sub>7A</sub>	> 10	> 10		


<sup>a</sup> [1,2-<sup>3</sup>H]5-Carboxamidotryptamine used for 5-HT<sub>1A,1D,5A,7A</sub>; [*N*-methyl-<sup>3</sup>H]LSD used for 5-HT<sub>2A,2C,6</sub>; [9-methyl-<sup>3</sup>H]BRL-43694 used for 5-HT<sub>3A</sub>; [*N*-methyl-<sup>3</sup>H]GR113808 used for 5-HT<sub>4</sub>. <sup>b</sup> Methysergide (5-HT<sub>2A,2C</sub>); tegaserod (5-HT<sub>4</sub>).

**Figure 1.** Chemical structures of baretin (1), 8,9-dihydrobaretin (2), and tegaserod (3).

greatly affected the affinity. It has proven difficult to develop selective ligands within the 5-HT<sub>2</sub> receptor subfamily. Baretin, showing a clear selectivity for the 5-HT<sub>2C</sub> receptor over the 5-HT<sub>2A</sub> receptor, could therefore be of interest for further investigation.


The cell lines were subjected to saturation analysis with a high-affinity radioligand to determine the receptor expression levels. Saturation of transfected HEK-293 cell membranes is due to receptors being occupied with the radioligand. When saturation occurs, all the receptors are occupied and no further specific binding site is available. The saturation experiments were performed in order to confirm that we have a robust cellular system that expresses the desired receptor and that this receptor binds the radioligand used in the competition-binding assays. The straight line in the three Scatchard analyses indicates that the radioligands [*N*-methyl-<sup>3</sup>H]-LSD and [*N*-methyl-<sup>3</sup>H]GR113808 bind to a single site on the transfected HEK-293 cell membrane as well as that the binding occurs with a single affinity to the receptor (Figures 2–4). *B*<sub>max</sub> expresses the number of available receptors/mg protein, which is illustrated by the Scatchard curve's intersection of the *x*-axis. *K*<sub>D</sub> is the value of free radioligand at half *B*<sub>max</sub> and expresses the affinity of the radioligand to the cell receptor: [*N*-methyl-<sup>3</sup>H]LSD bound to 5-HT<sub>2A</sub> receptors (*B*<sub>max</sub> = 6.17 fmol mg protein<sup>-1</sup> and *K*<sub>D</sub> = 21.17 nM) (Figure 2), [*N*-methyl-<sup>3</sup>H]LSD bound to 5-HT<sub>2C</sub> receptors (*B*<sub>max</sub> = 4.50 fmol mg protein<sup>-1</sup> and *K*<sub>D</sub> = 52.84 nM) (Figure 3), [*N*-methyl-<sup>3</sup>H]GR113808 bound to 5-HT<sub>4</sub> receptors (*B*<sub>max</sub> = 8.04 fmol mg protein<sup>-1</sup> and *K*<sub>D</sub> = 15.81 nM) (Figure 4). No specific radioligand binding was detected in the parental cell lines (data not shown).

We have thus shown that the marine natural product baretin specifically interacts with 5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, and 5-HT<sub>4</sub> receptors, while 8,9-dihydrobaretin interacts with the 5-HT<sub>2C</sub> receptor (Figures 2–4; Table 1). Among the selected 5-HT receptor subtypes, 8,9-


**Figure 2.** (A) Displacement of [*N*-methyl-<sup>3</sup>H]LSD on the serotonin subreceptor 5-HT<sub>2A</sub> by (●) baretin, (○) 8,9-dihydrobaretin, (■) 5-hydroxytryptamine, and (□) methysergide. (B) Saturation analysis of [*N*-methyl-<sup>3</sup>H]LSD binding to HEK-293 cell membranes expressing the human 5-HT<sub>2A</sub> receptor. Data points are triplicate values.

dihydrobaretin displayed affinity only for 5-HT<sub>2C</sub> receptors (Figure 2; Table 1), and the difference in affinity to the 5-HT<sub>2C</sub> receptor between baretin and 8,9-dihydrobaretin was approximately 10-fold (Table 1). We suggest that the small difference in chemical backbone between the two substances, a double bond in the tryptamine residue, may explain the considerable differences between them in affinity to the serotonin 5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, and 5-HT<sub>4</sub> receptors; the double bond in baretin causes a more rigid steric orientation of the bromotryptophan residue, resulting in a better fit of baretin in the receptor-binding pocket.

The bromine substituent is another chemical constituent that may play a profound role in the interaction between the cyclopeptides and the serotonin receptor. The importance of halogenated substituents for serotonin analogues has recently been highlighted in structure–activity studies. Halogenation not only accounted for drastic improvements in affinity but also produced changes in selectivity between serotonin subtypes.<sup>10,11</sup> However, further conformational and structure–activity studies of baretin and 8,9-dihydrobaretin are needed to elucidate how the double bond and the bromination influence the interactions with the 5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, and 5-HT<sub>4</sub> receptors.



**Figure 3.** (A) Displacement [N-methyl-<sup>3</sup>H]LSD on the serotonin subreceptor 5-HT<sub>2C</sub> by (●) baretтин, (○) 8,9-dihydrobaretтин, (■) 5-hydroxytryptamine, and (□) methysergide. (B) Saturation analysis of [N-methyl-<sup>3</sup>H]LSD binding to HEK-293 cell membranes expressing the human 5-HT<sub>2C</sub> receptor. Data points are triplicate values.



**Figure 4.** (A) Displacement of [N-methyl-<sup>3</sup>H]GR113808 on the serotonin subreceptor 5-HT<sub>4</sub> by (●) baretтин, (○) 8,9-dihydrobaretтин, (■) 5-hydroxytryptamine, and (□) tegaserod. (B) Saturation analysis of [N-methyl-<sup>3</sup>H]GR113808 binding to HEK-293 cell membranes expressing the human 5-HT<sub>4</sub> receptor. Data points are triplicate values.

The structure–activity relationship of baretтин and 8,9-dihydrobaretтин in the receptor–ligand binding assay, i.e., the affinities of the compounds for the 5-HT<sub>2C</sub> receptor, corresponds closely to the difference in activity found in the in vivo assay of settling larvae of *B. improvisus*.<sup>5</sup> Several serotonin receptors have been cloned and functionally characterized in invertebrates. Receptors related in DNA sequence, which activate intracellular signal pathways

identical to mammalian 5-HT<sub>1</sub> and 5-HT<sub>2</sub> receptors, have been found in *Drosophila melanogaster*, *Caenorhabditis elegans*, and the pond snail *Lymnaea stagnalis*.<sup>12</sup> In *D. melanogaster*, a receptor corresponding to the mammalian 5-HT<sub>7</sub> receptor has also been found.<sup>13,14</sup> The functional roles of 5-HT<sub>2</sub> receptors in invertebrates such as arthropods need to be further defined. It has been suggested that 5-HT<sub>2</sub> receptors are of particular importance in *D. melanogaster* embryogenesis.<sup>13</sup> Cypris larvae of the barnacle, *B. amphitrite*, show DNA sequences homologous to a human 5-HT<sub>1A</sub> receptor.<sup>15</sup> However, this barnacle serotonergic G protein-coupled receptor has not been functionally cloned, nor has its pharmacology been fully established. However, a 5-HT<sub>2</sub> receptor has recently been cloned and functionally characterized from another crustacean, the spiny lobster, *Panulirus interruptus*. This receptor is widely expressed in the gastrointestinal nervous system, where it is involved in the neurohormonal modulation of the gastrointestinal ganglion.<sup>16</sup> We suggest that the primary molecular target of the baretтins in cyprids of *B. improvisus* is the 5-HT<sub>2</sub> receptor, but further studies are needed to confirm the presence of this receptor in barnacles. It is plausible that the marine sponge *G. barretti* has evolved the ability to produce defense compounds that deter foulers and predators by interacting with serotonergic functions in these organisms; the sponge lacks a nervous system and so may not itself be affected by the release of these compounds.

In addition to interactions with 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> receptors, baretтин also displayed significant affinity for the 5-HT<sub>4</sub> receptor, with a  $K_i$  value of 1.91  $\mu$ M (Figure 4; Table 1). 8,9-Dihydrobaretтин on the other hand lacked affinity for the 5-HT<sub>4</sub> receptor. No invertebrate 5-HT<sub>4</sub> receptor has yet been cloned and functionally characterized, and so the importance of baretтин's affinity for 5-HT<sub>4</sub> receptors in the cyprid settlement assay remains to be elucidated.

Tegaserod (**3**) is a drug that has recently been approved for treatment of irritable bowel syndrome (IBS).<sup>17</sup> We selected it as a control due to a certain degree of similarity in chemical structure to baretтин and 8,9-dihydrobaretтин; in particular, the position of the double bond relative to the tryptophan ring in tegaserod is similar to that in baretтин. Tegaserod has a high affinity to 5-HT<sub>4</sub> as well as affinity to all the 5-HT<sub>2</sub> receptors.<sup>9,18</sup>

Neither baretтин nor 8,9-dihydrobaretтин displayed any significant interaction with the 5-HT<sub>3</sub> receptor. This was somewhat unexpected since England et al. (1998) have previously reported that  $\sigma$ -conotoxin from the cone snail *Conus geographus*, a peptide bearing a 6-bromotryptophan residue like the baretтins, caused an inactivation of 5-HT<sub>3</sub> receptors transfected in HEK-293 cells. Additionally,  $\sigma$ -conotoxin did not interact with the serotonin receptors 5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, and 5-HT<sub>4</sub>.<sup>8</sup> The similarities of the 6-bromo-substituted tryptophan residue in  $\sigma$ -conotoxin with baretтин and 8,9-dihydrobaretтин is not reflected in affinity for the 5-HT<sub>3</sub> receptor.

In conclusion, we have identified the marine cyclodipeptides baretтин and 8,9-dihydrobaretтин as selective serotonin receptor ligands. This may prove useful in further defining the functional roles of 5-HT receptors in invertebrates. Moreover, the small difference in chemical backbone between baretтин and 8,9-dihydrobaretтин results in striking differences in affinity to the human 5-HT receptors. This suggests a possible function for these compounds as templates to provide clues in drug discovery research aimed at disease states associated with the serotonergic system.

## Experimental Section

**Isolation of Baretтин and 8,9-Dihydrobaretтин.** The cyclodipeptides baretтин (cyclo[(6-bromo-8-tryptophan)arginine]) and 8,9-dihydrobaretтин (cyclo[(6-bromotryptophan)arginine]) were isolated from a specimen of *Geodia barretti* collected in the Swedish Koster Fjord, 1 nautical mile from Tjörnö Marine Biological Laboratory (58°53' N, 11°8' E). The isolation and structure elucidation of the compounds were conducted as outlined in our previous publication.<sup>5</sup>

**Chemicals.** [N-methyl-<sup>3</sup>H]LSD, [9-methyl-<sup>3</sup>H]BRL-43694, and [1,2-<sup>3</sup>H]5-carboxamidotryptamine were obtained from Perkin-Elmer (Boston,

MA). [*N*-methyl-<sup>3</sup>H]GR113808 was obtained from Amersham Bioscience (Buckinghamshire, UK). Tegaserod (**3**) was obtained from APIN Chemical Ltd. (Oxon, UK). All other chemicals used were obtained from Sigma-Aldrich (St. Louis, MO).

**Cell Membrane Preparation.** Membranes were prepared from human embryonic kidney-293 cells (HEK-293) transfected with human serotonin receptors. The following serotonin receptor subtypes were prepared from a glycerol stock (Guthrie, Sayre, PA): 5-HT<sub>1A</sub>, 5-HT<sub>1D</sub>, 5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, 5-HT<sub>3A</sub>, 5-HT<sub>4</sub>, 5-HT<sub>5A</sub>, 5-HT<sub>6</sub>, and 5-HT<sub>7A</sub>. Plasmids were purified according to QIAGEN Highspeed Plasmid Purification Maxi Kit protocol. HEK-293 cells were maintained in a 5% CO<sub>2</sub> humidified incubator (37 °C) in 10 cm Petri dishes with a growth medium consisting of Dulbecco's modified Eagles medium (Gibco, Paisley, UK) (DMEM) supplemented with d-glucose (4500 mg L<sup>-1</sup>), 10% FBS (Gibco, Paisley, UK), 100 μg/mL penicillin G (Gibco, Paisley, UK), and 100 μg/mL streptomycin (Gibco, Paisley, UK). The growth medium was replaced with 2 × 3 mL of OptiMEM (Gibco, Paisley, UK) prior to transfection. Six micrograms of receptor plasmid was transfected to each Petri dish, using Plus reagent (Gibco, Paisley, UK) and Lipofectamine reagent (Gibco, Paisley, UK). The transfection procedure was carried out according to the instructions given by the manufacturer. The HEK-293 cells and the transfection mixture were incubated for 4 h at 37 °C, 5% CO<sub>2</sub>. The cells were washed, cultured in growth medium, and incubated for 24 h, followed by a reculture of growth medium and then incubated for an additional 24 h. Medium was gently removed by suction, and cold phosphate buffer was added. The resuspended DNA was centrifuged at 4 °C (1000g for 5 min). The pellet was resuspended in 5 mL of cold 50 mM TRIS buffer (supplemented with 1 mM EDTA, pH 7.4). The suspension was mixed with a polytron and centrifuged at 4 °C (15000g for 20 min). The pellet was resuspended in 10 mM TRIS buffer (supplemented with 10% sucrose, 0.2 mM EDTA, pH 7.4) and mixed. The transfected HEK-293 cell membranes were stored at -80 °C. Protein determination was conducted according to Lowry et al. (1951).

**Radioligand Binding Assay.** Baretin and 8,9-dihydrobaretin were evaluated in an in vitro radioligand binding assay based on the displacement of radioligands from human 5-HT receptors expressed in HEK-293 cell membranes. [*N*-methyl-<sup>3</sup>H]LSD (86.8 Ci/mmol), [9-methyl-<sup>3</sup>H]BRL-43694 (69.50 Ci/mmol), [*N*-methyl-<sup>3</sup>H]GR113808 (81.0 Ci/mmol), and [1,2-<sup>3</sup>H]5-carboxamidotryptamine (29.70 Ci/mmol) were used as radioligands. Nonspecific radioligand binding was determined in the presence of 10 μM 5-hydroxytryptamine. Cell membrane suspension expressing the 5-HT receptor subtype (~10 μg protein) was suspended in a 50 mM TRIS buffer (supplemented with 3 mM MgCl<sub>2</sub>, pH 7.4) and incubated (60 min at 25 °C; total volume = 500 μL) with radioligand (0.8 mM [*N*-methyl-<sup>3</sup>H]LSD; 0.2 mM [9-methyl-<sup>3</sup>H]BRL-43694; 0.8 mM [1,2-<sup>3</sup>H]5-carboxamidotryptamine; or 0.05 mM [*N*-methyl-<sup>3</sup>H]GR113808) and concentrations ranging from 0.01 nM to 1 μM of the compound tested. The incubation was terminated by rapid filtration under vacuum through Whatman GF/B glass fiber filter sheets presoaked in washing buffer (50 mM TRIS, 1 mM EDTA, pH 7.4) using a Brandel cell harvester. The filters were washed with 3 × 2 mL of the same buffer, dissolved in 4 mL of EcoScint A (National Diagnostics, Atlanta, GA), and incubated overnight. The residual radioligand bound to the filter was determined by liquid scintillation counting (LS6000IC, Beckman, Fullerton, CA).

**Saturation Analysis.** Saturation studies were conducted on transfected HEK-293 cell membranes expressing 5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, or 5-HT<sub>4</sub> receptors. The radioligand [*N*-methyl-<sup>3</sup>H]LSD (86.8 Ci/mmol) was used for 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> and [*N*-methyl-<sup>3</sup>H]GR113808 (81.0 Ci/mmol) was used for 5-HT<sub>4</sub>. Total binding for each radioligand was determined with 10–12 concentrations of radioligand ranging from 6 to 300 nM. Nonspecific binding for each radioligand was determined in the same manner as total binding but with the addition of 400 μM 5-hydrox-

tryptamine to each concentration point (Figures 2, 3, 4). Specific binding was determined by subtracting nonspecific binding from total binding. Scatchard analysis was calculated by dividing bound radioligand by free radioligand. Approximately 7 μg of membrane protein was suspended together with radioligand in a 50 mM TRIS buffer (supplemented with 3 mM MgCl<sub>2</sub>, pH 7.4) and incubated (60 min at 25 °C; total volume = 100 μL). Scintillation was determined as described in the radioligand binding assay.

**Statistical Analysis.** Serotonin binding data were analyzed with nonlinear regression analysis using GraphPad Prism 4 (GraphPad Software, San Diego, CA). *K<sub>i</sub>* values were calculated based on *K<sub>D</sub>* values for [*N*-methyl-<sup>3</sup>H]LSD and [*N*-methyl-<sup>3</sup>H]GR113808. IC<sub>50</sub> values were converted to *K<sub>i</sub>* using the Cheng and Prusoff method.<sup>19</sup> The *B<sub>max</sub>* and *K<sub>D</sub>* values were obtained from three independent saturation experiments.

**Acknowledgment.** The authors wish to acknowledge the assistance of B.-M. Johansson for her skillful laboratory assistance. This project was financed by the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (FORMAS Marine), the Swedish Foundation for Strategic Research through the MASTEC program, the Swedish Research Council through contract 621-2002-4770 (P.R.J.), FORMAS through contract 210/2004-0285 (P.R.J.), and the Swedish Foundation for Strategic Environmental Research (MISTRA) through Marine Paint (M.D.).

## References and Notes

- Mayer, A. M.; Hamann, M. T. *Mar. Biotechnol.* **2004**, *6*, 37–52.
- Newman, D. J.; Cragg, G. M. *J. Nat. Prod.* **2004**, *67*, 1216–1238.
- Proksch, P. *Toxicol.* **1994**, *32*, 639–655.
- Sjögren, M.; Dahlström, M.; Göransson, U.; Jonsson, P. R.; Bohlin, L. *Biofouling* **2004**, *20*, 291–297.
- Sjögren, M.; Göransson, U.; Johnson, A. L.; Dahlström, M.; Andersson, R.; Bergman, J.; Jonsson, P. R.; Bohlin, L. *J. Nat. Prod.* **2004**, *67*, 368–372.
- Johnson, A. L.; Bergman, J.; Sjögren, M.; Bohlin, L. *Tetrahedron* **2004**, *60*, 961–965.
- Sölter, S.; Dieckmann, R.; Blumenberg, M.; Francke, W. *Tetrahedron Lett.* **2002**, *43*, 3385–3386.
- England, L. J.; Imperial, J.; Jacobsen, R.; Craig, A. G.; Gulyas, J.; Akhtar, M.; Rivier, J.; Julius, D.; Olivera, B. M. *Science* **1998**, *281*, 575–578.
- Beglinger, C. *Int. J. Clin. Pract.* **2002**, *56*, 47–51.
- Bozsing, D.; Simonek, I.; Simig, G.; Jakoczi, I.; Gacsalyi, I.; Levay, G.; Tihanyi, K.; Schmidt, E. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 3097–3099.
- Smith, B. M.; Smith, J. M.; Tsai, J. H.; Schultz, J. A.; Gilson, C. A.; Estrada, S. A.; Chen, R. R.; Park, D. M.; Prieto, E. B.; Gallardo, C. S.; Sengupta, D.; Thomsen, W. J.; Saldana, H. R.; Whelan, K. T.; Menzaghi, F.; Webb, R. R.; Beeley, N. R. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 1467–1470.
- Tierney, A. J. *Comp. Biochem. Physiol., Part A: Mol. Integr. Physiol.* **2001**, *128*, 791–804.
- Colas, J. F.; Choi, D. S.; Launay, J. M.; Maroteaux, L. *Ann. NY Acad. Sci.* **1997**, *812*, 149–153.
- Witz, P.; Amlai, N.; Plassat, J. L.; Maroteaux, L.; Borrelli, E.; Hen, R. *Proc. Natl. Acad. Sci. U.S.A.* **1990**, *87*, 8940–8944.
- Kawahara, H.; Isoai, A.; Shizuri, Y. *Gene* **1997**, *184*, 245–250.
- Clark, M. C.; Dever, T. E.; Dever, J. J.; Xu, P.; Rehder, V.; Sosa, M. A.; Baro, D. J. *J. Neurosci.* **2004**, *24*, 3421–3435.
- Johanson, J. F. *Aliment Pharmacol. Ther.* **2004**, *20*, Suppl. 7, 20–24.
- Beattie, D. T.; Smith, J. A.; Marquess, D.; Vickery, R. G.; Armstrong, S. R.; Pulido-Rios, T.; McCullough, J. L.; Sandlund, C.; Richardson, C.; Mai, N.; Humphrey, P. P. *Br. J. Pharmacol.* **2004**, *143*, 549–560.
- Cheng, Y.; Prusoff, W. H. *Biochem. Pharmacol.* **1973**, *22*, 3099–3108.

NP0601760