

Coumaroyl Iridoids and a Depside from Cranberry (*Vaccinium macrocarpon*)

Allison Turner,[†] Shao-Nong Chen,[†] Dejan Nikolic, Richard van Breemen, Norman R. Farnsworth, and Guido F. Pauli*

UIC/NIH Center for Botanical Dietary Supplements Research, Department of Medicinal Chemistry and Pharmacognosy (MC 781) and PCRPS, College of Pharmacy, University of Illinois at Chicago, 833 S. Wood Street, Chicago, Illinois 60612

Received June 7, 2006

Cranberry (*Vaccinium macrocarpon*) juice has been used for urinary tract infections for approximately 50 years. Recent research suggests that this botanical blocks adherence of pathogenic *E. coli* to urinary tract cells, thus preventing infection. While current evidence indicates that proanthocyanidins are responsible for this activity, these compounds may not reach the urinary tract; thus further investigation is warranted. Fractionation of cranberry juice concentrate was guided by a recently published antiadherence assay, and the resulting fractions were phytochemically characterized. Two new coumaroyl iridoid glycosides, 10-*p-trans*- (**1**) and 10-*p-cis*-coumaroyl-1*S*-dihydromonotropein (**2**), and a depside, 2-*O*-(3,4-dihydroxybenzoyl)-2,4,6-trihydroxyphenylmethylacetate (**3**), were isolated, and although these compounds did not have antiadherent activity in isolation, they might constitute a new group of marker compounds for this active fraction of cranberry.


Cranberry (*Vaccinium macrocarpon* Ait., Ericaceae) juice, generally in the form of cranberry juice cocktail consisting of a mixture of cranberry juice, water, and sugar, is a popular botanical dietary supplement^{1,2} primarily used for the treatment and prevention of urinary tract infections, with documented human use for at least 47 years.³ This use is supported by several clinical trials,^{4–11} although some trials, notably for children with neuropathic bladder and catheterization, did not find cranberry to be effective.^{12,13} Cranberry juice was originally believed to be active due to its acidifying effect on urine and/or the increased excretion of the cranberry urinary metabolite hippuric acid.^{14,15} However, subsequent research suggested that the bacteriostatic impact from acidification alone could not account for its demonstrated effects.^{16,17}

Inhibition of adherence of *Escherichia coli* to uroepithelial cells,^{18–20} rather than direct bacteriostatic or bactericidal activity, has been proposed as the mechanism of action. Specifically, there is support for inhibition of the papG fimbrial attachment of uropathogenic strains of *E. coli* to human cells^{21,22} by cranberry's A-type proanthocyanidin compounds, but not by a B-type dimer or the (–)-epicatechin monomer.^{18,23} Proanthocyanidins, however, may not be assimilated in the gut nor reach the urinary tract intact.²⁴ The putative active cranberry A-type proanthocyanidin oligomers, containing a second link (carbon–oxygen) between at least two of their epicatechin monomer units, are chemically similar to B-type proanthocyanidins, which contain only single, carbon–carbon links between units. Knowledge about the metabolic route in humans of A-type proanthocyanidins is extremely sparse. With very limited recent exceptions for small amounts of dimers and possibly a trimer,²⁵ evidence currently indicates that B-type proanthocyanidins, especially trimers and larger, are degraded in the gut, and/or not assimilated in any quantity, and do not reach the urinary tract intact.^{26–37} Whether this information can be applied to A-type proanthocyanidins is unknown and should be verified before abandoning the search for other active cranberry constituents. Additionally, while urine after cranberry ingestion has been shown to be antiadherent,^{16,17,19} and while proanthocyanidin metabolites³⁸ are possibly the active constituents, to date no researchers have elucidated any specific antiadherent cranberry compounds or metabolites thereof, proanthocyanidin or otherwise, found in urine after cranberry ingestion. It is also important to consider possible synergism between compounds, as seen for the antimutagenic activity of cranberry.³⁹ Relying on one class of isolated active

compounds and ignoring interactions from the remaining constituents, while tempting for reasons of feasibility, can generally be misleading in terms of explaining efficacy of herbals. Therefore, one aim of the present study was to remove proanthocyanidins and bactericidal benzoic acid and then further characterize the *in vitro* antiadherent fraction of cranberry juice. In this small but important step in the complete understanding of the use of this botanical for urinary tract infection, we present data on two new compounds that, while not active in isolation, are potential new (phytochemical) marker compounds for the bioactive cranberry fraction.

Results and Discussion

An assay using biologically relevant cells, and of sufficiently high throughput such that it can efficiently guide the fractionation of cranberry, was developed in our laboratory and recently published.⁴⁰ We have used this assay in the bioactivity-guided fractionation of cranberry juice and report here two new and one known compound (**1–3**) found in, and assisting in the identification of, the active fraction. Compounds **1** and **2** represent an isomeric pair of acylated dihydromonotropein iridoids and, together with previously reported congeners,⁴¹ underline the significance of this class of phytochemicals for the characterization of cranberry preparations. The third compound is the depside 2-*O*-(3,4-dihydroxybenzoyl)-2,4,6-trihydroxyphenylmethylacetate (**3**), a methylated form⁴² of another depside (**5**) so far only reported in *Papaver rhoeas*.⁴³



* To whom correspondence should be addressed. Tel: 312-355-1949. Fax: 312-355-2693. E-mail: gfp@uic.edu.

[†] Equally contributing authors.

Table 1. ^1H NMR [δ in ppm (mult, J in Hz)] Data of Compounds **1**, **2**, and **4** in CD_3OD^b and $\text{D}_2\text{O}^{a,c}$

pos.	1 ^b	1 ^c	2 ^b	2 ^c	4 ^d
1	5.514 (d, 5.3)	5.543 (d, 3.9)	5.466 (d, 5.4)	5.496 (d, 4.0)	5.54 (d, 3.6)
3	7.492 (d, 1.4)	7.526 (br s)	7.463 (d, 1.4)	7.527 (br s)	7.55 (br s)
5	2.933 (dddd, 15.2, 9.2, 1.7, 1.4, 0.5)	2.915 (m)	2.840 (dddd, 15.2, 9.4, 2.0, 1.4, 0.5)	2.828 (m)	2.95 (m)
6	Ha: 1.704 (m) Hb: 2.174 (m)	1.672 (m) 2.091 (m)	Ha: 1.630 (m) Hb: 2.113 (m)	1.674 (m) 2.073 (m)	2.08–1.63 (m w/ H7)
7	1.840–1.716 (2H, m)	1.831–1.710 (2H, m)	1.749–1.645 (2H, m)	1.794–1.670 (2H, m)	2.08–1.63 (m w/ H6)
9	2.319 (dd, 9.2, 5.3)	2.353 (dd 9.3, 4.0)	2.224 (dd, 9.4, 5.4)	2.270 (dd, 9.2, 4.2)	2.34 (dd, 9.3, 3.6)
10	4.182 (d, 11.3) 4.256 (d, 11.3)	4.159 (d, 11.1) 4.199 (d, 11.1)	4.114 (d, 11.2) 4.176 (d, 11.2)	4.194 (d, 11.2) 4.230 (d, 11.2)	3.56 (d, 17.5) 3.62 (d, 17.5)
1'	4.725 (d, 7.9)	4.797 (overlapped ^e)	4.711 (d, 7.9)	4.799 (overlapped ^e)	4.82 (d, 8.4)
2'	3.238 (dd, 9.0, 7.9)	3.285 (dd, 9.1, 8.6)	3.243 (dd, 9.0, 7.9)	3.291 (dd, 9.2, 8.1)	3.29 (t, 9.1, 8.1)
3'	3.329 (overlapped ^e)	3.487 (t, 9.1)	3.327 (overlapped ^e)	3.509 (dd, 9.5, 9.2)	3.51 (t, 9.1)
4'	3.352 (t, 9.0)	3.389 (t, 8.6)	3.349 (t, 9.0)	3.388 (t, 9.5)	3.41 (t, 9.1, 9.1)
5'	3.336 (overlapped ^e)	3.436–3.406 (m)	3.330 (overlapped ^e)	3.500–3.465 (m)	3.54–3.48 (m)
6'	3.662 (dd, 11.8, 5.2) 3.846 (dd, 11.8, 1.6)	3.638 (dd, 12.4, 5.2) 3.788 (dd, 12.4, 2.1)	3.667 (dd, 11.8, 5.1) 3.860 (dd, 11.8, 1.4)	3.708 (dd, 12.4, 5.5) 3.897 (dd, 12.4, 2.2)	3.74 (dd, 12.3, 5.7) 3.92 (br d, 12.3)
2''	6.385 (d, 15.9)	6.276 (d, 16.0)	5.858 (d, 12.6)	5.992 (d, 12.4)	
3''	7.668 (d, 15.9)	7.570 (d, 16.0)	6.914 (d, 12.6)	7.143 (d, 12.4)	
5'', 9''	7.487 (2H, br ddd, 8.7, 2.9, 2.0)	7.445 (d, 8.6)	7.616 (2H, br ddd, 8.7, 3.0, 2.1)	7.471 (d, 8.9)	
6'', 8''	6.804 (2H, br ddd, 8.7, 2.9, 2.0)	6.875 (d, 8.6)	6.750 (2H, br ddd, 8.7, 3.0, 2.1)	6.903 (d, 8.9)	

^a Some proton data were deduced from HMQC. ^b Data were recorded at 360 MHz in CD_3OD . The signals of CD_3OD at δ 3.305 ppm were used as reference. ^c Data were recorded at 500 MHz in D_2O . The signal of D_2O at δ 4.800 ppm was used as reference. ^d Data were recorded at 300 MHz in D_2O ; from Jensen et al. 2002.⁴¹ ^e Signals overlapped with solvent signals.

Cranberry juice, obtained as a frozen concentrate, was thawed, partially neutralized with aqueous ammonia (28% NH_4OH), and fractionated over HP-20 resin. Two more levels of fractionation, guided by positive activity in the antiadhesion assay, resulted in an active fraction, from which two new coumaroyl-iridoid glucosides (**1**, **2**) and a depside (**3**) were isolated using reversed-phase HPLC. These compounds were identified by spectroscopic analysis and by comparison of their data with literature values of the similar compounds dihydromonotropein (**4**) and a *Papaver* depside (**5**).^{41,43}

The deprotonated molecule of **1** at m/z 537.1629 ($[\text{M} - \text{H}]^-$ (high-resolution negative ion electrospray) corresponded to the molecular formula $\text{C}_{25}\text{H}_{30}\text{O}_{13}$ (calc 537.1608). The IR spectra showed the characteristic absorption bands of hydroxyl groups, free acid or ester groups, and alkenes at 3308, 1684, and 1138 cm^{-1} , respectively. UV absorptions at 215, 230, and 315 suggested that **1** had two independent conjugated systems, one of which was aromatic. In the ^1H NMR spectrum of **1** (Table 1), AM spin system resonances at δ_{H} 7.668 and 6.385 (each 1H, d, $J = 15.9$ Hz; H-3'', H-2'', respectively) revealed the presence of a *trans*-ethylene group conjugated with an aromatic ring and a typical AA'XX' spin system for 1,4-disubstituted aromatic ring protons at δ_{H} 7.487 and 6.804 (each 2H, brddd, $J = 8.7, 2.9, 2.0$ Hz; H-5''/H-9'', and H-6''/H-8'', respectively). The spectrum also revealed a six-carbon sugar moiety with all-axial ring protons at δ_{H} 4.725 (1H, d, $J = 7.9$ Hz; H-1'), 3.238 (1H, dd, $J = 9.0, 7.9$ Hz; H-2'), 3.329 (1H, overlapped with solvent signals; H-3'), 3.352 (1H, t, 9.0; H-4'), 3.336 (1H, overlapped with solvent signals; H-5'), 3.846 (1H, dd, $J = 11.8, 1.6$ Hz; H-6a'), and 3.662 (1H, dd, $J = 11.8, 5.2$ Hz; H-6b'). In addition to the observed resonances of characteristic spin systems and a sugar moiety, an olefinic proton was observed at δ_{H} 7.492 (1H, d, $J = 1.4$ Hz; H-3), a dioxymethine proton at δ_{H} 5.514 (1H, d, $J = 5.3$ Hz; H-1), methylene resonances at δ_{H} 4.256 and 4.182 (each 1H, d, $J = 11.3$ Hz; H-10), three proton resonances at δ_{H} 2.933 (1H, dddd, $J = 15.2, 9.2, 1.7, 1.4, 0.5$ Hz; H-5), 2.319 (1H, dd, $J = 9.2, 5.3$ Hz; H-9), and 2.174 (1H, m; H-6a), and three overlapped multiplets in the range δ_{H} 1.831–1.653 accounting for the C-7 methylene protons and one of the two C-6 protons. The shielded H-6b at δ_{H} 1.704 overlapped with H-7, as revealed in the HMQC spectrum. The ^{13}C and DEPT NMR spectra of **1** (Table 2) included resonances for a *trans*-ethylene group at δ_{C} 114.89 (d, C-2'') and 147.03 (d, C-3''), as well as two resonances representing four carbons of a 1,4-disubstituted aromatic ring at δ_{C} 131.32 (d,

Table 2. ^{13}C NMR Data [δ in ppm, mult] of Compounds **1**, **2**, and **4** in CD_3OH^a and D_2O^b

pos.	1 ^a	2 ^b	4 ^c
1	96.14 d	95.05 d	95.6 d
3	153.28 d	152.50 d	153.4 d
4	112.47 s	111.57 s	112.4 s
5	35.21 d	32.40 d	32.5 d
6	31.57 t	29.67 t	30.1 t
7	37.47 t	35.61 t	35.7 t
8	81.66 s	80.37 s	82.8 s
9	46.87 d	45.66 d	45.7 d
10	70.88 t	69.55 t	68.2 t
11	170.74 s	171.25 s	172.0 s
1'	100.88 d	98.88 d	99.4 d
2'	74.62 d	72.72 d	73.3 d
3'	77.88 d	75.72 d	76.3 d
4'	71.22 d	70.14 d	70.1 d
5'	78.40 d	76.38 d	77.0 d
6'	62.49 t	60.78 d	61.3 d
1''	169.23 s	169.22 s	
2''	114.89 d	116.89 d	
3''	147.03 d	144.32 d	
4''	127.13 s	127.31 s	
5'', 9''	131.32 d	131.51 d	
6'', 8''	116.85 d	115.34 d	
7''	161.39 s	156.83 s	

^a 90 MHz in CD_3OH . ^b 90 MHz in D_2O . ^c 75 MHz in D_2O ; from Jensen et al. 2002.⁴¹

C-5'' and 9'') and 116.85 (d, C-6'' and 8'') and six oxygenated carbon resonances belonging to the sugar substituent at δ_{C} 100.88 (d, C-1'), 74.62 (d, C-2'), 77.88 (d, C-3'), 71.22 (d, C-4'), 78.40 (d, C-5'), and 62.49 (t, C-6'). In addition, the spectra showed an acyl group and a carbonyl group at δ_{C} 169.23 (s, C-1'') and 170.74 (s, C-11), a pair of resonances for olefinic carbons at δ_{C} 153.28 (d, C-3) and 112.47 (s, C-4), three methine carbons at δ_{C} 96.14 (d, C-1), 35.21 (d, C-5), and 46.87 (d, C-9), three methylene carbons at δ_{C} 31.57 (t, C-6), 37.47 (t, C-7), and 70.88 (t, C-10), and an oxygenated quaternary carbon at δ_{C} 81.66 (s, C-8). These NMR data suggested that **1** was likely an iridoid glycoside, as they were similar to those of the known iridoid glucoside 6,7-dihydromonotropein (**4**) recently isolated from cranberry by Jensen et al.⁴¹

The primary NMR differences between **1** and **4**, in addition to the *trans*-coumaroyl signals in **1**, were that the protons at δ_{H} 3.56 (1H, d, $J = 17.5$ Hz; H-10b) and 3.62 (1H, d, $J = 17.5$ Hz;




Figure 1. Additional resonances (*/**) in the ^1H NMR spectrum of the parent fraction, from which **1** and **2** were isolated, revealed the presence of structurally related minor iridoids, indicated by the presence of pairs of doublets with the typical ~ 11 Hz geminal coupling of acylated C-10 oxymethylene resonances. For example, the signals at δ 4.329 and 4.205 (*a) likely belong to a third major monotropein-type iridoid. This indicates that these iridoids are distinct phytochemical markers of the antiadherence active fraction of *V. macrocarpon*.

H-10a) in **4** were shifted toward lower field to δ_{H} 4.182 (1H, d, $J = 11.3$) and 4.256 (1H, d, $J = 11.3$ Hz) in **1** and that the corresponding carbon at δ_{C} 68.2 (t, C-10) in **4** was also deshielded to δ_{C} 70.88 (t) in **1**. This evidence indicated that the coumaroyl group in compound **1** was connected to the C-10 hydroxyl function of the iridoid skeleton. Additionally, the shift in the geminal coupling constants for H-10a and H-10b from 17.5 Hz in **4** to 11.3 Hz in **1** suggested that the hydroxyl proton had been replaced by a bulky substituent.

In the HMBC spectrum, correlations of the C-10 methylene protons resonating at δ_{H} 4.182 and 4.256 with the quaternary carbon at δ_{C} 169.23 (C-1'') were observed. This quaternary carbon was also correlated with two other protons at δ_{H} 6.385 and 7.668 (H-2'' and H-3'', respectively). This evidence confirmed that the coumaroyl group must be attached at position 10. Two fragments of m/z 163.0344 and 373.1023 in the negative ion tandem mass spectrum supported this conclusion.

The coupling constants of $J_{1',2'}$ (7.9 Hz), $J_{2',3'}$ (9.0 Hz), and $J_{3',4'}$ and $J_{4',5'}$ (9.0 Hz in CD_3OD and 8.6 Hz in D_2O) indicated that all of these protons are pyranose ring protons in an axial position, confirming that the sugar substituent was glucose. In the ROESY spectrum of **1**, the correlations between H-1 and H-1' and one of two C-7 protons indicated that H-1 is in an α -orientation. Both H-5 and H-9 correlated with H-6 (δ_{H} 2.174) and with one of the two C-7 protons (δ_{H} 1.840–1.726). Moreover, correlations of H-10 (δ_{H} 4.182 and 4.256) with H-9 (δ_{H} 2.319), H-5 (δ_{H} 2.933), and H-7 (δ_{H} 1.840–1.716, 2H) were observed. NOE experiments provided evidence that these protons were in β -orientations; that is, **1** had the same relative configuration as **4**. Consequently, compound **1** was deduced to be 10-*trans*-coumaroyl-6,7-dihydromonotropein.

Compound **2** was very similar to compound **1**. Comparing the ^1H , ^{13}C , and DEPT NMR spectra of **2** with those of **1**, the only difference was for the proton and carbon resonances within the coumaroyl moiety. In the ^1H NMR spectra, the H-2'' and H-3'' signals were shifted upfield from δ_{H} 6.385 and 7.668 in **1** to δ_{H} 5.858 and 6.914 in **2**, respectively. The coupling constant between H-2'' and H-3'' differed markedly from 15.9 Hz in **1** to 12.6 Hz in **2**. Comparing the ^{13}C NMR spectra, the chemical shift of C-2'' was shifted downfield from δ_{C} 114.89 in **1** to 116.89 in **2**, while

the chemical shift of C-3'' moved to higher field from δ_{C} 147.03 in **1** to 144.32 in **2**. This evidence indicated that compound **2** had a *cis*-coumaroyl substituent, as opposed to the *trans*-coumaroyl seen in **1**. Therefore, compound **2** was determined to be 10-*cis*-coumaroyl-6,7-dihydromonotropein.

Additional iridoid compounds are likely to be present in cranberry. A ^1H NMR analysis of the parent fraction (Figure 1), from which the two iridoids **1** and **2** were isolated, reveals a total of four pairs of doublets between δ_{H} 4.6 and 4.1 with the typical coupling of C-10 acylated geminal protons ($J = 11.1$ to 11.4 Hz). While **1** and **2** account for one pair of doublets each, two pairs remain, for a possibility of at least two more compounds belonging to the same series of iridoids. At least one compound is highly likely to be a 10-*O*-substituted iridoid glycoside: a doublet can be clearly seen downfield from the H-10 resonances of both **1** and **2**, at δ_{H} 4.329 ($J = 11.4$ Hz); resonances for the second C-10 proton for this third iridoid are likely to be found at around 4.2, overlapping with the resonances of **1** and **2** (likely a $J = 11.4$ Hz doublet at 4.205). While the low available sample mass limited analysis in the present study, further analysis of this fraction with regard to the presence of additional iridoid glycosides is clearly warranted.

Compound **3** was isolated in a yield of 1.7 mg (13 ppm, dry weight) as an amorphous, colorless solid. The high-resolution negative ion electrospray mass spectrum of compound **3** revealed a deprotonated molecule at m/z 333.0606 $[\text{M} - \text{H}]^-$, corresponding to a molecular formula of $\text{C}_{16}\text{H}_{14}\text{O}_8$ (calc 333.0611). The negative ion tandem mass spectrum of compound **3** contained a fragment ion m/z 301, which corresponded to a loss of methanol and indicated the presence of a methoxyl group in **3**. The IR spectrum showed the characteristic absorption bands corresponding to a hydroxyl group, acyl or carboxyl groups, and alkenes at 3362, 1684, and 1138 cm^{-1} , respectively. In the ^1H NMR (Table 3) and ^1H - ^1H COSY spectra of **3**, only two aromatic spin systems were observed at lower field. One was a 1,3,4-trisubstituted ABX spin system with resonances at δ_{H} 7.529 (1H, dd, $J = 8.0, 2.0$ Hz; H-6'), 7.517 (1H, dd, $J = 2.0, 0.4$ Hz; H-2'), and 6.863 (1H, dd, $J = 8.0, 0.4$ Hz; H-5'), respectively. The other was a 1,2,4,6-tetrasubstituted AX spin system with resonances at δ_{H} 6.251 (1H, d, $J = 2.3$ Hz; H-3) and 6.135 (1H, d, $J = 2.3$ Hz; H-5). In addition to the above resonances, a methoxyl and a similarly shifted methylene resonance were

Table 3. NMR Data of Compounds **3** and **5**

pos.	δ_{H} (3) ^a	δ_{H} (5) ^b	δ_{C} (3) ^a	δ_{C} (5) ^b
1			107.14 s	107.7
2			158.63 s	158.5
3	6.251 (d, 2.3)	6.18 (d 2.2)	101.02 d	101.2
4			158.42 s	158.6
5	6.135 (d, 2.3)	6.27 (d 2.2)	102.12 d	102.1
6			152.40 s	152.4
7	3.479 (brs)	3.47 s (2H)	29.89 t	30.3
8			174.34 s	176.1
1'			121.65 s	121.8
2'	7.517 (dd 2.0, 0.4)	7.55 (d 8.2)	117.81 d	117.9
3'			146.45 s	146.4
4'			152.51 s	152.5
5'	6.863 (dd 8.0, 0.4)	6.86 (d 8.2)	116.09 d	116.1
6'	7.529 (dd, 8.0, 2.0)	7.56 (dd, 8.2, 2.2)	124.37 d	124.5
7'			166.33 s	166.4
MeO	3.550 s		52.33 q	

^a 360 MHz for ¹H and 90 MHz for ¹³C in CD₃OD. ^b 500 MHz for ¹H and 125 MHz for ¹³C in CD₃OD, from Hillenbrand et al.⁴²

observed at δ_{H} 3.550 (3H, s; OMe) and 3.479 (2H, brs; H-7a, H-7b), respectively. The ¹³C and DEPT NMR resonances could be ascribed to a 1,2,4,6-tetrasubstituted aromatic ring containing resonances at δ_{C} 107.14 (s, C-1), 158.63 (s, C-2), 101.02 (d, C-3), 158.42 (s, C-4), 102.12 (d, C-5), and 152.40 (s, C-6) and to a 1,3,4-trisubstituted aromatic ring with carbons resonating at δ_{C} 121.65 (s, C-1'), 117.81 (d, C-2'), 146.45 (s, C-3'), 152.51 (s, C-4'), 116.09 (d, C-5'), and 124.37 (d, C-6'), respectively. In addition, there were two carbonyl resonances at δ_{C} 174.34 (s, C-8) and 166.29 (s, C-7'), a methylene at δ_{C} 29.89 (t, C-7), and one methoxyl resonance at δ_{C} 52.33. The HMBC spectrum of **3** exhibited correlations between the respective protons and carbons, as illustrated in Figure 2. Compared to what is typical of a free carboxylic acid,⁴³ C-7' in **3** was shifted to a higher field, indicating that this is the site of esterification, which, by a process of elimination, must have occurred between the two rings of this depside. However, the C-7' shift supplied only indirect evidence; additional support for this ester configuration was provided by MS data. The positive ion electrospray tandem mass spectrum contained the dominant fragment ion of m/z 137.0239 (C₇H₅O₃) corresponding to the cleavage of the ester bond (see Figure 2). This ion further fragmented to lose CO, which further supported the assignment of the ester group. The NMR data were consistent with a similar compound recently discovered in *Papaver rhoeas* petals,⁴³ with the exception of the 8.2 Hz coupling constant reported for H-2'. It is our understanding that this is a typographical error on the part of Hillenbrand et al.,⁴³ because the presence of three *ortho* coupling constants is inconsistent with their *tri*- and *para*-substituted ring, and, additionally, a matching 2.2 Hz *meta* coupling constant (H-2' with H-6') is missing from their data. Compound **3** was thus deduced to be 2-*O*-(3,4-dihydroxybenzoyl)-2,4,6-trihydroxyphenylmethylacetate (**3**).⁴²

It should be noted that it is rare to find depsides in organisms that are not lichens,⁴⁴ although there is evidence that ionizing radiation, such as that used to preserve foods, can break down quercetin into various depsides.⁴⁵ Whether **3** is in fact a genuine

cranberry secondary metabolite or from some other biological source, related or unrelated to cranberries, deserves further investigation. Considering published evidence,⁴² however, the methyl ester group in **3** appears unlikely to be due to artifact formation in the presence of methanol/acid.

Experimental Section

General Experimental Procedures. Optical rotations were obtained with a Perkin-Elmer 241 polarimeter (Perkin-Elmer, Inc., MA). ¹H and ¹³C NMR were measured on Bruker Avance 360 and Avance 500 instruments (Bruker, Zürich, Switzerland). Chemical shifts (δ) were expressed in ppm with reference to the deuterated MeOH signals (¹H: 3.305, ¹³C: 49.000 ppm) or D₂O (4.8000 ppm). The digital resolution was always better than 0.1 Hz equivalent to 0.0002 ppm (e.g., 32K real data points, 8 ppm spectral width for ¹H NMR) in the ¹H and 1.2 Hz equivalent to 0.008 ppm (32K real data points, 250 ppm spectral width) in the ¹³C domain. HRESIMS data were recorded on a Micromass (Manchester, UK) Q-TOF-2 system. Infrared spectra were recorded from a thin film on a germanium ATR unit (JASCO FT/IR-410, JASCO Inc., MD). Thin-layer chromatography was performed on precoated TLC plates (250 μm thickness, KGF Si gel 60 and KGF RP-18 Si gel 60, EM Science, Germany, or 200 μm ALUGRAM SIL G/UV₂₅₄, Macherey-Nagel, Germany) with compounds visualized by spraying the dried plates with 5% H₂SO₄ in EtOH or 2% anisaldehyde and 5% H₂SO₄ in EtOH, followed by heating. Semipreparative HPLC was carried out on a Waters Delta 600 system with a Waters 996 photodiode array detector, Waters 717 plus autosampler, and Millennium32 chromatography manager (Waters Co., MA) on a GROM-Sil 120 ODS-4 HE (Watrex-International, Inc., CA) semipreparative column (5 μm , 300 \times 20 mm) with a flow rate of 6 mL/min. Diaion HP-20 (Supelco Co., PA) and lipophilic Sephadex LH-20 (25–100 μm , Sigma Chemical Co., MO) were used for column chromatography. A high-speed countercurrent chromatography system (Pharma-Tech CCC-1000, Pharma-Tech, MD) was also used for fractionation. All solvents were HPLC grade (Fisher Scientific Co., Hanover Park, IL).

Biological Activity. Fractionation was guided by an assay that measured inhibition of adherence of uropathogenic *E. coli* to human uroepithelial cells; for experimental details see our recent report.⁴⁰ In brief, P-fimbriated *E. coli* were grown on CFA agar to promote fimbrial growth. Human T-24 cells were grown to confluence in 96-well microplates. Substances to be tested were briefly incubated with *E. coli* and then poured over the T-24 cells and incubated at 35 °C for 1 h. Unadhered bacteria were carefully rinsed away, and new media was added to wells. Adherent bacteria were allowed to multiply for 4 to 6 h and detected with a microplate reader. Differences in initial quantities of adhered bacteria, compared with solvent controls, were calculated using a standard curve.

Botanical Source. Cranberry juice concentrate was obtained from Ocean Spray, Inc., in two separate lots (lot numbers T073101 and T11802). Because juice does not contain recognizable genetic material or cellular matter, botanical identification was not possible. An HPLC chromatogram for each lot, with peak matching for several cranberry metabolites, is available; see Supporting Information.

Fractionation and Isolation. Briefly, fractionation of cranberry juice consisted of four levels: initially following the general guidelines of Kandil et al.,⁴⁶ the first column was HP-20 reversed-phase resin. HP-20 active fractions were secondarily separated with high-speed coun-




Figure 2. HRMS fragmentation pattern and HMBC correlations of 2-*O*-(3,4-dihydroxybenzoyl)-2,4,6-trihydroxyphenylmethylacetate (**3**).




Figure 3. Bioactivity-guided fractionation of cranberry juice, leading to isolation of three compounds (1–3). Bold lines follow activity through the first three levels of fractionation. Longer lines from level three indicate further separation on HPLC.

tercurrent chromatography (HSCCC, conditions see below), followed, third, by a 4 m Sephadex LH-20 column, and the final fourth-level fractionation was accomplished on a semipreparative HPLC reversed-phase column (Figure 3). Bacterial antiadherence activity-guided fractionation was performed through the first three levels. The overall method was specifically designed to eliminate benzoic acid early in the fractionation scheme; this method primarily follows bioactivity and differs from more traditional schemes developed for the targeted isolation of polyphenolic compounds.

First-Level Fractionation. Two lots of cranberry juice concentrate, 31.8 kg total including H₂O content (approximately 6.4 kg dry weight, processed in two runs, approximately 12 and 19 kg, respectively), were adjusted on ice from pH 2.4 to approximately pH 5.5 with 28% ammonia (Fisher) and were fractionated over 2.25 kg Dianon HP-20 (Supelco/Sigma) that had been conditioned with deionized H₂O. Juice was allowed to flow through the column, which retained all colored anthocyanins. Effluent of this first fraction contained 95% of the mass (dry weight). The subsequent mobile phase sequence consisted of 100% deionized H₂O (10 and 14 L, first and second run, respectively), followed by 20% MeOH in deionized H₂O (8 and 9 L, respectively), then 50% MeOH (11 and 19 L, respectively), and finally 100% MeOH (11 and 13 L, respectively). Transitions in MeOH concentration were accomplished with a short gradient of approximately 1 L total volume. One-liter fractions were collected throughout and tested for antiadherent activity. Antiadherent fraction yield, from 100% MeOH fractions, was 34.6 g.

Second-Level Fractionation. Because the above active HP-20 first-level fractions contained large quantities of bactericidal benzoic acid, which interfered with antiadherence assessment, HSCCC separation on a Pharma-Tech CCC-1000 system was next used. A 12.9 g amount of the active fraction from level one was separated using a solvent system of CHCl₃–MeOH–H₂O (10:7:5), flow rate of 1.5 mL/min, UV detection (T2H mode, upper phase mobile). The resulting fractions were recombined per TLC into seven fractions. The first of these seven fractions, collected in 10 runs averaging 62.5 mL of solvent each (total 625 mL), with a yield of 5.05 g dry weight, contained antiadherent activity but no benzoic acid.

Third-Level Fractionation. The first HSCCC fraction (5.05 g, $K_D < 0.2$), containing all visible anthocyanins but no benzoic acid, was then fractionated over a 4 m Sephadex LH-20 column (300 mm × 25 mm i.d. precolumn; main column 4 m × 10 mm i.d., 108 g total LH-20; isocratic 100% MeOH mobile phase) in six injections of approximately 800 mg each (300 fractions at 8 mL each collected for each injection). Fractions obtained were recombined into 16 fractions per TLC.

Fourth-Level Fractionation. Fractions 6, 8–12, 14, and 16 from level three were further separated on a semipreparative HPLC reversed-phase column (Grom, SIL-120 ODS-4 HE, 300 × 20 mm, 5 μm) at 5 mL/min, with a mobile phase consisting of a mixture of MeOH (solvent A) and H₂O containing 0.1% TFA (solvent B). Compounds 1 and 2 were isolated using a gradient of 50:50 to 76:24 A:B over 35 min. Retention time for 1 was 23.6 min (6.8 mg), and retention time for 2 was 28.4 min (1.9 mg). For the isolation of compound 3, an isocratic method (55:45 A:B) was used. Retention time for 3 was 23.3 min (1.7 mg).

10-*p*-trans-Coumaroyl-1S-dihydromonotropein (1): colorless, amorphous solid; $[\alpha]_D^{25} -44.4$ (c 0.20, MeOH); UV (MeOH–H₂O) λ_{max} 215, 230, 310; IR (Ge ATR) ν_{max} 3308, 1684 br, 1634, 1604, 1515, 1169, and 1074 cm⁻¹; ¹H NMR and ¹³C NMR, see Tables 1 and 2; HRESIMS m/z 537.1629 (calcd for [C₂₅H₃₀O₁₃–H]⁻, 537.1608)

10-*p*-cis-Coumaroyl-1S-dihydromonotropein (2): colorless, amorphous solid; $[\alpha]_D^{25} +37.8$ (c 0.10, MeOH); UV (MeOH–H₂O) λ_{max} 215, 231, 314; IR (Ge ATR) ν_{max} 3308, 1695, 1684, 1558, 1507, 1457, 1188, 1160, and 1071 cm⁻¹; ¹H NMR and ¹³C NMR, see Tables 1 and 2; HRESIMS m/z 537.1649 (calcd for [C₂₅H₃₀O₁₃–H]⁻, 537.1608).

2-O-(3,4-Dihydroxybenzoyl)-2,4,6-trihydroxyphenylmethylacetate (3): colorless, amorphous solid; UV (MeOH–H₂O) λ_{max} 215, 230, 315; IR (Ge ATR) ν_{max} 3308, 1717, 1700, 1684, 1653, 1560, 1295, 1191, and 1138 cm⁻¹; ¹H NMR and ¹³C NMR, see Tables 1 and 2; HRESIMS m/z 333.0606 (calcd for [C₁₆H₁₄O₈–H]⁻, 333.0611).

Acknowledgment. Research was supported by an NRSA Fellowship for A.T. (F3 AT00623) from the National Center for Complementary and Alternative Medicine (NCCAM) and by grant number P50 AT00155 from the National Center for Complementary and Alternative Medicine (NCCAM), the Office of Dietary Supplements (ODS), the Office for Research on Women's Health (ORWH), and the National Institute of General Medical Sciences (NIGMS). The contents are solely the responsibility of the authors and do not necessarily represent official views of NCCAM, ODS, ORWH, or NIGMS. We thank Ocean Spray Cranberries, Inc. for the cranberry juice concentrate used in this study and for an unrestricted gift.

Supporting Information Available: HPLC chromatograms of the two lots of cranberry juice used in this research, including retention time for known cranberry compounds ideain chloride, resveratrol, and benzoic acid, are available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- Bennett, J.; Brown, C. M. *J. Am. Pharm. Assoc.* **2000**, *40*, 353–358.
- Blumenthal, M.; Brinckmann, J.; Wollschlaeger, B. *The ABC Clinical Guide to Herbs*, 1st ed.; American Botanical Council: Austin, TX, 2003; p 480.
- Bodel, P. T.; Cotran, R.; Kass, E. H. *J. Lab. Clin. Med.* **1959**, *54*, 881–888.
- Avorn, J.; Monane, M.; Gurwitz, J. H.; Glynn, R. J.; Choodnovskiy, I.; Lipsitz, L. A. *J. Am. Med. Assoc.* **1994**, *271*, 751–754.
- Dignam, R. R.; Ahmed, M.; Kelly, K. G.; Denman, S. J.; Zayon, M.; Kleban, M. *Ann. Long-Term Care* **1998**, *6*, 163–167.
- Gibson, L.; Pike, L.; Kilbourn, J. P. *J. Naturopathic Med.* **1991**, *2*, 45–47.
- Haverkorn, M. J.; Mandigers, J. *J. Am. Med. Assoc.* **1994**, *272*, 590.
- Howell, A. B.; Foxman, B. *J. Am. Med. Assoc.* **2002**, *287*, 3082–3083.
- Papas, P. N.; Bruschi, C. A.; Ceresia, G. C. *Southwest. Med.* **1966**, *47*, 17–20.
- Stothers, L. *Can. J. Urol.* **2002**, *9*, 1558–1562.
- Walker, E. B.; Barney, D. P.; Mickelsen, J. N.; Walton, R. J.; Mickelsen, R. A., Jr. *J. Fam. Pract.* **1997**, *45*, 167–168.
- Foda, M. M.; Middlebrook, P. F.; Gatfield, C. T.; Potvin, G.; Wells, G.; Schillinger, J. F. *Can. J. Urol.* **1995**, *2*, 98–102.
- Schlager, T. A.; Anderson, S.; Trudell, J.; Hendley, J. O. *J. Pediatr.* **1999**, *135*, 698–702.
- Kinney, A. B.; Blount, M. *Nurs. Res.* **1979**, *28*, 287–290.
- Schultz, A. *J. Community Health Nurs.* **1984**, *1*, 159–169.
- Schmidt, D. R.; Sobota, A. E. *Microbios* **1988**, *55*, 173–181.
- Sobota, A. E. *J. Urol.* **1984**, *131*, 1013–1016.
- Foo, L. Y.; Lu, Y.; Howell, A. B.; Vorsa, N. *J. Nat. Prod.* **2000**, *63*, 1225–1228.
- Habash, M. B.; Van der Mei, H. C.; Busscher, H. J.; Reid, G. *Can. J. Microbiol.* **1999**, *45*, 691–694.
- Howell, A. B.; Vorsa, N.; Der Marderosian, A.; Foo, L. Y. *N. Engl. J. Med.* **1998**, *339*, 1085–1086.
- Ahuja, S.; Kaack, B.; Roberts, J. *J. Urol.* **1998**, *159*, 559–562.
- Zafriri, D.; Ofek, I.; Adar, R.; Pocino, M.; Sharon, N. *Antimicrob. Agents Chemother.* **1989**, *33*, 92–98.
- Foo, L. Y.; Lu, Y.; Howell, A. B.; Vorsa, N. *Phytochemistry* **2000**, *54*, 173–181.
- Reid, G. *World J. Urol.* **1999**, *17*, 359–363.
- Tsang, C.; Auger, C.; Mullen, W.; Bornet, A.; Rouanet, J. M.; Crozier, A.; Teissedre, P. L. *Br. J. Nutr.* **2005**, *94*, 170–181.

- (26) Baba, S.; Osakabe, N.; Natsume, M.; Terao, J. *Free Radical Biol. Med.* **2002**, *33*, 142–148.
- (27) Deprez, S.; Brezillon, C.; Rabot, S.; Philippe, C.; Mila, I.; Lapierre, C.; Scalbert, A. *J. Nutr.* **2000**, *130*, 2733–2738.
- (28) Deprez, S.; Mila, I.; Huneau, J. F.; Tome, D.; Scalbert, A. *Antioxid. Redox. Signal.* **2001**, *3*, 957–967.
- (29) Donovan, J. L.; Manach, C.; Rios, L.; Morand, C.; Scalbert, A.; Remesy, C. *Br. J. Nutr.* **2002**, *87*, 299–306.
- (30) Downs, C. T.; McDonald, P. M.; Brown, K.; Ward, D. *J. Chem. Ecol.* **2003**, *29*, 845–858.
- (31) Duweler, K. G.; Rohdewald, P. *Pharmazie* **2000**, *55*, 364–368.
- (32) Gonthier, M.-P.; Donovan Jennifer, L.; Texier, O.; Felgines, C.; Remesy, C.; Scalbert, A. *Free Radical Biol. Med.* **2003**, *35*, 837–844.
- (33) Holt, R. R.; Lazarus, S. A.; Sullards, M. C.; Zhu, Q. Y.; Schramm, D. D.; Hammerstone, J. F.; Fraga, C. G.; Schmitz, H. H.; Keen, C. L. *Am. J. Clin. Nutr.* **2002**, *76*, 798–804.
- (34) Jimenez-Ramsey, L. M.; Rogler, J. C.; Housley, T. L.; Butler, L. G.; Elkin, R. G. *J. Agric. Food Chem.* **1994**, *42*, 963–967.
- (35) Nakamura, Y.; Tonogai, Y. *J. Agric. Food Chem.* **2003**, *51*, 7215–7225.
- (36) Spencer, J. P.; Chaudry, F.; Pannala, A. S.; Srail, S. K.; Debnam, E.; Rice-Evans, C. *Biochem. Biophys. Res. Commun.* **2000**, *272*, 236–241.
- (37) Spencer, J. P. E.; Schroeter, H.; Shenoy, B.; Srail, S. K. S.; Debnam, E. S.; Rice-Evans, C. *Biochem. Biophys. Res. Commun.* **2001**, *285*, 588–593.
- (38) Howell, A. B.; Leahy, M. M.; Kurowska, E.; Guthrie, N. *FASEB J.* **2001**, *15*, A284.
- (39) Vattem, D. A.; Jang, H. D.; Levin, R.; Shetty, K. *J. Food Biochem.* **2006**, *30*, 98–116.
- (40) Turner, A.; Chen, S. N.; Joike, M. K.; Pendland, S. L.; Pauli, G. F.; Farnsworth, N. R. *J. Agric. Food Chem.* **2005**, *53*, 8940–8947.
- (41) Jensen, H. D.; Krogfelt, K. A.; Cornett, C.; Hansen, S. H.; Christensen, S. B., *J. Agric. Food Chem.* **2002**, *50*, 6871–6874.
- (42) While this paper was under review, the discovery of the same compound, named jaboticabin, was published by: Reynertson, K. A.; Wallace, A. M.; Adachi, S.; Gil, R. R.; Yang, H.; Basile, M. J.; D'Armiento, J.; Weinstein, I. B.; Kennelly, E. J. *J. Nat. Prod.* **2006**, *69*, 1228–30.
- (43) Hillenbrand, M.; Zapp, J.; Becker, H. *Planta Med.* **2004**, *70*, 380–382.
- (44) Miao, V.; Coeffet-LeGal, M. F.; Brown, D.; Sinnemann, S.; Donaldson, G.; Davies, J. *Trends Biotechnol.* **2001**, *19*, 349–355.
- (45) Marfak, A.; Trouillas, P.; Allais, D.-P.; Champavier, Y.; Calliste, C.-A.; Duroux, J.-L. *J. Agric. Food Chem.* **2002**, *50*, 4827–4833.
- (46) Kandil, F. E.; Smith, M. A. L.; Rogers, R. B.; Pepin, M.-F.; Song, L. L.; Pezzuto, J. M.; Seigler, D. S. *J. Agric. Food Chem.* **2002**, *50*, 1063–1069.

NP060260F