

Highly Polar Spiroisoxazolines from the Sponge *Aplysina fulva*

Evan W. Rogers and Tadeusz F. Molinski*

Department of Chemistry and Biochemistry and Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92093

Received March 12, 2007

Two new highly polar brominated spiroisoxazolines, araplysinin *N*⁹-sulfamate (**1**) and an *N*-[5*S*,10*R*]-7,9-dibromo-10-hydroxy-8-methoxy-1-oxa-2-azaspiro[4.5]deca-2,6,8-triene-3-carboxy]-4-aminobutanoic acid (**2**), were isolated from a sample of *Aplysina fulva* collected in the Florida Keys. The absolute stereostructures of the new compounds were determined from analysis of MS, ¹H and ¹³C NMR, and CD spectroscopy. Compound **2** provides a structural clue that may unify the biosynthesis of brominated spiroisoxazolines.

Secondary metabolites from sponges of the order Verongidae¹ are typically highly oxidized compounds derived from bromotyrosine. Modified alkaloids derived from 3',5'-dibromotyrosine (Figure 1, *i*, R = H) have been found from various Verongid genera²—mainly *Aplysina*, *Psammaplysilla*, *Pseudoceratina*, and *Verongia*³—that are widely distributed throughout Mediterranean, Pacific, and Atlantic waters.⁴ Heterocycles based on brominated spiroisoxazolines (Figure 1, *iii*, R = H, (5*S*,10*R*)-7,9-dibromo-10-hydroxy-8-methoxy-1-oxa-2-azaspiro[4.5]deca-2,6,8-triene-3-carboxamide) are common natural products arising from bromotyrosine secondary metabolism.⁵ Since the first example of a spiroisoxazoline was reported by the Minale group from two Mediterranean species, *Aplysina aerophoba* and *Verongia thiona*,⁶ over 25 spiroisoxazoline analogues have been described.⁴

Recently, we reported geographic variability of the diastereomeric compositions of fistularin-3 and 11-*epi*-fistularin-3 in *Aplysina* species collected from Brazil and the United States (Florida Keys) and *Agelas* from Australia (the Great Barrier Reef).⁷ We have extended these investigations and now report two new polar spiroisoxazolines: the sulfamate **1** and carboxylic acid **2**. Both compounds occur in very low concentrations in the most polar fractions derived from column chromatography of methanol-soluble components of the sponge extract. Compound **2** is a lower homologue of purpuroceratic acid (**5**),⁸ reported by Kijoa and co-workers, and lacks the unusual aryl *C*-methyl group of the latter. Compound **1** is the *N*-sulfato derivative of the known compound araplysinin-1 (**3**) and is formally the decarboxylation product of the *N*-sulfato α -amino acid ianthesine D (**4**), reported by Okamoto et al. from *Ianthella* sp.⁹ Carboxylic acids **2** and **5** may provide a “missing link” that unifies the biosynthesis of several spiroisoxazoline alkaloids from Verongid sponges.

Moderately polar fractions obtained from silica chromatography of the CHCl₃–MeOH extract of *Aplysina fulva* contained 11-hydroxyfistularin-3 and 11-*epi*-fistularin-3 in variable proportions, as we have reported previously. The most polar fraction obtained from elution of the silica column (100% MeOH) was further separated by C₁₈ reversed-phase HPLC to give a 3:2 mixture of compounds **1** and **2**, respectively. Final purification of this mixture by preparative silica TLC gave pure samples of **1** and **2** as colorless solids.

Compound **1** is an optically active solid, [α]_D +100 (*c* 0.06, MeOH), with UV activity [λ]_{max} 206 nm (log ϵ 4.70), 278 (3.84)]. The compound did not dissolve in CHCl₃, but was partially soluble in MeOH and soluble in DMSO and CHCl₃–MeOH. Analysis of the formula of **1** by mass spectrometry was complicated by multiple pseudomolecular ions and neutral losses. The positive- and negative-ion ESIMS (*m/z* 841 [M – H + Na₂]⁺, 794 [M – Na][–], and 714





Figure 1. Putative biosynthesis of spiroisoxazoline carboxamide **iii** (R = H, (5*S*,10*R*)-7,9-dibromo-10-hydroxy-8-methoxy-1-oxa-2-azaspiro[4.5]deca-2,6,8-triene-3-carboxamide) from a 3',5'-dibromotyrosine residue.



[M – SO₃ – Na][–]) suggested a Na⁺ salt of sulfate-half acid. The formula of C₂₁H₂₃Br₄N₃O₈S for the neutral species, requiring 10 double-bond equivalents, was assigned from positive-ion MALDI HRMS (*m/z* 841.7643, [M – H + Na₂]⁺ Δm = 3.3 mmu). The presence of only 19 distinct signals in the ¹³C NMR spectrum suggested an element of symmetry in an aryl ring. The ¹H NMR spectrum of **1** (Table 1) indicated the presence of three vinyl and aromatic protons, comprising H-5 in the spiro-ring system (δ 6.31, s, 1H) and the two-proton signal in the second ring belonging to a symmetrical 3,5-dibromotyrosine (δ 7.41, s, 2H). The ¹H (CDCl₃–CD₃OD) and ¹³C NMR (DMSO-*d*₆) signals—in particular, the C=N signal (δ 154.6, s) and the characteristic AB quartet for H₂-7 (δ 3.01, d, *J* = 18.3 Hz; 3.79, d, *J* = 18.3 Hz)—were consistent with a 1-oxa-2-azaspirodeca-triene-ring system as seen in the structures of aeroplysinin-1,¹⁰ fistularin-3,¹² and related compounds. NMR

* Corresponding author. Tel: +1-858 534-7115. Fax: +1-858 822-0386. E-mail: tmolinski@ucsd.edu.

Table 1. NMR Data for Compound **1** (600 MHz)

no.	CD ₃ OD–CDCl ₃ (2:1) ^a				DMSO- <i>d</i> ₆			
	δ _C ^b	δ _H [mult., <i>J</i> (Hz)]	HMBC (H→C)	DQF-COSY	δ _C ^b	δ _H [mult., <i>J</i> (Hz)]	HMBC (H→C)	DQFCOSY
1	74.6 (CH)	4.16 s	3,4,5,6		73.6	3.91 d (7.5)	3,4,5,6	OH
2	122.2 (C) ^d				120.8			
3	148.5 (C)				147.3 ^c			
4	113.9 (C) ^d				113.1			
5	131.4 (CH)	6.31 s	1,2,3,4,6,7		131.2	6.58 s	1,2,3	
6	92.1 (C)				90.2			
7	39.7 (CH ₂)	3.02 d (18.3) 3.82 d (18.3)	1,5,6,8		39.3 ^c	3.21 d (18.6) 3.62 d (18.6)	1,5,6,8	
8	154.6 (C)				154.3 ^c			
9	160.6 (C)				158.9			
10	37.7 (CH ₂)	3.61 t (6.4)	9,11,12	11	36.2	3.39 td (6.6, 5.7)	9,11,12	11, CONH
11	29.9 (CH ₂)	2.09 pent (6.4)	10, 12	10, 12	29.4	1.98 p (6.6)	10,12	10,12
12	71.6 (CH ₂)	4.05 t (6.4)	10,11,13	11	71.2	3.95 t (6.6)	10,11	11
13	151.8 (C)				150.8 ^c			
14, 18	118.5 (C)				117.0			
15, 17	133.7 (CH)	7.41 s	13,14,18,19		133.0	7.49 s	13,14,18,19	
16	139.6 (C)				140.6			
19	35.3 (CH ₂)	2.79 t (7.3)	15,16,17,20	20	33.7	2.67 t (6.7)	15,16,17,20	20
20	45.4 (CH ₂)	3.20 t (7.3)	16,19	19	44.7	2.92 q (6.7)	16,19	19, SO ₃ NH
OMe	60.3 (CH ₃)	3.71 s	3		59.6	3.64 s	3	
OH					6.37 d (7.5)			1
CONH					8.55 t (5.7)	9		10
SO ₃ NH					4.14 t (6.7)			20

^a Run as a 3:2 mixture with compound **2**. ^b 500 MHz. ^c Assigned by HSQC and HMBC (*J* = 8 Hz) at 600 MHz. ^d Assignment based on calculated ¹³C chemical shifts (ChemDraw Ultra).

Table 2. NMR Data for Compound **2** (600 MHz)

no.	CD ₃ OD–CDCl ₃ (2:1) ^a				DMSO- <i>d</i> ₆			
	δ _C ^b	δ _H [mult., <i>J</i> (Hz)]	HMBC (H→C)	DQF-COSY	δ _C ^b	δ _H [mult., <i>J</i> (Hz)]	HMBC (H→C)	DQFCOSY
1	74.6 (CH)	4.15 s	3,4,5,6		73.4	3.93 s	3,4,5,6	
2	122.2 (C) ^d				120.8			
3	148.5 (C)				147.0			
4	113.9 (C) ^d				113.3			
5	131.4 (CH)	6.30 s	1,2,3,4,6,7		131.2	6.56 s	1,2,3,4,7	
6	91.9 (C)				90.1			
7	39.8 (CH ₂)	3.01 d (18.0) 3.79 d (18.0)	1,5,6,8		39.7 ^c	3.22 d (17.7) 3.63 d (17.7)	1,5,6,8	
8	154.7 (C)				154.8			
9	160.6 (C)				158.9			
10	40.2 (CH ₂)	3.30 t (6.9)		11	39.4 ^c	3.14 t (6.6)	9,11	11
11	26.1 (CH ₂) (CH ₂)	1.81 p (6.9)	10,12,13	10, 12	24.5	1.65 p (6.6)	10,12,13	10, 12
12	35.9 (CH ₂)	2.21 t (6.9)	10,11,13	11	34.5 ^c	2.09 bt (6.6)	10,11,13	11
13	181.5 (C)				174.7 ^c			
OMe	60.3 (CH ₃)	3.71 s	3		59.6	3.63 s	3	
OH								
CONH								
COOH								

^a Run as a 2:3 mixture with compound **1**. ^b 500 MHz. ^c Assigned by HSQC and HMBC (*J* = 8 Hz) at 600 MHz. ^d Assignment based on calculated ¹³C chemical shifts (ChemDraw Ultra).

signals were observed (Table 1) for other units including a 1,3-disubstituted propane chain terminated with an oxygen atom and an NH(CO) group, and an 1-aryl-2-ethylamine side chain.

A database survey (MarinLit, University of Canterbury) of known Verongid sponge compounds matched the expected formula (with the assumption of one SO₃ group) to a monosulfated derivative of araplyssillin-1 (**3**).¹¹ Verification of this assignment came from ¹³C NMR chemical shifts of **1** (Table 1), which matched closely those of **3**.⁹ (Table 1) The assembled structure **1** and complete chemical shifts assignments were obtained by linking substructures from interpretation of 2D NMR data (gCOSY, gDQFCOSY, gHMBC, and gHSQC).

The SO₃ group was located in **1** as follows. Three exchangeable signals were detected (ESIMS measured in CD₃OD); these appeared in the ¹H NMR spectrum (DMSO-*d*₆) and were attributable to an OH group (δ 6.37 d, *J* = 7.5 Hz), one amide NH (δ 8.55, t, *J* = 5.7 Hz), and an unidentified upfield NH signal (δ 4.14, t, *J* = 6.7 Hz). Since all other heteroatoms in the formula of **1** were accounted for, the SO₃ group was placed on a nitrogen atom that allowed us

to ascribe the latter NH signal to a sulfamate group (–NHSO₃[–]). The NH chemical shift is consistent with those observed for other sulfamates (e.g., δ 4.91 for ianthenisin C¹⁰).

Compound **2**, [α]_D +140 (*c* 0.04, MeOH), also showed an isotope pattern for the MS parent ions consistent with the presence of two Br atoms. The molecular mass of **2** was inferred as 466 amu from low-resolution positive- and negative-ion ESIMS (*m/z* 489 [M + Na]⁺, 465 [M – H][–]), but the formula C₁₄H₁₆Br₂N₂O₆ could be confirmed only by MALDI HRMS (*m/z* 488.9297 [M + Na]⁺, Δ*m* = 2.4 mmu) and required seven double-bond equivalents. Unlike **1**, the appearance of 14 distinct signals in the ¹³C NMR spectrum of **2** (Table 2) showed lack of symmetry. The ¹H NMR spectrum of compound **2** (Table 2) was much simplified compared to that of **1**, but retained signals due to the spiroisoxazoline unit and a 1,3-disubstituted propane side chain. The ¹³C NMR spectrum of **2** showed two C=O ¹³C NMR signals: one corresponding to a free carboxylic acid (δ 181.5 s) and a second due to the α-oximimo amide group (δ 158.9 s). Treatment of **2** with diazomethane gave the corresponding methyl ester **2a** with a new ¹H NMR signal (CD₃-





Figure 2. Proposed unified biosynthesis of 1–6 and aberrant S_N2 substitution of *S*-adenosylmethionine (SAM) by 3',5'-dibromotyrosine phenoxide (path b) by hypothetical “ C^3 -alanyl-methyltransferase”.

OD) due to a second OMe group (δ 3.66, s, 3H). Carboxylic acid **2** most closely resembles purpuroceranic acid **5** with the only difference being the absence of a methyl group at C-5.⁵

The relative configurations of **1** and **2** were addressed using ^{13}C NMR, ROESY, and comparisons of their circular dichroism (CD) spectra with those of known spiroisoxazolines. Since the ^{13}C NMR chemical shifts of the 1-oxa-2-azaspirodeca-triene-ring systems in **1** and **2** were essentially identical, we could assume the relative stereochemistry was the same in both and assign them by analysis of **1**. The ROESY spectrum (500 mS) of **1** (DMSO- d_6) showed dipolar coupling between the C-1 OH and the proximal diastereotopic proton of the C-7 methylene group (δ 3.62, d, $J = 18.0$ Hz)¹² that places both groups *syn* to each other. The absolute configurations of **1** and **2** were assigned by circular dichroism (CD). The CD spectra of **1** and **2** were very similar [CD MeOH, **1**: λ_{max} 244 ($\Delta\epsilon +8.3$), 285 (+8.0); **2**: λ_{max} 243 ($\Delta\epsilon +7.0$), 289 (+5.2)] and showed two prominent positive Cotton effects (CE) that were of the same sign as and similar magnitudes to those of (1*R*,6*S*)-spiroisoxazolines, particularly arothionin (**6a**).^{6,13} Thus, the absolute configurations are as depicted in structures **1** and **2**.



Compounds **2** and **5** contain a structural motif—a spiroisoxazoline carboxyl group *N*-acylated to 4-aminobutanoic acid—that shows a trend that may help explain their biosynthesis. We propose each dibromospiroisoxazoline derives from a *dipeptide* comprised of an *N*-terminal 3',5'-dibromotyrosine and another common α -amino acid (Figure 1, *i*, R = amino acid) that has undergone decarboxylation to the corresponding amine either before or after peptide bond formation (Figure 2). For example, if the 1,4-diaminobutane (putrescine) linker in **6a** derives from ornithine (or lysine, in the case of homoarothionin, **6b**¹⁴), it would appear the 4-aminobutanoic acid unit in **2** derives from glutamic acid. In contrast, the linker in **1**, **3**, **4**, fistularin-3,¹² and most other spiroisoxazolines is 3-amino-1-propanol, which does not have an obvious biogenesis unless it is considered as the decarboxylation product of homoserine, an uncommon amino acid that is an intermediate of one pathway to *S*-adenosylmethionine (SAM). Since the opposite end of the 3-amino-1-propanol linker in fistularin-3 is *O*-alkylated to the phenolic oxygen of a 3',5'-dibromotyrosine (DBT) unit, it is conceivable that the immediate precursor to the fistularin-3 linker is SAM, which participates in an aberrant S_N2 -type alkylation of DBT phenoxide at the more substituted *S*-CH₂ carbon (C-3) of the sulfonium ion instead of the *S*-Me carbon. After decarboxylation, the resultant amine **7** participates in amide bond formation with a spiroisoxazoline unit **ii** that is derived separately by “normal” *O*-methylation of DBT with SAM and arene oxidation (Figure 1). A putative “ C^3 -alanyl-methyltransferase” might catalyze the former transformation and rationalize the unusual *N,O*-substituted C₃ unit that links two of three DBT-derived groups in high molecular mass bromotyrosine natural products from Verongid sponges.

Insufficient quantities of **1** and **2** were available to assess their antifungal activity against *Cryptococcus neoformans* or *Candida albicans*. Further investigations are pending to identify antifungal active principles in *A. fulva* and other sponges from the Bahamas.

Experimental Section

General Experimental Procedures. Optical rotations were measured on a JASCO DIP370 or P1020 polarimeter. UV spectra were recorded on a Hewlett-Packard 8452A single-beam spectrometer, and CD spectra were measured using a JASCO 810 spectropolarimeter. IR

spectra were recorded on a Mattson Galaxy 3000 FTIR instrument. ESIMS was measured using a Finnigan LCQ Deca mass spectrometer, and high-resolution MALDI FTMS spectra were provided by the University of California, Riverside, MS laboratory. ^1H NMR and 2D NMR spectra were recorded on a Bruker 600 MHz DRX-600 equipped with a 5 mm cryoprobe, and ^{13}C NMR spectra were recorded on a Bruker Avance 500 MHz NMR spectrometer. Chemical shifts were referenced to CD_3OD ($\delta_{\text{H}} = 3.31$ ppm; $\delta_{\text{C}} = 49.0$ ppm) and DMSO ($\delta_{\text{H}} = 2.50$ ppm; $\delta_{\text{C}} = 39.5$ ppm). Solvents used were of HPLC grade.

Animal Material. Samples of *Aplysina fulva* (95-03-025, total ~1.4 kg) were collected by hand using scuba from Dry Rocks near Key Largo, Florida, in 1995 and immediately stored at -20 °C until needed. Voucher specimens are archived in the Department of Chemistry and Biochemistry, UC San Diego.

Extraction and Isolation. A CHCl_3 -MeOH-soluble fraction (3.14 g) of the MeOH extract of the sponge was separated by silica flash chromatography and eluted using a solvent gradient (1.5–100% MeOH- CHCl_3) to give an early eluting fraction (10% MeOH- CHCl_3) containing fistularin-3,¹² 11-epi-fistularin-3,¹⁵ and 11-oxoaerothionin¹⁶ as previously described.⁷ The crude fraction eluting with 100% MeOH was submitted to further purification by reversed-phase HPLC (Dynamax C₁₈, 5 μm , 10×250 mm, 3:1 H_2O - CH_3CN , 4.0 mL/min) to a single peak comprising a 3:2 mixture of **1** and **2** (4.8 mg, 0.0017% wet wt), respectively. Further purification of 3 mg of this mixture by silica TLC (4:21 MeOH- CHCl_3) provided **1** (1.2 mg) and **2** (0.7 mg) as white solids.

1: $[\alpha]_{\text{D}}^{23} +100$ (c 0.06, MeOH); UV (MeOH) λ_{max} 206 nm (log ϵ 4.70), 278 (3.84); CD (MeOH) λ 244 nm ($\Delta\epsilon$ +8.3), 285 (+8.0); IR (neat) ν_{max} 3288, 2939, 2864, 1664, 1593, 1544, 1458, 1310, 1218, 1044, 990, 933, 866, 739 cm^{-1} ; ^1H NMR (600 MHz) and ^{13}C NMR (125 MHz), see Table 1; ESIMS m/z 841 $[\text{M} - \text{H} + \text{Na}]^+$, 794 $[\text{M} - \text{Na}]^-$, 714 $[\text{M} - \text{SO}_3 - \text{Na}]^-$; MALDI HRMS m/z 841.7643 $[\text{M} - \text{H} + \text{Na}]^+$, calcd for $\text{C}_{21}\text{H}_{22}\text{N}_3\text{O}_8\text{Na}_2\text{S}^{81}\text{Br}_4$ 841.7616.

2: $[\alpha]_{\text{D}}^{23} +140$ (c 0.04, MeOH); UV (MeOH) λ_{max} 223 nm (log ϵ 3.93), 281 (3.63); CD (MeOH) λ 243 nm ($\Delta\epsilon$ +7.0), 289 (+5.2); IR (neat) ν_{max} 3321, 2959, 2930, 2850, 1666, 1581, 1407, 1310, 1271, 1218, 1048, 990, 921, 767, 739, 703 cm^{-1} ; ^1H NMR (600 MHz) and ^{13}C NMR (125 MHz), see Table 2; ESIMS m/z 489 $[\text{M} + \text{Na}]^+$, 465 $[\text{M} - \text{H}]^-$; MALDI HRMS m/z 488.9297 $[\text{M} + \text{Na}]^+$, calcd for $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_6\text{Na}^{79}\text{Br}_2$ 488.9273.

Methyl Ester of Carboxylic Acid 2. 2a. A solution of **2** (600 μg) in MeOH (~0.5 mL) at 0 °C was treated with an excess of ethereal solution CH_3N_2 (~0.2 M) and allowed to warm to rt over 20 min. The mixture was concentrated and separated on a pencil column (silica, 15:85 MeOH- CH_2Cl_2) to give **2a** as a colorless solid (310 μg): ^1H NMR (CD_3OD) δ 3.66 (s, 3H, OMe), 3.72 (s, 3H, OMe); HREIMS m/z 479.9542 $[\text{M}]^+$ calcd for $\text{C}_{15}\text{H}_{18}\text{O}_6\text{N}_2^{79}\text{Br}_2$ 479.9526.

Acknowledgment. We are grateful to J. Pawlik (University of North Carolina, Wilmington) and the National Underwater Research Center (NURC) for the opportunity to collect from the Key Largo field station. This research was supported by a grant from the National Institutes of Health (AI 39987).

References and Notes

- (1) Class, Demospongia; subclass, Ceractinomorph; order, Verongidae. Verongid sponges are easily identified in the field: upon tissue damage and exposure to air, their characteristic yellow pigmentation undergoes rapid aerial oxidation to blue, purple, and finally black pigments.
- (2) Sponges of the families Aplysinidae, Aplysinellidae, Ianthellidae, and Pseudoceratinidae are responsible for >90% of brominated compounds from Verongida.
- (3) The species “*Verongia*” is considered now by taxonomists to be synonymous with *Aplysina*. Hooper, J. N. A.; Wiedenmayer, F. In *Zoological Catalogue of Australia*; Wells, A., Ed.; CSIRO: Melbourne, 1994; Vol. 12.
- (4) (a) Masuno, M. N. Synthesis and Biological Evaluation of Brominated Marine Natural Product Analogs that Modulate the Skeletal Muscle FKBP12/Calcium Channel Complex. Ph.D. Thesis, University of California, Davis, 2006, Chapter 1. (b) Peng, J.; Li, J.; Hamann, M. T. The Marine Bromotyrosine Derivatives. In *The Alkaloids*; Cordell, G. A., Ed.; Elsevier: New York, 2005; Vol. 61, pp 59–261.
- (5) It is believed the spiroisoxazolines arise from oxidation of the α -amino group of a 4'-methoxy-3',5'-dibromotyrosine residue to a ketoxime that undergoes nucleophilic attack upon a putative arene epoxide (Figure 1, e.g., **ii**, arising from further oxidation of the phenyl ring to give the heterocyclic ring and a secondary alcohol. Anderson, R. J.; Faulkner, D. J. *Tetrahedron Lett.* **1973**, *14*, 1175–1178), although the order of reactions is speculative and identities of the responsible enzymes are presently unknown.
- (6) Fattorusso, E.; Minale, L.; Sodano, G.; Moody, K.; Thomson, R. J. *Chem. Soc., Chem. Commun. D* **1970**, *12*, 752–753.
- (7) Rogers, E. W.; Fernanda de Oliveira, M.; Berlinck, R. G. S.; König, G. M.; Molinski, T. F. *J. Nat. Prod.* **2005**, *68*, 891–896.
- (8) Kijjoa, A.; Bessa, J.; Wattanadilok, R.; Sawangwong, P.; Nascimento, M. S. J.; Pedro, M.; Silva, A. M. S.; Eaton, G. van Soest, R.; Herz, W. *Z. Naturforsch., B: Chem. Sci.* **2005**, *60*, 904–908.
- (9) Okamoto, Y.; Ojika, M.; Kato, S.; Sakagami, Y. *Tetrahedron* **2000**, *56*, 5813–5818.
- (10) Fattorusso, E.; Minale, L.; Sodano, G. *J. Chem. Soc., Perkin. Trans. I* **1972**, *1*, 16–18.
- (11) Longeon, A.; Guyot, M.; Vacelet, J. *Experientia* **1990**, *46*, 548–550.
- (12) Gopichand, Y.; Schmitz, F. J. *Tetrahedron Lett.* **1979**, *41*, 3921–3924.
- (13) McMillan, J. A.; Paul, I. C.; Goo, Y. M.; Rinehart, K. L.; Krueger, W. C.; Pschigoda, L. M. *Tetrahedron Lett.* **1981**, *22*, 39–42. The Cotton effects of the skewed diene in the CD spectrum of aerothionin (**6a**) were as follows: CD (MeOH) λ_{max} 284 ($\Delta\epsilon$ +21.3), 245 (+23.7) (after conversion, $[\theta] = 3300\Delta\epsilon$). Since **6a** is a C_2 symmetrical dimer, we estimated the following comparison values corresponding to the dichroism of one diene unit, assuming no interaction between the two chromophores: CD (MeOH) λ_{max} 284 ($\Delta\epsilon$ +10.7), 245 (+11.8).
- (14) Moody, K.; Thomson, R. H.; Fattorusso, E.; Minale, L.; Sodano, G. *J. Chem. Soc., Perkin Trans. I* **1972**, 18–24.
- (15) König, G. M.; Wright, A. D. *Heterocycles* **1993**, *36*, 1351–1358.
- (16) Acosta, A. L.; Rodriguez, A. D. *J. Nat. Prod.* **1992**, *55*, 1007–1012.

NP070109L