

Lindenane Sesquiterpene Dimers from *Chloranthus fortunei*Xia-Chang Wang,^{†,‡} Yi-Nan Zhang,[†] Li-Li Wang,[†] Shi-Ping Ma,[‡] Jing-Han Liu,[‡] and Li-Hong Hu^{*†}

Shanghai Research Center for Modernization of Traditional Chinese Medicine, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, 199 Guo Shou Jing Road, Zhangjiang Hi-Tech Park, Shanghai 201203, People's Republic of China, and College of Traditional Chinese Medicine, China Pharmaceutical University, 24 Tong Jia Xiang, Nanjing 210009, Jiangsu, People's Republic of China

Received December 30, 2007

Five new lindenane sesquiterpene dimers (**1–5**), named shizukaols K–O, and eight known sesquiterpene dimers were isolated from the roots of *Chloranthus fortunei*. The structures of **1–5** were elucidated using spectroscopic data, mainly 1D NMR, 2D NMR, and mass spectra.

In the course of searching for biologically active substances from traditional Chinese medicines, lindenane sesquiterpenoids were reported to exhibit antifungal activity^{1–3} and moderate cytotoxicity.⁴ A series of lindenane dimers have been isolated from *Chloranthus* spp.,^{1,5–12} and their complex structures and biological activities have been of interest to natural product chemists. Chloramultilide B, isolated from *Chloranthus spicatus*, was reported to show inhibitory activity against *Candida albicans* and *Candida parapsilosis*.¹ Chlorahololides A and B isolated from *Chloranthus holostegius* exhibited potent and selective inhibition of the delayed rectifier (I_K) K^+ current.⁵ Shizukaol B, cycloshizukaol A, and shizukaol F isolated from *Chloranthus japonicus* inhibited the expression of cell adhesion molecules.¹³

Chloranthus fortunei (A. Gray) Solms-Laub (Chloranthaceae) has been used in Chinese folk medicine for the treatment of bone fractures,¹⁴ but the chemical constituents of this plant have not been investigated. Thus, 13 compounds were isolated in our search for structurally unique lindenane dimers from the roots of *C. fortunei*. Five of the compounds were identified as new sesquiterpene dimers (**1–5**) and were named shizukaols K–O. The other eight known compounds were 13'-acetylshizukaol C¹² and shizukaols A,¹¹ B,¹⁰ D,¹⁰ E,⁸ F,⁸ I,⁸ and J.⁷ Their structures and relative configurations were elucidated using spectroscopic data, mainly 1D NMR, 2D NMR, and mass spectra.

Results and Discussion

Shizukaol K (**1**) was obtained as a white powder. The molecular formula was determined as $C_{38}H_{44}O_{11}$ from the HRESIMS (m/z 699.2784 [$M + Na$]⁺), which indicated 17 degrees of unsaturation. The IR spectrum revealed the presence of OH (3469.4 cm^{-1}) and carbonyl (1758.8 cm^{-1}) groups. The ¹³C NMR and DEPT experiments displayed 38 carbon resonances, which were ascribed to five carbonyl, eight olefinic, one methoxy, six methyl, six methylene, eight methine, and four quaternary carbons (Table 1). The characteristic high-field methylene signals of **1** at δ_H 0.28 (H-2 β , m) were diagnostic for the cyclopropane ring of a lindenane sesquiterpene.^{7–11} The ¹H–¹H COSY spectrum showed two sets of proton spin systems of a 1,2-disubstituted cyclopropane ring (δ_H 0.28, 0.98, 1.85, and 2.05; 0.69, 1.25, 1.50, and 1.55). Therefore, **1** appeared to be a lindenane dimer. Analysis of the ¹H and ¹³C NMR and mass spectra indicated that **1** was an isomer of 13'-acetylshizukaol C isolated from *Chloranthus japonicus*.¹² Their ¹³C and ¹H NMR data (Tables 1 and 2) were quite similar except that the olefinic proton of the tiglic acid residue in 13'-acetylshi-

Table 1. ¹³C NMR Data (δ) (100 MHz, CDCl₃) for Shizukaols K–O (**1–5**)

position	1	2	3	4	5
1	25.5, CH	26.3, CH	25.5, CH	25.8, CH	25.6, CH
2	15.7, CH ₂	15.8, CH ₂	15.7, CH ₂	15.8, CH ₂	15.7, CH ₂
3	24.5, CH	24.7, CH	24.5, CH	24.7, CH	24.6, CH
4	142.1, qC	142.3, qC	141.9, qC	142.3, qC	142.1, qC
5	131.5, qC	131.6, qC	132.0, qC	131.9, qC	131.9, qC
6	40.6, CH	40.5, CH	40.5, CH	40.8, CH	40.6, CH
7	131.1, qC	131.5, qC	131.8, qC	131.4, qC	131.6, qC
8	200.0, qC	200.7, qC	200.3, qC	200.4, qC	200.4, qC
9	80.0, CH	80.3, CH	79.8, CH	79.9, CH	79.8, CH
10	50.9, qC	51.1, qC	50.8, qC	50.9, qC	50.9, qC
11	147.5, qC	147.5, qC	146.3, qC	147.1, qC	146.6, qC
12	170.3, qC	170.9, qC	171.1, qC	171.5, qC	171.6, qC
13	20.2, CH ₃	20.2, CH ₃	19.8, CH ₃	20.3, CH ₃	20.1, CH ₃
14	15.0, CH ₃	14.3, CH ₃	15.1, CH ₃	15.3, CH ₃	15.1, CH ₃
15	25.1, CH ₂	25.2, CH ₂	25.1, CH ₂	25.2, CH ₂	25.1, CH ₂
1'	25.2, CH	25.4, CH	25.2, CH	25.4, CH	25.3, CH
2'	11.8, CH ₂	12.1, CH ₂	11.7, CH ₂	11.7, CH ₂	11.7, CH ₂
3'	28.1, CH	28.3, CH	28.0, CH	28.0, CH	27.6, CH
4'	76.9, qC	76.9, qC	76.9, qC	76.9, qC	77.0, qC
5'	60.0, CH	60.2, CH	60.4, CH	60.2, CH	60.8, CH
6'	22.4, CH ₂	22.0, CH ₂	22.2, CH ₂	22.3, CH ₂	23.0, CH ₂
7'	172.0, qC	165.0, qC	165.3, qC	168.5, qC	171.0, qC
8'	93.1, qC	92.5, qC	92.5, qC	93.2, qC	93.3, qC
9'	55.5, CH ₂	55.5, CH ₂	55.2, CH ₂	55.1, CH ₂	55.5, CH ₂
10'	44.4, qC	44.6, qC	44.6, qC	44.7, qC	44.6, qC
11'	123.4, qC	124.8, qC	124.3, qC	127.2, qC	123.1, qC
12'	171.1, qC	173.5, qC	173.5, qC	172.4, qC	172.3, qC
13'	55.0, CH ₂	8.7, CH ₂	8.4, CH ₂	54.5, CH ₂	55.7, CH ₂
14'	26.3, CH ₃	25.4, CH ₃	26.1, CH ₃	26.2, CH ₃	26.2, CH ₃
15'	69.7, CH ₂	70.8, CH ₂	70.8, CH ₂	71.6, CH ₂	71.1, CH ₂
OMe	52.3, CH ₃	52.7, CH ₃	52.6, CH ₃	52.7, CH ₃	52.7, CH ₃
1''	166.3, qC	165.5, qC	166.8, qC	171.1, qC	166.9, qC
2''	115.1, CH	144.5, qC	112.5, CH	20.7, CH ₃	112.6, CH
3''	158.4, qC	126.9, CH	159.5, qC		159.4, qC
4''	20.2, CH ₃	170.4, qC	66.7, CH ₂		66.6, CH ₂
5''	27.3, CH ₃	15.1, CH ₃	15.7, CH ₃		15.6, CH ₃
1'''	170.3, qC				172.4, qC
2'''	20.2, CH ₃				28.6, CH ₂
3'''					28.6, CH ₂
4'''					175.1, qC

zukaol C (δ_H 6.81, dd) was shifted upfield and sharpened in **1** (δ_H 5.73, br s). Furthermore, the two methyls (δ_H 1.79 and 1.80) of the tiglic acid residue in 13'-acetylshizukaol C were replaced by methyl signals at δ_H 1.94 and 2.19. In the HMBC spectrum of **1** (Figure 1), H-2'' (δ_H 5.73) correlated to C-1'' (δ_C 166.3), C-4'' (δ_C 20.2), and C-5'' (δ_C 27.3). The pair of signals at δ_H 1.94 and 2.19 (Me-4'' and Me-5'') showed correlations to C-1'', C-2'' (δ_C 115.1) and C-3'' (δ_C 158.4), indicating a senecioic acid moiety. A strong cross-peak of C-1'' (δ_C 166.3) with H₂-15' (δ_H 3.74 and 4.09) determined the senecioic acid residue to be at C-15'. Some key HMBC correlations were observed between H₂-13' (δ_H 4.80) and C-7' (δ_C

* To whom correspondence should be addressed. Tel: +86-21-50272221. Fax: +86-21-50272221. E-mail: simmhulh@mail.shcnc.ac.cn.


[†] Shanghai Research Center for Modernization of Traditional Chinese Medicine.

[‡] China Pharmaceutical University.

Table 2. ^1H NMR Data (300 MHz) for Shizukaols K–O (1–5)

position	1		2		3		4		5	
	δ_{H} (J in Hz) ^a	δ_{H} (J in Hz) ^a	δ_{H} (J in Hz) ^b	δ_{H} (J in Hz) ^a	δ_{H} (J in Hz) ^a	δ_{H} (J in Hz) ^a	δ_{H} (J in Hz) ^a	δ_{H} (J in Hz) ^a	δ_{H} (J in Hz) ^a	δ_{H} (J in Hz) ^a
1	2.05 m	2.02 m	2.39 m	2.05 m	2.05 m	1.98 m				
2 α	0.98 m	0.96 m	1.01 m	1.01 m	0.96 m	1.05 m				
2 β	0.28 m	0.25 m	0.40 m	0.28 m	0.31 m	0.25 m				
3	1.85 m	1.77 m	1.94 m	1.84 m	1.85 m	1.82 m				
6	3.91 bs.	3.91 bs.	4.18 bs.	3.88 d	3.91 d	3.86 d				
9	3.96 s	3.93 s	4.45 s	3.98 s	3.92 s	3.89 s				
13	1.88 s	1.79 s	2.01 s	1.80 s	1.91 s	1.82 s				
14	1.01 s	0.99 s	1.34 s	0.99 s	1.00 s	0.97 s				
15 α	2.75 m	2.72 d	2.95 d	2.75 d (16.5)	2.79 d (14.4)	2.72 m				
15 β	2.59 m	2.48 m	2.68 m	2.47 d (16.5)	2.55 m	2.39 m				
1'	1.55 m	1.55 m	1.54 m	1.59 m	1.60 m	1.53 m				
2' α	0.69 m	0.70 m	0.80 m	0.69 m	0.70 m	0.66 m				
2' β	1.25 m	1.24 m	1.62 m	1.26 m	1.28 m	1.25 m				
3'	1.50 m	1.51 m	1.74 m	1.44 m	1.44 m	1.40 m				
5'	1.75 m	1.68 m	2.15 m	1.86 m	1.89 m	1.71 m				
6' α	2.32 d (5.7)	2.53 m	2.66 m	2.55 dd	2.32 m	2.53 m				
6' β	2.38 d (5.7)	2.58 m	2.85 m	2.61 dd	2.69 m	2.62 m				
9'	1.83 m	1.74 m	1.93 m	1.93 m	1.80 m	1.75 m				
13'	4.80 dd	1.79 s	1.83 s	1.80 s	4.36 dd	4.87 dd				
14'	0.86 s	0.83 s	1.07 s	0.83 s	0.87 s	0.85 s				
15'	3.74 m	3.79 d	4.39 d (11.4)	3.78 d (12.0)	3.81 m	3.76 d				
	4.09 d (11.7)	4.03 d (11.4)	4.50 d (11.4)	4.21 d (12.0)	4.04 d (11.4)	4.20 d				
OMe	3.78 s	3.76 s	3.81 s	3.74 s	3.78 s	3.72 s				
2''	5.73 d			6.01 d	2.11 s	6.00 s				
3''		6.82 s	7.11 s							
4''	1.94 s			4.18 s		4.17 s				
5''	2.19 s	2.28 s	2.52 s	2.12 s		2.10 s				
2'''	2.09 s					2.62 m				
3'''						2.62 m				


^a Measured in CDCl_3 . ^b Measured in pyridine-*d*₅.

**Figure 1.** Key HMBC correlations of shizukaol K (1), shizukaol L (2), and shizukaol O (5).


172.0), C-11' (δ_{C} 123.4), C-12' (δ_{C} 171.1) and between H₃-2''' (δ_{H} 2.09) and C-1''' (δ_{C} 170.3), C-13' (δ_{C} 55.0). Thus, **1** has a senecioid acid residue attached to C-15' rather than tiglic acid as in 13'-acetylshizukaol C.

The relative configuration of **1** was determined by ROESY experiment (Figure 2). ROESY interactions found included H-1/H-9, H-6/H-14, H-6/H-13, H-9'/H-14', and H-9/H-5'. The ROESY correlations between H-3' and H₂-15' indicated that OH-4' was β -oriented. A ROESY interaction was also observed between H-2'' (δ_{H} 5.73) and H-4'' (δ_{H} 1.94) in the senecioid acid substructure. Hence, the structure of **1** was determined to be as shown, and it was named shizukaol K.

Shizukaol L (**2**) was obtained as a white powder, and the molecular formula was determined as $\text{C}_{36}\text{H}_{40}\text{O}_{11}$ from the HRES-IMS (m/z 671.2470 [$\text{M} + \text{Na}$]⁺). Compound **2** was clearly recognized as a sesquiterpene dimer from its ^{13}C and ^1H NMR data (Tables 1 and 2), which were quite similar to that of **1**. The significant differences observed in the NMR spectra were that the 13'-H₂ signal of **1** was absent and an allyl methyl (δ_{H} 1.80, 3H) appeared in **2**. The ^{13}C NMR of **2** showed a corresponding methyl signal in the high-field region (δ_{C} 8.7), characteristic of a 13'-methyl

**Figure 2.** Key ROESY correlations of shizukaol K (1).

group in lindenane sesquiterpenes.^{8,12–14} Moreover, the senecioid acid moiety of **1** disappeared, and five carbon signals (δ_{C} 170.4, 165.5, 144.5, 126.9, 15.1), an olefinic proton (δ_{H} 6.82, s), and a methyl (δ_{H} 2.28, s) appeared in the spectra of **2**. In the HMBC



spectrum (Figure 1), H-3'' (δ_{H} 6.82) correlated with C-1'' (δ_{C} 165.5), C-2'' (δ_{C} 144.5), C-4'' (δ_{C} 170.4), and C-5'' (δ_{C} 15.1), and H-5'' (δ_{H} 2.28) correlated with C-1'', C-2'', C-3'' (δ_{C} 126.9), and C-4''. A strong cross-peak of C-1'' (δ_{C} 165.5) with H₂-15' (δ_{H} 3.79 and 4.03) unambiguously placed the 2-methyl-2-butenedioic acid residue at C-15'. The relative configuration of **2** was also elucidated from correlations observed in the ROESY spectrum. Only the geometry of the double bond on the 2-methyl-2-butenedioic acid moiety could not be established from ROESY correlations, since no ROESY cross-peaks were observed between H-3'' and H-5''. This however suggested that the double bond had *E*-geometry. Pyridine-induced solvent shifts¹⁵ were therefore applied to confirm this result. The significant pyridine-induced solvent shifts (Table 2) for H-5'' ($\Delta\delta$ -0.24) from 4''-OH revealed that they were on the same side; the double bond possessed an *E*-geometry. The structure of **2** was therefore established, and it was named shizukaol L.

Shizukaol M (**3**) was obtained as a yellowish powder, and the molecular formula was determined to be C₃₆H₄₂O₁₀ from HRESIMS (m/z 657.2672 [M + Na]⁺). The similarity of the NMR data suggested the same skeleton for **2** and **3** but different substitution at C-15'. There were also signals assigned to five carbons (δ_{C} 166.8, 112.5, 159.5, 66.7, 15.7), an olefinic proton (δ_{H} 6.01, s), an oxygen-bearing methylene (δ_{H} 4.18, s), and a methyl group (δ_{H} 2.12, s) in the NMR spectra of **3**. Analysis of the key HMBC correlations and comparison of the data with that of Shizukaol I⁸ indicated that there was a γ -hydroxysenecioic acid attached to C-15'. Thus, **3** was concluded to be 13'-deoxyshizukaol I, and it was named shizukaol M.

Shizukaol N (**4**) was obtained as a white powder. The molecular formula was determined to be C₃₃H₃₈O₁₀ by HRESIMS (617.2366 [M + Na]⁺). Compound **4** was also identified as a sesquiterpene dimer, with the same skeleton as **1**–**3**, from its ¹³C and ¹H NMR data (Tables 1 and 2). By key HMBC correlations, the signal at δ_{H} 4.36 (dd, H₂-13') of **4** was assigned to a hydroxylated methylene at C-13'. The remaining two carbons at δ_{C} 171.1 (C-1'') and 20.7

(C-2'') and a methyl at δ_{H} 2.11 (H₃-2'', s) were confirmed to be an acetyl group attached to C-15' by the HMBC correlations observed between H₃-2'' (δ_{H} 2.11) and C-1'' (δ_{C} 171.1) and between H₂-15' (δ 3.81 and 4.04) and C-1''. Thus, the structure of **4** was established, and it was named shizukaol N.

Shizukaol O (**5**) was obtained as a yellowish powder, and the molecular formula was determined as C₄₀H₄₆O₁₄ from the HRESIMS (m/z 773.2783 [M + Na]⁺). The ¹³C and ¹H NMR data (Tables 1 and 2) were quite similar to those of **3**; both had a γ -hydroxysenecioic acid residue on C-15'. The main differences observed were the absence of the characteristic 13'-methyl signal in the high-field region (δ_{C} 8.4) and the addition of four carbons (δ_{C} 175.1, 172.4, 28.6, 28.6) and two methylene (δ_{H} 2.62, each two protons overlay) signals in the spectra of **5**. A succinic acid moiety was established by the key HMBC correlations (Figure 1): H₂-2''' and H₂-3''' (δ_{H} 2.62, each two protons overlay) both correlated with C-1''' (δ_{C} 172.4) and C-4''' (δ_{C} 175.1). A strong cross-peak of C-1''' (δ_{C} 172.4) with H₂-13' (δ_{H} 4.87) accounted for the succinic acid residue on C-13'. It seemed that the coexistence of the OH on γ -hydroxysenecioic acid and the carboxyl on succinic acid was uncommon. However, methylation of **5** with CH₂N₂ yielded a methyl ester that supported our conclusion. Hence the structure of **5** was determined, and it was named shizukaol O.

The eight known dimeric compounds were identified as 13'-acetylshizukaol C¹² and shizukaols A,¹¹ B,¹⁰ D,¹⁰ E,⁸ F,⁸ I,⁸ and J⁷ by comparison with data in the literature.

Experimental Section

General Experimental Procedures. Optical rotations were measured on a Perkin-Elmer 341 polarimeter. UV spectra were obtained on a Shimadzu UV-2450 UV-visible spectrophotometer. IR spectra were measured on a Nicolet FTIR 750 spectrophotometer. ¹H NMR, ¹³C NMR, ¹H–¹H COSY, DEPT, HSQC, HMBC, and ROESY spectra were recorded at 300 MHz for ¹H, at 100 MHz for ¹³C, and at 600 MHz for ROESY with Bruker AMX-300/400/600 instruments in CDCl₃ or pyridine-*d*₅ solution. HRESIMS was carried out using Micromass Q-ToF Global mass spectrometers. ESIMS were recorded on a Bruker Esquire 3000 Plus spectrometer. HPLC was performed with a Waters 2695 separation module equipped with a Waters 2996 photodiode array detector and a Kromacil C18 column (4.6 × 150 mm, 0.5 μ m). All solvents used were of chemical grade and purchased from the Shanghai Chemical Plant, Shanghai, China. Sephadex LH-20 (25–100 μ m) was purchased from Pharmacia. MCI gel CHP 20P (75–150 μ m) was purchased from Mitsubishi Chemical Ind., Tokyo, Japan. RP-18 (20–45 μ m) was purchased from Fuji Silysia Chemical Ltd. Silica gel (200–300 mesh) for column chromatography was purchased from Qingdao Marine Chemical Ltd., Qingdao, China. Silica gel plates (GF-254) for TLC were purchased from Yantai Huiyou Inc., Yantai, China.

Plant Material. The roots of *C. fortunei* were collected from Zhangzhu Town, Yixing City, Jiangsu Province, China, in May 2007 and identified by Prof. Gui-Xin Chou (Shanghai R&D Center for Standardization of Chinese Medicines). A voucher sample (20070530) was deposited at the Shanghai Research Center for Modernization of Traditional Chinese Medicine, Shanghai Institute of Materia Medica, Shanghai, China.

Extraction and Isolation Procedure. Dried and powdered roots of *C. fortunei* (5 kg) were extracted with MeOH (3 × 30 L) at room temperature. The extract was concentrated under reduced pressure to obtain a dark crude extract (470 g), which was suspended in H₂O, then partitioned with EtOAc to afford the EtOAc solubles (205 g). The EtOAc solubles were then subjected to a column of MCI gel eluted with 30%, 75%, and 90% aqueous MeOH, and 72 g of the 75% aqueous MeOH fraction (a major fraction containing the sesquiterpene dimers) was separated on a silica gel column eluted with petroleum ether–acetone (10:1–1:1) to yield eight fractions, I–VIII. Fraction III (1.2 g) was chromatographed on an RP-18 column, using 75% aqueous MeOH, to yield shizukaol A (31 mg). Fraction IV (11.4 g) was first subjected to a silica gel column using CHCl₃–acetone (10:1) as eluent, then separated further on Sephadex LH-20 (CHCl₃–MeOH (1:1)) and RP-18 (60% aqueous MeOH) columns to yield 13'-acetylshizukaol C (3.6 g), shizukaol E (12 mg), and shizukaol J (6 mg). Fraction V (1.9 g) was subjected to a RP-18 column (65% aqueous MeOH) to yield

compound **1** (30 mg). Fraction VI (32 g) was first subjected to a silica gel column using CHCl₃–acetone (8:2) as eluent, then purified on Sephadex LH-20 (CHCl₃–MeOH (1:1)) and RP-18 (55% aqueous MeOH) columns to yield shizukaol F (10.3 g), shizukaol B (285 mg), shizukaol D (406 mg), compound **3** (460 mg), and compound **4** (32 mg). Fraction VII (2.2 g) was first subjected to a Sephadex LH-20 column (CHCl₃–MeOH (1:1)), then further separated on a RP-18 column (55% aqueous MeOH) to yield shizukaol I (120 mg) and compound **2** (11 mg). Fraction VIII (1.7 g) was subjected to a Sephadex LH-20 column (CHCl₃–MeOH (1:1)), then separated further on a RP-18 column (50% aqueous MeOH) to yield compound **5** (320 mg).

Shizukaol K (1): white powder; $[\alpha]_D^{20}$ –150 (c 0.3, MeOH); UV (MeOH) λ_{\max} (log ϵ) 224 (4.54); IR (KBr) ν_{\max} 3469.4, 2935.2, 1758.8, 1695.1, 1648.9, 1438.7, 1376.9, 1228.5, 1141.7, 1085.7, 1031.7, 993.2 cm⁻¹; ¹³C and ¹H NMR data, see Tables 1 and 2; ESIMS m/z 699.4 [M + Na]⁺; HRESIMS m/z 699.2784 (calcd for C₃₈H₄₄O₁₁Na, 699.2781).

Shizukaol L (2): white powder; $[\alpha]_D^{20}$ –118.7 (c 0.3, MeOH); UV (MeOH) λ_{\max} (log ϵ) 224 (4.37); IR (KBr) ν_{\max} 3446.2, 2950.6, 1731.8, 1646.9, 1436.7, 1376.9, 1278.6, 1195.7, 1124.3, 1085.7, 995.1 cm⁻¹; ¹³C and ¹H NMR data, see Tables 1 and 2; ESIMS m/z 671.2 [M + Na]⁺, 647.5 [M – H]⁺; HRESIMS m/z 671.2470 (calcd for C₃₆H₄₀O₁₁Na, 671.2468).

Shizukaol M (3): yellowish powder; $[\alpha]_D^{20}$ –173.1 (c 0.3, MeOH); UV (MeOH) λ_{\max} (log ϵ) 224 (4.65); IR (KBr) ν_{\max} 3453.9, 2948.7, 1735.6, 1604.5, 1436.7, 1280.5, 1224.6, 1139.7, 1085.7, 995.1 cm⁻¹; ¹³C and ¹H NMR data, see Tables 1 and 2; ESIMS m/z 1291.3 [2M + Na]⁺, 633.4 [M – H]⁺; HRESIMS m/z 657.2672 (calcd for C₃₆H₄₂O₁₀Na, 657.2676).

Shizukaol N (4): white powder; $[\alpha]_D^{20}$ –128.0 (c 0.3, MeOH); UV (MeOH) λ_{\max} (log ϵ) 224 (4.20); IR (KBr) ν_{\max} 3434.7, 2933.2, 2250.6, 1735.6, 1602.6, 1436.7, 1378.9, 1263.2, 1085.7, 991.2 cm⁻¹; ¹³C and ¹H NMR data, see Tables 1 and 2; ESIMS m/z 617.2 [M + Na]⁺, 593.2 [M – H]⁺; HRESIMS m/z 617.2366 (calcd for C₃₃H₃₈O₁₀Na, 617.2363).

Shizukaol O (5): yellowish powder; $[\alpha]_D^{20}$ –116.8 (c 0.3, MeOH); UV (MeOH) λ_{\max} (log ϵ) 224 (4.67); IR (KBr) ν_{\max} 3448.2, 2935.2, 1731.8, 1604.5, 1436.7, 1280.5, 1224.6, 1157.1, 1085.7, 991.2 cm⁻¹; ¹³C and ¹H NMR data, see Tables 1 and 2; ESIMS m/z 773.1 [M + Na]⁺, 749.5 [M – H]⁺; HRESIMS m/z 773.2783 (calcd for C₄₀H₄₆O₁₄Na, 773.2785).

Methylation of 5. Freshly prepared CH₂N₂ in Et₂O was added to an ethereal solution of **5** (20 mg) at 0 °C, and the reaction mixture was stirred overnight, allowing the temperature to rise to RT. The solvent was evaporated to obtain the C-4''' methyl ester of **5** (15 mg): yellowish powder; ¹H NMR (300Mz, CDCl₃) δ_H 5.96 (H-2'', s), 4.77 (H-13', dd), 4.16 (H-4'', s), 3.97 (H-9, s), 3.91 (H-6, d), 3.72 (CH₃O-12, s), 3.65

(CH₃O-4''', s), 2.11 (H₃-5'', s), 1.87 (H₃-13, s), 0.98 (H₃-14, s), 0.85 (H₃-14', s), 0.71 (H-2'α, m), 0.28 (H-2β, m); ESIMS m/z 787.3 [M + Na]⁺, 763.6 [M – H]⁺.

Acknowledgment. Financial support by the Chinese National Natural Science Foundation of China (NNSF; No. 30371679, 30623008), Chinese National High-tech R&D Programs (2006AA020602, 2006AA02Z156, 2007AA02Z147), and Shanghai Science and Technology Commission (06DZ22028) is gratefully acknowledged.

Supporting Information Available: ¹H NMR, ¹³C NMR, HMBC, ROESY, and ESIMS spectra of shizukaols K–O (**1–5**) and the methyl ester of **5** are available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- Xu, Y. -J.; Tang, C. -P.; Ke, C. -Q.; Zhang, J. -B.; Weiss, H. -C.; Gesing, E. -R.; Ye, Y. *J. Nat. Prod.* **2007**, *70*, 1987–1990.
- Kawabata, J.; Tahara, S.; Mizutani, J. *Agric. Biol. Chem.* **1981**, *45*, 1447–1453.
- Tahara, S.; Fukushi, Y.; Kawabata, J.; Mizutani, J. *Agric. Biol. Chem.* **1981**, *45*, 1511–1512.
- Masaaki, U.; Yutaka, K.; Genjiro, K.; Yoshikazu, K.; Shigeo, N.; Chizuko, K.; Tsunematsu, T. *Chem. Pharm. Bull.* **1980**, *28*, 92–102.
- Yang, S.-P.; Gao, Z.-B.; Wang, F.-D.; Liao, S.-G.; Chen, H.-D.; Zhang, C.-R.; Hu, G.-Y.; Yue, J.-M. *Org. Lett.* **2007**, *9*, 903–906.
- Yang, S.-P.; Yue, J.-M. *Tetrahedron Lett.* **2006**, *47*, 1129–1132.
- Kawabata, J.; Fukushi, E.; Mizutani, J. *Phytochemistry* **1998**, *47*, 231–235.
- Kawabata, J.; Fukushi, E.; Mizutani, J. *Phytochemistry* **1995**, *39*, 121–125.
- Kawabata, J.; Fukushi, E.; Mizutani, J. *Phytochemistry* **1993**, *32*, 1347–1349.
- Kawabata, J.; Mizutani, J. *Phytochemistry* **1992**, *31*, 1293–1296.
- Kawabata, J.; Fukushi, Y.; Tahara, S.; Mizutani, J. *Phytochemistry* **1990**, *29*, 2332–2334.
- Kim, S. U.; Lee, Y. M.; Park, K. D.; Kang, T. H.; Kim, S. E.; Rho, M.-C.; Moon, J. S.; Choi, W. S. U.S. Pat. Appl. Publ. 2007154574 A1, 2007.
- Kwon, O. E.; Lee, H. S.; Lee, S. W.; Bae, K.; Kim, K.; Hayashi, M.; Rho, M.-C.; Kim, Y.-K. *J. Ethnopharm.* **2006**, *104*, 270–277.
- Editorial Committee of the Administration Bureau of Traditional Chinese Medicine. In *Chinese Materia Medica (Zhonghua Bencao)*; Shanghai Science & Technology: Shanghai, 1998; Vol. 3, pp 449–450.
- Demarco, P. V.; Farkas, E.; Doddrell, D.; Mylari, B. L.; Wenkert, E. *J. Am. Chem. Soc.* **1968**, *90*, 5480–5486.

NP7007544