

Tyrosinase-Inhibitory Constituents from the Twigs of *Cinnamomum cassia*Tran Minh Ngoc,[†] IkSoo Lee,[†] Do Thi Ha,[†] HongJin Kim,[†] ByungSun Min,[‡] and KiHwan Bae^{*†}

College of Pharmacy, Chungnam National University, Daejeon 305-764, South Korea, and College of Pharmacy, Catholic University of Daegu, Gyeongsan 712-702, South Korea

Received January 19, 2009

A methanol extract of the twigs of *Cinnamomum cassia* was found to possess inhibitory activity against tyrosinase. Purification of the MeOH extract afforded four new phenolics, cassiferaldehyde (**6**), icaricide DC (**9**), cinnacassinol (**10**), and dihydrocinnacasside (**13**), together with 10 known compounds (**1–5**, **7–12**, and **14**). The structures of the new compounds were determined by spectroscopic data interpretation. Compounds **1–6** and **8–13** showed strong inhibitory activity against tyrosinase, with IC₅₀ values ranging from 0.24 to 0.94 mM.


Cinnamomum cassia Blume (Lauraceae) is distributed in the southern part of mainland China, Myanmar, Laos, and Vietnam. The twigs are used in traditional Chinese medicine for treating dyspepsia, gastritis, blood circulation disturbances, diabetes, and inflammatory diseases.¹ Chemical and pharmacological investigations on *C. cassia* have resulted in the isolation of several bioactive compounds such as cinnamaldehyde and cinnamic acid, as well as coumarins, diterpenoids, and polyphenols,^{2,3} which have been found to exhibit antifungal, cytotoxic, antipyretic, antioxidant, and antimicrobial activities.^{4–9} In the course of a program to screen for tyrosinase inhibitors from plants, it was found that a MeOH extract of the twigs of *C. cassia* exhibited a strong inhibitory activity (>85% inhibition at 100 μg/mL). Since it has been reported that the tyrosinase inhibitors may have antimelanin synthesis activity and offer a potential treatment for Parkinson's disease,¹⁰ a phytochemical investigation was carried out. The present report describes the isolation of 14 phenolics (**1–14**) from the MeOH extract of the twigs of *C. cassia*, including the characterization of four new compounds (**6**, **9**, **10**, and **13**). The inhibitory activity of compounds **1–14** against tyrosinase has been evaluated.

chromatography of these fractions resulted in the purification of compounds **1–14**. The known compounds were identified as cinnamaldehyde (**1**),¹¹ 2-methoxycinnamaldehyde (**2**),¹² 2-hydroxycinnamaldehyde (**3**),¹³ cinnamic acid (**4**),¹⁴ coniferaldehyde (**5**),¹⁵ cinnamic alcohol (**7**),¹⁶ *o*-coumaric acid (**8**),¹⁷ dihydromelilotoside (**11**),¹⁸ methyl dihydromelilotoside (**12**),¹⁹ and rosavin (**14**).²⁰

Compound **6** was obtained as a colorless, amorphous powder, and its positive HREIMS showed a molecular ion peak [M]⁺ at *m/z* 178.0631, consistent with the molecular formula C₁₀H₁₀O₃. Its IR spectrum showed absorption bands for a hydroxy group and a carbonyl aldehyde, at ν_{\max} 3280 and 1673 cm⁻¹, respectively. The ¹H NMR spectrum displayed a singlet methoxy resonance and signals for two *trans*-olefinic protons and three aromatic protons. The ¹³C and DEPT NMR data (Table 1) of **6** were very similar to those of coniferaldehyde (**5**)¹⁵ except for signals indicating the locations of a methoxy group at C-2 (δ_C 148.9) and a hydroxy group at C-3 (δ_C 152.1). This was supported by the long-range correlations in the HMBC spectrum from the methoxy protons and the H-7 olefinic proton to C-2 and from the aromatic protons H-4 and H-5 to C-3 (Figure 1). Therefore, **6** was determined as a new cinnamaldehyde derivative and has been named cassiferaldehyde.

Compound **9** was obtained as a colorless gum, with $[\alpha]_D^{25} +22.5$. Its IR spectrum showed an absorption band at 1458 cm⁻¹ due to an aromatic function and strong absorption bands at 3399 and 1075 cm⁻¹ suggestive of a glycosidic structure. The molecular formula was established as C₁₉H₂₈O₁₀ from a molecular ion peak at *m/z* 416.1672 [M]⁺ in the HREIMS. The ¹H and ¹³C NMR data (Table 1) of **9** indicated the presence of a β -D-glucopyranosyl moiety, a β -D-apiofuranosyl moiety, and ethylphenyl resonances. The ¹H and ¹³C NMR data were very similar to those of icaricide D₁,²¹ with a downfield shift of C-4' (δ_C 78.8) indicating the linkage to C-1'' of an apiofuranosyl moiety. Analysis of the HBMBC spectrum of **9** showed long-range correlations between the H-4' proton and C-1'' carbon and between the H-1'' proton and the C-4' carbon (Figure 1). Consequently, the structure of icaricide DC (**9**) was determined as phenylethyl-1-*O*- β -D-apiofuranosyl-(1→4)- β -D-glucopyranoside.

Compound **10** was purified as an amorphous solid. The molecular formula, C₃₉H₅₄O₂₀, was observed from a molecular ion peak at *m/z* 842.8362 [M]⁺ in the HREIMS. Combined analysis of the ¹H, ¹³C, DEPT, and HMQC NMR spectra (Table 2) of **10** revealed the presence of an *o*-hydroxycinnamic alcohol skeleton possessing an oxygenated methylene signal, two *trans*-olefinic resonances, and *o*-hydroxyphenyl and phenyl methanol moieties. The ¹H and ¹³C NMR spectra also suggested the presence of four sugar moieties of two β -D-glucopyranosyl units, one β -D-apiofuranosyl unit, and one α -L-rhamnopyranosyl unit. The HMBC spectrum displayed key correlations of the H-1' aromatic proton to the C-2 carbon and the H-7'''' oxymethine protons to the C-1'' carbon, showing the connectivities of the β -D-glucopyranosyl to the cinnamic alcohol



The MeOH extract of the twigs of *C. cassia* was partitioned into hexane-, EtOAc-, and BuOH-soluble fractions. Repeated column

* To whom correspondence should be addressed. Tel: 82-42-821-5925. Fax: 82-42-823-6566. E-mail: baekh@cnu.ac.kr.

[†] Chungnam National University.

[‡] Catholic University of Daegu.

Table 1. NMR Spectroscopic Data for **6**, **9**, and **13** (in MeOD)

position	6		9		13	
	δ_C , mult.	δ_H (J in Hz)	δ_C , mult.	δ_H (J in Hz)	δ_C , mult.	δ_H (J in Hz)
1	129.5, qC		140.2, qC		131.5, qC	
2	148.9, qC		129.5, CH	7.26 d (1.6, 7.6)	157.1, qC	
3	152.1, qC		130.1, CH	7.26 d (1.6, 7.6)	123.5, CH	6.91 dt (2.8, 7.2)
4	120.9, CH	6.95 dd (1.5, 7.8)	127.3, CH	7.17 m	128.8, CH	7.14 d (7.2) overlap
5	125.9, CH	7.00t (7.8)	130.1, CH	7.26 d (1.6, 7.6)	116.4, CH	7.13 d (7.2) overlap
6	120.1, CH	7.17 dd(1.5, 7.8)	129.6, CH	7.26 d (1.6, 7.6)	131.1, CH	7.12 d (7.2) overlap
7	150.1, CH	7.90 d (16.2)	71.7, CH ₂	3.74 dt (9.6, 7.2)	27.1, CH ₂	2.97 m
8	130.6, CH	6.79 dd (7.8, 16.2)	37.4, CH ₂	2.92 t (7.2)	35.5, CH ₂	2.62 t (7.6)
9	196.7, CHO	9.67 d (7.8)			177.7, qC	
Glc-1'			103.4, CH	4.35 d (7.6)	102.7, CH	4.92 d (7.6)
2'			78.7, CH	3.51 m	75.1, CH	3.51 m
3'			78.0, CH	3.42 m	78.3, CH	3.42 m
4'			78.8, CH	3.41 m	71.5, CH	3.41 m
5'			78.0, CH	3.49 m	78.2, CH	3.49 m
6'			62.8, CH ₂	3.96 d (12.0)	62.6, CH ₂	3.89 d (12.0) 3.71 dd (5.2, 12.0)
Apio-1''				3.85 dd (2.4, 12.0)		
2''			110.6, CH	5.38 d (1.6)		
3''			77.9, CH	3.93 d (1.6)		
4''			80.8, qC			
5''			75.4, CH ₂	3.96 d (9.6)		
3.68 d (9.6)			66.3, CH ₂	3.59 dd (11.2, 14.2)		
OCH ₃	61.8, CH ₃	3.85 s				

at C-2, and another β -D-glucopyranosyl to phenylmethanyl at C-7'''. The correlations from H-1''' to C-6' and from H-1''' to C-6'' indicated the linkages of the α -L-rhamnopyranosyl and the β -D-apiofuranosyl to C-6' and C-6'', respectively. The $^3J_{C,H}$ correlations from H-2' to C-2'' and H-2'' to C-2' were supportive of the (2-2) linkage of two the β -D-glucopyranosyl units (Figure 1). Accordingly, the structure of cinnacassiol (**10**) was assigned as 2-[1-O-(phenylmethanyl)-O- β -D-apiofuranosyl-(1-6)- β -D-glucopyranosyl]-(2-2)(O- α -L-rhamnopyranosyl)-(1-6)- β -D-glucopyranosyl-cinnamic alcohol.

Compound **13** was isolated as a white amorphous powder, with $[\alpha]_D^{25} +29.1$, and showed a $[M]^+$ ion at m/z 328.1168 in the HREIMS, corresponding to the molecular formula C₁₅H₂₀O₈. The IR spectrum of **13** showed a glycosidic nature from absorptions at 3354 and 1087 cm⁻¹, and a band at 1735 cm⁻¹ for a carbonyl ester group. The ¹H NMR spectrum displayed the following representative signals: a propanoyl proton, an *o*-hydroxyphenyl resonance, and an aromatic proton of a β -D-glucopyranosyl moiety. The ¹H and ¹³C NMR data (Table 1) of **13** were closely comparable with those of dihydromelilotoside (**11**).¹⁸ Analysis of the 2D NMR data

allowed the assignment of the esterification location of the β -D-glucopyranosyl group in **13** at C-9, which was supported by the key HMBC correlation between the H-1' and the carbonyl carbon C-9 (δ_C 177.7) (Figure 1). Accordingly, the structure of dihydrocinnacasside (**13**) was elucidated as *o*-hydroxyphenylpropanoyl-O- β -D-glucopyranoside.

Compounds **1-14**, together with kojic acid and hydroquinone as positive controls, were examined for their inhibitory effects on tyrosinase activity in an in vitro assay.²² The results are presented in Table 3, and it may be seen that compounds **1-6** and **8-13** exhibited strong inhibition of tyrosinase with IC₅₀ values ranging from 0.24 to 0.94 mM. Of the positive controls used, hydroquinone is used as a commercial whitening agent in cosmetics.²³ The other compounds (**7** and **14**) were inactive (IC₅₀ >10 mM).

Experimental Section

General Experimental Procedures. Specific rotations were measured on a JASCO DIP-370 polarimeter. UV spectra were obtained using a Beckman DU-650 recording spectrophotometer. FT-IR spectra were carried out using a JASCO Report-100 infrared spectrometer. FT-NMR spectra were recorded on a Bruker DRX-300 spectrometer (¹H NMR, 300 MHz; ¹³C NMR, 75 MHz) using CD₃OD as the solvent and tetramethylsilane (TMS) as an internal standard. Chemical shifts (δ) are expressed in ppm with reference to the TMS signals. Two-dimensional (2D) NMR (HMQC, HMBC) experiments were performed on a Bruker Avance 500 spectrometer. HRMS was measured on a JMS-700 Mstation mass spectrometer. Semipreparative HPLC was conducted on Trilution LC with a UV/vis-151 detector, a 321 pump, a 402 syringe pump, and a GX-271 liquid handler (Gilson, Inc.), using a YMC-pack Pro C₁₈ (250 × 20 mm, i.d.) column. Column chromatography was performed using silica gel (Kieselgel 60, 70-230 mesh and 230-400 mesh, Merck). Thin-layer chromatography (TLC) was performed on Merck precoated silica gel 60 F₂₅₄ and/or RP-18 F_{254s} plates (0.25 mm), and compounds were observed under UV light at 254 and 365 nm or visualized by spraying the dried plates with 10% H₂SO₄, followed by heating at 180 °C. Optical density (OD) values in the tyrosinase inhibitory activity assays were read on an Emax Precision microplate reader.

Plant Material. The dried twigs of *C. cassia* were purchased from a pharmacy in Daejeon, Korea, in November 2006, and were identified by one of us (K.B.). A voucher specimen (CN 95125) was deposited at the herbarium in the College of Pharmacy, Chungnam National University.




Figure 1. Selected HMBC correlations of compounds **6**, **9**, **10**, and **13**.

Table 2. NMR Spectroscopic Data for **10** (in MeOD)

position	δ_C , mult.	δ_H (J in Hz)	position	δ_C , mult.	δ_H (J in Hz)
1	128.5, qC		1'''	111.7, CH	4.97 d (2.5)
2	156.3, qC		2'''	78.2, CH	3.90 d (2.5)
3	117.4, CH	7.18 dd (1.5, 7.5)	3'''	80.6, qC	
4	129.9, CH	7.21 dd (1.5, 7.5)	4'''	75.3, CH	3.95 d (10.0) 3.74 d (10.0)
5	123.7, CH	6.98 br t (7.5)	5'''	65.8, CH ₂	3.58 dd (1.5, 11.0)
6	127.4, CH	7.48 dd (1.5, 7.5)	1''''	102.5, CH	4.78 d (1.0)
7	126.6, CH	7.07 d (16.0)	2''''	72.4, CH	3.87 m
8	130.6, CH	6.34 dd (5.5, 16.0)	3''''	72.6, CH	3.70 m
9	64.2, CH ₂	4.25 dd (1.5, 5.5)	4''''	74.2, CH	3.39 m
1'	103.0, CH	4.48 d (8.0)	5''''	70.0, CH	3.70 m
2'	75.1, CH	3.51 m	6''''	18.2, CH ₃	1.27 d (6.3)
3'	78.2, CH	3.45 m	1'''''	139.0, qC	
4'	71.9, CH	3.29 m	2'''''	129.5, CH	7.42 brd (7.5)
5'	77.2, CH	3.58 m	3'''''	129.4, CH	7.33 d (7.5)
6'	68.3, CH ₂	3.67 m	4'''''	128.9, CH	7.27 br t (7.5)
		4.01 dd (2.0, 11.5)			
1''	103.3, CH	4.33 d (7.5)	5'''''	129.4, CH	7.33 d (7.5)
2''	75.1, CH	3.23 m	6'''''	129.5, CH	7.42 brd (7.5)
3''	78.3, CH	3.30 m	7'''''	71.9, CH ₂	4.87 d (12.5) 4.46 d (12.5)
4''	71.8, CH	3.36 m			
5''	77.1, CH	3.38 m			
6''	68.9, CH	3.61 m			
		4.02 dd (1.5, 11.0)			

Table 3. Inhibitory Activity of Isolated Compounds **1–14** against Tyrosinase

compound	inhibitory activity IC ₅₀ (mM) ^a
cinnamaldehyde (1)	0.52 ± 0.03
2-methoxy cinnamaldehyde (2)	0.42 ± 0.02
2-hydroxy cinnamaldehyde (3)	0.47 ± 0.03
cinnamic acid (4)	0.41 ± 0.01
coniferaldehyde (5)	0.24 ± 0.03
cassiferaldehyde (6)	0.26 ± 0.04
cinnamic alcohol (7)	>10
<i>O</i> -coumaric acid (8)	0.67 ± 0.03
icaraside DC (9)	0.71 ± 0.03
cinnacassinol (10)	0.94 ± 0.04
dihydromelilotoside (11)	0.57 ± 0.01
methyl dihydromelilotoside (12)	0.63 ± 0.02
dihydrocinnacasside (13)	0.67 ± 0.03
rosavin (14)	>10
kojic acid ^b	0.14 ± 0.02
hydroquinone ^b	4.21 ± 0.05

^a Values present mean ± SD of triplicate experiments. ^b Positive control.

Extraction and Isolation. The sliced twigs of *C. cassia* (30 kg) were extracted with hot MeOH (60 L × 3 times) for two days. The MeOH extracts were filtered, combined, and concentrated in vacuo, resulting in the production of a residue (1650 g). This residue was suspended in H₂O and then fractionated successively with hexane, EtOAc, and *n*-BuOH, producing a hexane-soluble (410 g), EtOAc-soluble (230 g), and *n*-BuOH-soluble (135 g) fraction, respectively. The hexane-soluble fraction (410 g) was subjected to silica gel column chromatography, eluted with hexane and EtOAc gradient mixtures (100:1–0:100), to give seven fractions, H1–H7. Fraction H1 was chromatographed on silica gel, eluting with hexane–EtOAc (25:1), as well as ODS silica gel with MeOH–H₂O (55:45), to afford **1** (6.0 g) and **2** (1.4 g). Fraction H6 was subjected to silica gel column chromatography, eluting with hexane–EtOAc (10:1), and then purification by medium-pressure liquid chromatography (MPLC) (YMC-ODS 15 × 30 mm, 5 μm), eluted with MeOH–H₂O (60:40), to give **3** (1.2 g). Compound **4** (2.2 g) was isolated and purified from fraction H7 using YMC-ODS column chromatography, with MeOH–H₂O (60:40) as eluant, and MPLC (YMC-ODS 15 × 300 mm, 5 μm), eluted with MeOH–H₂O (50:50).

The EtOAc-soluble fraction (230 g) was chromatographed on a silica gel column with a stepwise gradient of CHCl₃ and MeOH (60:1 to 0:1), to separate eight fractions, E1–E8. Subfraction E2 was chro-

matographed on a silica gel column, with mixtures of hexane–EtOAc (10:1 to 2:1) as the eluting solvent systems, and semipreparative HPLC (YMC-ODS, 10 × 250 mm, 5 μm), eluted with MeOH–H₂O (40:60), to afford **5** (516 mg) and **6** (4.2 mg). Compound **7** (120 mg) was purified from fraction E4 using silica gel column chromatography, eluted with hexane–EtOAc (3:1), and then purified by MPLC (YMC-ODS 15 × 300 mm, 5 μm), with MeOH–H₂O (60:40) as mobile phase. Subfraction E5 was subjected to silica gel column chromatography, eluted with CHCl₃–MeOH–H₂O (3:1:0.1), to afford **8** (52 mg).

The *n*-BuOH-soluble fraction (135 g) was suspended in H₂O and then applied to a Diaion HP-20 column with a H₂O and MeOH gradient eluent to give fractions B1–B14. Fraction B5 was subjected to passage over silica gel, using CH₂Cl₂–MeOH–H₂O (15:1:0.1 to 1:1:0.1) for elution, to yield nine subfractions, B5.1–B5.9. Subfraction B5.5 was applied to a Sephadex LH-20 column, eluted with MeOH–H₂O (1:5), and then YMC-ODS column chromatography with the mobile phase system MeOH–H₂O (1:4) to give **9** (87 mg) and **10** (2.7 mg). Fraction B6 was chromatographed on a Sephadex LH-20 column with MeOH–H₂O (1:5) as solvent and then purified by reversed-phase HPLC (YMC-ODS, 10 × 250 mm, 5 μm), with MeOH–H₂O (1:5) as mobile phase, yielding **11** (15 mg), **12** (10 mg), and **14** (43 mg).

Cassiferaldehyde (6): colorless, amorphous powder; mp 78–80 °C; UV (MeOH) λ_{\max} (log ϵ) 214 (3.30) and 350 (3.24) nm; IR (KBr) ν_{\max} 3280, 1673, 1586, 1331, 1206, 1106, and 727 cm⁻¹; ¹H and ¹³C NMR data, see Table 1; HREIMS m/z 178.0631 [M]⁺ (calcd for C₁₀H₁₀O₃, 178.0630).

Icariside DC (9): colorless gum; [α]_D²⁵ +22.5 (*c* 0.24, MeOH); UV (MeOH) λ_{\max} (log ϵ) 214 (3.00) nm; IR (KBr) ν_{\max} 3399, 2936, 1458, and 1066 cm⁻¹; ¹H and ¹³C NMR data, see Table 1; HREIMS m/z 416.1672 [M]⁺ (calcd for C₁₉H₂₈O₁₀, 416.1682).

Cinnacassinol (10): white, amorphous powder; [α]_D²⁵ -57.2 (*c* 0.35, MeOH); UV (MeOH) λ_{\max} (log ϵ) 204 (2.31), 248 (1.82), and 284 (1.33) nm; IR (KBr) ν_{\max} 3506, 3354, 2900, 1735, 1240, and 1087 cm⁻¹; ¹H and ¹³C NMR data, see Table 2; HREIMS m/z 842.8362 [M]⁺ (calcd for C₃₉H₅₄O₂₀, 842.8368).

Dihydrocinnacasside (13): colorless, amorphous powder; mp 174–175 °C; [α]_D²⁵ -29.1 (*c* 0.4, MeOH); UV (MeOH) λ_{\max} (log ϵ) 214 (2.55) nm; IR (KBr) ν_{\max} 3354, 2900, 1735, 1240, and 1087 cm⁻¹; ¹H and ¹³C NMR data, see Table 1; HREIMS m/z 328.1168 [M]⁺ (calcd for C₁₉H₂₈O₁₀, 328.1158).

Acid Hydrolysis of Compounds 9, 10, and 13. Solutions of **9** (5.2 mg), **10** (1.2 mg), and **13** (2.5 mg) in 0.5 N H₂SO₄ (dioxane–H₂O, 1:1, 2 mL) were each heated at 100 °C for 2 h. Each reaction mixture was diluted with H₂O (3 mL) and extracted with CHCl₃ (5 mL × 3 times). The H₂O layer was dried in vacuo after neutralization with 1 N NaOH and passed through a Sep-Pak C₁₈ cartridge (Waters, Milford,

MA). Sugars were identified as glucose (R_f 0.16), rhamnose (R_f 0.32), and apiose (R_f 0.38) by TLC in CHCl_3 -MeOH- H_2O (12:6:1), using the authentic samples D-glucose, L-rhamnose, and D-apsiose. The remaining eluate was concentrated to dryness, and the residue was stirred with D-cysteine methyl ester hydrochloride, hexamethyldisilazane, and trimethylsilylchloride in pyridine using the same procedures as in a previous report.²⁴ After the reaction, the supernatant was analyzed by GC [column: capillary column BD-5, 0.25 mm \times 30 m, detector: FID, detector temperature: 300 °C, injector temperature: 270 °C, carrier gas, N_2 ; column temperature, 210 °C]. The peaks corresponding to the D-glucosyl, L-rhamnosyl, and D-apsiosyl derivatives appeared with t_R 8.55, 5.31, and 4.03 min, respectively.

Tyrosinase Inhibitory Activity Assay. Mushroom tyrosinase and L-dopa used for the bioassay were purchased from Sigma Chemical Co. Antityrosinase activity was measured by spectrophotometry, according to the method of Mason and Peterson²⁵ with minor modifications. Each test substance was dissolved in 0.1 mL of 10% DMSO in aqueous solution and incubated with 0.1 mL of 135 U/mL mushroom tyrosinase in phosphate buffer solution (PBS, pH 6.8) at 25 °C for 10 min, and then 0.1 mL of L-dopa (0.5 mM, PBS pH 6.8) was added. The reaction mixture was incubated for 5 min. The amount of dopachrome in the mixture was determined by the optical density (OD) at 475 nm using an Emax Precision microplate reader. Kojic acid and hydroquinone (Sigma) were used as positive control agents. The inhibitory percentage of tyrosinase was calculated as follows: % inhibition = $\{[(A - B) - (C - D)] / (A - B)\} \times 100$ (A: OD at 475 nm without test substance; B: OD at 475 nm without test substance and tyrosinase; C: OD at 475 nm with test substance; D: OD at 475 nm with test substance, but without tyrosinase).

Acknowledgment. This work was supported by a grant from the Korea Food and Drug Administration (08182 Crude Drugs 257).

Supporting Information Available: ^1H , ^{13}C , HMQC, and HMBC spectra of the four new compounds (**6**, **9**, **10**, and **13**). This material is available free charge via the Internet at <http://pubs.acs.org>.

References and Notes

- Loi, D. T. *Vietnamese Medicinal Plants and Ingredients*; Medical Publishing House: Hanoi, 2004; p 862.
- Nohaza, T.; Kashiwada, Y.; Murakami, K.; Tomimasu, T.; Kido, M.; Yagi, A.; Hishioka, I. *Chem. Pharm. Bull.* **1981**, *29*, 2451–2459.

- Yazaki, K.; Okudu, T. *Phytochemistry* **1990**, *29*, 1559–1562.
- Singh, H. B.; Srivastava, M.; Singh, A. B.; Srivastava, A. K. *Allergy* **1995**, *50*, 995–999.
- Kwon, B. M.; Lee, S. H.; Choi, S. U.; Park, S. H.; Lee, C. O.; Cho, Y. K.; Sung, N. D.; Bok, S. H. *Arch. Pharm. Res.* **1998**, *21*, 147–152.
- Kurokawa, M.; Kumeda, C. A.; Yamamura, T.; Shiraki, K. *Eur. J. Pharmacol.* **1998**, *348*, 45–51.
- Lin, C. C.; Wu, S. J.; Chang, C. H.; Ng, L. T. *Phytother. Res.* **2003**, *17*, 726–730.
- Kim, H. O.; Park, S. W.; Park, H. D. *Food Microbiol.* **2004**, *21*, 105–110.
- Cheng, S. S.; Liu, J. Y.; Tsai, K. H.; Chen, W. J.; Chang, S. T. *J. Agric. Food Chem.* **2004**, *52*, 4395–4400.
- Tessari, I.; Bisaglia, M.; Valle, F.; Samori, B.; Bergantino, E.; Mammì, S.; Bubacco, L. *J. Biol. Chem.* **2008**, *283*, 16808–16817.
- Choi, J.; Lee, K. T.; Ka, H.; Jung, W. T.; Jung, H. J.; Park, H. J. *Arch. Pharm. Res.* **2001**, *24*, 418–423.
- Geirsson, J. K. F.; Arnadottir, L.; Jonsson, S. *Tetrahedron* **2004**, *60*, 9149–9153.
- Byoung, M. K.; Young, K. C.; Seung, H. L.; Ji, Y. N.; Song, H. B.; Soo, K. C.; Jeong, A. K.; Ihn, R. L. *Planta Med.* **1996**, *62*, 183–184.
- Huung, D. T. L.; Jo, Y. S.; Lee, M. K.; Bae, K. H.; Kim, Y. H. *Nat. Prod. Sci.* **2000**, *6*, 16–19.
- Sy, L. K.; Brown, G. D. *Phytochemistry* **1999**, *50*, 781–785.
- Mohammad, M.; Mojtahedi, E. A.; Sharifi, R.; Abaee, M. S. *Org. Lett.* **2007**, *9*, 2791–2793.
- Aldrich Library of ^{13}C and ^1H FT NMR Spectra*; Sigma-Aldrich: Milwaukee, 1992; Vol. 2, 1046A, 1046B.
- Taskova, R. M.; Gotfredsen, C. H.; Jensen, S. R. *Phytochemistry* **2005**, *66*, 1440–1447.
- Malakov, P. Y.; Papanov, G. Y.; De La Torre, M. C.; Rodriguez, B. *Fitoterapia* **1998**, *69*, 552–554.
- Kishida, M.; Akita, H. *Tetrahedron: Asymmetry* **2005**, *16*, 2625–2630.
- Miyase, T.; Ueno, A.; Takizawa, N.; Kobayashi, H.; Oguchi, H. *Chem. Pharm. Bull.* **1987**, *35*, 3713–3719.
- Mason, H. S.; Peterson, E. W. *Biochim. Biophys. Acta* **1965**, *111*, 134–146.
- Bernard, P.; Berthon, J. Y. *Int. J. Cosmet. Sci.* **2000**, *22*, 219–226.
- Hara, S.; Okabe, H.; Mihashi, K. *Chem. Pharm. Bull.* **1987**, *35*, 501–506.
- Mason, H. S.; Peterson, E. W. *Biochim. Biophys. Acta* **1965**, *111*, 134–146.

NP900031Q