

Indole Derivatives from a Marine Sponge-Derived Yeast as DPPH Radical Scavengers

Yasumasa Sugiyama,*[†] Yuki Ito,[†] Motofumi Suzuki,[‡] and Akira Hirota[†]

Laboratory of Applied Microbiology, School of Food and Nutritional Sciences, University of Shizuoka, 52-1 Yada, Suruga-ku, Shizuoka 422-8526, Japan, and Microbe Division/Japan Collection of Microorganisms, RIKEN BioResource Center, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

Received August 5, 2009

Two new indole derivatives (**3**, **4**) and three known compounds (**1**, **2**, **5**) were isolated as radical scavengers from the culture filtrate of a marine sponge-derived yeast. Their structures were determined to be tyrosol (**1**), tryptophol (**2**), 2-(1*H*-indol-3-yl)ethyl 2-hydroxypropanoate (**3**), 2-(1*H*-indol-3-yl)ethyl 5-hydroxypentanoate (**4**), and cyclo(L-Pro-L-Tyr) (**5**) on the basis of their spectroscopic data. The absolute configurations of compounds **3** and **5** were determined by chiral HPLC analysis combined with synthesis and Marfey's method, respectively. Each obtained compound was evaluated for DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging activity, and all compounds exhibited weak activities.


Excessive radicals are one of the risk factors for organisms, because they damage DNA and react with other biomolecules such as lipids and proteins. This unfavorable oxidative stress consequently induces various diseases including arteriosclerosis, cardiovascular disease, and cancer. Therefore, antioxidants are expected to be useful tools to combat and prevent those diseases.^{1,2}

In our screening program for antioxidants, we have reported several new compounds from soil-derived microorganisms^{3–7} and fermented foods^{8,9} in the past. We continued to investigate new antioxidants from microorganisms, and recently a yeast strain (USF-HO25) was found as a producer of the DPPH (2,2-diphenyl-1-picrylhydrazyl)¹⁰ radical scavenging compounds. The yeast was separated from the marine sponge *Halichondria okadai* collected at the west coast of Izu Peninsula in Shizuoka Prefecture, Japan. The USF-HO25 strain produced several indole derivatives and phenol derivatives in the culture filtrate. In this paper, we report the fermentation of the USF-HO25 strain and the isolation, structural elucidation, and radical scavenging activities of five compounds, including two novel indole derivatives.

The marine sponge-derived yeast, HO-25 strain, was identified as *Pichia membranifaciens* from the observation of cell and ascospore morphology under the light microscope and the molecular phylogenetic analysis based on the sequence of the D1/D2 region of the 26S rRNA gene. The cultivation of the USF-HO25 strain was conducted on a reciprocal shaker at 30 °C for 4 days. An EtOAc extract from the culture filtrate was successively purified by silica gel column chromatography, repeated Sephadex LH20 column chromatography, and reversed-phase HPLC to yield compounds **1–5**.

Compounds **1** and **2**, respectively, were identified as tyrosol¹¹ and tryptophol¹² on the basis of their spectroscopic data. It had been reported that both compounds were produced by a yeast, and tyrosol showed radical scavenging activity for DPPH.¹³

The molecular formula of **3** was established as C₁₃H₁₅NO₃ from the HR-FABMS data. The ¹H and ¹³C NMR spectra of **3** were similar to those of **2**, except that additional signals of a methyl group (δ_{H} 1.31, δ_{C} 20.6), an oxymethine group (δ_{H} 4.22, δ_{C} 67.9), and a carbonyl carbon (δ_{C} 176.5) were observed in the NMR spectra of **3** (Table 1). Hence, it was supported that compound **3** was a derivative of indole-3-ethanol. In addition, it was clear that a methyl group is connected to an oxymethine group, as exhibited by their



coupling constants in the ¹H NMR spectrum of **3**. The connection of three partial structures was established by HMBC correlations from both a methyl proton (δ_{H} 1.31 (H-3'')) and a methylene proton (δ_{H} 4.38 (H-2'')) to a carbonyl carbon (δ_{C} 176.5 (C-1'')) (Figure 1). Thus, the planar structure of **3** was determined to be 2-(1*H*-indol-3-yl)ethyl 2-hydroxypropanoate, a novel compound.

The absolute structure of **3** was decided by comparison of the chromatogram of a natural product with those of synthetic compounds. At first, we synthesized (*S*)-2-(1*H*-indol-3-yl)ethyl 2-hydroxypropanoate (**3a**) and (*R*)-2-(1*H*-indol-3-yl)ethyl 2-hydroxypropanoate (**3b**), which were reaction products from tryptophol with *L*-lactic acid and *D*-lactic acid, respectively. Subsequently, the synthesized compounds (**3a** and **3b**) and the natural product (**3**) were analyzed by HPLC using a chiral column. Two peaks were detected in the chromatogram of **3**, and they were assigned to **3a** and **3b** on the basis of their retention times. Moreover, **3b** was 1.6 times larger than that of **3a**. These results indicate that **3** is a mixture of the *S*-form and the *R*-form, and the composition ratio is 5:8 (*S*/*R*). Each of synthetic compounds **3a** and **3b** did not convert spontaneously to the opposite configuration when they were purified in a similar manner to the natural product. This phenomenon might indicate that the mixed configuration of **3** was derived from the biosynthetic process.

The molecular formula of **4** was determined to be C₁₅H₁₉NO₃ by the HR-FABMS spectrum. The ¹H and ¹³C NMR spectra of **4** were comparable to those of **2**, and it was suggested that **4** was an analogue of **3**. The signals of an oxymethylene moiety (δ_{H} 3.51, δ_{C} 62.4), three methylene groups (δ_{H} 2.33, 1.63, and 1.49, δ_{C} 34.9, 22.4, and 32.9), and a carbonyl carbon (δ_{C} 175.5) were observed besides the signals corresponding to an indole-3-ethanol moiety in the NMR spectra of **4** (Table 1). Four methylene groups represent an alkyl side chain, –CH₂(2'')–CH₂(3'')–CH₂(4'')–CH₂(5'')–O–,

* To whom correspondence should be addressed. Tel: +81-54-264-5555. Fax: +81-54-264-5099. E-mail: sugiyasu@u-shizuoka-ken.ac.jp.



[†] University of Shizuoka.

[‡] RIKEN BioResource Center.

Table 1. NMR Spectroscopic Data of Compounds **3** and **4** in CD₃OD

position	3			4		
	δ_C^a , mult.	δ_H^b (J in Hz)	HMBC ^c	δ_C , mult.	δ_H (J in Hz)	HMBC
2	123.8, CH	7.09, s	3a, 7a	123.7, CH	7.08, s	3a, 7a
3	111.6, qC			111.9, qC		
3a	128.7, qC			128.8, qC		
4	119.2, CH	7.56, d, (8.0)	6, 7a	119.3, CH	7.55, d (8.0)	6, 7a
5	119.7, CH	7.00, t (8.0, 7.5)	3a, 7	119.6, CH	7.00, t (8.0, 7.5)	3a, 7
6	122.3, CH	7.08, t ^d	4, 7a	122.3, CH	7.08, t ^e	4, 7a
7	112.2, CH	7.33, d (8.0)	3a, 5	112.2, CH	7.32, d (8.0)	3a, 5
7a	138.1, qC			138.1, qC		
1'	25.8, CH ₂	3.10, t (6.9)	2, 3, 3a, 2'	25.8, CH ₂	3.07, t (6.9)	2, 3, 3a, 2'
2'	66.4, CH ₂	4.38, m	3, 1', 1''	66.0, CH ₂	4.32, t (6.9)	3, 1', 1''
1''	176.5, qC			175.5, qC		
2''	67.9, CH	4.22, q (6.9)	1'', 3''	34.9, CH ₂	2.33, t (6.9)	1'', 3'', 4''
3''	20.6, CH ₃	1.31, d (6.9)	1'', 2''	22.4, CH ₂	1.63, quint	1'', 2'', 4'', 5''
4''				32.9, CH ₂	1.49, quint	2'', 3'', 5''
5''				62.4, CH ₂	3.51, t (6.9)	3'', 4''

^a ¹³C NMR spectrum was measured at 125 MHz. ^b ¹H NMR spectrum was measured at 500 MHz. ^c HMBC correlations optimized for 8 Hz are from proton started to the indicated carbons. ^d Coupling constants are unclear due to overlaps. ^e Coupling constants are unclear due to overlaps.

**Figure 1.** Partial structures and HMBC correlations of compound **3**.**Figure 2.** Partial structures and ¹H–¹H COSY and HMBC correlations of compound **4**.

as determined by analysis of the ¹H–¹H COSY spectrum of **4**. The key HMBC correlations were observed from the oxymethylene proton at δ_H 4.32 (H-2') and two methylene protons (δ_H 2.33 (H-2'') and 1.63 (H-3'')) of the alkyl chain to a carbonyl carbon (δ_C 175.5 (C-1'')) (Figure 2). For this reason, it was demonstrated that the alkyl chain was attached to an indole-3-ethanol moiety via a carbonyl carbon (δ_C 175.5). Therefore, the structure of **4** was determined to be 2-(1*H*-indol-3-yl)ethyl 5-hydroxypentanoate as a novel compound.

HR-FABMS, ¹H and ¹³C NMR, ¹H–¹H COSY, HMQC, and HMBC spectroscopic data showed that compound **5** was a cyclic dipeptide consisting of a proline and a tyrosine. Both chiral centers of **5** were determined to have *S* configurations by HPLC analysis of Marfey's derivatives¹⁴ from the hydrolysate. Therefore, the structure of **5** was identified as cyclo(L-Pro-L-Tyr).¹⁵

The five compounds (**1**–**5**) obtained in this study and BHT as a positive control were tested for DPPH radical scavenging activities at 300 μ M (final concentration), and their percentages of scavenged DPPH radical were calculated (Table 2). As a result, the activities of the indolic metabolites (**2**, **3**, and **4**) were slightly higher than the isolated phenolic compounds (**1** and **5**). However, between the three indolic derivatives (**2**, **3**, and **4**) there was no significant difference in their DPPH radical scavenging activities. Therefore, the presence of a side chain does not affect the antioxidative activity of an indolic derivative. In contrast, a high antioxidative activity

Table 2. DPPH Radical Scavenging Activities of Compounds **1**–**5** and BHT

compound ^a	scavenged DPPH radical (%) ^c
1	26
2	52
3	33
4	47
5	16
BHT ^b	86

^a Compounds were used at a final concentration of 300 μ M. ^b Butylhydroxytoluene; 2,6-di-*tert*-butyl-4-methylphenol, used as a positive control. ^c Percentage of scavenged DPPH radical after 48 h.

of tyrosol (**1**) was shown in the ORAC (oxygen radical absorbance capacity) assay, in spite of no activity for scavenging the DPPH radical.¹⁶ High antioxidative activity of tryptophol (**2**) was shown in the ABTS (2,2'-azinobis(3-ethylbenzthiazoline-6-sulfonic acid)) assay under hydrophilic conditions.¹⁷ Many indole-containing compounds are known to present various activities such as anticancer, antibiotic, and anti-inflammatory activities. Accordingly, the two novel indole derivatives (**3** and **4**) are expected to exhibit other biological activities along with their antioxidative properties.

Experimental Section

General Experimental Procedures. Optical rotation was recorded with a Horiba SEPA-200 spectrometer. UV spectra were recorded with a Shimadzu UV-160A spectrometer. NMR spectra were obtained with a JEOL JMN-ECA-500 spectrometer, using tetramethylsilane as an internal standard. High- and low-resolution mass spectra were measured by a JEOL JMS-700 spectrometer, using glycerol as a matrix. The HPLC analyses were carried out on a JASCO HPLC system: PU-2080 Plus and UV-2077 Plus. HPLC for preparation was done by JASCO 880-PU and JASCO UV-970 apparatus.

Fungal Material. Morphological characteristics of the yeast strain were examined by using standard methods described by Yarrow.¹⁸ Its nuclear DNA was isolated from colonies on a YM agar slant incubated at 25 °C for 4 days by using a MasterPure yeast DNA purification kit (Epicenter Technologies, Madison, WI) according to the manufacturer's instructions. The D1/D2 domain of the large-subunit (26S) rRNA gene from DNA was amplified and sequenced as described by Kurtzman and Robnett.¹⁹ A sequence-similarity search was performed by using the BLAST 2.2.21 (Altschul et al.)²⁰ in GenBank. The nomenclature of the yeast strain identified follows the classification of Kurtzman.²¹ The D1/D2 sequence of the yeast strain was identical to that of the type strain of *Pichia membranifaciens* (GenBank accession number U75725).

Fermentation. The USF-HO25 strain was inoculated into a medium (100 mL) that consisted of glucose 1%, polypeptone 0.2%, yeast extract 0.1%, meat extract 0.1%, and NaCl 3%, pH 7.2 in a 500 mL flask; 100 flasks were inoculated and incubated at 30 °C on a reciprocal shaker at 130 rpm for 4 days.

Isolation of DPPH Radical Scavengers from the Culture Broth of USF-HO25. The culture broth (10 L) was filtered, and the filtrate was adjusted to pH 3 and extracted with EtOAc (5 L \times 3). The combined EtOAc layer was dried over anhydrous Na₂SO₄ and filtered, and the solvent was removed under reduced pressure. The EtOAc extract (1.8 g) was applied to a silica gel column and eluted with *n*-hexane/EtOAc at 3:1, 1:1, and 1:3, then with EtOAc and MeOH. Two fractions that eluted with *n*-hexane/EtOAc (1:1) (fraction II, 568.0 mg) and MeOH (fraction V, 399.0 mg) showed DPPH radical scavenging activity in the TLC assay.²² Fractions II and V were further purified three times by Sephadex LH20 column chromatography using MeOH as an eluent. Subfractions 17 and 18 of fraction II obtained from the third Sephadex LH20 column were combined and further purified by preparative RP-HPLC (column: Capcell-Pak C₁₈ SG120, 15 \times 250 mm, Shiseido; mobile phase: 31% CH₃CN/H₂O), to yield compounds **1** (51.6 mg), **2** (44.3 mg), **3** (0.6 mg), and **4** (4.9 mg). Subfractions 6–11 of fraction V were combined and evaporated to dryness *in vacuo*, and the resulting residue was purified by preparative RP-HPLC (column: Capcell-Pak C₁₈ SG120, 15 \times 250 mm, Shiseido; mobile phase: 8% CH₃CN/H₂O) to yield **5** (60.7 mg).

2-(1H-Indol-3-yl)ethyl 2-hydroxypropanoate (3): colorless oil; UV (MeOH) λ_{\max} (log ϵ) 222 (4.48), 282 (3.73) nm; ¹H NMR, see Table 1; ¹³C NMR, see Table 1; HR-FABMS [M + H]⁺ *m/z* 234.1127 (calcd for C₁₃H₁₆NO₃, 234.1124).

2-(1H-Indol-3-yl)ethyl 5-hydroxypentanoate (4): colorless oil; UV (MeOH) λ_{\max} (log ϵ) 222 (4.33), 282 (3.57) nm; ¹H NMR, see Table 1; ¹³C NMR, see Table 1; HR-FABMS [M + H]⁺ *m/z* 262.1444 (calcd for C₁₅H₂₀NO₃, 262.1445).

Preparations of (S)-2-(1H-Indol-3-yl)ethyl 2-hydroxypropanoate (3a) and (R)-2-(1H-Indol-3-yl)ethyl 2-hydroxypropanoate (3b). L-Lactic acid (270 mg) was heated in thionyl chloride (30 mL) for 6 h at 55 °C. After the thionyl chloride was evaporated at low temperature, CH₂Cl₂ (20 mL) containing tryptophol (480 mg) and 4-dimethylaminopyridine (366 mg) was added, and the mixture was stirred for 6 h at room temperature. The reaction mixture was diluted with EtOAc, washed with 0.1 M HCl, and dried over anhydrous Na₂SO₄. The filtrate was evaporated to give crude materials, and they were applied to silica gel column chromatography. The fraction was eluted with *n*-hexane/EtOAc (2:1) and was further purified by preparative RP-HPLC (column: Capcell-Pak C₁₈ SG120, 15 \times 250 mm, Shiseido; mobile phase: 32% CH₃CN/H₂O) to yield **3a** (47.4 mg). Similarly, (R)-2-(1H-indol-3-yl)ethyl 2-hydroxypropanoate (**3b**, 4.0 mg) was prepared from D-lactic acid (45 mg) and tryptophol (81 mg).

(S)-2-(1H-Indol-3-yl)ethyl 2-hydroxypropanoate (3a): colorless oil; [α]_D²⁵ -8.8 (c 0.23, MeOH); ¹H NMR (CD₃OD, 500 MHz) δ 7.56 (1H, d, *J* = 8.0 Hz, H-4), 7.32 (1H, d, *J* = 8.0 Hz, H-7), 7.09 (1H, s, H-2), 7.08 (1H, t, H-6), 7.00 (1H, t, *J* = 8.0, 7.5 Hz, H-5), 4.38 (2H, m, H-2'), 4.22 (1H, q, *J* = 6.9 Hz, H-2''), 3.11 (2H, t, *J* = 6.9 Hz, H-1'), 1.31 (3H, d, *J* = 6.9 Hz, H-3''); ¹³C NMR (CD₃OD, 125 MHz) δ 176.5 (C-1'), 138.1 (C-7a), 128.7 (C-3a), 123.8 (C-2), 122.3 (C-6), 119.7 (C-5), 119.2 (C-4), 112.2 (C-7), 111.6 (C-3), 67.9 (C-2''), 66.4 (C-2'), 25.8 (C-1'), 20.6 (C-3''); HR-FABMS *m/z* 234.1137 [M + H]⁺ (calcd for C₁₃H₁₆NO₃, 234.1130).

(R)-2-(1H-Indol-3-yl)ethyl 2-hydroxypropanoate (3b): colorless oil; [α]_D²¹ +9.3 (c 0.11, MeOH); ¹H and ¹³C NMR data are identical to those for **3a**; HR-FABMS *m/z* 234.1142 [M + H]⁺ (calcd for C₁₃H₁₆NO₃, 234.1130).

Determination of the Configuration of Compound 3. The natural product (**3**) and the synthetic compounds (**3a** and **3b**) were analyzed by chiral HPLC (column: Ceramospher Chiral RU-2, 4.6 \times 250 mm, Shiseido; mobile phase: MeOH; flow rate: 0.5 mL/min; detection: 280 nm). The retention times of synthetic compounds **3a** and **3b** were 21.9 and 17.6 min, respectively. Two peaks that correspond to **3a** and **3b**

were detected in the chromatogram of the natural product (**3**). The ratio of the *S/R* enantiomers in the natural product mixture was calculated from the area of their peaks.

Measurement of DPPH Radical Scavenging Activity. DPPH radical scavenging assay was performed by the modified method presented previously.^{3–9} An ethanolic solution of the sample (750 μ M, 1 mL) was mixed with a 500 μ M DPPH ethanolic solution (0.5 mL) and 0.1 M acetate buffer (pH 5.5, 1 mL). The reaction mixture was left to stand for 48 h, at which time the absorbance of the mixture was measured at 517 nm. The DPPH radical scavenging activity was determined by comparison of the absorbance of sample with that of a blank control. BHT (butylhydroxytoluene; 2,6-di-*tert*-butyl-4-methylphenol) was used as a positive control at the same concentration as other samples.

Acknowledgment. We are grateful to M. Sugiyama for assistance in collecting the marine sponge.

Supporting Information Available: NMR spectra of **1–5** and spectroscopic data and additional information on the known compounds are available free of charge via the Internet at <http://pubs.acs.org>.

References and Notes

- Madamanchi, N. R.; Vendrov, A.; Runge, M. S. *Arterioscler. Thromb. Vasc. Biol.* **2005**, *25*, 29–38.
- Ames, B. N.; Gold, L. S.; Willett, W. C. *Proc. Natl. Acad. Sci. U.S.A.* **1995**, *92*, 5285–5265.
- Abe, N.; Murata, T.; Hirota, A. *Biosci. Biotechnol. Biochem.* **1998**, *62*, 661–666.
- Abe, N.; Murata, T.; Yamamoto, K.; Hirota, A. *Tetrahedron Lett.* **1999**, *40*, 5203–5206.
- Ito, C.; Abe, N.; Hirota, A. *Biosci. Biotechnol. Biochem.* **1999**, *63*, 1993–1996.
- Washida, K.; Abe, N.; Sugiyama, Y.; Hirota, A. *Biosci. Biotechnol. Biochem.* **2007**, *71*, 1052–1057.
- Sugiyama, Y.; Watanabe, K.; Hirota, A. *Biosci. Biotechnol. Biochem.* **2009**, *73*, 230–232.
- Hirota, A.; Inaba, M.; Chen, Y.-C.; Abe, N.; Taki, S.; Yano, M.; Kawaii, S. *Biosci. Biotechnol. Biochem.* **2004**, *68*, 1372–1374.
- Chen, Y.-C.; Sugiyama, Y.; Abe, N.; Kuruto-Niwa, R.; Nozawa, R.; Hirota, A. *Biosci. Biotechnol. Biochem.* **2005**, *69*, 999–1006.
- Blois, M. S. *Nature* **1958**, *181*, 1199–1200.
- SentheShanmuganathan, S.; Elsdén, S. R. *Biochem. J.* **1958**, *69*, 210–218.
- Kradolfer, P.; Niederberger, P.; Hütter, R. *Arch. Microbiol.* **1982**, *133*, 242–248.
- Valavanidis, A.; Nisiotou, C.; Papageorgiou, Y.; Kremli, I.; Satravelas, N.; Zinieris, N.; Zygalki, H. *J. Agric. Food Chem.* **2004**, *52*, 2358–2365.
- Marfey, P. *Carlsberg Res. Commun.* **1984**, *49*, 591–596.
- Tamura, S.; Suzuki, A.; Aoki, Y.; Otake, N. *Agric. Biol. Chem.* **1964**, *28*, 650–652.
- Fernández-Pachón, M. S.; Villaño, D.; Troncoso, A. M.; García-Parrilla, M. C. *Anal. Chim. Acta* **2006**, *563*, 101–108.
- Cano, A.; Alcaraz, O.; Arnao, M. B. *Anal. Bioanal. Chem.* **2003**, *376*, 33–37.
- Yarrow, D. *The Yeasts, A Taxonomic Study*, 4th ed.; Elsevier Science BV: Amsterdam, The Netherlands, 1998; pp 77–100.
- Kurtzman, C. P.; Robnett, C. J. *Antonie van Leeuwenhoek* **1998**, *73*, 331–371.
- Altschul, S. F.; Madden, T. L.; Schäffer, A. A.; Zhang, J.; Zhang, Z.; Miller, W.; Lipman, D. J. *Nucleic Acids Res.* **1997**, *25*, 3389–3402.
- Kurtzman, C. P. *The Yeasts, A Taxonomic Study*, 4th ed.; Elsevier Science BV: Amsterdam, The Netherlands, 1998; pp 273–352.
- Glavind, J.; Holmer, G. J. *Am. Oil Chem. Soc.* **1967**, *44*, 539–542.

NP900483G