

## Gymnochromes E and F, Cytotoxic Phenanthroperylenequinones from a Deep-Water Crinoid, *Holopus rangii*

Hilaire V. Kemami Wangun,<sup>†</sup> Alexander Wood,<sup>‡</sup> Catherine Fiorilla,<sup>‡</sup> John K. Reed,<sup>†</sup> Peter J. McCarthy,<sup>†</sup> and Amy E. Wright<sup>\*†</sup>


Harbor Branch Oceanographic Institute at Florida Atlantic University, Center for Marine Biomedical and Biotechnology Research, 5600 US 1 Fort Pierce, Florida 34946, and Novartis Institutes for Biomedical Research, Cambridge, Massachusetts 02139

Received August 27, 2009

Bioactivity-guided fractionation of metabolites from the crinoid *Holopus rangii* led to the discovery of two new phenanthroperylenequinone derivatives, gymnochromes E (**1**) and F (**2**). Gymnochrome E showed cytotoxic activity toward the NCI/ADR-Res with an  $IC_{50}$  of 3.5  $\mu$ M. It also inhibited histone deacetylase-1 with an  $IC_{50}$  of 3.3  $\mu$ M. Gymnochrome F was a moderate inhibitor of myeloid cell leukemia sequence 1 (MCL-1) binding to Bak. Two anthraquinone metabolites, emodic acid (**4**) and its new bromo derivative (**5**), were also isolated from the crinoid and show remarkable similarity to the phenanthroperylenequinone core, suggesting that these metabolites share the same polyketide biosynthetic pathway.

The past decade has seen a dramatic increase of marine organism derived anticancer lead compounds entering human clinical trials<sup>1</sup> resulting from recent technological advances in structure elucidation, organic synthesis, and biological assays. These lead compounds range in structural class from relatively simple linear peptides such as dolastatin 10,<sup>2</sup> to more complex polyketides such as discodermolide,<sup>3</sup> to very complex macrocyclic polyethers, such as halichondrin B.<sup>4</sup> Equally diverse are the molecular modes of action by which these molecules impart their biological activity and the increasing number of compounds working through novel modes of action. As part of our ongoing program to identify novel natural products with activity in target-directed oncology assays, materials from the HBOI Peak Library (generated by reversed-phase medium-pressure liquid chromatography) were assayed for their ability to inhibit the binding of MCL-1 to Bak using a FRET-based assay.<sup>5</sup> MCL-1 (an antiapoptotic member of the BCL-2 family) binds Bak (a pro-apoptotic BCL-2 member), which upon release from MCL-1 regulates apoptosis. A fraction derived from the crinoid *Holopus rangii* inhibited the binding of MCL-1 to Bak with an  $IC_{50}$  of 10  $\mu$ g/mL. NMR and MS analysis of the fraction suggested the presence of a series of highly unsaturated pigments. Bioassay- and spectroscopy-guided fractionation led to the isolation and characterization of two new members of the phenanthroperylenequinone family of natural products, which we have designated as gymnochromes E (**1**) and F (**2**), the known isogymnochrome B (**3**), and two anthraquinones, emodic acid (**4**) and its 7-bromo derivative (**5**). Here we report the isolation and structure elucidation of compounds **1**, **2**, and **5** as well as their biological activities.

A sample of freeze-dried crinoid was cut into small pieces and crushed. The resultant powder was extracted successively with EtOH and EtOAc/EtOH, 1:1. The combined extracts were concentrated by distillation under reduced pressure to give a dark green residue, which was fractionated on a C-18 stationary phase using vacuum column chromatography. Further purification using reversed-phase HPLC and monitoring by bioassay and mass spectrometry led to the isolation of **1** (3 mg), **2** (5 mg), **3** (1.5 mg), **4** (2.1 mg), and **5** (0.5 mg). The spectroscopic data observed for **3** were consistent with those reported for isogymnochrome B,<sup>6</sup> and **3** has tentatively been assigned as isogymnochrome B. The spectroscopic data observed for **4** were identical to those reported for emodic acid,<sup>7</sup> allowing for its identification.




Compound **1** was isolated as a dark brown oil. The IR spectrum of **1** shows an absorption band observed at 1634  $cm^{-1}$  characteristic of the carbonyl of a hydrogen-bonded quinone. The UV-vis spectrum of **1** showed maxima ( $\lambda_{max}$ ) at 311, 372, 524, and 596 nm that are similar to those of the well-known compound hypericin,<sup>8,9</sup> suggesting that **1** is a phenanthroperylenequinone derivative. The electrospray mass spectrum (ESIMS) of **1** detected in negative mode showed a complex multiplet of three peaks at  $m/z$  775, 777, and 779, suggesting that **1** is dibrominated. The <sup>1</sup>H NMR spectrum of **1** recorded in DMSO-*d*<sub>6</sub> showed the presence of two methyl resonances [ $\delta_H$  -0.09 (H<sub>3</sub>-22), 0.94 (H<sub>3</sub>-17)] as well as an aromatic proton resonance at  $\delta_H$  6.59 (H-9 and H-12) that also suggested that **1** is a dibromophenanthroperylenequinone derivative. In addition to the signals attributable to the dibromohydroxy phenanthroperylenequinone skeleton, analysis of the <sup>13</sup>C and HSQC spectra of **1** revealed the presence of two different aliphatic side chains. The first one consisted of three methylene carbons, an oxymethine carbon, and a methyl carbon. It was unambiguously identified as a 2-hydroxypropyl moiety on the basis of correlations observed in the COSY spectrum that revealed the sequential connectivity of H<sub>2</sub>-18 → H-19 → H<sub>2</sub>-20 → H<sub>2</sub>-21 → H<sub>3</sub>-22. For the second aliphatic side chain a methylene carbon, an oxymethine carbon, and a methyl carbon were identified. The COSY spectrum clearly established the connectivity of H<sub>2</sub>-15 → H-16 → H<sub>3</sub>-17 and assigned the presence of a 2-hydroxypropyl moiety in **1**. This assignment was further supported by the HMBC spectrum,

\* To whom correspondence should be addressed. Phone: 772-242-2459. E-mail: awright33@fau.edu.

<sup>†</sup> Harbor Branch Oceanographic Institute at Florida Atlantic University.

<sup>‡</sup> Novartis Institutes for Biomedical Research.




**Figure 1.** Key HMBC and COSY correlations observed for gymnochrome E (1).

which showed correlations between the methyl protons H<sub>3</sub>-17 ( $\delta$  0.94, d,  $J$  = 6.1) and both C-15 and C-16, as well as correlations between the methylene protons H<sub>2</sub>-15 (H-15a,  $\delta$ <sub>H</sub> 3.47 dd,  $J$  = 8.2, 13.0; H-15b,  $\delta$ <sub>H</sub> 3.64 dd,  $J$  = 13.0, 2.0) and the methyl carbon C-17 ( $\delta$ <sub>C</sub> 24.8). The two aliphatic fragments were connected to the dibromohydroxy phenanthroperylenequinone moiety with the aid of diagnostic correlations observed in the HMBC spectrum (Figure 1), in particular through three-bond connectivity observed between H<sub>2</sub>-18 (H-18a,  $\delta$ <sub>H</sub> 3.56 dd,  $J$  = 13.0, 8.9; H-18b,  $\delta$ <sub>H</sub> 3.73 dd,  $J$  = 13.0, 4.8) and C-5 ( $\delta$ <sub>C</sub> 115.4) and C-3b ( $\delta$ <sub>C</sub> 122.0) as well as connectivity between H<sub>2</sub>-15 and C-3a ( $\delta$ <sub>C</sub> 122.5) and C-2 ( $\delta$ <sub>C</sub> 115.0). On the basis of these data the structure of **1** was assigned as 1,6,8,10,11,13-hexahydroxy-2,5-dibromo-3-(2-hydroxypropyl)-4-(2-hydroxypentyl)phenanthroperylene-7,14-quinone, which we are designating gymnochrome E. Further support for this assignment came from the ESIMS/MS analysis of **1**, which showed a fragment ion at  $m/z$  645, representing the cleavage of the side chains (see Supporting Information). Gymnochrome E represents the 9 or 12 debromo derivative of isogymnochrome B (**3**).

Compound **2** was isolated as a dark brown oil. The UV-vis spectrum as well as the ESIMS/MS fragmentation pattern suggested that **2** is a homologue of **1**. The electrospray mass spectrum (ESIMS) of **2** indicated in negative mode a complex multiplet of five peaks at  $m/z$  959, 961, 963, 965, and 967, suggesting that **2** is tetrabrominated. The lack of an aromatic proton signal in the <sup>1</sup>H NMR spectrum of **2** suggested that the aromatic part of **2** was fully substituted. Furthermore, the <sup>1</sup>H NMR spectrum of **2** revealed the presence of a side chain, which could be ascribed to a 2-hydroxypentyl side chain, on the basis of the correlations observed in the HMBC and COSY spectra. The structure of **2** was thus elucidated as 1,6,8,10,11,13-hexahydroxy-2,5,9,12-tetrabromo-3,4-(2-hydroxypentyl)phenanthroperylene-7,14-quinone, which we have designated as gymnochrome F. Compound **2** was reported previously as the solvolysis product of its disulfated derivative isogymnochrome D.<sup>6</sup> As it was possible that compound **2** was an artifact from our isolation procedure in which H<sub>2</sub>O/TFA (0.05%) was used as the elution solvent during HPLC, the crude MeOH extract of the crinoid was analyzed by ESIMS. The detection of an ion with  $m/z$  963 (negative ionization mode) in the mass spectrum strongly supported the natural origin of **2**.

The configurations of the chiral centers in the side chains have been tentatively assigned using arguments similar to those described in detail by De Riccardis et al.<sup>6</sup> Briefly, it has been determined that the orientation of the side chain in perylenequinones is regulated by both the helicity of the ring system and the configurations of the chiral centers in the side chain. When the helicity of the ring and the configuration of the side chain chiral carbon are *M* and *R*, respectively, then the side chain is oriented above the aromatic ring system and is in the shielding zone of the aromatic ring system. When the helicity of the ring and the configuration of the side chain asymmetric carbon are *M* and *S*, respectively, then the side chain



**Figure 2.** CD spectra of **1** and **2**.

is directed away from the aromatic ring and is outside the shielding zone of the aromatic ring system. Therefore if one knows the helicity of the ring system, one can assign configurations to the chiral center of the side chain based upon the observation of shielding (or lack thereof) in the <sup>1</sup>H NMR data observed for the protons in the side chain.

Compounds **1** and **2** have CD spectra (Figure 2) that are virtually superimposable with that of isogymnochrome D, which has been reported previously to have *M* axial chirality.<sup>6</sup> For compound **1**, the propyl side chain methyl (H<sub>3</sub>-17) of compound **1** is not shielded ( $\delta$ <sub>H</sub> 0.94), which argues that it is directed away from the aromatic ring and that C-16 has an *S* configuration. The <sup>1</sup>H resonance of the methyl group of the pentyl moiety (H<sub>3</sub>-22) is shielded ( $\delta$ <sub>H</sub> -0.09) and therefore lies above the aromatic ring, which suggests an *R* configuration for the C-19 chiral center. In compound **2**, which has C<sub>2</sub> symmetry, the H<sub>3</sub>-19 methyl resonance is shielded, suggesting that it is oriented above the ring, and therefore C-16 has been assigned an *R* configuration. An additional datum to confirm this assignment is the close agreement in the <sup>1</sup>H NMR data observed for **2** with that reported for the desulfated isogymnochrome D produced during the course of the assignment of the structure of isogymnochrome D.<sup>6</sup> Gymnochrome F is therefore the desulfated form of isogymnochrome D.

Compound **5** was obtained as a yellow oil. The HR-ESIMS spectrum suggested a formula of C<sub>15</sub>H<sub>7</sub>BrO<sub>7</sub> for **5**. The UV-vis spectrum of **5** was similar to that observed for **4**, suggesting that they belong to the same family of natural products. The primary difference in the <sup>1</sup>H NMR spectra was the lack of the proton resonance observed at  $\delta$ <sub>H</sub> 6.51 (H-7) in **5**, which suggested that **5** is the 7-bromo derivative of **4**. Thus **5** was identified as 7-bromoemodic acid. The co-occurrences of **1**–**5** in the same organism strongly suggest a biogenetic relationship. Biosynthetic studies on the hypericin class of compounds indicate that the biosynthesis of the phenanthroperylenequinones and related natural products involves two consecutive reaction cascades, consisting of the biosynthesis of the two anthraquinones, followed by their condensation.<sup>10</sup>

Gymnochrome E (**1**) inhibited the proliferation of the NCI/ADR-Res (multi-drug-resistant ovarian cancer cell line) with an IC<sub>50</sub> value of 3.5  $\mu$ M and did not show significant inhibitory activity at a concentration of 6.4  $\mu$ M against the PANC-1 pancreatic carcinoma or DLD-1 human colorectal adenocarcinoma cell lines. Gymnochrome E also inhibited histone deacetylase-1 (HDAC-1) with an IC<sub>50</sub> of 10.9  $\mu$ M. Gymnochrome F (**2**) did not show significant inhibitory activity at a concentration of 5.2  $\mu$ M against the PANC-1, NCI/ADR-Res, or DLD-1 tumor cell lines but was a moderate inhibitor of MCL-1 binding to Bak with an IC<sub>50</sub> of 3.3  $\mu$ M. Gymnochrome E (**1**) exhibited minimum inhibition concentrations (MICs) of 25  $\mu$ g/mL against both *Staphylococcus aureus* and methicillin-resistant *S. aureus* (MRSA), while gymnochrome F (**2**) exhibited MICs of 12.5  $\mu$ g/mL against *S. aureus* and MRSA. Methicillin-resistant *S. aureus* has emerged as a serious threat due to its resistance to penicillin-class antibiotics. Compounds **1** and **2** did not show activity against either *Pseudomonas aeruginosa* or

**Table 1.** NMR Spectroscopic Data for Gymnochrome E (**1**) and Gymnochrome F (**2**)<sup>a</sup>

no.	<b>1</b>		HMBC <sup>c</sup>	<b>2</b>		
	$\delta_C^{b,e}$	$\delta_H$ (J in Hz) <sup>b</sup>		$\delta_C^{d,e}$	$\delta_H$ (J in Hz) <sup>d</sup>	HMBC
1, 6	158.9	158.9		159.6		
2, 5	115.0	115.4		116.4		
3, 4	145.6	143.7		144.0		
3a, 3b	122.5	122.0		121.5		
6a, 14a	109.5	109.4		109.2		
7, 14	183.0	183.0		185.0		
7a, 13a	102.5	102.5		103.0		
8, 13	168.9	168.9		163.7		
9, 12	106.3	106.2	6.59, s	102.0		
10, 11	175.7	175.7		168.3		
10a, 10b	120.3	120.3		120.7		
14b, 14c	127.5	127.6		125.1		
14d, 14g	121.7	121.6		121.5		
14e, 14f	124.7	124.9		125.6		
15	45.7	3.47, dd (13.0, 8.2) 3.64, dd (13.0, 2.0)	2, 3, 3a, 17 2, 3, 3a, 17	45.7	3.85, dd (13.7, 6.8) 3.98, dd (13.7, 5.5)	2, 3, 3a, 16, 17 2, 3, 3a, 16, 17
16	67.5	3.42, brs		70.8	3.54, brs	
17	24.8	0.94, d (6.1)	15, 16	37.0	0.43, brs	
18	45.4	3.56, dd (13.0, 8.9) 3.73, dd (13.0, 4.8)	3b, 4, 5, 19, 20 3b, 4, 5, 19, 20	17.5	0.51, brs 0.70, brs 0.85, brs	
19	69.5	3.29, brs		12.3	0.22, t (6.8)	17, 18
20	37.1	0.03, m 0.27, m				
21	17.3	0.58, m 0.81, m				
22	13.6	-0.09, t (6.8)	20, 21			

<sup>a</sup> <sup>1</sup>H and <sup>13</sup>C NMR data were measured at 600.17 and 150.9, respectively. <sup>b</sup> Recorded in DMSO-*d*<sub>6</sub> solution. <sup>c</sup> HMBC correlations are from the proton(s) stated to the indicated carbons. <sup>d</sup> Recorded in MeOH-*d*<sub>4</sub> solution. <sup>e</sup> Assignments made by interpretation of the HSQC and HMBC spectra as well as through comparison of the observed <sup>13</sup>C NMR spectrum with the <sup>13</sup>C spectrum calculated using ACDLab v 11 (see Supporting Information for calculated values).

*Candida albicans* at the concentrations tested. Emodic acid (**4**) and its 7-bromo derivative (**5**) showed no significant activity in any of the assays.

## Experimental Section

**General Experimental Procedures.** The UV spectra were collected on a Hitachi U-3010 spectrophotometer. The circular dichroism (CD) spectra were collected on a JASCO J-815 circular dichroism system. The IR spectra were collected on a Midac M-1200 with Galactic GRAMS/386 software. NMR data were collected on a JEOL ECA-600 spectrometer operating at 600.17 MHz for <sup>1</sup>H and 150.9 MHz for <sup>13</sup>C. The edited-g-HSQC spectrum was optimized for 140 Hz; the g-HMBC spectrum was optimized for 8 Hz; chemical shifts were referenced to solvent (CD<sub>3</sub>OD  $\delta_H$  observed at 3.31 ppm and  $\delta_C$  observed at 49.0 ppm; for DMSO-*d*<sub>6</sub>  $\delta_H$  observed at 2.46 and  $\delta_C$  observed at 40.0 ppm). The ESIMS were measured using a Finnigan LTQ mass spectrometer, and the HRESIMS were measured using a Kratos MS50TC mass spectrometer. HPLC was performed using a Hitachi LaChrom binary gradient HPLC system equipped with a diode array detector (L-7455) monitoring at 230 nm with a semipreparative C18 column (218TP1010, Grace Vydac). All solvents used were HPLC grade. Simulated <sup>13</sup>C NMR spectra were calculated for comparison to the observed spectra using ACD <sup>13</sup>C predictor version 11.

**Biological Material.** The specimen used in this study was identified as *Holopus rangii* d'Orbigni, 1837 (phylum Echinodermata, class Crinoidea, order Cyrtocrinida, family Holopodidae). Taxonomy was confirmed by Dr. David Pawson, Senior Research Scientist, National Museum of Natural History, Smithsonian Institution. A museum voucher specimen is cataloged in the Harbor Branch Oceanographic Museum (HBOI sample ID 8-V-00-1-002; catalog no. 070:00047). *H. rangii* is a stalked crinoid with an enlarged root that typically attaches to a hard substrate. Four arms attach to a thick pentagonal plate, and each arm divides in two, forming eight arms that are conical, tuberculated on the outer surface, and almost twice as long as the foot. The arms stretch out to feed, but retract into a tight ball when disturbed or to prevent predation. The live animal is dark green and remains green in alcohol. The museum specimen is approximately 4 cm in

length. The samples were collected with the *Johnson-Sea-Link* submersible (dive number JSL II-3203) from a depth of 358 m on a rocky slope on the south coast of Curacao (12°04.34' N, 68°53.62' W), May 8, 2000.

**Extraction and Isolation.** The frozen crinoid *Holopus rangii* (152 g) was freeze-dried and then extracted exhaustively by macerating with EtOH and EtOAc/EtOH (1:1 v/v) using a Waring blender (3 × 400 mL). The combined filtered extracts were concentrated by distillation under reduced pressure. The resulting dark brown oil (3.97 g) was separated under vacuum-column chromatographic conditions on a reversed-phase C18 stationary phase. A 150 mL Büchner funnel fitted with a medium-porosity fritted glass disk was used as the column. The stationary phase was packed to a total height of 4 cm. The crude extract was adsorbed onto a portion of the C18 stationary phase and applied as slurry to the column. Fractions were eluted using the following elution series: fraction 1: H<sub>2</sub>O (150 mL), fraction 2: ACN/H<sub>2</sub>O (8:2) (150 mL), fraction 3: IPA (2-propanol) (150 mL), fraction 4: ACN/H<sub>2</sub>O/TFA (8:2:0.1) (150 mL), and fraction 5: CH<sub>2</sub>Cl<sub>2</sub>/MeOH (1:1) (150 mL). Active components were isolated on the basis of bioassays and spectroscopic analysis. Final purification was achieved by semipreparative HPLC on fraction 2 using a Vydac C18 Protein and Peptide Column [(10 mm × 250 mm, 10 μm particle size), flow rate 2.5 mL/min; eluent: H<sub>2</sub>O containing 0.05% TFA/ACN (2:3 v/v), monitored by UV at 230 nm], yielding **1** (3 mg), **2** (5 mg), **3** (1.5 mg), **4** (2.1 mg), and **5** (0.5 mg).

**Gymnochrome E (1):** dark brown oil; UV (MeOH)  $\lambda_{max}$  (log  $\epsilon$ ) 293 (4.32), 311 (4.24), 372 (4.01), 524 (4.08), and 596 (4.32) nm; CD (MeOH, *c* 10 μg/mL) reported in Figure 2; IR (film) 3388, 2928, 1634, 1609, 1574, 1458, 1245, 1114, and 1032 cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) see Table 1; <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) see Table 1; ESIMS/MS (*m/z*) 718 and 645 [M - H]<sup>-</sup>; HRESIMS *m/z* 776.9966 [M + H]<sup>+</sup> (calcd for C<sub>36</sub>H<sub>27</sub>Br<sub>2</sub>O<sub>10</sub>, 776.9965).

**Gymnochrome F (2):** dark brown oil; UV (MeOH)  $\lambda_{max}$  (log  $\epsilon$ ) 300 (4.45), 333 (4.36), 498 (4.11), 554 (4.23), and 598 (4.46) nm; CD (MeOH, *c* 10 μg/mL) reported in Figure 2; IR (film) 3388, 2928, 1705, 1609, 1451, 1245, 1128, and 1004 cm<sup>-1</sup>; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) see Table 1; <sup>13</sup>C NMR (MeOH-*d*<sub>4</sub>) see Table 1; ESIMS/MS (*m/z*) 875 and

803 [M - H]<sup>-</sup>; HRESIMS *m/z* 960.8463 [M + H]<sup>+</sup> (calcd for C<sub>38</sub>H<sub>29</sub>Br<sub>4</sub>O<sub>10</sub>, 960.8489).

**7-Bromoemodic acid (5):** yellow oil; IR (film) 3402, 2921, 1684, 1629, 1210, and 1135 cm<sup>-1</sup>; UV (MeOH) λ<sub>max</sub> (log ε) 271 (4.08) and 437 (3.88) nm; <sup>1</sup>H NMR (MeOH-*d*<sub>4</sub>) δ 8.25 (H-1, s, 1H), 7.78 (H-3, s, 1H), and 7.27 (H-9, s, 1H); ESIMS/MS (*m/z*) 334, 297, 268, and 225; HRESIMS *m/z* 376.9273 [M - H]<sup>-</sup> (calcd for C<sub>15</sub>H<sub>6</sub>BrO<sub>7</sub>, 376.9296).

**Cytotoxicity Assays.** **1** and **2** were evaluated for their effects on proliferation of the PANC-1 human pancreatic carcinoma (ATCC No. CRL-1469), DLD-1 human colorectal adenocarcinoma (ATCC No. CCL-221), and NCI-ADR-RES (formerly MCF-7/ADR) human ovarian carcinoma cell lines. The PANC-1 and DLD-1 cell lines were obtained from the American Type Culture Collection (Rockville, MD). The NCI/ADR-RES cell line was obtained from the NCI-Frederick Cancer DCTD Tumor/Cell Line Repository (Bethesda, MD). Assays were run using protocols described previously.<sup>11</sup> All samples were assayed a minimum of three times to derive the final IC<sub>50</sub> value.

**HDAC Assays.** This assay was run as per the protocol described in Sambucetti et al.<sup>12</sup>

**MCL-1 Assays.** This assay was run as per the protocol described in Calcul et al.<sup>5</sup>

**Antimicrobial Assays.** The antimicrobial activities of **1–5** against *Staphylococcus aureus* (ATCC 29213), MRSA (methicillin-resistant *Staphylococcus aureus*, ATCC 700787), *Pseudomonas aeruginosa* (ATCC 27853), and *Candida albicans* (ATCC 44506) were determined as described in Park et al.<sup>13</sup>

**Acknowledgment.** The work described in this paper was funded by NIH grant 2U19-CA529955. We thank Prof. Dr. C. Hertweck and Dr. K. Ishida from the Leibniz Institute for Natural Product Research and Infection Biology (Germany) for the CD measurement. We thank G. Samples for sample preparation and D. Harmody and P. Linley for bioassays. We thank the governments of The Netherlands Antilles and Curacao for allowing us to operate in their territorial waters and collect the specimens used in this research. We thank A. Debrot and W. L. Bakhuis of the CARMABI Foundation (Caribbean Marine Biological Institute), Curacao, and L. Pors and K. van Dongen of the Curacao

Marine Park, who provided exceptional assistance in planning and collaboration during the expedition. This is Harbor Branch Oceanographic Institute at Florida Atlantic University Contribution No. 1799.

**Supporting Information Available:** LRMS, <sup>1</sup>H and <sup>13</sup>C NMR, and 2D NMR spectra for **1** and **2** as well as <sup>1</sup>H NMR spectra for **4** and **5**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References and Notes

- (1) Simmons, T. L.; Andrianasolo, E.; McPhail, K.; Flatt, P.; Gerwick, W. H. *Mol. Cancer Ther.* **2005**, *4*, 333–342.
- (2) Pettit, G. R.; Yoshiaki Kamano, Y.; Herald, C. L.; Tuinman, A. A.; Boettner, F. E.; Kizu, H.; Schmidt, J. M.; Baczynskyj, L.; Tomer, K. B.; Bontems, R. J. *J. Am. Chem. Soc.* **1987**, *109*, 6883–6885.
- (3) Gunasekera, S. P.; Gunasekera, M.; Longley, R. E.; Schulte, G. K. *J. Org. Chem.* **1990**, *55*, 4912–4945.
- (4) Bai, R.; Paull, K. D.; Herald, C. L.; Malspeis, L.; Pettit, G. R.; Hamel, E. *J. Biol. Chem.* **1991**, *266*, 15882–15889.
- (5) Calcul, L.; Chow, R.; Oliver, A. G.; Tenney, K.; White, K. N.; Wood, A. W.; Fiorilla, C.; Crews, P. *J. Nat. Prod.* **2009**, *72*, 443–449.
- (6) De Riccardis, F.; Iorizzi, M.; Minale, L.; Riccio, R.; De Forges, B. R.; Debitus, C. *J. Org. Chem.* **1991**, *56*, 6781–6787.
- (7) Alvi, K. A.; Nair, B.; Gallo, C.; Baker, D. *J. Antibiot.* **1997**, *50*, 264–266.
- (8) Skalkos, D.; Tatsis, E.; Gerathanassis, I. P.; Troganis, A. *Tetrahedron* **2002**, *58*, 4925–4626.
- (9) Pace, N.; Mackinney, G. *J. Am. Chem. Soc.* **1941**, *63*, 2570–2574.
- (10) Falk, H. *Angew. Chem., Int. Ed.* **1999**, *38*, 3116–3136.
- (11) Gunasekera, S. P.; Zuleta, I. A.; Longley, R. E.; Wright, A. E.; Pomponi, S. A. *J. Nat. Prod.* **2003**, *66*, 1615–1617.
- (12) (a) Sambucetti, L. C.; Fischer, D. D.; Zabudoff, S.; Kwon, P. O.; Chamberlin, H.; Trogani, N.; Xu, H.; Cohen, D. *J. Biol. Chem.* **1999**, *274*, 34940–34947. (b) Remiszewski, S. W.; Sambucetti, L. C.; Atadja, P.; Bair, K. W.; Cornell, W. D.; Green, M. A.; Howell, K. L.; Jung, M.; Kwon, P.; Trogani, N.; Walker, H. *J. Med. Chem.* **2002**, *45*, 753–757.
- (13) Park, Y. C.; Gunasekera, S. P.; Lopez, J. V.; McCarthy, P. J.; Wright, A. E. *J. Nat. Prod.* **2006**, *69*, 580–584.

NP900526Y