

Resorcylic Acid Lactones with Cytotoxic and NF- κ B Inhibitory Activities and Their Structure–Activity Relationships

Sloan Ayers,[†] Tyler N. Graf,[†] Audrey F. Adcock,[‡] David J. Kroll,[‡] Susan Matthew,[§] Esperanza J. Carcache de Blanco,^{§,⊥} Qi Shen,^{||} Steven M. Swanson,^{||} Mansukh C. Wani,[∇] Cedric J. Pearce,[#] and Nicholas H. Oberlies^{*,†}

[†]Department of Chemistry and Biochemistry, University of North Carolina at Greensboro, P.O. Box 26170, Greensboro, North Carolina 27402, United States

[‡]Department of Pharmaceutical Sciences, BRITE, North Carolina Central University, 1801 Fayetteville Street, Durham, North Carolina 27707, United States

[§]Division of Pharmacy Practice and Administration, and [⊥]Division of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, The Ohio State University, Columbus, Ohio 43210, United States


^{||}Department of Medicinal Chemistry and Pharmacognosy, University of Illinois at Chicago, Chicago, Illinois 60612, United States

[∇]Natural Products Laboratory, Research Triangle Institute, P.O. Box 12194, Research Triangle Park, North Carolina 27709, United States

[#]Mycosynthetix, Inc., 505 Meadowland Drive, Suite 103, Hillsborough, North Carolina 27278, United States

S Supporting Information

ABSTRACT: As part of our ongoing investigation of filamentous fungi for anti-cancer leads, an active fungal extract was identified from the Mycosynthetix library (MSX 63935; related to *Phoma* sp.). The initial extract exhibited cytotoxic activity against the H460 (human non-small cell lung carcinoma) and SF268 (human astrocytoma) cell lines and was selected for further study. Bioactivity-directed fractionation yielded resorcylic acid lactones (RALs) **1** (a new natural product) and **3** (a new compound) and the known RALs zeanol (**2**), (*SE*)-7-oxozeanol (**4**), (*SZ*)-7-oxozeanol (**5**), and LL-Z1640-1 (**6**). Reduction of (*SE*)-7-oxozeanol (**4**) with sodium borohydride produced **3**, which allowed assignment of the absolute configuration of **3**. Other known resorcylic acid lactones (**7**–**12**) were purchased and assayed in parallel for cytotoxicity with isolated **1**–**6** to investigate structure–activity relationships in the series. Moreover, the isolated compounds (**1**–**6**) were examined for activity in a suite of biological assays, including antibacterial, mitochondria transmembrane potential, and NF- κ B. In the latter assay, compounds **1** and **5** displayed sub-micromolar activities that were on par with the positive control, and as such, these compounds may serve as a lead scaffold for future medicinal chemistry studies.



In one component of our collaborative project to identify anticancer leads from diverse natural product study materials,^{1,2} extracts of filamentous fungi from the Mycosynthetix library, representing over 50 000 accessions, are being investigated systematically. Of the estimated 1.5 million species of fungi in the world, only about 75 000 have been described in the literature,³ and it is likely that an even smaller percentage have been pursued for bioactive secondary metabolites. Yet, many of the pharmaceutical breakthroughs of the 20th century originated in fungal cultures, including the well-known examples of antibiotics (i.e., penicillin),⁴ immunosuppressants (e.g., cyclosporin A),⁵ and cholesterol-lowering agents (e.g., compactin).⁶ The recently approved natural product analogue fingolimod,⁷ for the treatment of multiple sclerosis, and a clinical trial on psilocybin for terminally ill cancer patients⁸ are but two examples that demonstrate contemporary interest in fungal secondary


metabolites. Moreover, recent genomic data support the potential for fungi to biosynthesize a wealth of compounds.⁹ In short, and as is often reported in this journal,¹⁰ fungi from many different sources continue to be a valuable resource of bioactive secondary metabolites.¹¹

An extract of a filamentous fungus, MSX 63935, isolated from leaf litter collected in Nigeria, displayed promising cytotoxic activity against a human tumor panel.^{12,13} Previous research on this fungal isolate has examined a variety of biological targets, including assays for methicillin-resistant *Staphylococcus aureus* (MRSA), tuberculosis (TB), gene regulation, activity in the CNS, and various agricultural pests. However the only positive result was against MRSA, and no antibiotic compounds had been

Received: January 21, 2011

Published: April 22, 2011

isolated.¹⁴ Thus, these new cytotoxicity results spawned a bioactivity-directed fractionation study, and a series of resorcylic acid lactones (RALs; 1–6) were isolated and characterized, including a new natural product, 15-*O*-desmethyl-(*SZ*)-7-oxozeaenol (1), and a new compound, 7-*epi*-zeaenol (3). In conjunction with structurally related analogues purchased from commercial sources, these compounds were evaluated against the human tumor panel, yielding preliminary structure–activity relationship data. The isolated compounds (1–6) were examined also for antibacterial activity, as well as activity in mitochondrial transmembrane potential and NF- κ B inhibition assays. The NF- κ B inhibition assay revealed promising hits (compounds 1 and 5), both of which had sub-micromolar activities.



RESULTS AND DISCUSSION

Through the ongoing exploration of extracts of filamentous fungi for cytotoxicity,¹ fungus MSX 63935 showed promising activity (i.e., >96% inhibition of H460 cell growth at 20 μ g/mL). The solid-phase culture was scaled up to a 2.8 L Fernbach flask, and a 1:1 CHCl₃/MeOH extract was generated, which was defatted with a 3:3:4 CH₃CN/MeOH/hexanes partition. The resultant organic-soluble material was a white solid, which consisted almost exclusively of compounds 1–6 (see Figure S1 in the Supporting Information for an analytical HPLC of this solid). In fact, of all the anticancer leads examined from this library of filamentous fungi over the past three years, MSX 63935 was one of the most proficient producers of small molecules studied to date. This white solid was advanced directly to preparative scale RP-HPLC to isolate all six compounds in >95% purity, as measured by analytical HPLC. Compound 3 was a new compound, 1 was a new natural product, and 2, 4, 5, and 6 were known compounds. Of these, the production of 5 was quite high, in excess of 800 mg from a solid-phase culture grown in a 2.8 L Fernbach flask.

The molecular formula for the major compound (5) was C₁₉H₂₂O₇, as determined via HRESIMS. The NMR data, in conjunction with the HRMS data and UV maxima (from the PDA detector of the HPLC) at 234, 274, and 314 nm, indicated 5 to be the known compound (*SZ*)-7-oxozeaenol (also known as

LL-Z1640-2),^{15,16} first described in 1978 by Ellestad and co-workers.¹⁷ The $[\alpha]_D$ of 5 was -83 , compared to the literature value of -76 , indicating the absolute configuration of 5 to be the same as LL-Z1640-2.¹⁶ A reference standard was purchased, and the ¹H NMR data were identical to the isolated compound (Table S1). The absolute configuration of 5 had been established by synthesis.^{15,16}

The molecular formula for 1 was C₁₈H₂₀O₇, as determined via HRESIMS. The ¹H NMR spectra of 1 and 5 were very similar (see Tables 1 and S1). The major difference between the NMR data of 1 and 5 was the absence of the methoxy signals at δ_H/δ_C 3.80/55.5 in 1 that were present in 5. A new phenol peak was observed at δ_H 10.39 in 1, further supporting the absence of a methoxy moiety in 1 relative to 5. Compound 1 has been described previously as an intermediate in the synthesis of 5;¹⁶ however, it was ascribed the trivial name 15-*O*-desmethyl-(*SZ*)-7-oxozeaenol, since it had not been described from nature previously. The absolute configuration of 1 had been established previously by synthesis.¹⁶

The HRESIMS data of compound 2 corresponded to a formula of C₁₉H₂₄O₇, indicative of the addition of two protons relative to 5. The ¹H NMR spectrum of 2 suggested the identity of 2 as zeaenol, and comparison of the NMR data to literature confirmed this assignment.¹⁸ Moreover, the $[\alpha]_D$ of 2 was -93 , which matched literature data exactly.¹⁸ The absolute configuration of 2 had been established previously by X-ray diffraction.¹⁸

Compound 3 was isolated as a white powder, and the HRESIMS data also suggested a formula of C₁₉H₂₄O₇ as with 2. The ¹H NMR data of 3 were similar to 2; however, there were numerous signals that had shifted slightly compared to 2 (see Figures S4 and S5 for ¹H and ¹³C NMR spectra, respectively, of 3). The COSY spectrum of 3 indicated the same carbon skeleton and oxygenation pattern as 2, suggesting that 2 and 3 were diastereomers.

A recent report described a C-7 epimer (C-6' in their numbering scheme) of aigalomycin B, which differs from 2 by the presence of an epoxide at C-11/C-12 (C-1'/C-2' in their numbering scheme) instead of an alkene.¹⁹ The same report described both C-7 epimers of 5,6-dihydrozeaenol. Hence, given these literature precedents and the 2D-NMR data indicating that 3 and 2 shared an identical skeleton and oxygenation pattern, it was hypothesized that 3 was the C-7 epimer of 2. The reduction of the C-7 ketone of (*SE*)-7-oxozeaenol (4; described below) was performed using sodium borohydride. The major product of the reduction of 4 had the identical retention time and UV spectrum to those of 3. Moreover, after isolation of the major reduction product by preparative HPLC, the ¹H NMR data were identical to those of isolated 3, and thus, the configuration at C-7 for 3 was established as *R*. That 3 was the major product of the sodium borohydride reduction of 4 was somewhat surprising, as sodium borohydride is known to give mixtures of 1, 4- and 1,2-reductions of α,β -unsaturated ketones; only a trace amount of 2 was detected by HPLC in the sodium borohydride reduction of 4. Regardless, these data supported the assignment of 3 as 7-*epi*-zeaenol.

Compound 4 was isolated as a white powder. HRESIMS data indicated a formula identical to 5 of C₁₉H₂₂O₇. NMR data established 4 as the known compound (*SE*)-7-oxozeaenol.¹⁸ To the best of our knowledge, the NMR data for 4 have not been reported previously, and thus they are summarized in Table S1. The $[\alpha]_D$ value does not appear to have been reported either (see Experimental Section). Moreover, since the acetone of 4 has been reported, it was synthesized, and the ¹H NMR data were in excellent agreement with literature data.¹⁸

Table 1. NMR Data for Compounds **1** and **3** (700 MHz for ^1H , 176 MHz for ^{13}C ; chemical shifts in δ , coupling constants in Hz, DMSO- d_6)

position	15-O-desmethyl-(5Z)-7-oxozeaenol (1)		7- <i>epi</i> -zeaenol (3)	
	δ_{C}	δ_{H} mult (J in Hz)	δ_{C}	δ_{H} mult (J in Hz)
1	170.6		169.0	
3	73.6	5.17, qdd (5.7, 12.4, 2.1)	71.8	5.09, qdd (6.1, 12.4, 2.1)
4a	36.2	2.56, m	38.6	2.35, ddd (15.1, 9.6, 8.9)
4b	36.2	3.19, m	38.6	2.43, m
5	142.8	6.17, ddd (11.5, 11.5, 2.8)	127.1	5.54, ddd (15.8, 8.2, 4.1)
6	126.6	6.43, dd (11.5, 2.1)	133.1	5.60, dd (15.8, 6.9)
7	200.8		74.7	4.04, m
8	81.6	4.39, dd (4.4, 2.1)	77.6	3.44, m
9	72.6	3.84, m	72.8	3.57, m
10a	36.0	1.98, m	36.8	2.14, m
10b	36.0	2.02, m	36.8	2.52, m
11	131.6	5.95, ddd (15.3, 10.3, 4.8)	132.5	6.11, ddd (15.8, 7.6, 4.8)
12	131.3	6.76, d (15.3)	128.5	6.59, d (15.8)
13	143.8		139.5	
14	107.4	6.29, d (2.8)	102.5	6.43, d (2.3)
15	162.3		161.6	
16	101.5	6.19, d (2.8)	99.9	6.31, d (2.3)
17	164.1		158.8	
18	102.5		110.5	
19	20.0	1.36, d (5.7)	20.5	1.30, d (6.1)
20			55.3	3.74, s
7-OH				4.83, br s
8-OH		4.98, d (4.4)		4.57, br d (3.9)
9-OH		5.08, d (5.7)		4.89, br s
15-OH		10.39, s		
17-OH		11.82, s		10.53, s

Compound **6** was isolated as a white powder. The HRESIMS data indicated a formula identical to **2** and **3** of $\text{C}_{19}\text{H}_{24}\text{O}_7$. The NMR data for **6** matched that of LL-Z1640-1.¹⁸ The $[\alpha]_{\text{D}}$ of **6** was measured as -78 , which compared favorably with the literature value of -81 .¹⁸ There appears to be some confusion in the literature about the absolute configuration of **6**, which was determined initially by X-ray diffraction.¹⁷ Sugawara et al.¹⁸ later reported position C-8 as having an *R* configuration, whereas the X-ray studies¹⁷ indicated C-8 as *S*; the latter assignment seems to be consistent on the basis of biosynthetic comparisons to the other isolated compounds.

Compounds **1–6**, along with other known RALs hypothemycin (**7**), zearalenone (**8**), α -zearalenol (**9**), β -zearalenol (**10**), α -zearalanol (**11**), and β -zearalanol (**12**) were assayed against the three cancer cell lines MCF-7, H460, and SF268. The IC_{50} results for **1–12** are summarized in Table 2. From these, it was clear that the enone (i.e., compounds **1**, **4**, **5**, and **7**), but not exclusively a *cis*-enone (i.e., note **4**), was required for cytotoxic activity. A recent review highlights some of the biology of the resorcylic acid lactones,²⁰ and they are known to inhibit with nanomolar potency the ATPase activity of two critical classes of proteins: the HSP90 chaperone^{21,22} and a subset of 16 kinases possessing a crucial cysteine residue in their ATPase binding pocket.²³ HSP90 client proteins include several oncogenic proteins, such as Src kinase, and HSP90 inhibition results in the dissociation and degradation of these proteins. The activity of

the new members of this class of polyketides, described herein, toward these targets remains to be investigated.

Compounds **1–6** were examined for activity in a series of three other assays. With respect to NF- κ B inhibitory activity, compound **5** (Table 3) was the most potent, being approximately a half-order of magnitude less potent than the positive control (rocgamide). Previously, **5** was reported as a potent inhibitor of TAK1, which is a member of the mitogen-activated kinase kinase kinase (MEKK) family that participates in proinflammatory cellular signaling pathways by activating NF- κ B.²⁴ As with the cytotoxicity data, the enone moiety seemed to enhance potency, and in this case, the *cis* relationship (i.e., compounds **1** and **5**) was more potent than the *trans* relationship (i.e., compound **4**). In an assay for mitochondrial transmembrane potential (MTP), all compounds displayed IC_{50} values greater than $10 \mu\text{M}$. Since the positive control for this assay, staurosporine, had an IC_{50} value in the single digit nanomolar range, **1–6** were considered inactive. Finally, compounds **1–6** were examined and found inactive against *Escherichia coli*, *Staphylococcus aureus*, *Mycobacterium smegmatis*, and *Bacillus subtilis*; MIC values were all $>500 \mu\text{g}/\text{mL}$.

In summary, a series of six structurally related resorcylic acid lactones (**1–6**) were isolated from MSX 63935. Data in both the cytotoxicity assays and the NF- κ B assay suggest that compound **5** was the most promising. Fortunately, this fungus was a prolific producer of **5**, generating over 800 mg per solid-phase culture in a 2.8 L Fernbach flask, using standard techniques prior to any

Table 2. Cytotoxicity of Compounds Isolated from MSX 63935 (1–6) and Purchased Standards with Similar Structural Features (7–12) against a Panel of Human Tumor Cell Lines

compound	IC ₅₀ values (in μM) ^a				
	MCF-7	H460	SF268	HT-29	MDA-MB-435
15-O-desmethyl-(5Z)-7-oxozeaenol (1)	10.3	4.1	8.9	11.5	15.8
zeaenol (2)	>100	>100	>100	>25	>25
7- <i>epi</i> -zeaenol (3)	>100	>100	>100	>25	>25
5 <i>E</i> -7-oxozeaenol (4)	4.9	1.2	5.6	4.4	5.5
5 <i>Z</i> -7-oxozeaenol (5)	3.8	1.2	5.1	3.6	3.3
LL-Z1640-1 (6)	>50	>25	>50	>50	>50
hypothemycin (7)	16.3	4.3	12.7	nt ^c	nt
zearalenone (8)	>25	>25	>50	nt	nt
α -zearalenol (9)	>25	>25	>50	nt	nt
β -zearalenol (10)	>25	15.4	>50	nt	nt
α -zearalanol (11)	>50	>50	>50	nt	nt
β -zearalanol (12)	>25	9.5	>25	nt	nt
camptothecin ^b	0.07	<0.01	0.04	nt	nt
silvestrol ^b	nt	nt	nt	0.004	0.006

^aIC₅₀ values were determined as the concentration required to reduce cellular proliferation by 50% relative to untreated controls following 72 h of continuous exposure. ^{12,13} ^b Positive controls. ^c Indicates “not tested”.

Table 3. NF- κ B Inhibitory Activity of Compounds Isolated from MSX 63935 (1–6)

compound	IC ₅₀ (μM)
15-O-desmethyl-(5Z)-7-oxozeaenol (1)	0.75
zeaenol (2)	>50
7- <i>epi</i> -zeaenol (3)	>50
7-oxozeaenol (4)	11.5
(5Z)-7-oxozeaenol (5)	0.25
LL-Z1640-1 (6)	>50
rocaglamide (positive control)	0.075

growth optimization studies. As such, studies are ongoing to probe the structure–activity relationships further by using **5** as starting materials for medicinal chemistry studies and by modifying the fermentation procedures to generate various analogues.

EXPERIMENTAL SECTION

General Experimental Procedures. Optical rotations, UV spectra, and IR spectra were obtained on a Rudolph Research Autopol III polarimeter, a Varian Cary 100 Bio UV–vis spectrophotometer, and a Perkin-Elmer Spectrum One with Universal ATR attachment, respectively. NMR experiments were conducted in either DMSO-*d*₆ or CDCl₃ with TMS as reference. NMR instrumentation was a Bruker Ultrashield Plus with Avance III console, Topspin software version 2.1, and a QNP style Cryoprobe (operating at 700.13 MHz for ¹H, 176.05 MHz for ¹³C). A JEOL ECA-500 (operating at 500 MHz for ¹H, 125 MHz for ¹³C) was also used for some experiments. HRESIMS was performed on a Waters SYNAPT MS system. HPLC was carried out on Varian Prostar HPLC systems equipped with Prostar 210 pumps and a Prostar 335 photodiode array detector (PDA), with data collected and analyzed using Galaxie Chromatography Workstation software (version 1.9.3.2). For preparative HPLC, a Gemini-NX C18 (5 μm ; 250 \times 21.2 mm) column was used at a 21.2 mL/min flow rate, while for analytical HPLC, a Gemini-NX C18 (5 μm ; 250 \times 4.6 mm) column was used with a 1 mL/min flow rate

(both from Phenomenex, Inc.). Reference standards of (5Z)-7-oxozeaenol (**5**) and zeaenol (**2**) were obtained from EMD Chemicals. Zearalenone (**8**), α -zearalenol (**9**), β -zearalenol (**10**), α -zearalanol (**11**), and β -zearalanol (**12**) were obtained from Sigma-Aldrich. Hypothemycin (**7**) was obtained from Enzo Life Sciences. All other reagents and solvents were obtained from Fisher Scientific and were used without further purification.

Producing Organism and Fermentation. Mycosynthetix fungal strain 63935 was isolated in 1992 by Dr. Barry Katz of MYCOsearch from leaf litter collected at the Agricultural Farms, University of Nigeria, Ede. DNA analyses were performed by MIDI Laboratories, Inc. (Newark, DE), and the D2 variable region of the large subunit (LSU) rRNA was sequenced and compared to their database; the closest match could determine only that this fungus was related to *Phoma* sp.; these data were deposited in Genbank (accession no. JF767207). The culture was stored on a malt extract slant and was transferred periodically. A fresh culture was grown on a similar slant, and a piece was transferred to a medium containing 2% soy peptone, 2% dextrose, and 1% yeast extract (YESD media). Following incubation (7 d) at 22 °C with agitation, the culture was used to inoculate 50 mL of a rice medium, prepared using rice to which was added a vitamin solution and twice the volume of rice with H₂O, in a 250 mL Erlenmeyer flask. This was incubated at 22 °C until the culture showed good growth (approximately 14 d). The scale-up culture was grown in a 2.8 L Fernbach flask containing 150 g of rice and 300 mL of H₂O and was inoculated using a seed culture grown in YESD medium. This was incubated at 22 °C for 14 d.

Extraction and Isolation. To the large-scale solid fermentation was added 500 mL of 1:1 MeOH/CHCl₃. The mixture was shaken for 16 h, then filtered, and the solvent was evaporated (3.7 g of waxy, white solid). The extract was then partitioned between CH₃CN/MeOH/hexanes (150/150/200 mL), the CH₃CN/MeOH solubles were separated, and the solvent was evaporated (1.7 g of white powder, Figure S1, Supporting Information). As such, it was dissolved in 5.5 mL of DMSO and purified via seven separate injections by preparative HPLC using a gradient that initiated with 35:65 and increased linearly to 45:55 CH₃CN/H₂O over 40 min. Compound **1** eluted at \sim 7.6 min (5.2 mg), **2** at \sim 9 min (90.7 mg), **3** at \sim 11.8 min (5.6 mg), and **4** at \sim 14 min (73.1 mg), and **5** and **6** eluted together from 16 to 20 min.

The mixture of **5** and **6** was subjected to two further rounds of shave-and-recycle preparative HPLC under the same conditions to yield >95% pure **5** (885.9 mg) and **6** (80.5 mg).

15-O-Desmethyl-(5Z)-7-oxozeaenol (1): white solid; $[\alpha]_D^{23}$ -59 (c 0.07, MeOH); UV (MeOH) λ_{\max} (log ϵ) 232 (4.35), 271 (3.89), 313 (3.62) nm; IR (diamond) ν_{\max} 3348, 3146, 2951, 1690, 1646, 1602, 1467, 1350, 1312, 1259, 1216, 1167, 1021, 965 cm^{-1} ; ^1H NMR (DMSO- d_6 , 700 MHz) and ^{13}C NMR (DMSO- d_6 , 175 MHz), see Table 1; HRESIMS m/z 371.1100 $[\text{M} + \text{Na}]^+$ and 347.1122 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{18}\text{H}_{20}\text{O}_7\text{Na}$, 371.1107; $\text{C}_{18}\text{H}_{19}\text{O}_7$, 347.1131).

Zeaenol (2): white solid; $[\alpha]_D^{23}$ -91.9 (c 1.00, MeOH); ^1H and ^{13}C NMR data were in good agreement with the literature;¹⁸ HRESIMS m/z 387.1413 $[\text{M} + \text{Na}]^+$ and 363.1442 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{19}\text{H}_{24}\text{O}_7\text{Na}$, 387.1420; $\text{C}_{19}\text{H}_{23}\text{O}_7$, 363.1443).

7-epi-Zeaenol (3): white solid; $[\alpha]_D^{23}$ -90.4 (c 0.25, MeOH); UV (MeOH) λ_{\max} (log ϵ) 235 (4.30), 270 (3.94), 312 (3.64) nm; IR (diamond) ν_{\max} 3376, 2965, 2930, 2899, 1636, 1602, 1571, 1353, 1312, 1248, 1212, 1157, 1041, 971 cm^{-1} ; ^1H NMR (DMSO- d_6 , 700 MHz) and ^{13}C NMR (DMSO- d_6 , 175 MHz), see Table 1; HRESIMS m/z 387.1410 $[\text{M} + \text{Na}]^+$ and 363.1446 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{19}\text{H}_{24}\text{O}_7\text{Na}$, 387.1420; $\text{C}_{19}\text{H}_{23}\text{O}_7$, 363.1443).

5E-7-Oxozeaenol (4): white solid; $[\alpha]_D^{23}$ -27.0 (c 0.33, MeOH); ^1H and ^{13}C NMR (DMSO- d_6 , 500 MHz) data for **4**, see Table 1; HRESIMS m/z 385.1257 $[\text{M} + \text{Na}]^+$ and 361.1281 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{19}\text{H}_{22}\text{O}_7\text{Na}$, 385.1263; $\text{C}_{19}\text{H}_{21}\text{O}_7$, 361.1287, respectively).

5Z-7-Oxozeaenol (5): white solid; $[\alpha]_D^{23}$ -83.3 (c 0.34, MeOH); ^1H and ^{13}C NMR data were in good agreement with the literature;^{15,16} HRESIMS m/z 385.1252 $[\text{M} + \text{Na}]^+$ and 361.1278 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{19}\text{H}_{22}\text{O}_7\text{Na}$, 385.1263; $\text{C}_{19}\text{H}_{21}\text{O}_7$, 361.1287).

LL-Z1640-1 (6): white solid; $[\alpha]_D^{23}$ -78.3 (c 0.33, MeOH); ^1H and ^{13}C NMR data were in good agreement with the literature;¹⁸ HRESIMS m/z 387.1410 $[\text{M} + \text{Na}]^+$ and 363.1424 $[\text{M} - \text{H}]^-$ (calcd for $\text{C}_{19}\text{H}_{24}\text{O}_7\text{Na}$, 387.1420; $\text{C}_{19}\text{H}_{23}\text{O}_7$, 363.1443).

Acetonide of (5E)-7-Oxozeaenol (4a). To 1.54 mg of **4** were added 100 μL of 2,2-dimethoxypropane and 100 μL of acetone. A trace amount of pyridinium *p*-toluenesulfonate was added, and the solution was stirred for 2 h at 40 $^\circ\text{C}$. H_2O (200 μL) was added to the solution, which was then injected directly onto semipreparative HPLC (ODS-A, 250 \times 10 mm i.d., 5 μm ; YMC). The elution proceeded at 4 mL/min via a gradient that initiated with 50:50 and increased linearly over 20 min to 100:0 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$. The acetonide eluted from 14.7 to 15.3 min and was collected to yield 1.47 mg of **4a** (86% yield). The ^1H NMR data of **4a** were in excellent agreement with literature.¹⁸

Reduction of 4. (5E)-7-Oxozeaenol (**4**; 4.6 mg) was weighed into a 2 mL reaction vial, and 0.25 mL of both 2-propanol and THF were added. The compound dissolved after brief stirring. NaBH_4 (10.6 mg) was added with continued stirring. After 30 min, the solution had turned cloudy, and by analytical HPLC the presence of **4** was not discernible. H_2O (0.5 mL) was added to this reaction product to dissolve all solids, and this solution was purified in a single step via preparative HPLC using the aforementioned conditions. The major peak was collected between 10 and 13 min (2.53 mg; 57% yield) and was confirmed by ^1H and ^{13}C NMR to be identical to isolated **3**.

Cytotoxicity Assay. The cytotoxicity measurements against the MCF-7²⁵ human breast carcinoma (Barbara A. Karmanos Cancer Center), NCI-H460²⁶ human large cell lung carcinoma (HTB-177, American Type Culture Collection (ATCC)), and SF-268²⁷ human astrocytoma (NCI Developmental Therapeutics Program) cell lines were performed as described previously.¹³ Moreover, a second cytotoxicity assay was performed on only the isolated compounds using the HT-29²⁸ human colorectal adenocarcinoma (HTB-38, ATCC) and the MDA-MB-435²⁹ human melanoma (HTB-129, ATCC) cell lines. All cells were propagated in RPMI 1640 medium supplemented with fetal bovine serum (10%), penicillin (100 units/mL), and streptomycin (100 $\mu\text{g}/\text{mL}$). Cells were plated at either 2500

(HT-29) or 5000 (MCF-7, MDA-MB-435) cells per well in 96-well microtiter plates and incubated for 16 h in a humidified, 5% CO_2 atmosphere at 37 $^\circ\text{C}$. Test substance was then added at the following final concentrations: 25, 5, 1, 0.2, and 0.04 $\mu\text{g}/\text{mL}$. After a 72 h incubation, an MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium, inner salt) assay was performed using a commercially available kit according to the manufacturer's instructions (CellTiter 96 AQueous One Solution Cell Proliferation Assay, Promega Corp.).

Antimicrobial Assay. Antibiotic activity was examined using an agar plate diffusion assay. Overnight cultures of *Escherichia coli*, *Staphylococcus aureus*, *Mycobacterium smegmatis*, and *Bacillus subtilis* were used to inoculate molten LB media or Middlebrook 7H9 media (Difco) with 1% glycerol, containing 1.5% agar and kept at 50 $^\circ\text{C}$; these were then used to prepare assay plates. Samples (dissolved in 10 μL of MeOH) were applied to the surface of the assay dish, and positive controls were treated in a similar manner (penicillin G, novobiocin, and streptomycin; all from Sigma). The bioassay plates were incubated overnight at 37 $^\circ\text{C}$. Biological activity of the standards could be detected to 1 $\mu\text{g}/\text{mL}$ (except that penicillin G was active against *E. coli* at 100 $\mu\text{g}/\text{mL}$ only), whereas the test compounds showed no activity at 500 $\mu\text{g}/\text{mL}$ and were thus deemed inactive.

NF- κB p65 Assay. An ELISA-based NF- κB inhibitory assay was performed as described previously.³⁰ Briefly, HeLa cells (ATCC CCL-2) were treated with various concentrations of test compounds, positive control, or solvent control, and their nuclei extracted using the NE-PER kit (Pierce Biotechnology). The specific binding ability of activated p65 subunits of NF- κB in the nucleus was detected according to the manufacturer's protocol (EZ-Detect Transcription Factor Assay System ELISA kit, Pierce). Rocaglamide (Enzo Life Sciences International, Inc.) was used as a positive control (IC_{50} value of 0.075 μM).

Mitochondria Transmembrane Potential ($\Delta\Psi$) Assay. The mitochondrial transmembrane potential assay kit (Cayman Chemical Company) was adapted to detect the $\Delta\Psi$ using a procedure published previously.³¹ $\Delta\Psi$ is used to represent mitochondrial membrane transition events. Briefly, HT-29 cells were treated with various concentrations of the test compounds or positive control. Then, cells were incubated with the lipophilic cationic dye 5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolylcarbocyanide (JC-1; Cayman Chemical Company). After incubation, cells were rinsed with a wash buffer to remove unbound staining reagent. Samples were analyzed by a FLUOstar Optima fluorescence plate reader (BMG Labtech, Inc.) using an excitation wavelength of 485 nm and emission wavelength of 530 nm for JC-1 monomers from apoptotic cells and an excitation wavelength of 560 nm and emission wavelength of 595 nm for JC-1 aggregates from healthy cells. Measurements were performed in triplicate and are representative of at least two independent experiments. For this assay, staurosporine (Cayman Chemical Company, Ann Arbor, MI) was used as a positive control (IC_{50} value of 2.5 nM).

■ ASSOCIATED CONTENT

Supporting Information. ^1H NMR spectra for compounds **1–6**, ^{13}C NMR spectra for compounds **1** and **3**, and an HPLC chromatogram for the crude extract of MSX 63935. This information is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: Nicholas_Oberlies@uncg.edu. Tel: 336-334-5474.

■ ACKNOWLEDGMENT

This research was supported by P01 CA125066 from the National Cancer Institute/National Institutes of Health,

Bethesda, MD, USA. The Golden LEAF Foundation (Rocky Mount, NC) provided partial support to D.J.K. Mycology technical support was provided by B. Darveaux and M. Lawrence. The authors thank Dr. K. Knagge and M. Su, both of the David H. Murdock Research Institute, Kannapolis, NC, for some NMR and mass spectrometry data and Dr. J. Fuchs (Ohio State University) for helpful suggestions.

REFERENCES

- (1) Kinghorn, A. D.; Carache de Blanco, E. J.; Chai, H. B.; Orjala, J.; Farnsworth, N. R.; Soejarto, D. D.; Oberlies, N. H.; Wani, M. C.; Kroll, D. J.; Pearce, C. J.; Swanson, S. M.; Kramer, R. A.; Rose, W. C.; Fairchild, C. R.; Vite, G. D.; Emanuel, S.; Jarjoura, D.; Cope, F. O. *Pure Appl. Chem.* **2009**, *81*, 1051–1063.
- (2) Orjala, J.; Oberlies, N. H.; Pearce, C. J.; Swanson, S. M.; Kinghorn, A. D. In *Bioactive Compounds from Natural Sources*, 2nd ed.; Tringali, C., Ed.; Taylor & Francis: London, accepted.
- (3) Hawksworth, D. L.; Rossman, A. Y. *Phytopathology* **1997**, *87*, 888–891.
- (4) Fleming, A. *Penicillin, Its Practical Application*; Butterworth & Co. Ltd.: London, 1946; p 380.
- (5) Dreyfus, M.; Harri, E.; Hofmann, H.; Pache, W.; Tschertter, H. *Eur. J. Appl. Microbiol.* **1976**, *3*, 125–133.
- (6) Brown, A. G.; Smale, T. C.; King, T. J.; Hasenkamp, R.; Thompson, R. H. *J. Chem. Soc., Perkin Trans. 1* **1976**, *11*, 1165–1170.
- (7) Brinkmann, V.; Billich, A.; Baumruker, T.; Heining, P.; Schmouder, R.; Francis, G.; Aradhye, S.; Burtin, P. *Nat. Rev. Drug Discovery* **2010**, *9*, 883–897.
- (8) Griffiths, R.; Richards, W.; Johnson, M.; McCann, U.; Jesse, R. *J. Psychopharmacol.* **2008**, *22*, 621–632.
- (9) Misiek, M.; Hoffmeister, D. *Planta Med.* **2007**, *73*, 103–115.
- (10) The use of the term “fungus” in titles of articles published in the *Journal of Natural Products* has increased steadily over the last 10 years, from appearing 12 times in 2000 to appearing 31 times in 2009; at the time of manuscript submission, the term was on target to increase to nearly 40 times in 2010.
- (11) Pearce, C.; Eckard, P.; Gruen-Wollny, I.; Hanske, F. G. In *Natural Product Chemistry for Drug Discovery*; Buss, A. D.; Butler, M. S., Eds.; The Royal Society of Chemistry: Cambridge, 2010; pp 215–244.
- (12) Alali, F. Q.; El-Elimat, T.; Li, C.; Qandil, A.; Alkofahi, A.; Tawaha, K.; Burgess, J. P.; Nakanishi, Y.; Kroll, D. J.; Navarro, H. A.; Falkinham, J. O., III; Wani, M. C.; Oberlies, N. H. *J. Nat. Prod.* **2005**, *68*, 173–178.
- (13) Li, C.; Lee, D.; Graf, T. N.; Phifer, S. S.; Nakanishi, Y.; Riswan, S.; Setyowati, F. M.; Saribi, A. M.; Soejarto, D. D.; Farnsworth, N. R.; Falkinham, J. O., III; Kroll, D. J.; Kinghorn, A. D.; Wani, M. C.; Oberlies, N. H. *J. Nat. Prod.* **2009**, *72*, 1949–1953.
- (14) These previous studies were retrieved from the Mycosynthetix database, which covers previous research on these organisms going back over 30 years.
- (15) Tatsuta, K.; Takano, S.; Sato, T.; Nakano, S. *Chem. Lett.* **2001**, 172–173.
- (16) Dakas, P. Y.; Jogireddy, R.; Valot, G.; Barluenga, S.; Winssinger, N. *Chemistry* **2009**, *15*, 11490–11497.
- (17) Ellestad, G. A.; Lovell, F. M.; Perkinson, N. A.; Hargreaves, R. T.; McGahren, W. J. *J. Org. Chem.* **1978**, *43*, 2339–2343.
- (18) Sugawara, F.; Kim, K. W.; Kobayashi, K.; Uzawa, J.; Yoshida, S.; Murofushi, N.; Takahashi, N.; Strobel, G. A. *Phytochemistry* **1992**, *31*, 1987–1990.
- (19) Xu, L. X.; He, Z. X.; Xue, J. H.; Chen, X. P.; Wei, X. Y. *J. Nat. Prod.* **2010**, *73*, 885–889.
- (20) Winssinger, N.; Barluenga, S. *Chem. Commun.* **2007**, 22–36.
- (21) Schulte, T. W.; Akinaga, S.; Soga, S.; Sullivan, W.; Stensgard, B.; Toft, D.; Neckers, L. M. *Cell Stress Chaperones* **1998**, *3*, 100–108.
- (22) Sharma, S. V.; Agatsuma, T.; Nakano, H. *Oncogene* **1998**, *16*, 2639–2645.
- (23) Schirmer, A.; Kennedy, J.; Murli, S.; Reid, R.; Santi, D. V. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 4234–4239.
- (24) Ninomiya-Tsuji, J.; Kajino, T.; Ono, K.; Ohtomo, T.; Matsumoto, M.; Shiina, M.; Mihara, M.; Tsuchiya, M.; Matsumoto, K. *J. Biol. Chem.* **2003**, *278*, 18485–18490.
- (25) Soule, H. D.; Vazquez, J.; Long, A.; Albert, S.; Brennan, M. *J. Natl. Cancer Inst.* **1973**, *51*, 1409–1416.
- (26) Carney, D. N.; Gazdar, A. F.; Bunn, P. A., Jr.; Guccion, J. G. *Stem Cells* **1982**, *1*, 149–164.
- (27) Rosenblum, M. L.; Gerosa, M. A.; Wilson, C. B.; Barger, G. R.; Pertuiset, B. F.; de Tribolet, N.; Dougherty, D. V. *J. Neurosurg.* **1983**, *58*, 170–176.
- (28) Fogh, J.; Trempe, G. In *Human Tumor Cells In Vitro*; Plenum: New York, 1975; pp 115–159.
- (29) Rae, J. M.; Creighton, C. J.; Meck, J. M.; Haddad, B. R.; Johnson, M. D. *Breast Cancer Res. Treat.* **2007**, *104*, 13–19.
- (30) Salim, A. A.; Pawlus, A. D.; Chai, H. B.; Farnsworth, N. R.; Douglas Kinghorn, A.; Carcache-Blanco, E. J. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 109–112.
- (31) Deng, Y.; Balunas, M. J.; Kim, J. A.; Lantvit, D. D.; Chin, Y. W.; Chai, H.; Sugiarsa, S.; Kardono, L. B.; Fong, H. H.; Pezzuto, J. M.; Swanson, S. M.; de Blanco, E. J.; Kinghorn, A. D. *J. Nat. Prod.* **2009**, *72*, 1165–1169.