

Cannabinomimetic Lipid from a Marine Cyanobacterium

Marcelino Gutiérrez,^{*,†,‡} Alban R. Pereira,[†] Hosana M. Deboni,^{†,§} Alessia Ligresti,[⊥] Vincenzo Di Marzo,[⊥] and William H. Gerwick^{*,†}

[†]Center for Marine Biotechnology and Biomedicine, Scripps Institution of Oceanography, and Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California, San Diego, La Jolla, California 92093, United States


[‡]Instituto de Investigaciones Científicas y Servicios de Alta Tecnología, Ciudad del Saber, Clayton 0843-01103, Panamá

[§]Departamento de Física e Química, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, São Paulo 14040903, Brazil

[⊥]Institute of Biomolecular Chemistry, National Research Council, 80078 Pozzuoli, Italy

S Supporting Information

ABSTRACT: NMR-guided fractionation of two independent collections of the marine cyanobacteria *Lyngbya majuscula* obtained from Papua New Guinea and *Oscillatoria* sp. collected in Panama led to the isolation of the new lipids serinolamide A (3) and propenediester (4). Their structures were determined by NMR and MS data analysis. Serinolamide A (3) exhibited a moderate agonist effect and selectivity for the CB₁ cannabinoid receptor ($K_i = 1.3 \mu\text{M}$, >5-fold) and represents the newest addition to the known cannabinomimetic natural products of marine origin.



An important development in neuropharmacological research has been the discovery of specific cannabinoid receptors (CB₁ and CB₂) in central and peripheral mammalian tissues.^{1–3} Both receptor subtypes have amino acid sequences characteristic of G-protein-coupled receptors (GPCR).^{2,4} CB₁ receptors exhibit a widespread distribution in the brain and are responsible for the psychological and anticonvulsive effects produced by marijuana.^{3,4} CB₂ receptors are most abundant in the immune system, and they appear to be involved in the anti-inflammatory and possibly other therapeutic effects of cannabis.^{3–5} Thus, CB₂ represents an attractive pharmacological target for developing cannabinoid-based therapeutic agents in the treatment of several human conditions (e.g., cancer, chemotherapy relief, neuropathic pain, neurotrauma, and appetite regulation).⁶


The characterization of the CB₁ and CB₂ receptors has launched the quest for their corresponding endogenous ligands, which led in 1992 to the isolation of the lipid derivative anandamide (1).^{2,7} This endogenous metabolite binds to both CB₁ and CB₂ receptors and is found in nearly all tissues in a wide range of animals.² Besides 1, several other endogenous fatty acid-derived compounds have been reported that are capable of binding one or both of the cannabinoid receptors.^{6,8} Among them, 2-arachidonoylglycerol (2) is more abundant and efficacious than 1, but as a monoglyceride, it takes part in several other pathways of lipid metabolism that minimizes its availability for endocannabinoid signaling.^{2,9}

Marine organisms in general, and cyanobacteria in particular, have been a rich source of unusual alkyl amides that resemble structurally some aspects of anandamide and other endocannabinoids.¹⁰ Our group has previously reported cannabinomimetic activity for the chlorosulfolipid malhamensilipin A,^{10c} as well as the fatty acid-based metabolites grenadadiene^{10d} and semiplenamides A, B, and G.^{10e}

As part of an ongoing program aimed at characterizing the bioactive natural products of various marine organisms, we investigated the extracts of two independent collections of marine cyanobacteria that shared similar NMR characteristics. The first extract, prepared from a specimen of *Lyngbya majuscula* collected in Papua New Guinea, was fractionated chromatographically to afford the new cannabinoid-active lipid, serinolamide A (3), bearing a strong structural similarity to anandamide (1). Extracts from this same collection of *L. majuscula* were reported previously to produce the potent cancer cell cytotoxin apratoxin D, a cyclodepsipeptide metabolite.¹¹ The second extract, derived from a *Oscillatoria* sp. specimen, showed NMR resonances analogous to those of serinolamide A (3), and its ¹H NMR-guided fractionation led to the discovery of the new cyanobacterial lipid propenediester (4). Herein, we report the isolation, structure elucidation, and biological activity of serinolamide A (3) and propenediester (4), two new endocannabinoid-like lipids isolated from marine cyanobacteria.

Received: July 22, 2011

Published: October 14, 2011



The marine cyanobacterium *L. majuscula* was collected by hand using scuba from New Ireland in Papua New Guinea. The sample was extracted with CH₂Cl₂–MeOH (2:1), and 8.8 g of the resulting extract was fractionated by silica gel vacuum-liquid chromatography (VLC) to obtain nine fractions (A–I). Further purification of this material using reversed-phase HPLC yielded 2.9 mg (0.03%) of serinolamide (3). HRESITOFMS analysis of 3 showed a pseudomolecular ion [M + H]⁺ at *m/z* 384.3479 and its sodium adduct [M + Na]⁺ at *m/z* 406.3295, corresponding with the molecular formula, C₂₃H₄₅NO₃, and indicating that this compound possesses two degrees of unsaturation. IR spectroscopy showed absorptions at 3351 and 1622 cm⁻¹, suggesting the presence of hydroxy and amide-carbonyl groups. The ¹H NMR spectrum of 3 showed resonances for three methyl groups, two of which were consistent with being directly attached to oxygen and nitrogen atoms, respectively [δ 3.33 (s, H-23) and 3.01 (s, H-19)] (Table 1). The third methyl group

Table 1. NMR Spectroscopic Data (500 MHz, CDCl₃) for Serinolamide A (3)

carbon	δ_C , mult. ^a	δ_H , mult. (<i>J</i> in Hz) ^b	HMBC ^c	COSY
1	174.3, C			
2	34.3, CH ₂	2.40, m	1, 3, 4	3
3	28.1, CH ₂	2.32, m	1, 2, 4, 5	2, 4
4	128.4, CH	5.43, m	3	3, 5
5	131.6, CH	5.47, m	6	4, 6
6	32.6, CH ₂	1.96, dd (6.5, 13.5)	4, 5, 7	5
7	29.5, CH ₂	1.25, m		
8 ^d	29.2, CH ₂	1.25, m		
9 ^d	29.3, CH ₂	1.25, m		
10 ^d	29.64, CH ₂	1.25, m		
11–15 ^d	29.68, 5 × CH ₂	1.25, m		
16	31.9, CH ₂	1.25, m		
17	22.7, CH ₂	1.28, m	14, 16	18
18	14.1, CH ₃	0.88, t (6.5)	16, 17	17
19	33.8, CH ₃	3.01, s	1, 20	
20	57.8, CH	4.39, m	1, 19, 21, 22	21, 22
21	62.3, CH ₂	3.78, m	20, 22	20
22	71.0, CH ₂	3.66, dd (7.0, 10.0) 3.56, dd (4.5, 10.0)	20, 21, 23 20, 21, 23	20
23	58.9, CH ₃	3.33, s		

^aRecorded at 125 MHz. ^bRecorded at 500 MHz. ^cFrom proton to the indicated carbon. ^dAssignments may be interchanged.

at δ 0.88 (t, *J* = 6.5 Hz, H-18) showed a chemical shift and multiplicity typical of a terminal methyl group of a fatty acid chain. Furthermore, resonances for 16 methylenes were observed, some of which were deshielded to shifts corresponding to oxymethylenes [δ 3.78 (m, H-21), 3.66 (dd, *J* = 7.0, 10.0 Hz, H-22a), and 3.56 (dd, *J* = 4.5, 10.0 Hz, H-22b)]. Other methylene groups appeared at high field [δ 2.40 (m,

H-2), 2.32 (m, H-3), 1.96 (m, H-6), and 1.28 (m, H-17)], which, in combination with a 20H signal at δ 1.25 for 10 methylenes and two methine resonances at δ 5.43 (m, H-4) and 5.47 (m, H-5), defined a monounsaturated alkyl chain. The remaining resonance in the spectrum of 3 was a deshielded methine resonance at δ 4.39 (m, H-20) and, by HSQC, corresponded to a nitrogen-bonded methine.

The ¹³C NMR spectrum of 3 showed resonances for only 19 carbon atoms. However, the carbon at δ 29.7 was of larger intensity, indicating a superposition of several carbon atoms, as is typical for chemically equivalent methylenes in a fatty acid chain. With 23 carbon atoms assigned to 3 by HRMS analysis, the ¹³C NMR and gHSQC spectra were interpreted to contain three methyl groups, 16 methylenes, three methines, and one quaternary carbon (Table 1).

Overall, analysis of the NMR data of compound 3 (Table 1) allowed the construction of two substructures (Figure 1A).




Figure 1. Assembly of compounds 3 (A) and 4 (B) from their 2D NMR data.

In the first, a prominent singlet at δ 1.25 ppm (20 H) and a triplet at δ 0.88 (3H) suggested the presence of a mostly saturated fatty acid moiety. COSY and HMBC correlations allowed the assignment of the spin system formed by H-2, H-3, H-4, H-5, and H-6 (I, Figure 1A), and the *trans*-geometry of the C-4/C-5 double bond was assigned on the basis of the δ_C values of C-3 and C-6, which are typical for allylic methylenes of a *trans* double bond.¹² The group of 10 overlapped methylenes observed at δ 1.25 ppm was localized between C-6 and C-17 on the basis of COSY and HMBC correlations, completing the assignment of substructure I (Figure 1A).

The second substructure, II, consisted of a dimethyl serinol unit that was elucidated as follows. The spin system formed by H-20, H-21, and H-22 was assigned by COSY and ^{2,3}J HMBC correlations. The chemical shifts of carbons C-21 and C-22 were typical for oxygen-bonded methylenes, while the chemical shift of the carbon C-20 indicated that it is attached to nitrogen. Methyl groups at δ 3.01 (H-19) and 3.33 (H-23) were attached to the serinol nitrogen and the oxygen at C-22, respectively, on the basis of diagnostic HMBC correlations, thus completing the assignment of substructure II. Substructures I and II were connected by HMBC correlations observed between the N-Me protons with carbons C-1 and C-20 as well as correlations observed between H-20 and carbons C-1, C-21, and C-22 (Figure 1A).

The absolute configuration at C-20 of the serinol unit was established via chiral derivatization of **3** with (*R*)- and (*S*)-2-(anthracen-9-yl)-2-methoxyacetic acid (9-AMA), according to the methodology reported by Riguera and co-workers for determining the absolute configuration of β -chiral primary alcohols.¹³ Examination of $\Delta\delta$ values obtained from the *R*- and *S*-ester derivatives **3a** and **3b** led to an *S*-configuration assigned at C-20. For the natural product **3** the priority rules for the substituents change, and thus, serinolamide (**3**) possesses the *R*-configuration at C-20.




Figure 2. $\Delta\delta^{RS}$ values (ppm, DMSO- d_6) for the 9-AMA esters **3a** and **3b**. The priority order of the substituents at C-20 changes when $R_2 = \text{H}$; thus the natural product **3** possesses an *R*-configuration at C-20.

In the case of *Oscillatoria* sp., a black mat of cyanobacterial biomass was collected by hand from Isla Canales de Afuera, Panamá. A similar extraction and fractionation strategy produced nine subfractions (A–I), for which fractions A (1.3 mg) and B (22.0 mg) displayed similar intriguing ^1H NMR resonances and LC-MS data to those for compound **3**. Thus, fractions A and B were combined and subjected to further silica gel column chromatography to afford metabolite **4** (6.6 mg, 0.1%).

HRESIMS of **4** yielded an $[\text{M} + \text{Na}]^+$ peak at m/z 417.2978, which when combined with the NMR data (Table 2), was consistent with the molecular formula $\text{C}_{24}\text{H}_{42}\text{O}_4$, possessing four degrees of unsaturation (calcd for $\text{C}_{24}\text{H}_{42}\text{O}_4\text{Na}$, 417.2975). Intense IR absorptions at 1746 and 1136 cm^{-1} suggested the presence of double and single carbon–oxygen bonds of an ester functional group. The ^1H NMR spectrum of **4** (Table 2) displayed resonances assignable to a deshielded vinylic proton at δ 7.23 and its coupling partner at δ 5.08, a complex multiplet for two deshielded methylenes between δ 4.76 and 4.70, six methylene multiplets at δ 2.43, 2.06, 1.99, 1.81, 1.40, and 1.28, nine isochronous methylenes centered at δ 1.25, a methyl singlet at δ 2.07, and a final methyl triplet at δ 0.88. In agreement with these assignments, the ^{13}C NMR spectrum of **4** showed two sp^2 methine carbons at δ 136.9 and 107.4, a methylene carbon at δ 57.9 (adjacent to oxygen), two methyl resonances at δ 20.9 and 14.1, and a series of 12 methylene carbons ranging from δ 35.8 to 22.4. This spectrum was completed by deshielded quaternary carbonyl-type resonances at δ 170.8 and 170.1, as well as a quaternary sp^2 carbon at δ 148.5 and a sp^2 methylene resonance at δ 109.7. Extensive analysis by 2D NMR methods, including the HSQC, HMBC, and COSY spectra, revealed the planar structure of propenediester (**4**) as described below.

The most prominent resonance in the ^1H NMR spectrum of **4** was an apparent singlet centered at δ 1.25, which accounted for a total of 18 protons assigned as methylenes via the HSQC

Table 2. NMR Spectroscopic Data (600 MHz, CDCl_3) for Propenediester (**4**)

carbon	δ_{C} , mult. ^a	δ_{H} , mult. (J in Hz) ^b	HMBC ^c	COSY
1	170.1, C			
2	33.3, CH ₂	2.43, t (7.4)	1, 3, 4	3, 4
3	22.4, CH ₂	1.81, tt (7.4, 7.6)	1, 2, 5	2, 4
4	35.1, CH ₂	2.06, t (7.6)	2, 3, 5, 6, 19	2, 3, 6, 19
5	148.5, C			
6	35.8, CH ₂	1.99, t (7.6)	5, 7, 19	4, 7, 19
7	27.7, CH ₂	1.40, tt (6.8, 7.6)	8	6, 8
8	29.7, CH ₂	1.25, m		7
9–12 ^d	29.6, 4 × CH ₂	1.25, m		
13 ^d	29.5, CH ₂	1.25, m		
14 ^d	29.4, CH ₂	1.25, m		
15 ^d	29.3, CH ₂	1.25, m		
16	31.9, CH ₂	1.25, m		17
17	22.7, CH ₂	1.28, m	16	16, 18
18	14.1, CH ₃	0.88, t (7.1)	16, 17	17
19	109.7, CH ₂	4.73, d (16.6)	4, 6	4, 6
20	136.9, CH	7.23, dt (1.5, 6.5)	1, 21, 22	21, 22
21	107.4, CH	5.08, dt (6.5, 7.1)	20	20, 22
22	57.9, CH ₂	4.72, dd (1.5, 7.1)	20, 21, 23	20, 21
23	170.8, C			
24	20.9, CH ₃	2.07, s	23	

^aRecorded at 125 MHz. ^bRecorded at 600 MHz. ^cFrom proton to the indicated carbon. ^dAssignments may be interchanged.

spectrum, and was indicative of a saturated fatty acid chain. COSY data allowed definition of the termini flanking this alkyl chain. At one end an ethyl group was found as revealed by a methyl triplet at δ 0.88 (H-18, $J = 7.1$ Hz, δ_{C} 14.1) and a methylene multiplet at δ 1.28 (H-17, δ_{C} 22.7). The other end was composed of a 1,1-disubstituted double bond, as indicated by a deshielded sp^2 methylene group at δ 4.73 (H₂-19, apparent d, $J = 16.6$ Hz, δ_{C} 109.7) and a sp^2 quaternary carbon at δ 148.5 (C-5). The protons for this unusual branching methylene showed both COSY and HMBC correlations with flanking methylenes at δ 1.99 (H-6, apparent t, $J = 7.6$ Hz, δ_{C} 35.8) and 2.06 (H-4, apparent t, $J = 7.6$ Hz, δ_{C} 35.1). The former was separated from the C₉ saturated alkyl chain mentioned above by an additional methylene at δ 1.40 (H-7, m, δ_{C} 27.7), whereas the latter methylene was found to comprise a new spin system with methylenes at δ 1.81 (H-3, tt, $J = 7.4, 7.6$ Hz, δ_{C} 22.4) and 2.43 (H-2, t, $J = 7.4$ Hz, δ_{C} 33.3). HMBC correlations from H-3 and H-2 were used to situate an ester-type carbonyl (δ 170.1, C-1) at the end of this spin system, thus completing the elucidation of the major portion of compound **4** (fragment III, Figure 1B).

A second spin system in **4** contained a polarized vinylic pair at δ 7.23 (H-20, dt, $J = 1.5, 6.5$ Hz, δ_{C} 136.9) and 5.08 (H-21, dt, $J = 6.5, 7.1$ Hz, δ_{C} 107.4), followed by a deshielded methylene (adjacent to oxygen) at δ 4.72 (H-22, dd, $J = 1.5, 7.1$ Hz, δ_{C} 57.9), exhibiting vicinal and allylic coupling with H-21 and H-20, respectively (fragment IV). The configuration of the polarized double bond was determined as *Z* on the basis of a $^3J_{\text{H}20,\text{H}21}$ value of 6.5 Hz. A HMBC correlation from H-20 to C-1 enabled substructures III and IV to be connected. Likewise, a HMBC correlation from H-22 with a carbonyl carbon at δ 170.8 (C-23) was used to join IV with V, identified as an acetyl group comprising C-23 and the methyl singlet at δ 2.07 (H-24, δ_{C} 20.9). This completed the structure elucidation

of propenediester (**4**), a compound with a uniquely branched fatty acid chain.

Due to the similarity in structural features of metabolites **3** and **4** to those of known endocannabinoids (e.g., **1** and **2**), these were evaluated for cannabimimetic properties in radioligand binding assays specific for human CB₁ and CB₂ receptors. Compound **3** exhibited moderate affinity for the CB₁ cannabinoid receptor with a K_i value of 1.3 μM , whereas **4** was completely inactive (Table 3). More importantly, **3** appears

Table 3. CB₁ and CB₂ Receptor Affinity of Compounds **3** and **4**

compound	CB ₁ IC ₅₀ (μM)	CB ₂ IC ₅₀ (μM)
Δ^9 -THC ^a	0.0028	0.0095
3	2.3 ± 0.1^b	>10
4	>10	>10

^aPositive control. ^b $K_i = 1.3 \mu\text{M}$.

to be selective for CB₁ (>5-fold). Although other CB₁ agonists (e.g., Δ^9 -THC, HU-210, ACEA) exhibit much higher affinity and selectivity over CB₂ receptors, only a handful of marine natural products are reported to show such potency and selectivity for CB₁ receptors.¹⁴ Furthermore, chemical entities exhibiting K_i values in the micromolar range such as **3** potentially can exert cannabimimetic effects in vivo, making them attractive from a therapeutic development perspective. Metabolite **3** was cytotoxic when evaluated in vitro against the H-460 human lung cell line (50 $\mu\text{g}/\text{mL}$, 0% cell survival; 5 $\mu\text{g}/\text{mL}$, 60% cell survival), whereas compound **4** was inactive at all tested concentrations (up to 20 $\mu\text{g}/\text{mL}$). These effects might be ascribed to the potential CB₁ agonist activity of **3**, since CB₁ agonists have been shown to exert cytotoxic effects via either induction of apoptosis or inhibition of proliferation.¹⁵ The lack of cannabinoid-type activity for propenediester (**4**) in these binding assays suggests that moieties containing a hydrogen bond donor at the hydrophilic end of their molecule are required to elicit cannabinoid activity. *N*-Arachidonoyl-L-serine, an endogenous mediator found in mammalian tissues and sharing perhaps a greater chemical similarity with **3** than **1**,¹⁶ was found to be completely inactive in both human CB₁ and CB₂ binding assays ($K_i > 10 \mu\text{M}$), despite containing a secondary amide group in its chemical structure, and possibly because of the presence of a terminal carboxyl group instead of the methoxy group of **3**.

Biogenetically, the C-1 to C-18 substructure of compounds **3** and **4** is probably derived from a PKS pathway, responsible for the production of fatty acids. The serinol unit in **3** likely originates from serine, which has been reported to be biosynthesized by cyanobacteria from phosphoglycerate through the phosphorylated pathway.¹⁷ Incorporation of *N*-Me and *O*-Me groups in the serinol residue of compound **3** occurs presumably via SAM-dependent methyltransferases.¹⁸ In the case of compound **4**, its intriguing 1,1-disubstituted double bond branch at C-5 resides at a predicted carbonyl site in the nascent polyketide and, thus, is probably produced by a β -branch mechanism involving an HMG-CoA synthase cassette¹⁹ followed by dehydration to generate the double bond. Although highly speculative, it is tempting to attribute fragment **IV** in **4** to glycerol, which would undergo elimination of its middle substituent (C21) through a more stable secondary intermediate (elimination at C-20 or C-22 would involve a primary intermediate).

EXPERIMENTAL SECTION

General Experimental Procedures. Optical rotations were measured on a JASCO P-2000 polarimeter, whereas UV data were obtained using a Beckman DU800 spectrophotometer. IR spectra were recorded on a Nicolet 100 FT-IR spectrometer. ¹H, ¹³C, and 2D NMR spectra were collected at a ¹H resonance frequency of either 600 MHz (Bruker Avance III DRX600 equipped with a 1.7 mm TCI cryoprobe) or 500 MHz (Varian INOVA 500). Chemical shifts were calibrated internally to the residual signal of the solvent in which the sample was dissolved (CDCl₃, δ_{H} 7.26, δ_{C} 77.0; DMSO-*d*₆, δ_{H} 2.50, δ_{C} 39.5). High-resolution mass spectra were obtained on a ThermoFinnigan MAT900XL mass spectrometer. HPLC was carried out using a dual Waters 515 pump system equipped with a Waters 996 photodiode array detector. Vacuum and flash chromatographic separations were performed using silica gel type H (10–40 μm , Aldrich) and silica gel 60 (40–63 μm , EMD), respectively. Merck TLC sheets (silica gel 60 F254) were used for analytical TLC (aluminum-supported, layer thickness 200 μm) and preparative TLC (glass-supported, layer thickness 250 μm). All solvents were purchased as HPLC grade. Cytotoxicity to H-460 cells was determined as previously reported.²¹

Cyanobacterial Collections and Taxonomic Identification.

The marine cyanobacterium *Lyngbya majuscula* (voucher specimen available from W.H.G. as collection code PNG 5-22-05-3) was collected by scuba at a depth of 15 m from New Ireland, Papua New Guinea, in May 2005 (S 3°49.626', E 152°26.017'). Specimens of *Oscillatoria* sp. (voucher specimen available from W.H.G. as collection code PAC-17-FEB-10-3) were collected by hand at a depth of 1–1.5 m on flat rocks at Isla Canales de Afuera, Coiba National Park, Republic of Panama, in February 2010 (07°41.723' N, 81°38.025' W). The samples were stored in 1:1 EtOH–H₂O and frozen at –20 °C prior to extraction. Morphological and molecular characterizations as well as phylogenetic analyses on *L. majuscula* (PNG 5-22-05-3) are described elsewhere.¹¹ Morphological characterization on *Oscillatoria* sp. was performed using an Olympus IX51 epifluorescent microscope (1000 \times) equipped with an Olympus U-CMAD3 camera. Morphological comparison and putative taxonomic identification of the cyanobacterial specimen were performed in accordance with modern classification systems.²⁰ These results were consistent with a phylogenetic analysis of this organism using the 16S rRNA subunit, as detailed elsewhere.²²

Extraction and Isolation. *L. majuscula* and *Oscillatoria* sp. were extracted separately with 2:1 CH₂Cl₂–MeOH and concentrated to dryness in vacuo to give 8.8 g and 91.0 mg of crude extract, respectively. Separate VLC prefractionation of the crude extracts using a gradient with 0–100% EtOAc in hexanes followed by 0–100% of MeOH in EtOAc yielded nine fractions (A–I). VLC fraction H from *L. majuscula* eluted with 25% MeOH in EtOAc (1.5 g) and was further fractionated by flash column chromatography on silica gel using a stepwise gradient of MeOH in CH₂Cl₂ to yield nine sub-fractions (H1–H9). Fraction H2 was purified by reversed-phase HPLC (continuous gradient from 60:40 MeOH–H₂O to 100% MeOH in 65 min at 2 mL/min) to give 2.9 mg of serinolamide A (**3**) as a pale yellow oil (0.03%). VLC fractions A (eluted with 100% hexanes, 1.3 mg) and B (eluted with 10% EtOAc in hexanes, 22.0 mg) from *Oscillatoria* sp. were combined according to their ¹H NMR, LCMS, and TLC profiles and subjected to ¹H NMR-guided fractionation using silica gel column chromatography (10% EtOAc in hexanes) to yield 6.6 mg (0.1%) of propenediester (**4**).

Serinolamide A (3): pale yellow, oil [α]_D²⁵ +2.6 (*c* 0.28, CHCl₃); IR (film) λ_{max} 3351 (br), 2930, 2853, 1622, 1457, 1404, 1099, 1047 cm⁻¹; ¹H and ¹³C NMR, see Table 1; ESIMS *m/z* 384 (75) [M + H]⁺, 767 (100) [2 M + H]⁺; HRESITOFMS *m/z* [M + H]⁺ 384.3479 (calcd for C₂₃H₄₆NO₃, 384.3472).

Propenediester (4): colorless oil; IR (neat) 3005, 2926, 2855, 1746, 1371, 1224, 1136, 1029, 895 cm⁻¹; ¹H, ¹³C, and 2D NMR data, see Table 2; HRESITOFMS *m/z* [M + Na]⁺ 417.2978 (calcd for C₂₄H₄₂O₄Na, 417.2975).

Derivatization of Serinolamide (3) with (R)- and (S)-9-AMA Acids. Esters **3a** and **3b** were prepared separately by treatment of

serinolamide (**3**) (1 mg) with the corresponding (*R*)- and (*S*)-9-AMA acids (1.1 equiv) in the presence of EDC (1.1 equiv) and DMAP (catalytic) in dry CH₂Cl₂, under a N₂ atmosphere. The reaction was stirred at room temperature for 12 h. The organic layer was washed sequentially with H₂O, HCl (1 M), H₂O, NaHCO₃ (sat), and H₂O, then dried (Na₂SO₄) and concentrated under reduced pressure to obtain the corresponding ester. Final purification was achieved by flash column chromatography on silica gel 230–400 mesh (elution with hexanes–ethyl acetate mixtures) to yield 2.2 mg of **3a** and 1.4 mg of compound **3b**.

(*R*)-9-AMA ester derivative (**3a**): $[\alpha]_D^{25} -10.1$ (c 1.4, CHCl₃); ¹H NMR (500 MHz, DMSO-*d*₆) δ 4.37 (1H, m, H-21), 4.08 (1H, dd, *J* = 7.2, 11.4 Hz, H-20a), 4.0 (1H, dd, *J* = 5.3, 11.4 Hz, H-20b), 3.08 (1H, m, H-22a), 2.98 (1H, dd, *J* = 5.5, 10.1 Hz, H-22b), 2.94 (3H, s, H-23), 2.16 (3H, s, H-19); HRESITOFMS *m/z* [M + H]⁺ 632.4313 (calcd for C₄₀H₅₈NO₅, 632.4310).

(*S*)-9-AMA ester derivative (**3b**): $[\alpha]_D^{25} +10.1$ (c 0.93, CHCl₃); ¹H NMR (500 MHz, DMSO-*d*₆) δ 4.46 (1H, m, H-21), 4.07 (1H, dd, *J* = 8.5, 11.6 Hz, H-20a), 3.94 (1H, dd, *J* = 4.55, 11.5 Hz, H-20b), 2.89 (3H, s, H-23), 2.88 (1H, bs, H-22a), 2.87 (1H, bs, H-22b), 1.97 (3H, s, H-19); HRESITOFMS *m/z* [M + H]⁺ 632.4304 (calcd for C₄₀H₅₈NO₅, 632.4310).

CB₁ and CB₂ Receptor Binding Assays. Membranes from HEK-293 cells, stably transfected with either the human recombinant CB₁ receptor (*B*_{max} = 2.5 pmol/mg protein) or human recombinant CB₂ receptor (*B*_{max} = 4.7 pmol/mg protein), were incubated with [³H]-CP-55,940 (0.14 nM/*K*_d = 0.18 nM and 0.084 nM/*K*_d = 0.31 nM, respectively, for CB₁ and CB₂ receptors) as the high-affinity ligand. This ligand was then displaced with 10 μM WIN 55212-2 as the heterologous competitor for nonspecific binding (*K*_i values 9.2 and 2.1 nM respectively for CB₁ and CB₂ receptors). The two compounds were tested in a preliminary screening following the procedure described by the manufacturer (Perkin-Elmer, Milano, Italy). The compound that displaced [³H]CP-55,940 by more than 50% at 10 μM was further analyzed by carrying out a complete concentration–response curve (using 10, 100, 500, 1000, 2500, 10 000 nM concentrations). Displacement curves were generated by incubating test compounds for 90 min at 30 °C with [³H]CP-55,940 (0.14 nM and 0.084 μM in CB₁ and CB₂ binding assays, respectively). *K*_i values were calculated by applying the Cheng–Prusoff equation to the IC₅₀ values (obtained by GraphPad) for the displacement of the bound radioligand by increasing concentrations of the test compounds. Data are means ± SEM of *n* = 3 experiments.

■ ASSOCIATED CONTENT

● Supporting Information

¹H NMR, ¹³C NMR, and 2D NMR spectra in CDCl₃ of compounds **3** and **4**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Tel: (507)-5170732. Fax: (507)-5170701. E-mail: mgutierrez@indicat.org.pa and Tel: (858)-534-0576. Fax: (858)-534-0529. E-mail: wgerwick@ucsd.edu.

■ ACKNOWLEDGMENTS

We gratefully acknowledge the governments of Panamá and Papua New Guinea for permission to make the organism collections. M.G. thanks SENACYT, Panamá, for financial support; H.M.D. thanks FAPESP, Brazil, for providing a fellowship. Special thanks are due to Dr. R. Riguera for kindly providing the (*R*)- and (*S*)-9-AMA acids for stereochemical analysis. We also thank K. Tidgewell and C. Sorrels for help with the cyanobacterial collections, N. Engene for identification of the source cyanobacterium, Y. Su (UCSD) for the MS data acquisition, and F. Freire and J. Seco for comments on the

stereochemical analysis. This work was supported by NIH CA 10085, CA 52955, and the Fogarty International Center's International Cooperative Biodiversity Groups program (grant no. TW006634).

■ REFERENCES

- (1) Felder, C. C.; Glass, M. *Annu. Rev. Pharmacol. Toxicol.* **1998**, *38*, 179–200.
- (2) Lambert, D. M.; Fowler, C. J. *J. Med. Chem.* **2005**, *48*, 5059–5087.
- (3) Stott, C. G.; Guy, G. W. *Euphytica* **2004**, *140*, 83–93.
- (4) Howlett, A. C.; Barth, F.; Bonner, T. I.; Cabral, G.; Casellas, P.; Devane, W. A.; Felder, C. C.; Herkenham, M.; Mackie, K.; Martin, B. R.; Mecholaum, R.; Pertwee, R. G. *Pharmacol. Rev.* **2002**, *54*, 161–202.
- (5) Marzo, V. D.; Petrocellis, L. D. *Annu. Rev. Med.* **2006**, *57*, 553–574.
- (6) Pacher, P.; Batkai, S.; Kunos, G. *Pharmacol. Rev.* **2006**, *58*, 389–462.
- (7) Devane, W. A.; Hanus, L.; Breuer, A.; Pertwee, R. G.; Stevenson, L. A.; Griffin, G.; Gibson, D.; Mandelbaum, A.; Etinger, A.; Mechoulam, R. *Science* **1992**, *258*, 1946–1949.
- (8) Gonsiorek, W.; Lunn, C.; Fan, X.; Narula, S.; Lundell, D.; Hipkin, W. *Mol. Pharmacol.* **2000**, *57*, 1045–1050.
- (9) Mecholaum, R.; Shabat, S. B.; Hanus, L.; Ligumsky, M.; Kaminski, N. E.; Schatz, A. R.; Gopher, A.; Almong, A.; Martin, B. R.; Compton, D. R.; Pertwee, R. G.; Griffin, G. *Biochem. Pharmacol.* **1995**, *50*, 83–90.
- (10) (a) Soderstrom, K.; Murray, T. F.; Yoo, H. D.; Ketchum, S.; Milligan, K.; Gerwick, W.; Ortega, M. J.; Salva, J. *Adv. Exp. Med. Biol.* **1997**, *433*, 73–77. (b) Pereira, A.; Pfeifer, T. A.; Grigliatti, T. A.; Andersen, R. J. *ACS Chem. Biol.* **2009**, *4*, 139–144. (c) Bisogno, T.; Melck, D.; De Petrocellis, L.; Bobrov, M. Y.; Gretskeya, N. M.; Bezuglov, V. V.; Sitachitta, N.; Gerwick, W. H.; Di Marzo, V. *Biochem. Biophys. Res. Commun.* **1998**, *248*, 515–522. (d) Han, B.; McPhail, K. L.; Ligresti, A.; Di Marzo, V.; Gerwick, W. H. *J. Nat. Prod.* **2003**, *66*, 1364–1368.
- (11) Gutiérrez, M.; Suyama, T. L.; Engene, N.; Wingerd, J. S.; Matainaho, T.; Gerwick, W. H. *J. Nat. Prod.* **2008**, *71*, 1099–1103.
- (12) Gunstone, F. D.; Pollard, M. R.; Scrimgeour, C. M.; Vedanayagam, H. S. *Chem. Phys. Lipids* **1977**, *18*, 115–129.
- (13) Freire, F.; Seco, J. M.; Quiñoá, E.; Riguera, R. *Chem. Commun.* **2007**, 1456–1458.
- (14) Tidgewell, K.; Clark, B. T.; Gerwick, W. H. In *Comprehensive Natural Products Chemistry*, 2nd ed.; Moore, B.; Crews, P., Eds.; Pergamon Press: Oxford, UK, 2010; pp 141–188.
- (15) Malfitano, A. M.; Ciaglia, E.; Gangemi, G.; Gazzerzo, P.; Laezza, C.; Bifulco, M. *Expert Opin. Ther. Targets* **2011**, *15*, 297–308.
- (16) Milman, G.; Maor, Y.; Abu-Lafi, S.; Horowitz, M.; Gallily, R.; Batkai, S.; Mo, F. M.; Offertaler, L.; Pacher, P.; Kunos, G.; Mechoulam, R. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 2428–2433.
- (17) Colman, B.; Norman, E. *Physiol. Plant.* **1997**, *100*, 133–136.
- (18) Jones, A. C.; Monroe, E. A.; Eisman, E. B.; Gerwick, L.; Sherman, D.; Gerwick, W. H. *Nat. Prod. Rep.* **2010**, *27*, 1048–1065.
- (19) Geders, T. W.; Gu, L.; Mowers, J. C.; Liu, H.; Gerwick, W. H.; Hakansson, K.; Sherman, D. H.; Smith, J. L. *J. Biol. Chem.* **2007**, *282*, 35954–35963.
- (20) (a) Geitler, L. In *Cyanophyceae*; Rabenhorst, L., Ed.; Akademische Verlag: Leipzig, 1932; reprint: Koeltz Scientific Books: Königstein, 1985; pp 1060–1061. (b) Komárek, J.; Anagnostidis, K. In *Süßwasserflora Von Mitteleuropa*; Büdel, B.; Gartner, G.; Krienitz, L.; Schagerl, M., Eds.; Gustav Fischer: Jena, 2005; *19/2*, pp 483–606.
- (21) Mevers, E.; Liu, W. T.; Engene, N.; Mohimani, H.; Byrum, T.; Pevzner, P. A.; Dorrestein, P. C.; Spadafora, C.; Gerwick, W. H. *J. Nat. Prod.* **2011**, *74*, 928–936.
- (22) Malloy, K. L.; Suyama, T. L.; Engene, N.; Debonsi, H.; Cao, Z.; Matainaho, T.; Di Marzo, V.; Spadafora, C.; Murray, T. F.; Gerwick, W. H. *J. Nat. Prod.* (submitted).