

Chemokine receptors and neural function

Charlene Cho¹ and Richard J Miller²

¹Department of Neurobiology, Pharmacology and Physiology, University of Chicago, Chicago, Illinois, USA; and
²Department of Molecular Pharmacology and Biological Chemistry, Northwestern University Medical School, Chicago, Illinois, USA

Numerous studies have demonstrated that chemokines play an integral role in diseases marked by inflammation. Recently, it has also been shown that chemokines and their receptors are widely expressed in the central nervous system by all types of cells, including neurons. The functions of neuronal chemokine receptors have yet to be fully defined. However, there are indications that neuronal chemokine receptors play an integral role in the development of the nervous system, in the regulation of neuronal excitability and in the signal transduction pathways that regulate neuronal survival. This review explores these topics and discusses the overall impact that chemokines may have on neuronal function. *Journal of NeuroVirology* (2002) **8**, 573–584.

Keywords: calcium; GPCR; neuronal survival; neurotransmission; signal transduction

Introduction

Survival of the organism requires that the host response to injury, insult, and/or infection be swift and accurate. In vertebrates, that function is carried out by the immune system and initially characterized by the trafficking (i.e., circulation, homing, and extravasation) of the appropriate immune system cells to the site of damage. One of the most important classes of molecules involved in directing this process are the chemokines (CHEMOtaxic cytoKINES), a family of low-molecular-weight (8 to 10 kDa), basic proteins whose production can be up-regulated in the presence of inflammatory stimuli (Rollins, 1997; Baggiolini, 1998; Luster, 1998).

As local mediators of immune cell infiltration, chemokines are believed to play a prominent role in a variety of disease states characterized by inflammation. For example, arthritis (Barnes *et al*, 1998), asthma (Kita and Gleich, 1996; Lukacs *et al*, 1996; Griffiths-Johnson *et al*, 1997), atherosclerosis (Boring

et al, 1998; Gu *et al*, 1998; Goslin *et al*, 1999), inflammatory bowel syndrome (Katsuta *et al*, 2000), and glomerulonephritis (Brown *et al*, 1996; Segerer *et al*, 2000) are all believed to be mediated in some part by chemokines and their effects on target cells. In addition, multiple sclerosis (MS), amyotrophic lateral sclerosis (ALS), Parkinson's disease (PD), and Alzheimer's disease (AD), four neurological disorders that are discussed in this special issue, are all marked by inflammation at some point during disease onset or progression and thus are also very likely to include a chemokine-mediated component.

Chemokines and their receptors are also involved in certain virally mediated diseases, such as acquired immunodeficiency syndrome (AIDS) (Cocchi *et al*, 1995; Feng *et al*, 1996; Berger, 1997; Rucker *et al*, 1997). Studies have shown that two chemokine receptors, CCR5 and CXCR4, play a crucial role in human immunodeficiency virus (HIV)-1 infection (Alkhatib *et al*, 1996; Choe *et al*, 1996; Deng *et al*, 1996; Doranz *et al*, 1996; Dragic *et al*, 1996; Feng *et al*, 1996; Oberlin *et al*, 1996; Berson and Doms, 1998; Berger *et al*, 1999). This finding has been extended to studies of the central nervous system (CNS) and resulted in data implicating chemokine receptors in the neuropathogenesis of AIDS. For example, CCR5 expression has been found to be elevated in microglia of patients with HIV-1-associated dementia (HAD) (Rottman *et al*, 1997; Sanders *et al*, 1998;

Address correspondence to Richard J Miller, PhD, Department of Molecular Pharmacology and Biological Chemistry, Northwestern University Medical School, 303 East Chicago Avenue, Ward Building, 8-296, Chicago, IL 60611, USA. E-mail: r-miller10@northwestern.edu

Received 14 March 2002; revised 22 April 2002; accepted 15 May 2002.

Vallat *et al*, 1998; Wang *et al*, 2002). In addition, chemokine receptor expression by cells in the CNS has been shown to facilitate infection by HIV-1 (Nuovo *et al*, 1994; Saito *et al*, 1994; Gabuzda *et al*, 1998; Brack-Werner, 1999; Sabri *et al*, 1999), a topic elaborated upon in this issue by Zheng and Benveniste. These reports, along with studies demonstrating that infected microglia and perivascular macrophages, the only types of cells in the CNS that allow for productive viral replication (Koenig *et al*, 1986; Wiley *et al*, 1986; Watkins *et al*, 1990; Lee *et al*, 1993; Takahashi *et al*, 1996), synthesize and secrete into the extracellular milieu soluble toxic products that elicit neuronal death, have led to the currently held belief that chemokine receptors play a major role in the molecular mechanisms by which HIV-1 induces neuronal loss (Giulian *et al*, 1990; Pulliam *et al*, 1991; Genis *et al*, 1992; Gelbard *et al*, 1994a, 1994b; Kaul *et al*, 2001). Indeed, such a mechanism of action could account for the neuropathological symptoms observed in AIDS (Sanders *et al*, 1998; Miller and Meucci, 1999; Zheng *et al*, 1999a, 1999b; Gabuzda and Wang, 2000; Pandey and Bolsover, 2000).

Although studies reporting the involvement of chemokines and their receptors in various disease states underscore their importance in the propagation of disease, the expression of chemokines and their cognate receptors in the CNS under normal conditions suggests that these molecules subservise physiological functions in the brain. This review will discuss that possibility, with particular emphasis on chemokine-mediated effects on neurons.

Expression of chemokine receptors by neurons

It was initially believed that in the CNS, neurons were the sites of chemokine receptor expression, whereas astrocytes and microglia were the major sites of chemokine synthesis. However, numerous reports have made it increasingly clear that all of the major cell types in the brain are capable of expressing both chemokines and chemokine receptors, especially in the presence of proinflammatory stimuli (Lavi *et al*, 1998; Ascensio and Campbell, 1999; Hesselgesser and Horuk, 1999; Mennicken *et al*, 1999; Bajetto *et al*, 2001).

Chemokine receptors are members of the G-protein-coupled receptor (GPCR) superfamily (Murphy, 1994, 1996; Premack and Schall, 1996), a class of membrane receptors distinguished structurally by the presence of seven transmembrane domains and functionally by their ability to signal intracellularly through heterotrimeric GTP-binding proteins (Gether and Kobilka, 1998; Ji *et al*, 1998; Lefkowitz, 1998). Comparative analysis of the primary sequence of chemokine receptors has yielded several observations: (1) the total number of amino acids can range from 340 to 370; (2) the N-terminal segment is relatively short and acidic; (3) the pres-

ence of a cysteine at the N-terminus and in each of the four extracellular loops; (4) the conservation of the DRYLAIV sequence (or some slight variation thereof) in the second intracellular loop; and (5) the existence of numerous serines and threonines in the C-terminal tail that serve as targets for phosphorylation following receptor activation (Murphy, 1994, 1996; Baggiolini *et al*, 1997; Berson and Doms, 1998; Ward and Westwick, 1998; Bajetto *et al*, 2001). It has also been reported that chemokine receptors undergo dimerization upon ligand binding, and that such a consequence serves a pivotal role in transducing the signal within the cell (Rodriguez-Frade *et al*, 1999a, 2001; Vila-Coro *et al*, 1999).

At present, chemokine receptors are subdivided according to the chemokine ligands to which they bind. Because the κ (CXC) and κ (CC) subfamilies comprise the largest groups of chemokine ligands, their cognate receptors also constitute the largest groups of chemokine receptors (Table 1). Notably, a new classification system for chemokines and their receptors has been approved (Murphy *et al*, 2000; Zlotnik and Yoshie, 2000). However, for purposes of clarity and consistency with the large body of literature that was published prior to approval of the new classification system, this review will utilize the traditional nomenclature to describe chemokines and their receptors.

Neuronal expression of κ chemokine receptors

Expression of all of the κ chemokine receptors by neurons has been reported. CXCR1, although originally believed to be lacking in the CNS, has been found at the level of mRNA in cholinergic septal neurons (Puma *et al*, 2001). CXCR2 expression, as revealed by immunohistochemistry, has been found on the surface of projection neurons in the hippocampus, dentate nucleus of the cerebellum, and

Table 1 The κ (CXC) and κ (CC) chemokines and their cognate receptors constitute the largest groups within the chemokine family

Chemokine receptors	Chemokine ligands
CXCR-1	IL-8, GCP-2
CXCR-2	IL-8, GRO- κ , - κ , - κ , NAP-2, ENA-78, GCP-2
CXCR-3	IP-10, MIG, I-TAC
CXCR-4	SDF-1 κ , - κ , - κ
CXCR-5	BLC/BCA-1
CCR-1	MIP-1 κ , -1 κ , RANTES, MCP-2, -3, -4, HCC-1, MIPF-1
CCR-2	MCP-1, -2, -3, -4
CCR-3	Eotaxin-1, -2, -3, RANTES, MIP-3, MIP-1 κ , MCP-2, -3, -4
CCR-4	RANTES, MDC, TARC
CCR-5	RANTES, MCP-2, MIP-1 κ , -1 κ
CCR-6	LARC
CCR-7	SLC, ELC
CCR-8	I-309, TARC, MIP-1 κ
CCR-9	TECK
CCR-10	ESkine
CCR-11	MCP-1, -2, -4
CX ₃ CR-1	Fractalkine
XCR-1	Lymphotactin κ , κ

locus coeruleus (Horuk *et al*, 1997), as well as on hippocampal neurons from embryonic weeks 17 to 22 fetal brains (Hesselgesser *et al*, 1997). Studies of CXCR3 have revealed protein expression on neurons residing in the cortical and subcortical regions of an AD brain (Coughlan *et al*, 2000). Similar results have been found in studies of human fetal neurons and NT2.N neurons (Xia *et al*, 2000), a clonally derived cell line that possesses multiple features consistent with a neuronal phenotype, including glutamate receptors, all three classes of neurofilaments, acetylcholinesterase, muscarinic acetylcholine receptors, and polarized axonal and dendritic processes (Lee and Andrew, 1986; Pleasure *et al*, 1992; Llanes *et al*, 1995). Expression of CXCR4, a topic of particular interest given its role in brain development and AIDS-related neuropathology, has been reported at the level of mRNA and protein in various subpopulations of cortical, hippocampal, and cerebellar neurons in adult (Wong *et al*, 1996; Lavi *et al*, 1997; Westmoreland *et al*, 1998; Zhang *et al*, 1998) and embryonic (Jazin *et al*, 1997; Klein *et al*, 1999; McGrath *et al*, 1999; Boutet *et al*, 2001) brains. In addition, CXCR4 expression may follow a developmental pattern, as indicated by immunohistochemical and immunofluorescence data showing an increase in neuronal expression from birth to 9 months of age in rhesus macaque brain (Westmoreland *et al*, 2002). CXCR5 transcripts have been shown to be present in neurons localized to the granule and Purkinje cell layers of the cerebellum (Kaiser *et al*, 1993) as well as other brain regions and several neuronal cell lines (Kouba *et al*, 1993).

Neuronal expression of κ chemokine receptors

Reports of neuronal expression of κ chemokine receptors have focused mainly on CCR1, CCR2, CCR3, CCR4, and CCR5, with CCR3 and CCR5 being of particular interest due to their role in molecular mechanisms of HIV-1 infection. Expression of CCR5 has been detected at the protein level in human hippocampal and cerebellar neurons (Rottman *et al*, 1997), whereas CCR5 transcripts and protein have been reported in human embryonic neurons (Boutet *et al*, 2001). Human brain cell cultures have also been reported to express CCR5 and CCR3, as revealed by immunocytochemistry (Sanders *et al*, 1998). Studies in nonhuman primates have extended these observations to certain populations of neurons in the hippocampus, dentate gyrus, and cerebral cortex (Westmoreland *et al*, 1998, 2002; Zhang *et al*, 1998). These results were further elaborated upon in studies of normal brain that reported CCR3 staining in pyramidal neurons of the frontal cortex and entorhinal cortex as well as in large neurons of the dentate nucleus of the cerebellum (Sanders *et al*, 1998). Rat primary hippocampal neurons have also been reported to express transcripts of CCR1, CCR4, CCR5, CCR9, and CCR10 (Meucci *et al*, 1998). Interestingly, hNT neurons, a human cell line that has been reported

to possess morphological, neurochemical, and physiological properties characteristic of mature neurons (Andrews, 1984; Pleasure and Lee, 1993; Hartley *et al*, 1999; Saporta *et al*, 2000), have been shown to express CXCR2, CXCR4, CCR1, and CCR5 (Hesselgesser *et al*, 1997). The CX₃CR1 receptor has been reported to be expressed by cultured hippocampal neurons and dorsal root ganglia neurons (Meucci *et al*, 1998; Oh *et al*, 2001). The C receptor has not yet been reported to be expressed on cells of the nervous system.

Effects of chemokines on neurons

Much of the data reporting chemokine responsiveness by neurons indicate that they serve in a developmental and trophic capacity in the CNS and may also influence neuronal excitability and synaptic communication. Studies aimed at identifying the function of chemokines in the developing brain have reported that chemokines can influence the migration of several different populations of neurons and neuronal progenitors. For example, Released upon Activation Normal T-Cell Expressed and Secreted (RANTES) has been reported to induce migration of mouse dorsal root ganglion (DRG) neurons (Bolin *et al*, 1998). Other *in vitro* studies have demonstrated that stromal cell derived factor (SDF)-1 can elicit migration of cerebellar granule neurons (Bajetto *et al*, 1999) and mouse cerebellar granule precursor cells (Klein *et al*, 2001). Rat E15 neuronal progenitor cells have also been reported to exhibit the same response in a dose-dependent manner (Lazarini *et al*, 2000). These *in vitro* results were extended to the *in vivo* level in studies of mice lacking the genes for SDF-1 or CXCR4. Compared to wild-type animals, severe abnormalities in cerebellar development are observed in the brains of these animals. The specific defects reported included a marked distortion of the laminar structure, premature inward migration of external granule cells, and the absence of foliation. These results suggest strongly that chemokines are intimately involved in normal brain development (Ma *et al*, 1998; Zou *et al*, 1998). In addition, recent studies in CXCR4 knockout mice have revealed that other areas of brain development are also affected. The dentate gyrus of the hippocampus was found to be smaller than normal, and this effect was attributed not only to deficits in granule cell migration but also to a reduction in the number of dividing neuronal precursors (manuscript submitted).

The idea that chemokine receptors can affect the development and survival of neuronal precursors suggests possible trophic effects of chemokines. Indeed, studies exploring the effects of chemokines on cells of the immune system cells have yielded data indicating chemokines can promote survival and/or proliferation. These reports, combined with the mechanistic similarities of hematopoiesis and neurogenesis (Mehler and Kessler, 1997; Quesenberry

et al, 1999; Terskikh *et al*, 2001), would suggest that an analogous effect of chemokines on cells in the CNS is very likely to exist. In support of this are several studies demonstrating that interleukin (IL)-8 can enhance the survival of cultured hippocampal pyramidal neurons (Araujo and Cotman, 1993; Horuk *et al*, 1997). Growth-related oncogene (GRO)- κ has also been shown to promote survival of cerebellar granule cells (Limatola *et al*, 2000a). Fractalkine, the sole member of the κ family of chemokines, has also been reported to promote survival of hippocampal pyramidal neurons in the presence of neurotoxic stimuli (Meucci *et al*, 2000). All of these reports serve as striking examples of how regulated expression of chemokines could provide survival promoting effects in the brain.

Although studies on IL-8 and fractalkine support a trophic role for chemokines in the CNS, there are reports of other chemokines, namely SDF-1 κ , inducing the opposite effect. Several studies by independent investigators have reported that exposure of neurons to SDF-1 κ results in apoptotic death (Hesselgesser *et al*, 1998; Kaul and Lipton, 1999; Zheng *et al*, 1999a, 1999b). However, other investigators have reported that SDF-1 κ can actually promote neuronal survival (Meucci *et al*, 1998) and, in one case, proliferation of cerebellar granule precursor cells in the presence of the neuronal mitogen, sonic hedgehog (Klein *et al*, 2001). These contradictory results and similar observations of anti- (Suzuki *et al*, 2001) and proapoptotic (Colamussi *et al*, 2000) effects of SDF-1 κ in cells of the immune system suggest that the effect of SDF-1 κ on cell survival may be a context dependent one.

Cellular effects of chemokines on neurons

One well-established event that follows chemokine receptor activation is the rapid dissociation of the respective heterotrimeric G-protein into the κ and β/γ subunits. In virtually every regard, this event is a critical one, for it provides a significant number of options for inducing cellular events (Gether, 2000; Dhanasekaran and Gutkind, 2001), including mobilization of intracellular calcium (Ca²⁺) and modulation of ion channel activity (Miller, 1998). This has been demonstrated in studies of cells of the immune system known to be responsive to chemokines. For example, natural killer cells exhibit Ca²⁺ transients following treatment with RANTES, SDF-1 κ , and macrophage-derived chemokine (MDC) (Damaj *et al*, 1996). The same effect has been reported in neutrophils treated with IL-8 (Schorr *et al*, 1999). Other studies have shown that chloride channels and Ca²⁺-activated potassium (K⁺) channels in primary human macrophages open in the presence of SDF-1 κ and macrophage inflammatory protein (MIP)-1 (Liu *et al*, 2000). Eosinophils have also been reported to respond to chemokines in a similar manner, as demonstrated by the opening of Ca²⁺-activated

K⁺ channels following RANTES exposure (Saito *et al*, 1996). Meanwhile, studies carried out in heterologous expression systems (e.g., *Xenopus laevis* oocytes) have shown that large currents due to the opening of G-protein-coupled inwardly rectifying K⁺ channels (Kir 3.1) are observed following stimulation of CXCR4 and CCR5 (Madani *et al*, 1998). These results, especially given the important roles played by Ca²⁺ and ion channels in neuronal function, have prompted investigators to look for similar effects in neurons.

Consistent with observations made of cells of the immune system, chemokine receptor activation in neurons has also been found to induce Ca²⁺ transients and modulate ion channel activity. For example, mouse granule and Purkinje cells exhibit Ca²⁺ transients following stimulation with IL-8 and GRO- κ (Giovannelli *et al*, 1998). Similar effects have been observed in hippocampal neurons exposed to MDC-1, RANTES, SDF-1 κ , and fractalkine (Meucci *et al*, 1998). The response of rat Purkinje neurons to chemokines appears to be quite varied, as Ca²⁺ transients are observed in the presence of a wide variety of chemokines (Gillard *et al*, 2002). Interestingly, rat cerebellar granule cells exhibit responsiveness to a more narrow range of chemokines (Limatola *et al*, 2000b; Klein *et al*, 2001; Gillard *et al*, 2002). Studies on neuronal ion channels have reported that voltage-dependent calcium channels (VDCCs) are subject to inhibition by chemokines. For example, hippocampal pyramidal neurons in the presence of fractalkine exhibit a reduction in the voltage-sensitive Ca²⁺ current (Meucci *et al*, 1998). Cholinergic septal neurons treated with IL-8 also respond in a similar manner (Puma *et al*, 2001). In addition, recordings of G1A1 cells (HEK293 cells stably expressing N-type VDCCs) treated with SDF-1 κ show a significant reduction in the Ca²⁺ current (Oh *et al*, 2002). Interestingly, chemokines have also been shown to enhance the generation of action potentials in cultured rat dorsal root ganglia (Oh *et al*, 2001) as well as induce excitatory effects in rat Purkinje neurons (Gillard *et al*, 2002).

Another aspect of neuronal function that has been explored concerns the effects of chemokines on neurotransmitter release. Studies on granule cells and Purkinje cells in cerebellar slices have suggested that stimulation with SDF-1 κ probably results in the release of glutamate (Limatola *et al*, 2000b), whereas GRO- κ may provoke the release of gamma-aminobutyric acid (GABA) (Giovannelli *et al*, 1998) at the synapse. Glutamate is the major excitatory neurotransmitter in the brain, and has been shown to play an important role in mediating synaptic plasticity (a topic which is discussed in this issue by Ragozzino). In addition, excess glutamate at the synapse can result in neurotoxicity (Choi and Rothman, 1990), an observation that may be consistent with the SDF-1 κ -induced neuronal cell death previously described. Overall, these results suggest

that chemokines can regulate neuronal activity not only at the level of ion channel activity but also neurotransmitter release and synaptic communication.

Molecular effects of chemokines on neurons

Because the response of neurons and other cell types to chemokines is the result of various intracellular events, many investigators have sought to examine the signaling pathways activated by chemokines. Most of these studies have been conducted in endothelial cells and in cells of the immune system and have reported the activation of a number of intracellular proteins. However, because attempts to undertake those same types of investigations in neurons are relatively recent, few reports are available regarding the effects of chemokines on intracellular signaling pathways in neurons. Nevertheless, the available data are consistent with the possibility that activation of signaling proteins in neurons may allow chemokines to regulate neuronal excitability and synaptic plasticity as well as the fate and development of neuronal progenitors, as described above.

Chemokines activate a number of signaling pathways, and virtually all of them have been found to involve Ca^{2+} . This property is consistent with the preceding discussion describing the Ca^{2+} transients observed following chemokine receptor stimulation in neurons and other cell types. Molecules such as calcium/calmodulin-dependent kinase (CaMK) (Zheng *et al*, 1999a), protein kinase C (PKC) (Laudanna *et al*, 1998; Guinamard *et al*, 1999), and proline-rich nonreceptor tyrosine kinase 2 (Pyk2) (Del Corno *et al*, 2001) have all been shown to be activated by chemokines. Furthermore, the identification of Pyk2, a protein that has been shown to interact directly with c-Src (Dikic *et al*, 1996), has prompted investigators to examine members of the mitogen-activated protein kinase (MAPK) family as potential targets of chemokine receptor stimulation.

All three members of the MAPK family (e.g., extracellular signal-regulated kinases 1 and 2 [ERK1/2], c-Jun N-terminal kinase/stress-activated protein kinase [JNK/SAPK], and p38) have been observed to undergo phosphorylation following chemokine receptor activation in cells of the immune system. Phosphorylation of ERK1/2 has been reported following stimulation of CXCR1, CXCR2 (Venkatakrishnan *et al*, 2000), CXCR3 (Bonnachi *et al*, 2001), CXCR4 (Ganju *et al*, 1998), CCR1, CCR3 (Kampen *et al*, 2000), and CCR5 (Sato *et al*, 2001). Similarly, stimulation of CXCR3 with IP-10 and monokine induced by interferon gamma (MIG) in mouse cortical neurons results in the activation of ERK1/2 (Xia *et al*, 2000). The same effect has been observed following stimulation of CXCR2 in mouse primary cortical neurons (Xia and Hyman, 2002). Treatment of rat E15 neuronal progenitors and cortical neuronal cultures with SDF-1 κ also elicits the activation of ERK1/2 (Lazarini *et al*, 2000).

The same observation has been made in hippocampal neurons stimulated with fractalkine (Meucci *et al*, 1998). Activation of JNK/SAPK has also been observed in immune system cells following exposure to RANTES, MIP-1 κ , and IL-8 (Yang *et al*, 2001) and as a result of open reading frame (ORF) 74 signaling (also known as HHV8 chemokine receptor, the constitutively active Kaposi's sarcoma-associated herpesvirus GPCR) (Bais *et al*, 1998). A similar effect has been reported in rat cerebellar granule neurons in which JNK1 activation was observed following stimulation of CXCR2 with GRO- κ (Limatola *et al*, 1999). Activation of p38 has been reported in immune system cells treated with RANTES (Sato *et al*, 2001; Wong *et al*, 2001) and GRO- κ (Kampen *et al*, 2000) and suggested in a study of SDF-1-induced neuronal death (Kaul and Lipton, 1999). Signaling by ORF74 has also been reported to provoke activation of p38 (Bais *et al*, 1998). As indicated by these reports, MAPK family members are clearly targeted for activation following chemokine receptor stimulation in neurons.

Protein kinase B (PKB)/c-Akt, a serine/threonine kinase widely recognized for its involvement in cell survival, has also been identified as an intracellular target of chemokine receptor stimulation. Investigations of c-Akt in immune system cells have revealed that activation is observed following stimulation of CXCR3 (Bonnachi *et al*, 2001) and CXCR4 (Kijowski *et al*, 2001). The same effect has been observed in mouse primary cortical neurons treated with mouse homologue of human Gro- α (Gro- κ /KC) (Xia and Hyman, 2002) and in rat primary hippocampal neurons treated with fractalkine (Meucci *et al*, 2000). Consistent with this is the report that showed that phosphatidylinositol-3-kinase (PI3K), a key enzyme involved in the regulation of PKB/c-Akt, is activated following treatment with monocyte chemoattractant protein (MCP)-1 and RANTES in HEK293 cells expressing CCR2 and CCR5 (Mellado *et al*, 2001).

Studies of chemokine receptor activation in immune system cells have also reported the activation of a family of proteins known as janus kinases (JAKs) and signal transducers and activators of transcription (STATs). Investigations in immune cells have revealed that exposure to the κ chemokines RANTES and MIP-1 κ results in the activation of STAT1 and STAT3 (Wong and Fish, 1998). These results were further investigated with RANTES alone and led to the finding that the activation of both JAK2 and JAK3 accompany STAT1 and STAT3 in transducing the signal from CCR5 (Wong *et al*, 2001). In contrast, activation of other JAK/STAT family members, namely STAT5b and JAK1, was observed in the presence of RANTES in HEK293 cells engineered to express CCR5. This study also reported that the activation of STAT5b was found to depend heavily on its physical association with CCR5 (Rodriguez-Frade *et al*, 1999b). Another study has reported that T cells stimulated with the κ

chemokine SDF-1 κ results in activation of four members of the STAT family, STAT1, STAT2, STAT3, and STAT5b, and two members of the JAK family, JAK 1 and JAK2. Similar to the report on CCR5, activation of JAK1 and JAK2 was reported to be tightly associated with their physical association with CXCR4 (Vila-Coro *et al*, 1999). The activation of JAKs and STATs have yet to be reported in neurons exposed to chemokines.

Obviously, the activation of transcription factors such as STATs is an important end point in signaling pathways, because the proteins produced as a result of their activity significantly impact the cell's ability to respond appropriately to stimuli. Attempts to examine this concept in greater detail, particularly in the context of chemokine regulation of neurons, have led to the identification of several other well-studied transcription factors such as nuclear factor (NF)- κ B and cAMP response element binding (CREB) protein.

NF- κ B seems to be constitutively active in cells of the cortex and hippocampus, regions of the brain reported to be very active metabolically (Kaltschmidt *et al*, 1994). This observation is believed to indicate that activation of NF- κ B is associated with protective mechanisms (e.g., antioxidant production) operative in the presence of noxious stimuli (i.e., lipopolysaccharides, hypoxia, etc.) (Schutze *et al*, 1992; Kaltschmidt *et al*, 1994). In support of this are reports showing that NF- κ B activation can be induced in neurons following depolarization or treatment with glutamate (Guerrini *et al*, 1995), tumor necrosis factor- α (TNF- κ), or other proinflammatory cytokines (O'Neill and Kaltschmidt, 1997). Moreover, NF- κ B has been implicated in the regulation of neuronal survival and plasticity (Mattson *et al*, 2000) and transcriptional responses evoked by GPCRs (Ye, 2001). These reports, reminiscent of the effects of chemokines on immune cells and neurons, have led investigators to examine whether chemokine receptor stimulation could result in the activation of NF- κ B. This hypothesis has been confirmed by several recent reports. For example, ORF74 signaling has been reported to activate NF- κ B in endothelial cells (Pati *et al*, 2001). This same effect has been observed in hippocampal neurons treated with fractalkine (Meucci *et al*, 2000).

A third transcription factor that has been identified in studies of signal transduction pathways activated by chemokines is CREB protein. Transactivation of CREB occurs via phosphorylation at serine 133, a modification that has been shown to be inducible by increases in cAMP or Ca²⁺ (Shaywitz and Greenberg, 1999). Although the former event is not observed in activation of chemokine receptors, the latter is, and hence investigators have explored the possibility that CREB is part of the signaling network targeted by chemokine receptor activation. Activation of CREB is observed in T cells treated with the CXC chemokine SDF-1 κ (Suzuki *et al*, 2001). The same effect has been reported in primary hippocampal neurons follow-

ing treatment with SDF-1 κ , RANTES, and fractalkine (Meucci *et al*, 1998). Interestingly, another study reported that although occupation of the calcium response element (CRE) was observed in human myeloid cells treated with MIP-1 κ , it was not by CREB, but rather by a heterodimer composed of JunD and ATF-2. This was believed to result in part from the extremely close similarity of the consensus sequence of ATF-2 (5'-T(G/T)ACGTCA-3') to that of CRE (5'-TGACGTCA-3') as well as the report that ATF-2 DNA-binding affinity could be enhanced through phosphorylation by activated JNK, an event that was observed to occur in these cells following stimulation with MIP-1 κ (Yang *et al*, 2001).

As indicated by the preceding discussion, stimulation of chemokine receptors results in the activation of diverse proteins in the cell. These proteins are not always found within the same signaling pathways, and as a result, attributing the cellular response to the activation of a particular signal transduction pathway poses quite a challenge. One way investigators have attempted to explain how and why cells might experience such effects is by looking for "crosstalk" between pathways. More specifically, it is thought that activation of proteins conventionally associated with disparate pathways might be a consequence of enzymatic proteins (e.g., kinases) that interact with multiple substrates. For example, the classic, linear pathway leading to the activation of ERK1/2 is the Ras/Raf/MEK pathway. However, phosphorylation of ERK1/2 has also been reported to be a function of activated PKC (Ueda *et al*, 1996; Grammer and Blenis, 1997; Axmann *et al*, 1998). Interestingly, ERK1/2 can also serve as upstream regulators of CREB (Sato *et al*, 1997; Cammarota *et al*, 2001) and STAT5 phosphorylation (Pircher *et al*, 1997, 1999); yet STAT5 phosphorylation is believed to be mediated directly by JAK2 (Flores-Morales *et al*, 1998; Saharinen *et al*, 2000), a protein tyrosine kinase that can apparently activate NF- κ B indirectly by promoting its dissociation from I κ B (Digicaylioglu and Lipton, 2001). Another degree of complexity is added by the recent report that NF- κ B is likely to be a downstream target of PKC (McAllister-Lucas *et al*, 2001; Ruland *et al*, 2001), an enzyme that, depending on the particular isoform, can be regulated by both the production of diacylglycerol (DAG) and/or a rise in the level of intracellular Ca²⁺ (Way *et al*, 2000). As evidenced by reports in the literature as well as within this review, all of these events occur following stimulation of chemokine receptors. Collectively then, these reports suggest that chemokines, in activating multiple signal transduction pathways, may foster the engagement of atypical effector:effector relationships such that "crosstalk" can occur in the generation of the cell's response.

Without dismissing the fact that the expression and function of proteins is a context sensitive issue, reports that certain proteins interact and regulate each other should be considered valuable information. The cell's response to extracellular stimuli is a

function of not a few but many proteins, and the identification of each of these proteins will help to develop a comprehensive explanation of how the cell utilizes different components of the intracellular machinery to achieve that response. In addition, because the controlled activity of proteins is the fundamental basis of signal propagation within the cell, individual proteins found to regulate the activity of other proteins can lead to the identification of "crosstalk" mechanisms utilized by the cell. Finally, as new findings are reported from studies linking signal to target, preexisting relationships between proteins will be referenced and evaluated for their relevance to and use in other cell systems. With respect to chemokine-mediated effects in the nervous system, this approach has been inarguably productive, for chemokine- (and cytokine)-mediated effects in the immune system have provided intriguing clues about parallel mechanisms operative in the CNS.

Conclusion

The study of chemokines and their receptors has only recently been undertaken and as a result, knowledge of the details governing their existence and function are not yet complete. However, as investigations on

chemokines continue to be reported, data from these studies will surely provide further insight about several issues that concern cell biology as whole. First, the ability of chemokines to induce directional migration of cells is an important phenomenon in normal and disease physiology, and the study of chemokines at the molecular and structural level will reveal how such a complex process can be controlled. Second, chemokines and their receptors have been reported to affect brain development, neuronal survival, and synaptic plasticity. However, the specific sequence of events and molecular mechanisms by which these effects are achieved are not yet known in their entirety. Admittedly, the data generated from further investigation of these topics will aid in the general understanding of development, apoptosis, and neuronal behavior. Third, most chemokines exhibit very low or no fidelity for a single receptor, and thus present an interesting model for how specificity can be achieved, both at the extracellular and intracellular level. With respect to the latter, the identification of signaling intermediates and transcription factors activated by chemokines should provide important information about how chemokines induce their cellular effects, and, by extension, provide leads for therapeutic agents in the treatment of diseases that are characterized by inflammation.

References

- Alkhatib G, Combadiere C, Broder CC, Feng Y, Kennedy PE, Murphy PM, Berger EA (1996). CC CKR5: RANTES, MIP-1 κ , MIP-1 κ receptor as a fusion cofactor for macrophage-tropic-HIV-1. *Science* **272**: 1955–1958.
- Andrews PW (1984). Retinoic acid induces neuronal differentiation of a cloned human embryonal carcinoma cell line *in vitro*. *Dev Biol* **103**: 285–293.
- Araujo DM, Cotman CW (1993). Trophic effects of interleukin-4, -7, and -8 on hippocampal neurons cultures: potential involvement of glial-derived factors. *Brain Res* **600**: 49–55.
- Ascensio VC, Campbell IL (1999). Chemokines in the CNS: plurifunctional mediators in diverse states. *Trends Neurosci* **22**: 504–512.
- Axmann A, Seidel D, Reimann T, Hempel U, Wenzel KW (1998). Transforming growth factor-beta1-induced activation of the Raf-MEK-MAPK signaling pathway in rat lung fibroblasts via a PKC-dependent mechanism. *Biochem Biophys Res Commun* **249**: 456–460.
- Baggiolini M (1998). Chemokines and leukocyte traffic. *Nature* **392**: 565–568.
- Baggiolini M, Dewald B, Moser B (1997). Human chemokines: an update. *Annu Rev Immunol* **15**: 675–705.
- Bais C, Santomaso B, Coso O, Arvanitakis L, Geras Raaka E, Gutkind JS, Asch AS, Cesarman E, Gershengorn E, Mesri EA (1998). G-protein coupled receptor of Kaposi's sarcoma-associated herpesvirus is a viral oncogene and angiogenesis activator. *Nature* **392**: 86–89.
- Bajetto A, Bonavia R, Barbero S, Piccioli P, Costa A, Florio T, Schettini G (1999). Glial and neuronal cells express functional chemokine receptor CXCR4 and its natural ligand stromal cell-derived factor 1. *J Neurochem* **73**: 2348–2357.
- Bajetto A, Bonavia R, Barbero S, Florio T, Schettini G (2001). Chemokines and their receptors in the central nervous system. *Front Neuroendocrinol* **22**: 147–184.
- Barnes DA, Tse J, Kaufhold M, Owen M, Hesselgesser J, Streiter R, Horuk R, Perez H (1998). Polyclonal antibody directed against human RANTES ameliorates disease in the Lewis rat adjuvant-induced arthritis model. *J Clin Invest* **101**: 2910–2919.
- Berger EA (1997). HIV entry and tropism: the chemokine receptor connection. *AIDS* **11 (Suppl A)**: S3–S16.
- Berger EA, Murphy PM, Farber JM (1999). Chemokine receptors as HIV-1 coreceptors: roles in viral entry, tropism, and disease. *Annu Rev Immunol* **17**: 657–700.
- Berson JF, Doms RW (1998). Structure-function studies of the HIV-1 coreceptors. *Semin Immunol* **10**: 237–248.
- Bolin LM, Murray R, Lukacs NW, Streiter RM, Kunkel SL, Schall TJ, Bacon KB (1998). Primary sensory neurons migrate in response to the chemokine RANTES. *J Neuroimmunol* **81**: 49–57.
- Bonnachi A, Romagnani P, Romanelli RG, Efsen E, Annunziato F, Lasagni L, Francalanci M, Serio M, Laffi G, Pinzani M, Gentilini P, Marra F (2001). Signal transduction by the chemokine receptor CXCR3. *J Biol Chem* **276**: 9945–9954.
- Boring L, Gosling J, Cleary M, Charo IF (1998). Decreased lesion formation in CCR2 O/O mice reveals a role for chemokines in the initiation of atherosclerosis. *Nature* **394**: 894–897.

- Boutet A, Salim H, Leclerc P, Tardieu M (2001). Cellular expression of functional chemokine receptor CCR5 and CXCR4 in human embryonic neurons. *Neurosci Lett* **311**: 105–108.
- Brack-Werner R (1999). Astrocytes: HIV cellular reservoirs and important participants in neuropathogenesis. *AIDS* **13**: 1–22.
- Brown Z, Robson RL, Westwick J (1996). Regulation and expression of chemokines: potential role in glomerulonephritis. *J Leukoc Biol* **59**: 75–80.
- Cammarota M, Bevilacqua LR, Dunkley PR, Rostas JA (2001). Angiotensin II promotes the phosphorylation of cyclic AMP-responsive element binding protein (CREB) at Ser133 through an ERK1/2-dependent mechanism. *J Neurochem* **79**: 1122–1128.
- Choe H, Farzan M, Sun Y, Sullivan N, Rollins B, Ponath PD, Wu L, Mackay CR, LaRosa G, Newman W, Gerard N, Gerard C, Sodroski J (1996). The κ -chemokine receptors CCR3 and CCR5 facilitate infection by HIV-1 primary isolates. *Cell* **85**: 1135–1148.
- Choi DW, Rothman SM (1990). The role of glutamate neurotoxicity in hypoxic-ischemic neuronal death. *Annu Rev Neurosci* **13**: 171–182.
- Cocchi F, De Vico AL, Garzino-Demo A, Arya SK, Gallo RC, Lusso P (1995). Identification of RANTES, MIP-1 alpha, and MIP-1 beta as the major HIV-1 suppressive factors produced by CD8+ T cells. *Science* **270**: 1811–1815.
- Colamussi ML, Secchiero P, Zella D, Curreli S, Mirandola P, Capitani S, Zauli G (2000). Stromal derived factor-1alpha induces apoptosis in activated primary CD4+ T cells. *AIDS* **14**: 748–750.
- Coughlan CM, McManus CM, Sharron M, Gao Z, Murphy D, Jaffer S, Choe W, Chen W, Hesselgesser J, Gaylord H, Kalyuzhny A, Lee VM, Wolf B, Doms RW, Kolson DL (2000). Expression of multiple functional chemokine receptors and monocyte chemoattractant protein-1 in human neurons. *Neuroscience* **97**: 591–600.
- Damaj BB, McColl SR, Neote K, Hebert CA, Naccache PH (1996). Diverging signal transduction pathways activated by interleukin 8 (IL-8) and related chemokines in human neutrophils. IL-8 and Gro-alpha differentially stimulate calcium influx through IL-8 receptors A and B. *J Biol Chem* **271**: 20540–20544.
- Del Corno M, Liu Q-H, Schols D, de Clercq E, Gessani S, Freedman BD, Collman RG (2001). HIV-1 gp120 and chemokine activation of Pyk2 and mitogen-activated protein kinases in primary macrophages mediated by calcium-dependent, pertussis toxin-insensitive chemokine receptor signaling. *Blood* **98**: 2909–2916.
- Deng HK, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhart M, Di Marzio P, Marmon S, Sutton RE, Hill CM, Davis CB, Peiper SC, Schall TJ, Littman DR, Landau NR (1996). Identification of a major HIV-1 co-receptor for primary isolates of HIV-1. *Nature* **381**: 661–666.
- Dhanasekaran N, Gutkind JS (2001). Signaling by G protein coupled receptors and G proteins: a perspective. *Oncogene* **20**: 1530–1531.
- Digicaylioglu M, Lipton SA (2001). Erythropoietin-mediated neuroprotection involves cross-talk between Jak2 and NF- κ B signalling cascades. *Nature* **412**: 641–647.
- Dikic I, Tokiwa G, Lev S, Courtneidge SA, Schlessinger J (1996). A role for Pyk2 and Src in linking G-protein-coupled receptors with MAP kinase activation. *Nature* **383**: 547–550.
- Doranz BJ, Rucker J, Yi Y, Smyth RJ, Samson M, Peiper SC, Parmentier M, Collman RG, Doms RW (1996). A dual-tropic primary HIV-1 isolate that uses fusin and the beta-chemokine receptors CKR-5, CKR-3 and CKR-2b as fusion cofactors. *Cell* **85**: 1149–1158.
- Dragic T, Litwin V, Allaway GP, Martin SR, Huang YX, Nagashima KA, Cayanan C, Maddon PJ, Koup RA, Moore JP, Paxton WA (1996). HIV-1 entry into CD4+ cells is mediated by the chemokine receptor CC-CKR-5. *Nature* **381**: 667–673.
- Feng Y, Broder CC, Kennedy PE, Berger EA (1996). HIV-entry cofactor: functional cDNA cloning of a seven-transmembrane, G-protein-coupled receptor. *Science* **272**: 872–877.
- Flores-Morales A, Pircher TJ, Silvernoinen O, Gustafsson JA, Sanchez-Gomez M, Norstedt G, Haldosen LA, Wood TJ (1998). In vitro interaction between STAT 5 and JAK 2; dependence upon phosphorylation status of STAT 5 and JAK 2. *Mol Cell Endocrinol* **138**: 1–10.
- Gabuzda D, He J, Ohagen A, Vallat A-V (1998). Chemokine receptors in HIV-1 infection of the central nervous system. *Semin Immunol* **10**: 203–213.
- Gabuzda D, Wang J (2000). Chemokine receptors and mechanisms of cell death in HIV neuropathogenesis. *J Neuro Virol* **6** (Suppl 1): S24–S32.
- Ganju RK, Brubaker SA, Meyer J, Dutt P, Yang Y, Qin S, Newman W, Groopman JE (1998). The κ -chemokine, stromal cell-derived factor-1 κ , binds to the transmembrane G-protein-coupled CXCR-4 receptor and activates multiple signal transduction pathways. *J Biol Chem* **273**: 23169–23175.
- Gelbard HA, Dzenko K, Diloreto D, Delcerro C, Delcerro M, Epstein LG (1994a). Neurotoxic effects of tumor necrosis factor in primary human neuronal cultures are mediated by activation of the AMPA receptor subtype: implications for AIDS neuropathogenesis. *Dev Neurosci* **15**: 418–422.
- Gelbard HA, Nottet HSLM, Swindells S, Jett M, Dzenko KA, Genis P, White R, Wang L, Choi Y-B, Zhang D, Lipton SA, Tourtellotte WW, Epstein LG, Gendelman HE (1994b). Platelet activating factor: a candidate HIV-1 induced neurotoxin. *J Virol* **68**: 4628–4635.
- Genis P, Jett M, Bernton EW, Boyle T, Gelbard HA, Dzenko K, Keane RW, Resnick L, Mizrachi Y, Volsky DJ, Epstein LG (1992). Cytokines and arachidonic metabolites produced during human immunodeficiency virus (HIV)-infected macrophage-astroglia interactions: implications for the neuropathogenesis of HIV disease. *J Exp Med* **176**: 1703–1718.
- Gether U (2000). Uncovering molecular mechanisms involved in activation of G-protein-coupled receptors. *Endocr Rev* **21**: 90–113.
- Gether U, Kobilka BK (1998). G-protein coupled receptors: mechanism of agonist activation. *J Biol Chem* **273**: 17979–17982.
- Gillard SE, Lu M, Mastracci RM, Miller RJ (2002). Expression of functional chemokine receptors by rat cerebellar neurons. *J Neuroimmunol* **124**: 16–28.
- Giovannelli A, Limatola C, Ragozzino D, Mileo AM, Ruggieri A, Ciotti MT, Mercanti D, Santoni A, Eusebi F (1998). CXC chemokines interleukin-8 (IL-8) and growth-related gene product alpha (GROalpha) modulate Purkinje neuron activity in mouse cerebellum. *J Neuroimmunol* **92**: 122–132.

- Giulian D, Vaca K, Noonan CA (1990). Secretion of neurotoxins by mononuclear phagocytes infected with HIV-1. *Science* **250**: 1593–1596.
- Goslin J, Slymaker S, Gu L, Tseng S, Zlot CH, Young SG, Rollins BJ, Charo IF (1999). MCP-1 deficiency reduces susceptibility to atherosclerosis in mice that overexpress human apolipoprotein B. *J Clin Invest* **103**: 773–778.
- Grammer TC, Blenis J (1997). Evidence for MEK-independent pathways regulating the prolonged activation of the ERK-MAP kinases. *Oncogene* **14**: 1635–1642.
- Griffiths-Johnson DA, Collins PD, Jose PJ, Williams TJ (1997). Animal models of asthma: role of chemokines. *Methods Enzymol* **288**: 241–266.
- Gu L, Okada Y, Clinton SK, Gerard C, Sukhova GK, Libby P, Rollins BJ (1998). Absence of monocyte chemoattractant protein-1 reduces atherosclerosis in low density lipoprotein receptor deficient mice. *Mol Cell* **2**: 275–281.
- Guerrini L, Blasi F, Denis-Donini S (1995). Synaptic activation of NF- κ B by glutamate in cerebellar granule neurons *in vitro*. *Proc Natl Acad Sci USA* **92**: 9077–9081.
- Guinamard R, Signoret N, Masamichi I, Marsh M, Krosaki T, Ravetch JV (1999). B cell antigen receptor engagement inhibits stromal cell-derived factor (SDF)-1 alpha chemotaxis and promotes protein kinase C (PKC)-induced internalization of CXCR4. *J Exp Med* **189**: 1461–1466.
- Hartley RS, Margulis M, Fishman PS, Lee VM, Tang CM (1999). Functional synapses are formed between human NTER2 (NT2N, hNT) neurons grown on astrocytes. *J Comp Neurol* **407**: 1–10.
- Hesselgesser J, Halks-Miller M, DelVecchio V, Peiper SC, Hoxie J, Kolson DL, Taub D, Horuk R (1997). CD4-independent association between HIV-1 gp120 and CXCR4: functional chemokine receptors are expressed in human neurons. *Curr Biol* **7**: 112–121.
- Hesselgesser J, Horuk R (1999). Chemokine and chemokine receptor expression in the central nervous system. *J NeuroVirol* **5**: 13–26.
- Hesselgesser J, Taub D, Baskar P, Greenberg M, Hoxie J, Kolson DL, Horuk R (1998). Neuronal apoptosis induced by HIV-1 gp120 and the chemokine SDF-1 alpha is mediated by the chemokine receptor CXCR4. *Curr Biol* **8**: 595–598.
- Horuk R, Martin AW, Wang Z, Schweitzer L, Gerassimides A, Guo H, Lu Z, Hesselgesser J, Perez HD, Kim J, Parker J, Hadley TJ, Peiper SC (1997). Expression of chemokine receptors in subsets of neurons in the central nervous system. *J Immunol* **158**: 2882–2890.
- Jazin EE, Soderstrom S, Ebendal T, Larhammar D (1997). Embryonic expression of the mRNA for the rat homologue of the fusin/CXCR-4 HIV-1 co-receptor. *J Neuroimmunol* **79**: 148–154.
- Ji TH, Grossman M, Ji I (1998). G-protein coupled receptors: diversity of receptor-ligand interactions. *J Biol Chem* **273**: 17299–17302.
- Kaiser E, Forster R, Wolf I, Ebensperger C, Kuehl WM, Lipp M (1993). The G protein-coupled receptor BLR1 is involved in murine B cell differentiation and is also expressed in neuronal tissues. *Eur J Immunol* **23**: 2532–2539.
- Kaltschmidt C, Kaltschmidt B, Neumann H, Wekerle H, Baeuerle PA (1994). Constitutive NF- κ B activity in neurons. *Mol Cell Biol* **14**: 3981–3992.
- Kampen GT, Stafford S, Adachi T, Jinquan T, Quan S, Grant JA, Skov PS, Poulsen LK, Alam R (2000). Eotaxin induces degranulation and chemotaxis of eosinophils through the activation of ERK2 and p38 mitogen-activated protein kinases. *Blood* **95**: 1911–1917.
- Katsuta T, Lim C, Shimoda K, Shibuta K, Mitra P, Banner BF, Mori M, Barnard GF (2000). Interleukin-8 and SDF1-alpha mRNA expression in colonic biopsies from patients with inflammatory bowel disease. *Am J Gastroenterol* **95**: 3157–3164.
- Kaul M, Garden G, Lipton SA (2001). Pathways to neuronal injury and apoptosis in HIV-associated dementia. *Nature* **410**: 988–994.
- Kaul M, Lipton SA (1999). Chemokines and activated macrophages in HIV gp120-induced neuronal apoptosis. *Proc Natl Acad Sci USA* **96**: 8212–8216.
- Kijowski J, Baj-Krzworzeka M, Majka M, Reza R, Marquez LA, Christofidou-Solomidou M, Janowska-Wieczorek A, Ratajczak MZ (2001). The SDF-1-CXCR4 axis stimulates VEGF secretion and activates integrins but does not affect proliferation and survival in lymphohematopoietic cells. *Stem Cells* **19**: 453–466.
- Kita H, Gleich GJ (1996). Chemokines active on eosinophils: potential roles in allergic inflammation. *J Exp Med* **183**: 2421–2426.
- Klein RS, Rubin JB, Gibson HD, DeHaan EN, Alvarez-Hernandez X, Segal RA, Luster AD (2001). SDF-1 α induces chemotaxis and enhances Sonic hedgehog-induced proliferation of cerebellar granule cells. *Development* **128**: 1971–1981.
- Klein RS, Williams KC, Alvarez-Hernandez X, Westmoreland S, Force T, Lackner AA, Luster AD (1999). Chemokine receptor expression and signaling in macaque and human fetal neurons and astrocytes: implications for the neuropathogenesis of AIDS. *J Immunol* **163**: 1636–1646.
- Koenig S, Gendelman HE, Orenstein JM, Canto MCD, Pezeshkpour GH, Yungbluth M, Janotta F, Aksamit A, Martin MA, Fauci AS (1986). Detection of AIDS virus in macrophages in brain tissue from AIDS patients with encephalopathy. *Science* **233**: 1089–1093.
- Kouba M, Vanetti M, Wang X, Schafer M, Holt V (1993). Cloning of a novel putative G-protein-coupled receptor (NLR) which is expressed in neuronal and lymphatic tissue. *FEBS Lett* **321**: 173–178.
- Laudanna C, Mochly-Rosen D, Liron T, Constantin G, Butcher EC (1998). Evidence of zeta protein kinase C involvement in polymorphonuclear neutrophil integrin-dependent adhesion and chemotaxis. *J Biol Chem* **273**: 30306–30315.
- Lavi E, Kolson DL, Ulrich AM, Fu L, Gonzalez-Scarano F (1998). Chemokine receptors in the human brain and their relationship to HIV infection. *J NeuroVirol* **4**: 301–311.
- Lavi E, Strizki JM, Ulrich AM, Zhang W, Fu L, Wang Q, O'Connor M, Hoxie JA, Gonzales-Scarano F (1997). CXCR-4 (Fusin), a co-receptor for the type 1 human immunodeficiency virus (HIV-1), is expressed in the human brain in a variety of cell types, including microglia and neurons. *Am J Pathol* **151**: 1035–1042.
- Lazarini F, Casanova P, Tham TN, De Clercq E, Arenzana-Seisdedos F, Baleux F, Dubois-Dalq M (2000). Differential signalling of the chemokine receptor CXCR4 by stromal cell-derived factor 1 and the HIV glycoprotein in rat neurons and astrocytes. *Eur J Neurosci* **12**: 117–125.
- Lee SC, Hatch WC, Liu W, Kress Y, Lyman WD, Dickson DW (1993). Productive infection of human fetal microglia by HIV-1. *Am J Pathol* **143**: 1032–1039.

- Lee VM-Y, Andrew PW (1986). Differentiation of NTERA-2 clonal human embryonal carcinoma cells into neurons involves the induction of all three neurofilament proteins. *J Neurosci* **6**: 514–521.
- Lefkowitz RJ (1998). G protein coupled receptors: new roles for receptor kinases and κ -arrestins in receptor signaling and desensitization. *J Biol Chem* **273**: 18677–18680.
- Limatola C, Ciotti MT, Mercanti D, Vacca F, Ragozzino D, Giovannelli A, Santoni A, Eusebi F, Mildedi R (2000a). The chemokine growth-related gene product beta protects rat cerebellar granule cells from apoptotic cell death through alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate receptors. *Proc Natl Acad Sci USA* **97**: 6197–6201.
- Limatola C, Giovannelli A, Maggi L, Ragozzino D, Castellani L, Ciotti MT, Mercanti D, Santoni A, Eusebi F (2000b). SDF-1 α -mediated modulation of synaptic transmission in rat cerebellum. *Eur J Neurosci* **12**: 2497–2504.
- Limatola C, Mileo AM, Giovannelli A, Vacca F, Ciotti MT, Mercanti D, Santoni A, Eusebi F (1999). The growth-related gene product beta induces sphingomyelin hydrolysis and activation of c-Jun N-terminal kinase in rat cerebellar granule neurones. *J Biol Chem* **274**: 36537–36543.
- Liu QH, Williams DA, McManus C, Baribaud F, Doms RW, Schols D, De Clercq E, Kotlikoff MI, Collman RG, Freedman BD (2000). HIV-1 gp120 and chemokines activate ion channels in primary macrophages through CCR5 and CXCR4 stimulation. *Proc Natl Acad Sci USA* **97**: 4832–4837.
- Llanes C, Collman RG, Hrin R, Kolson DL (1995). Acetylcholinesterase expression in NTERA 2 human neuronal cells: a model for developmental expression in the nervous system. *J Neurosci Res* **42**: 791–802.
- Lukacs NW, Streiter RM, Chensue SW, Kunkel SW (1996). Activation and regulation of chemokines in allergic airway inflammation. *J Leukoc Biol* **59**: 13–17.
- Luster A (1998). Chemokines-chemotactic cytokines that mediate inflammation. *N Engl J Med* **338**: 436–445.
- Ma Q, Jones D, Borghesani PR, Segal RA, Nagasawa T, Kishimoto T, Bronson RT, Springer TA (1998). Impaired B-lymphopoiesis, myelopoiesis, and derailed cerebellar neuron migration in CXCR4- and SDF-1-deficient mice. *Proc Natl Acad Sci USA* **95**: 9448–9453.
- Madani N, Kozak SL, Kavanaugh MP, Kabat D (1998). gp120 envelope glycoproteins of human immunodeficiency viruses competitively antagonize signaling by coreceptors CXCR4 and CCR5. *Proc Natl Acad Sci USA* **95**: 8005–8010.
- Mattson M, Culmsee C, Yu Z, Camandola S (2000). Roles of nuclear factor kappaB in neuronal survival and plasticity. *J Neurochem* **74**: 443–456.
- McAllister-Lucas LM, Inohara N, Lucas PC, Ruland J, Benito A, Li Q, Chen S, Chen FF, Yamaoka S, Verma IM, Mak TW, Nunez G (2001). Bimp1, a MAGUK family member linking protein kinase C activation to Bcl10-mediated NF-kappaB induction. *J Biol Chem* **276**: 30589–30597.
- McGrath KE, Koniski AD, Maltby KM, McGann JK, Palis J (1999). Embryonic expression and function of the chemokine SDF-1 and its receptor, CXCR4. *Dev Biol* **213**: 442–456.
- Mehler MF, Kessler JA (1997). Hematolymphopoietic and inflammatory cytokines in neural development. *Trends Neurosci* **20**: 357–365.
- Mellado M, Rodriguez-Frade JM, Vila-Coro AJ, Fernandez S, de Ana AM, Jones DR, Toran JL, Martinez AC (2001). Chemokine receptor homo- or heterodimerization activates distinct signaling pathways. *EMBO J* **20**: 2497–2507.
- Mennicken F, Maki R, de Souza EB, Quirion R (1999). Chemokines and chemokine receptors in the CNS: a possible role in neuroinflammation and patterning. *Trends Pharmacol Sci* **20**: 73–78.
- Meucci O, Fatatis A, Simen AA, Bushell TJ, Gray PW, Miller RJ (1998). Chemokines regulate hippocampal neuronal signaling and gp120 neurotoxicity. *Proc Natl Acad Sci USA* **95**: 8075–8080.
- Meucci O, Fatatis A, Simen AA, Miller RJ (2000). Expression of CX3CR1 chemokine receptors on neurons and their role in neuronal survival. *Proc Natl Acad Sci USA* **97**: 8075–8080.
- Miller RJ (1998). Presynaptic receptors. *Annu Rev Pharmacol Toxicol* **38**: 201–227.
- Miller RJ, Meucci O (1999). AIDS and the brain: is there a chemokine connection? *Trends Neurosci* **22**: 471–479.
- Murphy PM (1994). The molecular biology of leukocyte chemoattractant receptors. *Annu Rev Immunol* **12**: 593–633.
- Murphy PM (1996). Chemokine receptors: structure, function and role in microbial pathogenesis. *Cytokine Growth Factor Rev* **7**: 47–64.
- Murphy PM, Baggiolini M, Charo IF, Herbert CA, Horuk R, Matsushima K, Miller LH, Oppenheim JJ (2000). International union of pharmacology. XXII. Nomenclature for chemokine receptors. *Pharmacol Rev* **52**: 145–176.
- Nuovo GJ, Gallery F, MacConnell P, Braun A (1994). In situ detection of polymerase chain reaction-amplified HIV-1 nucleic acids and tumor necrosis factor- κ RNA in the central nervous system. *Am J Pathol* **144**: 659–666.
- Oberlin E, Amara A, Bachelier F, Bessia C, Virelizier JL, Arenzara-Seisdedos F, Schwartz O, Heard JM, Clark-Lewis I, Legler DF, Loetscher M, Baggiolini M, Moser B (1996). The CXC chemokine SDF-1 is the ligand for LESTR/fusin and prevents infection by T-cell-line-adapted HIV-1. *Nature* **382**: 833–835.
- Oh SB, Endoh T, Simen AA, Ren D, Miller RJ (2002). Regulation of calcium currents by chemokines and their receptors. *J Neuroimmunol* **123**: 66–75.
- Oh SB, Tran PB, Gillard SE, Hurley RW, Hammond DL, Miller RJ (2001). Chemokines and glycoprotein120 produce pain hypersensitivity by directly exciting primary nociceptive neurons. *J Neurosci* **21**: 5027–5035.
- O'Neill LAJ, Kaltschmidt C (1997). NF- κ B: a crucial transcription factor for glial and neuronal cell function. *Trends Neurosci* **20**: 252–258.
- Pandey V, Bolsover SR (2000). Immediate and neurotoxic effects of HIV protein gp120 act through CXCR4 receptor. *Biochem Biophys Res Commun* **274**: 212–215.
- Pati S, Cavrois M, Guo HG, Foulke J, Kim J, Feldman RA, Reitz M (2001). Activation of NF-kappaB by the human herpesvirus 8 chemokine receptor ORF74: evidence for a paracrine model of Kaposi's sarcoma pathogenesis. *J Virol* **18**: 8660–8673.
- Pircher TJ, Flores-Morales A, Mui AL, Saltiel AR, Norstedt G, Gustafsson JA, Haldosen LA (1997). Mitogen-activated protein kinase inhibition decreases growth hormone stimulated transcription mediated by STAT5. *Mol Cell Endocrinol* **133**: 169–176.

- Pircher TJ, Petersen H, Gustafsson JA, Haldosen LA (1999). Extracellular signal-regulated kinase (ERK) interacts with signal transducer and activator of transcription (STAT)5a. *Mol Endocrinol* **13**: 555–565.
- Pleasure SJ, Lee VM (1993). NTera 2 cells: a human cell line which displays characteristics expected of a human committed neuronal progenitor cell. *J Neurosci Res* **90**: 2174–2178.
- Pleasure SJ, Page C, Lee VM-Y (1992). Pure, postmitotic polarized human neurons derived from NTera 2 cells provide a system for expressing exogenous proteins in terminally differentiated neurons. *J Neurosci* **12**: 1802–1815.
- Premack BA, Schall TJ (1996). Chemokine receptors: gateways to inflammation and infection. *Nat Med* **2**: 1174–1178.
- Pulliam L, Herndier BG, Tang NM, McGrath MS (1991). HIV-1 infected macrophages produce soluble factors that cause histological and neurochemical alterations in cultured human brains. *J Clin Invest* **87**: 503–512.
- Puma C, Danick M, Quirion R, Ramon F, Williams S (2001). The chemokine interleukin-8 acutely reduces Ca²⁺ currents in identified cholinergic septal neurons expressing CXCR1 and CXCR2 receptor mRNAs. *J Neurochem* **78**: 960–971.
- Quesenberry PJ, Hulspar R, Joly M, Benoit B, Engstrom C, Rielly J, Savarese T, Pang L, Recht L, Ross A, Stein G, Stewart M (1999). Correlates between hematopoiesis and neurogenesis: neural stem cells. *J Neurotrauma* **16**: 661–666.
- Rodriguez-Frade JM, Mellado M, Martinez AC (2001). Chemokine receptor dimerization: two are better than one. *Trends Immunol* **22**: 612–617.
- Rodriguez-Frade JM, Vila-Coro AJ, De Ana AM, Albar AJ, Martinez AC, Mellado M (1999a). The chemokine monocyte chemoattractant protein-1 induces functional responses through dimerization of its receptor CCR2. *Proc Natl Acad Sci USA* **96**: 3628–3633.
- Rodriguez-Frade JM, Vila-Coro AJ, Martin A, Nieto M, Sanchez-Madrid F, Proudfoot AE, Wells TN, Martinez AC, Mellado M (1999b). Similarities and differences in RANTES- and (AOP)-RANTES-triggered signals: implications for chemotaxis. *J Cell Biol* **144**: 755–765.
- Rollins BJ (1997). Chemokines. *Blood* **90**: 909–928.
- Rottman J, Ganley KP, Williams K, Wu L, Mackay CR, Ringler DJ (1997). Cellular localization of the chemokine receptor CCR5. *Am J Pathol* **151**: 1341–1351.
- Rucker J, Edinger AL, Sharron M, Samson M, Lee B, Berson JF, Yi Y, Margulies B, Collman RG, Doranz BJ, Parmentier M, Doms RW (1997). Utilization of chemokine receptors, orphan receptors, and herpesvirus-encoded receptors by diverse human and simian immunodeficiency virus. *J Virol* **71**: 8999–9007.
- Ruland J, Dunca GS, Eila A, del Barco Barrantes I, Nguyen L, Plyte S, Millar DG, Bouchard D, Wakeham A, Ohashi PS, Mak TW (2001). Bcl10 is a positive regulator of antigen receptor-induced activation of NF-kappaB and neural tube closure. *Cell* **104**: 33–42.
- Sabri F, Tresoldi E, Di Stefano M, Polo S, Monaco MC, Verani A, Fiore JR, Lusso P, Major E, Chiodi F, Scarlatti G (1999). Nonproductive human immunodeficiency virus type 1 infection of human fetal astrocytes: independence from CD4 and major chemokine receptors. *Virology* **264**: 370–384.
- Saharinen P, Takaluoma K, Silvennoinen O (2000). Regulation of the Jak2 tyrosine kinase by its pseudokinase domain. *Mol Cell Biol* **20**: 3387–3395.
- Saito M, Sato R, Hisatome I, Narahashi T (1996). RANTES and platelet-activating factor open Ca²⁺-activated K⁺ channels in eosinophils. *FASEB J* **10**: 792–798.
- Saito Y, Sharer LR, Epstein LG, Michaels J, Mintz M, Louder M, Golding K, Cvetkovich TA, Blumberg BM (1994). Overexpression of nef as a marker for restricted HIV-1 infection of astrocytes in postmortem pediatric central nervous system. *Neurology* **44**: 474–481.
- Sanders VJ, Pittman CA, White MG, Wang G, Wiley CA, Achim CL (1998). Chemokines and receptors in HIV encephalitis. *AIDS* **12**: 1021–1026.
- Saporta S, Willing AE, Colina LO, Zigova T, Milliken M, Daadi MM, Sanberg PR (2000). In vitro and in vivo characterization of hNT neuron neurotransmitter phenotypes. *Brain Res Bull* **53**: 263–268.
- Sato K, Kawasaki H, Nagayama H, Enomoto M, Morimoto C, Tadokoro K, Juji T, Takahashi TA (2001). Signaling events following chemokine receptor ligation in human dendritic cells at different developmental stages. *Int Immunol* **13**: 167–179.
- Sato N, Kamino K, Tateishi K, Satoh T, Nishiwaki Y, Yoshiwa A, Miki T, Ogihara T (1997). Elevated amyloid beta protein(1–40) level induces CREB phosphorylation at serine-133 via p44/42 MAP kinase (Erk1/2)-dependent pathway in rat pheochromocytoma PC12 cells. *Biochem Biophys Res Commun* **232**: 637–642.
- Schorr W, Swandulla D, Zeilhofer HU (1999). Mechanisms of IL-8-induced Ca²⁺ signaling in human neutrophil granulocytes. *Eur J Immunol* **29**: 897–904.
- Schutze S, Potthoff K, Machleidt T, Berkovic D, Wiegmann K, Kronke M (1992). TNF activates NF-kB by phosphatidylcholine-specific phospholipase C-induced “acidic” sphingomyelin breakdown. *Cell* **71**: 765–776.
- Seeger S, Regele H, Mack M, Kain R, Cartron JP, Colin Y, Kerjaschki D, Schlondorff D (2000). The Duffy antigen receptor for chemokines is up-regulated during acute renal transplant rejection and crescentic glomerulonephritis. *Kidney Int* **58**: 1546–1556.
- Shaywitz AJ, Greenberg ME (1999). CREB: a stimulus induced transcription factor induced by a diverse array of extracellular signals. *Annu Rev Biochem* **68**: 821–861.
- Suzuki Y, Rahman M, Mitsuya H (2001). Diverse transcriptional response of CD4(+) T cells to stromal cell-derived factor (SDF)-1: cell survival promotion and priming effects of SDF-1 on CD4(+) T cells. *J Immunol* **167**: 3064–3073.
- Takahashi K, Wesselingh SL, Griffin DE, McArthur JE, Johnson RT, Glass JD (1996). Localization of HIV-1 in the brain using polymerase chain reaction/in situ hybridization and immunocytochemistry. *Ann Neurol* **39**: 705–711.
- Terskikh AV, Easterday MC, Li L, Hood L, Kornblum HI, Geschwind DH, Weissman IL (2001). From hematopoiesis to neurogenesis: evidence of overlapping genetic programs. *Proc Natl Acad Sci USA* **98**: 7934–7939.
- Ueda Y, Hirai S, Osada S, Suzuki A, Mizuno K, Ohno S (1996). Protein kinase C activates the MEK-ERK pathway in a manner independent of Ras and dependent on Raf. *J Biol Chem* **271**: 23512–23519.
- Vallat AV, De Girolami U, He J, Mhashilkar A, Marasco W, Shi B, Gray F, Bell J, Keohane C, Smith TW, Gabuzda D

- (1998). Localization of HIV-1 co-receptors CCR5 and CXCR4 in the brain of children with AIDS. *Am J Pathol* **152**: 167–178.
- Venkatakrishnan G, Salgia R, Groopman JE (2000). Chemokine receptors CXCR-1/2 activate mitogen activated protein kinase via the epidermal growth factor receptor in ovarian cancer cells. *J Biol Chem* **275**: 6868–6875.
- Vila-Coro AJ, Rodriguez-Frade JM, De Ana AM, Moreno-Ortiz MC, Martinez AC, Mellado M (1999). The chemokine SDF-1 α triggers CXCR4 receptor dimerization and activates the JAK/STAT pathway. *FASEB J* **13**: 1699–1710.
- Wang J, Crawford K, Menglan Y, Wang H, Gorry PR, Gabuzda D (2002). Regulation of CC chemokine receptor 5 and CD4 expression and human immunodeficiency virus type 1 replication in human macrophages and microglia by T-helper type 2 cytokines. *J Infect Dis* **185**: 885–897.
- Ward SG, Westwick J (1998). Chemokines: understanding their role in T-lymphocyte biology. *Biochem J* **333**: 457–470.
- Watkins BA, Dorn HH, Kelly WB, Armstrong RC, Potts BJ, Michaels F, Kufta CV, Dubois-Dalcq M (1990). Specific tropism of HIV-1 for microglial cells in primary human brain cultures. *Science* **249**: 549–553.
- Way KJ, Chou E, King GL (2000). Identification of PKC-isoform-specific biological actions using pharmacological approaches. *Trends Pharmacol Sci* **21**: 181–187.
- Westmoreland SV, Alvarez X, deBakker C, Aye P, Wilson ML, Williams KC, Lackner AA (2002). Developmental expression patterns of CCR5 and CXCR4 in the rhesus macaque brain. *J Neuroimmunology* **122**: 146–158.
- Westmoreland SV, Rottman JB, Williams KC, Lackner AA, Sasseville VG (1998). Chemokine receptor expression on resident and inflammatory cells in the brain of macaques with simian immunodeficiency virus encephalitis. *Am J Pathol* **152**: 659–665.
- Wiley CA, Schrier RD, Nelson JA, Lampert PW, Oldstone MBA (1986). Cellular localization of human immunodeficiency virus within the brains of acquired immunodeficiency patients. *Proc Natl Acad Sci USA* **83**: 7089–7093.
- Wong M, Fish E (1998). RANTES and MIP-1 κ activate STAT's in T cells. *J Biol Chem* **273**: 309–314.
- Wong M, Uddin S, Majchrzak B, Huynh T, Proudfoot AEI, Plataniotis LC, Fish EN (2001). RANTES activates Jak2 and Jak3 to regulate engagement of multiple signaling pathways in T cells. *J Biol Chem* **276**: 11427–11431.
- Wong ML, Xin WW, Dumas RS (1996). Rat LCR1: cloning and cellular distribution of a putative chemokine receptor in brain. *Mol Psychiatr* **1**: 133–140.
- Xia M, Bacskai BJ, Knowles RB, Qin SX, Hyman BT (2000). Expression of the chemokine receptor CXCR3 on neurons and the elevated expression of its ligand IP-10 in reactive astrocytes: *in vitro* ERK1/2 activation and role in Alzheimer's disease. *J Neuroimmunol* **108**: 227–235.
- Xia M, Hyman BT (2002). Gro α /KC, a chemokine receptor CXCR2 ligand, can be a potent trigger for neuronal ERK 1/2 and PI-3K pathways and for tau hyperphosphorylation—a role in Alzheimer's disease? *J Neuroimmunol* **122**: 55–64.
- Yang Y-M, Hatch WC, Liu Z-Y, Du B, Groopman JE (2001). κ -Chemokine induction of activation protein-1 and cyclic AMP responsive element activation in human myeloid cells. *Cell Growth Differ* **12**: 211–221.
- Ye RD (2001). Regulation of nuclear factor κ B activation by G-protein coupled receptors. *J Leukoc Biol* **70**: 839–848.
- Zhang L, He T, Talal A, Wang G, Frankel SS, Ho DD (1998). In vivo distribution of the human immunodeficiency virus/simian immunodeficiency virus coreceptors: CXCR4, CCR3, and CCR5. *J Virol* **72**: 5035–5045.
- Zheng J, Ghorpade A, Niemann D, Cotter RL, Thylin MR, Epstein L, Swartz JM, Shepard RB, Liu X, Nukuna A, Gendelman HE (1999a). Lymphotropic virions affect chemokine receptor-mediated neural signaling and apoptosis: implications for human immunodeficiency virus type 1-associated dementia. *J Virol* **73**: 8256–8267.
- Zheng J, Thylin MR, Ghorpade A, Xiong H, Persidsky Y, Cotter R, Niemann D, Che M, Zheng Y-C, Gelbard HA, Shepard RB, Swartz JM, Gendelman HE (1999b). Intracellular CXCR4 signaling, neuronal apoptosis and neuropathogenic mechanisms of HIV-1-associated dementia. *J Neuroimmunol* **98**: 185–200.
- Zlotnik A, Yoshie O (2000). Chemokines: a new classification system and their role in immunity. *Immunity* **12**: 121–127.
- Zou YR, Kottman AH, Kuroda M, Taniuchi I, Littman DR (1998). Function of the chemokine receptor CXCR4 in haematopoiesis and in cerebellar development. *Nature* **393**: 595–599.