

Short Communication

NKG2D contributes to efficient clearance of picornavirus from the acutely infected murine brain

Chandra Deb¹ and Charles L Howe^{1,2,3,4,5}

Departments of ¹Neurology, ²Neuroscience, and ³Immunology, ⁴The Molecular Neuroscience Program, and ⁵The Translational Immunovirology and Biodefense Program, Mayo Clinic College of Medicine, Rochester, Minnesota, USA

Activated murine cytotoxic T cells express the NKG2D natural cytotoxicity receptor. This receptor recognizes major histocompatibility complex (MHC) class I-like molecules expressed on the surface of infected cells and serves to augment T cell-mediated cytotoxicity. The role of NKG2D-mediated augmentation in the clearance of central nervous system viral infections has not been explored. Using the Theiler's murine encephalomyelitis virus model, the authors found that NKG2D-positive CD8+ cytotoxic T cells enter the brain, that NKG2D ligands are expressed in the brain during acute infection, and that interruption of NKG2D ligand recognition via treatment with a function-blocking antibody attenuates the efficacy of viral clearance from the central nervous system. *Journal of NeuroVirology* (2008) 14, 261–266.

Keywords: cytotoxic T cell; enterovirus; H60; Mult1; neurovirulence; Rae1; TMEV

Members of the picornavirus family of small, nonenveloped, positive-stranded RNA viruses, which includes enterovirus 71, poliovirus, hepatitis A virus, the coxsackieviruses, and the rhinoviruses, are a frequent cause of infection worldwide. Despite the fact that members of this family infect more humans than any other group of viruses (Rotbart, 2002), host factors that control picornaviral replication are poorly understood. Even less is known about host factors that control picornaviral neurovirulence and clearance of picornaviruses from the central nervous system (CNS) (Buenz and Howe, 2006). We and others have used the Theiler's murine encephalomyelitis virus (TMEV) as a mouse model of picornavirus infection of the CNS (Buenz *et al*, 2006; Jin *et al*, 2007; Rubio *et al*, 2006; Tsunoda

et al, 2006). Previous studies have extensively characterized the role of antiviral CD8+ cytotoxic T lymphocytes (CTLs) in the clearance of TMEV from the brain during acute infection (Mendez-Fernandez *et al*, 2003), and it is clear in mice of the H-2^b major histocompatibility complex (MHC) class I haplotype that CTLs specific for the VP2_{121–130} TMEV peptide are responsible for viral clearance during the first 2 weeks of CNS infection (Myoung *et al*, 2007). However, the role of additional host factors such as costimulatory molecules has not been thoroughly explored.

NKG2D is a natural cytotoxicity receptor expressed on natural killer cells (NKC) and on activated CTLs in mice (Eagle and Trowsdale, 2007). This receptor recognizes stress- and infection-regulated MHC class I-like molecules such as Rae1, Mult1, and H60 expressed on the surface of murine cells (Raulet, 2003). Although NKG2D is a direct mediator of cytotoxicity for NKCs, it serves as a costimulatory receptor for CTLs and augments cytotoxicity downstream from T-cell receptor recognition of peptides presented on MHC class I (Markiewicz *et al*, 2005). Because the mechanisms that direct CTL-mediated clearance of infected neural cells are unclear, we asked whether NKG2D-positive immune effectors are present in the CNS during acute TMEV infection, whether NKG2D ligands are expressed in the brain following infection,

Address correspondence to Charles L. Howe, Department of Neurology, Guggenheim 442-C, Mayo Clinic College of Medicine, 200 First Street SW, Rochester, MN 55905, USA. E-mail: howec@mayo.edu.

The authors are grateful to Reghann LaFrance-Corey for expert technical assistance and to Nikilyn Kinzel for conceptual, textual, and experimental guidance. This work was supported by a grant from the National Multiple Sclerosis Society (RG3636), by an early career development award from the Mayo Clinic (CLH), and by Donald and Frances Herdrich.

Received 9 January 2008; revised 14 March 2008; accepted 21 March 2008.

and whether interruption of NKG2D-mediated costimulation of CTLs alters viral clearance from the CNS.

Using flow-cytometric analysis of a previously characterized preparation of brain-infiltrating leukocytes (BILs) (Howe *et al.*, 2007), we measured the number of NKG2D-positive immune cells present in

the CNS at 3 and 7 days after TMEV infection (2×10^5 plaque-forming units [PFU] intracerebrally [i.c.]; 4- to 6-week-old C57Bl/6J mice; all experiments adhered to Mayo Institutional Animal Care and Use Committee [IACUC] guidelines). We found that at 3 days postinfection (d.p.i.), $6\% \pm 0.4\%$ of $CD45^{hi}$ BILs were CD8-positive, of which $20\% \pm 0.6\%$ were also




Figure 1 NKG2D-positive antiviral $CD8^+$ T cells infiltrate the brain during acute infection with TMEV. Brain-infiltrating leukocytes (BILs) were isolated from mice at 3 (A–C) and 7 (D–F) days postinfection (dpi) and analyzed by flow cytometry. The number of $CD45^{hi}CD8^+$ cells increased between 3 (A) and 7 (D) dpi. Likewise, the number of $CD45^{hi}$ cells positive for both CD8 and NKG2D increased from 20% at 3 dpi (B) to 76% at 7 dpi (E). Moreover, the intensity of NKG2D surface labeling increased from a mean fluorescence intensity (MFI) of 35 at 3 dpi (shaded histogram in C) to 89 at 7 dpi (shaded histogram in F). The MFI for the isotype control was 4 at 3 dpi (open histogram in C) and 5 at 7 dpi (open histogram in F). BILs were further analyzed for antiviral specificity by costaining with an MHC class I tetramer loaded with the VP2_{121–130} H-2D^b-specific viral peptide (G–I). Of the cells that were both $CD8^+$ and tetramer-positive (box “H” in G), 81% were also NKG2D-positive, with an MFI of 136 (H). In contrast, only 12% of cells that were $CD8^+$ but tetramer-negative (box “I” in G) were also NKG2D-positive, with an MFI of only 18 (I).

positive for the NKG2D costimulatory receptor. By 7 d.p.i., the BILs were comprised of $24\% \pm 2\%$ CTLs, with $76\% \pm 2\%$ of these cells positive for NKG2D ($n = 4-6$ mice per time point; mean \pm 95% CI). Thus, between 3 and 7 d.p.i., the phenotype of immune cells infiltrating the CNS shifted toward a population that was enriched in NKG2D⁺CD8⁺CD45^{hi} CTLs (Figure 1A to F). The presence of NKG2D-positive CTLs within the infected hippocampus (Wada and Fujinami, 1993) was confirmed by simultaneously immunostaining cryosections from mice at 7 d.p.i. with anti-CD8 (53-6.7; 1:100) and anti-NKG2D (CX5; 1:50) (Figure 2A to E) (Howe *et al*, 2004).

In order to determine the viral specificity of the NKG2D-positive CTLs at 7 d.p.i., we costained BILs with anti-CD8, anti-NKG2D, and the TMEV VP2₁₂₁₋₁₃₀-specific tetramer (Beckman Coulter Immunomics, San Diego, CA; 1:100) (Howe *et al*, 2007). Flow cytometry revealed that $74\% \pm 2\%$ of CD45^{hi}CD8⁺ cells were VP2-specific (Figure 1G), whereas none of these cells were stained by an irrelevant D^b/E7 tetramer (data not shown) (Mendez-Fernandez *et al*, 2005). Moreover, $81\% \pm 2\%$ of the VP2-specific CTLs were also NKG2D-positive (Figure 1H), whereas only $12\% \pm 2\%$ of the VP2 tetramer-negative CTLs were positive for NKG2D (Figure 1I) ($n = 4$ mice; mean \pm 95% CI). We conclude that a robust population of NKG2D-positive VP2-specific

CTLs are present within the brain at 7 days after infection with TMEV.

Others have reported the up-regulation of NKG2D ligands following infection in a variety of peripheral tissues (Eagle and Trowsdale, 2007). However, the up-regulation of these ligands by picornavirus infection has not been previously explored. We found that Rae1, Mult1, and H60 were all up-regulated within the hippocampus during acute TMEV infection, as determined by reverse transcriptase polymerase chain reaction (RT-PCR) analysis of RNA isolated from the excised hippocampus (see Table 1 for conditions). Of the three ligands we analyzed, Rae1 showed the most robust up-regulation, with a peak induction of 10 ± 1 -fold over the uninfected hippocampus at 7 d.p.i. ($F(4, 9) = 4.845$, $P = .017$ by one-way analysis of variance [ANOVA]; $q(9, 5) = 5.841$, $P = .012$ with the SNK pairwise comparison between 0 and 7 d.p.i.; mean \pm 95% CI) (Figure 3A). H60 and Mult1 were also up-regulated 2 ± 0.1 -fold and 2 ± 0.2 -fold over uninfected, respectively, at 7 d.p.i. ($n = 3$ mice; mean \pm 95% CI) (Figure 3A).

In order to identify the cellular locus of NKG2D ligand expression in the hippocampus, we attempted, unsuccessfully, to immunostain fresh frozen or paraffin-embedded sections of hippocampus after fixation with paraformaldehyde, methanol, or ethanol:acetone. We tested specific anti-Rae1 (clone




Figure 2 Hippocampal sections were stained with anti-NKG2D (A) and anti-CD8 (B). Colocalization (C–E) confirmed the presence of CD8⁺ (green) and NKG2D-positive (red) cells in the hippocampus at 7 dpi. DAPI is shown in blue in (C). Arrowheads in (D) indicate cells that are double-positive for CD8 and NKG2D. A representative double-positive cell is shown in (E). Scale bar in C is 200 microns and refers to A–C; scale bar in D is 100 microns; scale bar in E is 5 microns.

Table 1 RT-PCR conditions for NKG2D ligands

Gene	Forward primer	Reverse primer	UPL probe
Rae-1 γ	ATACACCAACGGGCTGGAT	CTTCGCTTCATACCAGAGAGG	cccagcag
H-60	ACAGCATAGCATCTACTTTTATCCAC	TCCATGGCACTGCTGTTATC	cctggaga
Mult-1	AGCTCATGTTGCACTGGAAA	TCATCAAGGTACTGAAAAGATCCTG	tctggagc
GAPDH	AGCTTGTTCATCAACGGGAAG	TTTGATGTTAGTGGGGTCTCG	catcacca

Note. Excised hippocampus was processed for RNA purification using the Qiagen RNeasy Lipid-Tissue kit. cDNA was prepared using the Roche 1st Strand cDNA Synthesis kit and analyzed using the Roche Universal Probe Library taqman probes identified above and the Roche TaqMan Master kit. Samples were amplified for 45 cycles using 95°C melt for 10 s, 55°C anneal for 30 s, and 72°C elongate for 5 s. Primers are given as 5' to 3'.

186107; 1:25; R&D Systems) and anti-H60 (clone 205326; 1:100; R&D Systems) antibodies as well as a pan-specific NKG2D ligand antibody (goat anti-Rae1; 1:100; Santa Cruz Biotechnology) and the pan-specific NKG2D-Fc chimeric molecule (R&D Systems

139-NK; 1:10). Likewise, we attempted to confirm the expression of the NKG2D ligands at the protein level within the hippocampus by Western blot analysis of hippocampal lysates. Of the numerous antibodies tested (same as immunostaining, above), only the anti-H60 antibody (clone 205326; 1:100; R&D Systems) successfully identified a prominent band that was present within the hippocampus only after infection and which was also present in lysate from the YAC1 NK cell target line (Figure 3B). The apparent molecular weight of this prominent band (ca. 50 kDa) matches that described by Hasan and colleagues (Hasan *et al*, 2005). The relative difference in mobility between the putative H60 band in the hippocampal lysate and the YAC-1 cell lysate may reflect differential expression of one of the three recently identified isoforms of H60 that share high sequence homology but differ in molecular weight (Takada *et al*, 2008). We conclude that NKG2D ligands such as H60 are up-regulated at both the transcriptional and translational level within the hippocampus following acute TMEV infection, but we are unable to identify the cells that express these ligands with the currently available immunologic tools.




Figure 3 NKG2D ligands are upregulated in the hippocampus during acute infection with TMEV and interference with NKG2D recognition of these ligands increases viral load in the CNS. qRT-PCR analysis of brain mRNA revealed a 10-fold upregulation of Rae1 at 7 dpi (A) as compared to the uninfected brain. Likewise, H60 and Mult1 were also upregulated throughout the acute phase of infection (A). Western blot analysis of hippocampal protein lysates showed the translational upregulation of H60, with peak expression at 7 dpi (B). The 50 kDa band (arrow) upregulated in hippocampus was also present in YAC-1 cell lysates (B). Treatment with the CX5 anti-NKG2D function blocking antibody throughout the acute phase of infection resulted in a 10-fold increase in viral load within the brain at 7 dpi (C). Bar graph shows mean \pm 95% confidence interval. Five mice were treated in each group and individual viral titers are shown as black circles in (C). All CX5-treated mice showed increased viral titer.

Finally, we asked whether interruption of NKG2D-mediated recognition of NKG2D ligands was sufficient to alter viral load within the hippocampus. We used the CX5 function blocking antibody following a therapeutic regime adapted from Lanier and colleagues (100 μ g/mouse intraperitoneally [i.p.] on -1, +1, and +4 d.p.i.; kindly provided by Dr. Lanier, University of California San Francisco) (Ogasawara *et al*, 2003). Control mice received an equivalent dose of polyclonal rat immunoglobulin G (IgG) (Kim *et al*, 2007). Plaque assay (Pavelko *et al*, 1998) of freshly excised brain homogenates at 7 d.p.i. revealed that CX5 treatment led to 9 ± 2 times more infectious virus as compared to control IgG-treated mice ($U(5, 5) = 15$, $P = .008$, Mann-Whitney rank sum test; mean \pm 95% CI) (Figure 3C). Based on these observations, we propose that VP2-specific CTLs expressing NKG2D recognize NKG2D ligands such as Rae1 and H60 expressed on infected cells within the brain and thereby engage an enhanced cytotoxic response (Markiewicz *et al*, 2005) that contributes to the clearance of picornaviruses from the CNS.

In addition to expression on activated CTLs, NKG2D is also a primary cytotoxicity receptor for NKC and NKC activity is certainly modulated by viral infection. We chose to specifically focus on the role of NKG2D on antiviral CTLs because the number of NKC present within the brain during the peak of viral clearance is low compared to CTLs (ca. 10% of CD45^{hi} cells are NK1.1-positive at 7 d.p.i.; data not shown). However, we certainly cannot rule out a role for NKC-mediated recognition of NKG2D ligands on infected neural cells as part of an early response required for viral clearance. Likewise, NKG2D is expressed by subpopulations of CD4 T cells (Saez-Borderias *et al*, 2006), $\gamma\delta$ T cells (Dandekar *et al*,

2005), and macrophages (Baba *et al*, 2006), suggesting that multiple cell types may be inhibited by CX5 treatment. Regardless, our observations provide evidence that NKG2D-mediated recognition of virus-infected targets by CTLs participates in the clearance of picornaviruses from the CNS. These findings extend previous studies regarding the immunovirology of NKG2D ligand expression, which have largely focused on viral evasion of immune surveillance (Lodoen *et al*, 2003, 2004; Vilarinho *et al*, 2007), and add to the potential repertoire of therapeutic strategies for reducing and controlling CNS picornaviral infections (Buenz and Howe, 2006; Buenz *et al*, 2006).

References

- Baba T, Ishizu A, Iwasaki S, Suzuki A, Tomaru U, Ikeda H, Yoshiki T, Kasahara M (2006). CD4⁺/CD8⁺ macrophages infiltrating at inflammatory sites: a population of monocytes/macrophages with a cytotoxic phenotype. *Blood* **107**: 2004–2012.
- Buenz EJ, Howe CL (2006). Picornaviruses and cell death. *Trends Microbiol* **14**: 28–36.
- Buenz EJ, Rodriguez M, Howe CL (2006). Disrupted spatial memory is a consequence of picornavirus infection. *Neurobiol Dis* **24**: 266–273.
- Dandekar AA, O'Malley K, Perlman S (2005). Important roles for gamma interferon and NKG2D in gammadelta T-cell-induced demyelination in T-cell receptor beta-deficient mice infected with a coronavirus. *J Virol* **79**: 9388–9396.
- Eagle RA, Trowsdale J (2007). Promiscuity and the single receptor: NKG2D. *Nat Rev Immunol* **7**: 737–744.
- Hasan M, Krmpotic A, Ruzsics Z, Bubic I, Lenac T, Halenius A, Loewendorf A, Messerle M, Hengel H, Jonjic S, Koszinowski UH (2005). Selective down-regulation of the NKG2D ligand H60 by mouse cytomegalovirus m155 glycoprotein. *J Virol* **79**: 2920–2930.
- Howe CL, Bieber AJ, Warrington AE, Pease LR, Rodriguez M (2004). Antiapoptotic signaling by a remyelination-promoting human antimyelin antibody. *Neurobiol Dis* **15**: 120–131.
- Howe CL, Ure D, Adelson JD, LaFrance-Corey R, Johnson A, Rodriguez M (2007). CD8⁺ T cells directed against a viral peptide contribute to loss of motor function by disrupting axonal transport in a viral model of fulminant demyelination. *J Neuroimmunol* **188**: 13–21.
- Jin YH, Mohindru M, Kang MH, Fuller AC, Kang B, Gallo D, Kim BS (2007). Differential virus replication, cytokine production, and antigen-presenting function by microglia from susceptible and resistant mice infected with Theiler's virus. *J Virol* **81**: 11690–11702.
- Kim J, Chang CK, Hayden T, Liu FC, Benjamin J, Hamerman JA, Lanier LL, Kang SM (2007). The activating immunoreceptor NKG2D and its ligands are involved in allograft transplant rejection. *J Immunol* **179**: 6416–6420.
- Lodoen M, Ogasawara K, Hamerman JA, Arase H, Houchins JP, Mocarski ES, Lanier LL (2003). NKG2D-mediated natural killer cell protection against cytomegalovirus is impaired by viral gp40 modulation of retinoic acid early inducible 1 gene molecules. *J Exp Med* **197**: 1245–1253.
- Lodoen MB, Abenes G, Umamoto S, Houchins JP, Liu F, Lanier LL (2004). The cytomegalovirus m155 gene product subverts natural killer cell antiviral protection by disruption of H60-NKG2D interactions. *J Exp Med* **200**: 1075–1081.
- Markiewicz MA, Carayannopoulos LN, Naidenko OV, Matsui K, Burack WR, Wise EL, Fremont DH, Allen PM, Yokoyama WM, Colonna M, Shaw AS (2005). Costimulation through NKG2D enhances murine CD8⁺ CTL function: similarities and differences between NKG2D and CD28 costimulation. *J Immunol* **175**: 2825–2833.
- Mendez-Fernandez YV, Hansen MJ, Rodriguez M, Pease LR (2005). Anatomical and cellular requirements for the activation and migration of virus-specific CD8⁺ T cells to the brain during Theiler's virus infection. *J Virol* **79**: 3063–3070.
- Mendez-Fernandez YV, Johnson AJ, Rodriguez M, Pease LR (2003). Clearance of Theiler's virus infection depends on the ability to generate a CD8⁺ T cell response against a single immunodominant viral peptide. *Eur J Immunol* **33**: 2501–2510.
- Myoung J, Hou W, Kang B, Lyman MA, Kang JA, Kim BS (2007). The immunodominant CD8⁺ T cell epitope region of Theiler's virus in resistant C57BL/6 mice is critical for anti-viral immune responses, viral persistence, and binding to the host cells. *Virology* **360**: 159–171.
- Ogasawara K, Hamerman JA, Hsin H, Chikuma S, Bour-Jordan H, Chen T, Pertel T, Carnaud C, Bluestone JA, Lanier LL (2003). Impairment of NK cell function by NKG2D modulation in NOD mice. *Immunity* **18**: 41–51.
- Pavelko KD, van Engelen BG, Rodriguez M (1998). Acceleration in the rate of CNS remyelination in lysolecithin-induced demyelination. *J Neurosci* **18**: 2498–2505.
- Raulet DH (2003). Roles of the NKG2D immunoreceptor and its ligands. *Nat Rev Immunol* **3**: 781–790.
- Rotbart HA (2002). Clinical significance, diagnosis, and treatment of picornavirus infections. In: *Molecular biology of Picornaviruses*. Semler BL, Wimmer E (eds). Washington, DC: ASM Press, pp 357–365.
- Rubio N, Sanz-Rodriguez F, Lipton HL (2006). Theiler's virus induces the MIP-2 chemokine (CXCL2) in astrocytes from genetically susceptible but not from resistant mouse strains. *Cell Immunol* **239**: 31–40.
- Saez-Borderias A, Guma M, Angulo A, Bellosillo B, Pende D, Lopez-Botet M (2006). Expression and function

- of NKG2D in CD4+ T cells specific for human cytomegalovirus. *Eur J Immunol* **36**: 3198–3206.
- Takada A, Yoshida S, Kajikawa M, Miyatake Y, Tomaru U, Sakai M, Chiba H, Maenaka K, Kohda D, Fugo K, Kasahara M (2008). Two Novel NKG2D ligands of the mouse H60 family with differential expression patterns and binding affinities to NKG2D. *J Immunol* **180**: 1678–1685.
- Tsunoda I, Libbey JE, Kobayashi-Warren M, Fujinami RS (2006). IFN-gamma production and astrocyte recognition by autoreactive T cells induced by Theiler's virus infection: role of viral strains and capsid proteins. *J Neuroimmunol* **172**: 85–93.
- Vilarinho S, Ogasawara K, Nishimura S, Lanier LL, Baron JL (2007). Blockade of NKG2D on NKT cells prevents hepatitis and the acute immune response to hepatitis B virus. *Proc Natl Acad Sci U S A*. **104**: 18187–18192.
- Wada Y, Fujinami RS (1993). Viral infection and dissemination through the olfactory pathway and the limbic system by Theiler's virus. *Am J Pathol* **143**: 221–229.