

Erratum

Erratum to ‘Detailed structure analysis of deposit layer in TEXTOR by means of TEM techniques’ [J. Nucl. Mater. 290–293 (2001) 295–298] ☆

S. Muto <sup>a,\*</sup>, N. Yokoya <sup>b</sup>, T. Tanabe <sup>a</sup>

<sup>a</sup> Center for Integrated Research in Science and Engineering, Nagoya University, Furo-cho, Chikusa-ku, Nagoya 464-8603, Japan

<sup>b</sup> Department of Nuclear Engineering, Graduate School of Engineering, Nagoya University, Chikusa-ku, Nagoya 464-8603, Japan

Unfortunately, in the final stages of publication, the incorrect Figs. 3–5 were inserted into this paper. The correct figures are given below. The Publisher apologises for the error.




Fig. 3. Electron energy-loss raw spectrum from the deposit in 100–600 eV.




Fig. 4. Comparison of EEL spectra in 150–350 eV from the deposit layer, crystalline boron and B<sub>4</sub>C from top to bottom. The pre-edge backgrounds are subtracted.

☆ PII of original article S0022-3115(00)00576-6.

\* Corresponding author. Tel.: +81-52 789 5157; fax: +81-52 789 3791.

E-mail addresses: muto@cirse.nagoya-u.ac.jp (S. Muto), tanabe@cirse.nagoya-u.ac.jp (T. Tanabe).




Fig. 5. Change in the intensity ratio of  $\pi^*$  and  $\sigma^*$  core-loss spectra as a function of angle,  $\theta$ , between the sample surface normal and incident electron direction (solid circle). The broken line illustrates  $\cos \theta$ .