



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Journal of Nuclear Materials 321 (2003) 118

Journal of  
nuclear  
materials

[www.elsevier.com/locate/jnucmat](http://www.elsevier.com/locate/jnucmat)

## Erratum

# Erratum to: 'Enthalpy and Gibbs energy of formation of cerium dicarbide'

[J. Nucl. Mater. 317 (2003) 102–108] ☆

R. Vidhya <sup>a</sup>, M.P. Antony <sup>a</sup>, P.R. Vasudeva Rao <sup>a,\*</sup>, B. Viswanathan <sup>b</sup>

<sup>a</sup> *Fuel Chemistry Division, Chemical Group, Indira Gandhi Centre for Atomic Research, Kalpakkam 603 102, Tamil Nadu, India*

<sup>b</sup> *Department of Chemistry, Indian Institute of Technology, Madras 600 036, India*

The Publisher regrets that in the original article several minus signs were not printed. Below are the correct notations:

1. Abstract, the last sentence should read: The recommended Gibbs energy and enthalpy of formation of CeC<sub>2</sub>(s) at 298 K are  $(-103.0 \pm 6.0)$  and  $(-120.1 \pm 11.0)$ .
2. Section 4.3, the fifth last sentence should read: The average of the results obtained by using the Gibbs energy functions of UC<sub>1.94</sub> and ThC<sub>1.94</sub>,  $(-120.1 \pm 11.0)$  kJ mol<sup>-1</sup>, is chosen to be the recommended value for  $\Delta_f H_{298}^0$  of CeC<sub>2</sub>(s).
3. Section 5, the last sentence should read: The enthalpy of formation and the Gibbs energy of formation of CeC<sub>2</sub>(s) at 298 K calculated from the present study are:  $(-120.1 \pm 11.0)$  and  $(-103.0 \pm 6.0)$  kJ mol<sup>-1</sup>, respectively.

☆ doi of original article: [10.1016/S0022-3115\(03\)00071-0](https://doi.org/10.1016/S0022-3115(03)00071-0).

\* Corresponding author. Tel.: +91-4114-280229; fax: +91-4114-280065.

E-mail address: [vasu@igcar.ernet.in](mailto:vasu@igcar.ernet.in) (P.R. Vasudeva Rao).