

Contents

A selection of papers from the Spent Fuel Workshop 2004

Preface	vii	
Contents	ix	
Gamma radiolysis of NaCl brine: Effect of dissolved radiolysis gases on the radiolytic yield of long-lived products, <i>M. Kelm and E. Bohnert</i>	1	
The effects of alpha-radiolysis on UO ₂ dissolution determined from batch experiments with ²³⁸ Pu-doped UO ₂ , <i>S. Stroes-Gascoyne, F. Garisto and J.S. Betteridge</i>	5	
The inhibiting effects of hydrogen on the corrosion of uranium dioxide under nuclear waste disposal conditions, <i>M.E. Broczkowski, J.J. Noël and D.W. Shoesmith</i>	16	
Radionuclide release from high burnup spent fuel during corrosion in salt brine in the presence of hydrogen overpressure, <i>A. Loida, V. Metz, B. Kienzler and H. Geckeis</i>	24	
		Leaching behaviour of unirradiated high temperature reactor (HTR) UO ₂ -ThO ₂ mixed oxides fuel particles, <i>C. Alliot, B. Grambow and C. Landesman</i> 32
		Radolytic modelling of spent fuel oxidative dissolution mechanism. Calibration against UO ₂ dynamic leaching experiments, <i>J. Merino, E. Cera, J. Bruno, J. Quiñones, I. Casas, F. Clarens, J. Giménez, J. de Pablo, M. Rovira and A. Martínez-Esparza</i> 40
		Enhanced diffusion under alpha self-irradiation in spent nuclear fuel: Theoretical approaches, <i>C. Ferry, P. Lovera, C. Poinssot and P. Garcia</i> 48
		Spent fuel radionuclide source-term model for assessing spent fuel performance in geological disposal. Part I: Assessment of the instant release fraction, <i>L. Johnson, C. Ferry, C. Poinssot and P. Lovera</i> 56
		Spent fuel radionuclide source term model for assessing spent fuel performance in geological disposal. Part II: Matrix alteration model and global performance, <i>C. Poinssot, C. Ferry, P. Lovera, C. Jegou and J.-M. Gras</i> 66