

# Synthesis of 3-(2-Oxo-2,3-dihydrobenzo[*b*]furan-3-ylidenehydrazono)-2,3-dihydrobenzo[*b*]furan-2-one

N. A. Pulina<sup>a</sup>, V. V. Zalesov<sup>b</sup>, and S. S. Kataev<sup>c</sup>

<sup>a</sup> Perm State Pharmaceutical Academy, Federal Agency of Health Protection and Social Development,  
ul. Lenina 48, Perm, 614900 Russia  
e-mail: pulina\_jane@mail.ru

<sup>b</sup> Perm State University, Perm, Russia

<sup>c</sup> Perm Regional Forensic Medical Examination Bureau, Perm, Russia

Received April 4, 2006

**Abstract**—Reactions of benzophenone and fluoren-9-one hydrazones with (2-hydroxyphenyl)oxoacetic acid gave [carboxy(2-hydroxyphenyl)methylidenehydrazono](2-hydroxyphenyl)acetic acid which underwent intramolecular cyclization with formation of 3-(2-oxo-2,3-dihydrobenzo[*b*]furan-3-ylidenehydrazono)-2,3-dihydrobenzo[*b*]furan-2-one. The symmetric azine was also obtained by reactions of (2-hydroxyphenyl)oxoacetic acid with triphenylphosphoranylidenehydrazones derived from benzophenones, fluoren-9-one, and 1-methyl-2,3-dihydro-1*H*-indole-2,3-dione.

**DOI:** 10.1134/S1070428007060103

The synthesis of hydrazono(2-hydroxyphenyl)acetic acids **Ia–Ie** and their cyclization into 3-hydrazono-2,3-dihydrobenzo[*b*]furan-2-ones **IIa–IIe** on heating in glacial acetic acid or concentrated hydrochloric acid (Scheme 1) were described in [1–3]. Since early 1970s these transformations remained beyond the scope of researchers' interest. Taking into account that compounds **I** and **II** may be important from the preparative viewpoint [4], we performed a study in this field.


We found that reactions of (2-hydroxyphenyl)oxoacetic acid (**III**) with benzophenone and fluoren-9-one hydrazones (reactant ratio 1:1 or 2:1) in anhydrous toluene at 20–25°C give no expected compounds **I**. Instead, the products are (2-hydroxyphenyl)[carboxy(2-hydroxyphenyl)methylidenehydrazono]acetic acid (**IV**, yield 67–72%) and azines **Va–Vd** (Scheme 2). Compounds **Va–Vd** were described in [5–7].

The IR spectrum of **IV** contained a broadened absorption band at 1744 cm<sup>-1</sup> due to stretching vibrations


of the acid carbonyl group and bands in the region 1617–1599 cm<sup>-1</sup>, belonging to C=N and C=C bonds. Compound **IV** displayed in the <sup>1</sup>H NMR spectrum a multiplet centered at δ 7.20 ppm from aromatic protons and a broadened signal at δ 11.00 ppm from the acid OH protons. In the mass spectrum of **IV**, the molecular ion peak (*m/z* 328) had a relative intensity of 1.9%, and fragment ion peaks did not contradict the assumed structure.

Presumably, the reaction includes intermediate formation of unsymmetrical azine like **A**, which reacts with the second molecule of acid **III** to produce azine **IV**. This reaction path is confirmed by the presence of the corresponding ketones (benzophenones or fluoren-9-one) in the reaction mixture. Acid **III** also reacted under analogous conditions with triphenylphosphoranylidenehydrazones derived from benzophenones, fluoren-9-one, and 1-methyl-2,3-dihydro-1*H*-indole-2,3-dione to give azine **IV**, but the yield of


Scheme 1.



R<sup>1</sup> = R<sup>2</sup> = H, R<sup>3</sup> = Ph (**a**), NH<sub>2</sub>C(O) (**b**), NH<sub>2</sub>S(O) (**c**); R<sup>1</sup> = Me, R<sup>2</sup> = H, R<sup>3</sup> = Ph (**d**), PhC(O) (**e**).



V, R = R' = Ph (a), R = Ph, R' = 4-BrC<sub>6</sub>H<sub>4</sub> (b); RR' = 9*H*-fluoren-9-ylidene (c), 1-methyl-2-oxo-2,3-dihydro-1*H*-indol-3-ylidene (d).



the latter was considerably smaller. Apart from triphenylphosphine oxide, the corresponding symmetric azines **Va–Vd** were isolated from the reaction mixture. Here, as in the above reaction, compound **IV** is likely to be formed through intermediate **A**. Thus, the processes leading to symmetric azines **IV** and **V** are concurrent, and the product ratio is determined most probably by thermodynamic factors.

On heating in acetic anhydride, compound **IV** underwent intramolecular cyclization to 3-(2-oxo-2,3-dihydrobenzo[*b*]furan-3-ylidenehydrazono)-2,3-dihydrobenzo[*b*]furan-2-one (**VI**). The IR spectrum of **VI** contained an absorption band at 1812 cm<sup>-1</sup> due to the lactone carbonyl groups, and C=N and C=C vibrations appeared in the region 1609–1598 cm<sup>-1</sup>. The aromatic protons gave a four-proton multiplet centered at δ 7.30 ppm and two two-proton multiplets at δ 7.65 and 7.83 ppm. No molecular ion peak was present in the mass spectrum of **VI**, but the fragment ion peaks (see Experimental) were consistent with its structure.

## EXPERIMENTAL

The IR spectra were recorded in mineral oil on FSM-1201 and Specord M80 instruments. The <sup>1</sup>H NMR spectra were measured on RYa-2310 (60 MHz) and Bruker DRX-500 (500 MHz) spectrometers from solutions in DMSO-*d*<sub>6</sub> relative to TMS as internal reference. The mass spectra (electron impact, 70 eV) were obtained on a Varian MAT-311 instrument with direct sample admission into the ion source. The progress of reactions and the purity of products were monitored by

TLC on Silufol UV-254 plates (diethyl ether–benzene–acetone, 10:9:1).

**[Carboxy(2-hydroxyphenyl)methylidenehydrazono](2-hydroxyphenyl)acetic acid (IV).** *a.* A suspension of 0.01 mol of benzophenone or fluoren-9-one hydrazone in 10 ml of anhydrous toluene was added to a solution of 1.66 g (0.01 mol) of acid **III** in 10 ml of the same solvent, and the mixture was stirred at room temperature until it became homogeneous. The resulting solution was cooled to 0°C, and the precipitate was filtered off and recrystallized from toluene. Yield 67–72%, yellow crystals, mp 183–185°C (decomp.). Mass spectrum, *m/z* (*I*<sub>rel</sub>, %): 310 (2.4), 292 (22.6), 264 (12.3), 238 (15.5), 192 (20.1), 164 (5.7), 146 (7.3), 120 (43.6), 102 (100), 93 (28.6), 76 (21.4). Found, %: C 58.82; H 3.58; N 8.60. C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>6</sub>. Calculated, %: C 58.54; H 3.66; N 8.54.

*b.* A suspension of 0.01 mol of benzophenone or fluoren-9-one triphenylphosphoranylidenehydrazono in 10 ml of anhydrous toluene was added to a solution of 1.66 g (0.01 mol) of acid **III** in 10 ml of the same solvent, and the mixture was stirred for 24 h at room temperature. The mixture was cooled to 0°C, and the precipitate was filtered off and recrystallized from toluene. Yield 42–58%.

**3-(2-Oxo-2,3-dihydrobenzo[*b*]furan-3-ylidenehydrazono)-2,3-dihydrobenzo[*b*]furan-2-one (VI).** A mixture of 3.28 g (0.01 mol) of compound **IV** and 20 ml of acetic anhydride was kept at room temperature until it became homogeneous. It was then cooled to 0°C, and the precipitate was filtered off and recryst-

tallized from acetonitrile. Yield 2.59 g (89%), orange crystals, mp 265–267°C (decomp.). Mass spectrum,  $m/z$  ( $I_{\text{rel}}$ , %): 264 (63.8), 236 (100), 208 (20.5), 102 (5.9), 76 (11.3). Found, %: C 65.68; H 2.78; N 9.55.  $\text{C}_{16}\text{H}_8\text{N}_2\text{O}_4$ . Calculated, %: C 65.75; H 2.74; N 9.59.

## REFERENCES

1. Tomchin, A.B., Ioffe, I.S., and Rusakov, E.A., *Zh. Obshch. Khim.*, 1971, vol. 41, pp. 1791, 1797; *Zh. Org. Khim.*, 1972, vol. 8, p. 1295; *Zh. Org. Khim.*, 1974, vol. 10, p. 604.
2. Fries, K. and Pfaffendorf, W., *Chem. Ber.*, 1912, vol. 45, p. 154.
3. Ranti, C. and Collino, F., *Farmaco, Ed. Sci.*, 1969, vol. 24, p. 577.
4. Zalesov, V.V. and Rubtsov, A.E., *Khim. Geterotsikl. Soedin.*, 2004, p. 163.
5. Schweizer, E.E., O'Neil, G.J., and Wemple, J.N., *J. Org. Chem.*, 1964, vol. 29, p. 1744.
6. DeMore, W.B., Pritchard, H.O., and Davidson, N., *J. Am. Chem. Soc.*, 1959, vol. 81, p. 5874.
7. Mariconi, E.J. and Murray, J.J., *J. Org. Chem.*, 1964, vol. 29, p. 3577.