

mother liquor amounted to 0.14 g., m.p. 95.5–99°, making the total yield 1.09 g. (68%). Recrystallization of a sample of the first crop material from methanol raised the m.p. to 106.5–107.5° corr.

A 0.50-g. sample of *trans*-stilbene was similarly oxidized with 15.4 mg. of osmium tetroxide and 1.25 g. of sodium metaperiodate (addition time 70 minutes). A subsequent stirring period of only 25 minutes was required to effect decolorization of the dark mixture. The first crop of 2,4-dinitrophenylhydrazone, obtained as described above, amounted to 1.35 g. (85% yield) of bright orange needles, m.p. 240–241.5° corr. undepressed on admixture with authentic benzaldehyde 2,4-dinitrophenylhydrazone, m.p. 241.5–242.5° corr.

*Oxidation of cyclohexene and of cyclopentene.* Ether-water method. A mixture of 15 ml. of ether, 15 ml. of water, 0.405 g. of cyclohexene, and 65.4 mg. of osmium tetroxide was stirred while a total of 2.32 g. of finely powdered sodium metaperiodate was added over a 40-minute period. The temperature was maintained at 24–26° during the addition and for an 80-minute period thereafter during which period the initially dark reaction mixture changed to pale yellow and considerable sodium iodate separated. The mixture was extracted thoroughly with ethyl acetate, and the combined organic layers were filtered through a little sodium sulfate. The solution then was treated with 2.5 g. of 2,4-dinitrophenylhydrazine and 5 drops of concentrated hydrochloric acid which produced an immediate precipitation. After standing for 1 hour the yellow product was separated and washed with ethyl acetate and ether; yield 1.81 g. (77%), m.p. 234–235° corr. (dec.). Repeated recrystallization of a sample from nitromethane gave *adipaldehyde bis-2,4-dinitrophenylhydrazone*, as yellow needles, m.p. 241.2–241.6° corr. (dec.).

*Anal.* Calc'd for:  $C_{18}H_{18}N_8O_8$ : C, 45.57; H, 3.82. Found: C, 45.33; H, 3.71.

A 0.387-g. specimen of cyclopentene was similarly oxidized with 74.2 mg. of osmium tetroxide and 2.83 g. of sodium metaperiodate (addition time 40 minutes). The subsequent stirring period to dispel the dark color was 25 minutes. The first crop of 2,4-dinitrophenylhydrazone isolated as described above amounted to 1.85 g., m.p. 187.5–189° corr. (reported 186–187°<sup>8</sup> 192.5–193.3°<sup>9</sup>). A second crop amounting to 0.14 g., m.p. 181.5–183°, was isolated. A specimen of the first crop material, recrystallized from nitromethane, melted at 192–193°.

DEPARTMENT OF CHEMISTRY  
UNIVERSITY OF OTTAWA  
OTTAWA, ONTARIO

DEPARTMENT OF CHEMISTRY  
UNIVERSITY OF WISCONSIN  
MADISON, WISCONSIN

(8) R. H. Hall and B. K. Howe, *J. Chem. Soc.*, 2480 (1951).

(9) A. C. Cope, H. L. Dryden, Jr., C. G. Overberger, and A. A. D'Addieco, *J. Am. Chem. Soc.*, **73**, 3416 (1951).

## Reductive Cleavage of Tosylates

DONALD B. DENNEY AND BERNARD GOLDSTEIN

Received January 12, 1956

In connection with some other work, it became necessary to convert several *p*-toluenesulfonic acid esters to the original alcohols. Normal hydrolytic

procedures were not acceptable, as a further requirement was that the alkyl-oxygen bond of the alcohol remain intact during the cleavage. Several reductive methods for performing such an operation are available, and they have been used with varying degrees of success. Recently, Kenner and Murray<sup>1</sup> have utilized Raney nickel for such cleavages; however, large excesses of nickel are required and in our hands the reactions were very slow. It has now been found that sodium in liquid ammonia smoothly reduces *p*-toluenesulfonates to the parent alcohol and toluene. The fate of the sulfur has not been determined. The general procedure is to add the tosylate in ether to a stirred solution of sodium in ammonia. After the reaction is complete, as is evidenced by the discharge of the blue color, water is added and the product is extracted with ether and then is distilled. The yields of alcohol were about 55%. No attempt was made to determine conditions for maximum yields. It was found by experiment that *ca.* six atoms of sodium were required per mole of tosylate reduced. This method is clearly advantageous for work in which the alkyl-oxygen bond must not be broken, and it may also be of value for tosylates which eliminate easily under hydrolytic conditions, since no olefin has been detected in any of these reductions.

## EXPERIMENTAL

*Reductive cleavage of cyclohexyl p-toluenesulfonate.* To a stirred solution of 13.8 g. (0.60 mole) of sodium in 300 ml. of liquid ammonia, there was added over a period of 40 minutes 26.4 g. (0.104 mole) of cyclohexyl *p*-toluenesulfonate in 100 ml. of anhydrous ether. Soon after the addition was completed, the solution turned orange, and solid ammonium chloride then was added until the orange color was discharged. Water, 50 ml., was added and the ammonia was allowed to evaporate. After evaporation, 60 ml. of water was added and the aqueous mixture was extracted with four 50-ml. portions of ether. The ether was dried over sodium sulfate. Distillation gave a small fraction, b.p. 26–28° (9 mm.) followed by a main fraction, b.p. 60–61° (9 mm.),  $n_D^{25}$  1.4675. The infrared spectra of the main fraction was identical in all respects with that of a known sample of cyclohexanol. The yield was 5.57 g. (54%).

*Reductive cleavage of n-octyl p-toluenesulfonate.* In a manner entirely analogous to that described above, 18.31 g. (0.0645 mole) of *n*-octyl *p*-toluenesulfonate was reduced with 6.9 g. (0.30 mole) of sodium. Distillation afforded a small amount of toluene followed by a main fraction b.p. 86–88° (8 mm.),  $n_D^{25}$  1.4230, identical in all respects with that of a known sample of *n*-octyl alcohol (infrared spectra). The yield was 4.82 g. (56%).

SCHOOL OF CHEMISTRY  
RUTGERS UNIVERSITY  
NEW BRUNSWICK, NEW JERSEY

(1) Kenner and Murray, *J. Chem. Soc.*, 5178 (1949).