



scribed previously and would appear to be quite difficult to obtain by any other means. It seems clear that we have available an excellent method for preparing certain substituted naphthalene compounds from selected  $\beta, \gamma$ -unsaturated lactones. The scope and limitations of the method are being explored further.

Other striking differences in chemical behavior between I and II have been observed and these will be discussed in a forthcoming paper.

#### EXPERIMENTAL<sup>8</sup>

*Reaction of  $\alpha$ -benzylidene- $\gamma$ -phenyl- $\Delta^{\beta, \gamma}$ -butenolide (II) with benzene.* In a 2-l., round bottomed flask, fitted with a mechanical stirrer, dropping funnel, and reflux condenser, were placed 19.8 g. (0.15 mol.) of anhydrous aluminum chloride in 250 ml. of dry, thiophene-free benzene. The mixture was cooled to 10–20° and stirred for 1 hr. To this mixture was added dropwise with stirring a solution containing 12.4 g. (0.05 mol.) of II in 250 ml. of dry benzene, the temperature being maintained at 10–20° during the addition. The mixture turned brick-red. When all of the butenolide had been added, the mixture was stirred for an additional 3 hr. at room temperature. The complex was decomposed with dilute hydrochloric acid and the resulting mixture extracted with ether. The ether layer was washed with dilute sodium bicarbonate solution and water and then dried over anhydrous magnesium sulfate. Upon evaporation of the solvent on steam bath, a yellowish white solid was obtained. Crystallization from 95% ethanol gave 8.5 g. (70.8%) of white needles of 4-phenyl-2-naphthoic acid (III), m.p. 262.5–263.5°.

*Anal.* Calcd. for  $C_{17}H_{12}O_2$ : C, 82.24; H, 4.83. Found: C, 82.52; H, 4.81. Mol. wt. calcd.: 248. Found: 248 (neut. equiv.).

$\lambda_{\max}^{EOH}$  245 m $\mu$  ( $\epsilon$  31,000), 290 m $\mu$  ( $\epsilon$  7990). Infrared absorption at 1695  $cm^{-1}$  (s) and 2600  $cm^{-1}$  (w).

*Reaction of II with anisole.* In a 500-ml. round bottomed flask, fitted with a mechanical stirrer, dropping funnel, and reflux condenser, were placed 4.7 g. (0.036 mol.) of anhydrous aluminum chloride and 3.8 g. (0.036 mol.) of anisole in 65 ml. of methylene chloride. The mixture was stirred for 1 hr. at 10°. To this mixture was added dropwise with stirring a solution containing 3 g. (0.012 mol.) of II in 100 ml. methylene chloride, the temperature being maintained at 10–20° during the addition. When all of the butenolide had been added, the mixture was stirred for an additional 3 hr. at room temperature. The complex was decomposed with 250 ml. of dilute (1:15) hydrochloric acid and the two layers which formed were separated, the water layer was washed with methylene chloride and the combined extracts were washed with dilute acid and water until neutral to litmus. Excess methylene chloride was removed by evaporation on a steam bath. The product was crystallized from 95% ethanol to give 1.5 g. (50%) of white needles, m.p. 258–260°.

*Anal.* Calcd. for  $C_{17}H_{12}O_2$ : C, 82.24; H, 4.83. Found: C, 82.57; H, 5.17.

Mixed melting point with the product obtained with benzene showed no depression. The ultraviolet and infrared spectra were also identical.

*Decarboxylation of 4-phenyl-2-naphthoic acid (III).* III [8.13 g. (0.033 mol.)] and 0.648 g. (0.0028 mol.) of copper chromite in 10 ml. of quinoline were heated under slow reflux for 7 hr. The mixture was treated with 100 ml. of ether and washed with several portions of dilute hydrochloric acid followed by sodium bicarbonate solution and water. The ether layer was dried over anhydrous magnesium sulfate.

The ether was removed by evaporation and 1-phenylnaphthalene was obtained as a viscous oil, b.p. 333–335° at

748 mm.,  $n_D^{23}$  1.6654 (reported b.p. 334–336°,  $n_D^{18}$  1.6692<sup>10</sup>).

*Nitration of 1-phenylnaphthalene.* To 3 ml. of a 50-50 mixture of fuming nitric acid and glacial acetic acid, cooled in an ice bath, was added dropwise 0.25 ml. of 1-phenylnaphthalene. The solution was mixed and poured over ice. The 4-nitro-1-phenylnaphthalene thus obtained was crystallized from petroleum ether to give yellow crystals melting at 129–130° (reported<sup>11</sup> m.p. 129–130°).

*Spectral measurements and analyses.* Infrared spectra were obtained on a Perkin-Elmer 21 spectrophotometer, using chloroform as solvent. Ultraviolet spectra were measured in 95% ethanol using a Beckman DK-2 spectrophotometer. Microanalyses were conducted by Micro-Tech Laboratories, Skokie, Ill.

*Acknowledgment.* The authors gratefully acknowledge the financial assistance of the Research Corp. and the Philip D. Armour Cancer Fund.

DEPARTMENT OF CHEMISTRY  
ILLINOIS INSTITUTE OF TECHNOLOGY  
CHICAGO 16, ILL.

(9) R. Weiss and K. Woidich, *Monatsh.*, **46**, 455 (1925).

(10) W. Davies, N. W. Gamble, and W. E. Savage, *J. Chem. Soc.*, 4678 (1952).

(11) R. T. Arnold, C. Collins, and W. Zenk, *J. Am. Chem. Soc.*, **62**, 983 (1940).

#### Aluminum Iodide as a Friedel-Crafts Catalyst<sup>1</sup>

ERNEST R. KLINE, BRUCE N. CAMPBELL, JR.,<sup>2</sup>  
AND EARL C. SPAETH<sup>3</sup>

Received April 1, 1959

A comparison of aluminum iodide with aluminum chloride and aluminum bromide as catalysts in typical Friedel-Crafts alkylations and acylations seems not to have been reported. The absence of such an investigation may be due, in part, to the well known sensitivity of aluminum iodide to air and moisture.<sup>4</sup>

Aluminum iodide was prepared by two methods which minimize contact with air and moisture. Comparisons have been made with aluminum chloride and aluminum bromide as catalysts in the isopropylation, succinoylation, and benzoylation of benzene. The results of these experiments are summarized in Table I. The reactions using aluminum chloride reported in this table are each the best of several made to determine conditions which give moderately good yields. The alkylations with aluminum bromide and aluminum iodide (Method B) represent the average of two runs. Although aluminum iodide acts as a catalyst in all these reactions,

(1) Based on part of the Ph.D. thesis of B.N.C., University of Connecticut, 1958.

(2) To whom inquiries should be sent. Present address: Department of Chemistry, MacMurray College, Jacksonville, Ill.

(3) Present address: Apt. 203, 1600 N. Quinn St., Arlington, Va.

(4) G. W. Watt and J. L. Hall, *Inorganic Syntheses*, McGraw-Hill Book Company, Inc., New York, N. Y., 1953, Vol. IV, p. 119.

(8) M.p.'s are corrected.