

adducts were not detected in these cases, even though complete disappearance of TPPO was indicated in some cases by the absence of oxygen evolution on warming. This result suggests that biradicals like II are very short lived and decay to stable products before they can be spin trapped. This observation is in agreement with the suggestion of Bartlett and Chu<sup>7</sup> that hindered phenols act as electron donors to TPPO rather than intercepting biradicals like II, since radicals that can be intercepted by phenols might also be expected to form spin adducts.

### Conclusion

The decomposition of TPPO mainly gives singlet oxygen and the production of III probably is a minor pathway. This is evidenced by the brisk oxygen evolution when TPPO-PBN mixtures are rapidly warmed to room temperature. Since the peroxy spin adduct is unstable and since PBN reacts with singlet oxygen to produce nonradical products, an exact estimate of the ratio of the yields of singlet oxygen and spin adducts is not possible. However, our results combined with the kinetic evidence of Stephenson and McClure<sup>2</sup> confirm the biradical mechanism by which TPPO decomposes to singlet oxygen. The formation of PBNOx in many previous photooxidation studies<sup>12,13</sup> can be explained by our finding that nitrene spin traps are oxidized to acyl nitroxides by singlet oxygen at low temperatures.

### Experimental Section

**General Procedures.** ESR spectra were recorded on a Varian E-109 ESR spectrometer. Low-temperature experiments were done with a quartz Dewar insert. The temperature was controlled by varying the flow of nitrogen (precooled by passing through a stainless steel coil immersed in liquid nitrogen) through the insert.

Ozone was generated by using a Welsbach Model T-23 ozonator. Triphenyl phosphite (Aldrich) was washed with base and distilled in vacuo. Commercially available PBN (Eastman) was recrystallized from hexane before use. DMPO obtained from Spin Trap Producers (Guelph, Canada) was used without further purification. Dichloromethane was washed with sulfuric acid and distilled water, dried over CaCl<sub>2</sub>, distilled from CaH<sub>2</sub>, and stored over molecular sieves.

**Spin Trapping: TPPO-PBN Experiments.** TPPO was made by the slow addition of 8 mg of triphenyl phosphite dissolved in 0.1 mL of dichloromethane to a saturated solution of ozone in dichloromethane (0.3 mL) at -78 °C, with continued bubbling of ozone. After the addition was complete, excess ozone was flushed out with nitrogen and a precooled solution of 5 mg of PBN in 0.1 mL of dichloromethane was added. The sample was then warmed in a -15 °C bath for 10-15 min and cooled back to -70 °C, and the ESR spectrum was recorded after deoxygenation. To establish the stability of the peroxy radical spin adduct, we added methanol and pyridine (0.1 mL, 1:1) at -78 °C to a TPPO-PBN sample that had been warmed at -5 °C until oxygen evolution almost stopped.

**Quenching Experiments with  $\beta$ -Carotene.** A blank experiment was done as follows. To a TPPO-PBN mixture prepared as described above was added 0.1 mL of precooled methanol and pyridine. The mixing of the reagents was effected by a stream of nitrogen and the sample was transferred to a -65 °C bath. After 30 min, the sample was deoxygenated with N<sub>2</sub> again and the ESR spectrum was recorded at -70 °C. In the  $\beta$ -carotene experiments, 0.1 mL of a solution of  $\beta$ -carotene of the required concentration was added before the addition of methanol and pyridine and treated exactly as the blank. Volumes were made equal by addition of dichloromethane.

**Acknowledgment.** This work was supported in part by grants from the NSF and NIH (HL-16029).

**Registry No.** 1, 79101-59-0; ozone, 10028-15-6; TPPO, 29833-83-8; (PhO)<sub>3</sub>P, 101-02-0.

## Enzymatic Conversion of $\alpha$ -Keto Aldehydes to Optically Active $\alpha$ -Hydroxy Acids Using Glyoxalase I and II<sup>1</sup>

Mark A. K. Patterson,<sup>2</sup> Richard P. Szajewski,<sup>3</sup> and Geroge M. Whitesides\*

Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Received May 18, 1981

$\alpha$ -Keto aldehydes (RCOCHO, R = CH<sub>3</sub>, CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>, Ph, *p*-CH<sub>3</sub>Oph, *p*-ClPh) have been converted to optically active  $\alpha$ -hydroxy acids (RCHOHCO<sub>2</sub>H) by the combined action of glutathione and the (immobilized) enzymes glyoxalase I (GX-I, EC 4.4.1.5) and glyoxalase II (GX-II, EC 3.1.2.6). The reaction seems to provide a practical if specialized method for synthesizing 1-10-g quantities of product with enantiomeric excesses (ee) in the range 75-99%. Efforts to increase the scale of the reaction are accompanied by decreases in the ee of the product and in the turnover number reached by the enzymes.

### Introduction

This paper describes the conversion of  $\alpha$ -keto aldehydes to optically active acids, using a catalytic system comprised of two enzymes, glyoxalase I<sup>4-13</sup> (GX-I, EC 4.4.1.5) and

glyoxalase II<sup>14,15</sup> (GX-II, EC 3.1.2.6), and the cysteine-containing cofactor glutathione (GSH,  $\gamma$ -L-Glu-L-Cys-Gly) (eq 1). GX-I accepts a number of sterically unhindered  $\alpha$ -keto aldehydes as substrates but does not accept substances that have large substituents close to the keto

(1) Supported by the National Institutes of Health, Grant GM-26543.

(2) NSF Predoctoral Fellow, 1977-1980.

(3) NIH Postdoctoral Fellow, 1976-1978.

(4) Alexander, N. M.; Boyer, J. L. *Anal. Biochem.* 1971, 41, 29-38.

(5) Vander Jagt, D. L.; Han, L.-P. B.; Lehman, C. H. *Biochemistry* 1972, 11, 3735-3740.

(6) Vander Jagt, D. L.; Han, L.-P. B. *Biochemistry* 1973, 12, 5161-5167.

(7) Ekwall, K.; Mannervik, B. *Biochim. Biophys. Acta* 1973, 297, 297-299.

(8) Uotila, L.; Koivusalo, M. *Eur. J. Biochem.* 1975, 52, 493-503.

(9) Kester, M. V.; Norton, S. J. *Biochim. Biophys. Acta* 1975, 391, 212-221.

(10) Vander Jagt, D. L.; Daub, E.; Krohn, J. A.; Han, L.-P. B. *Biochemistry* 1975, 14, 3669-3675.


(11) Han, L.-P. B.; Davison, L. M.; Vander Jagt, D. L. *Biochim. Biophys. Acta* 1976, 445, 486-499.

(12) Hall, S. S.; Doweiko, A. M.; Jordan, F. J. *Am. Chem. Soc.* 1978, 100, 5934-5939.

(13) Marmstål, E.; Aronsson, A.-C.; Mannervik, B. *Biochem. J.* 1979, 183, 23-30.

(14) Wieland, T.; Pfeleiderer, G.; Lau, H. *Biochem. Z.* 1956, 327, 393-406. Uotila, L. *Biochemistry* 1973, 12, 3944-3951. Ball, J. C.; Vander Jagt, D. L. *Anal. Biochem.* 1979, 98, 472-477.

(15) Ball, J. C.; Vander Jagt, D. L. *Biochemistry* 1981, 20, 899-905.



group.<sup>5,8,10,11</sup> Its action appears to involve an enediol-proton transfer mechanism,<sup>12</sup> with the hemithioacetal formed between GSH and  $\alpha$ -keto aldehydes as substrate.<sup>6,10</sup> GX-II has been studied in less detail but seems to function primarily as a catalyst for the hydrolysis of thioesters of GSH.<sup>14,15</sup> The combination of GX-I and GX-II has been reported to convert methylglyoxal to lactic acid, having >99% the D(-) configuration,<sup>4,7</sup> and phenylglyoxal to mandelic acid (>95% D(-)).<sup>10</sup> GX-I and GX-II are commercially available.

### Results

GX-I and GX-II were immobilized in a cross-linked gel derived from poly(acrylamide-co-N-(acryloxy)succinimide) (PAN)<sup>16</sup> in yields of ~51% and 31%, respectively. Immobilized GX-I and GX-II are stable at pH 7 and 25 °C for periods of several months in the absence of reactants. In the presence of  $\alpha$ -keto aldehydes, both lose activity, presumably as a result of modification of arginine residues.<sup>15,17-19</sup> This instability of the enzymes (especially GX-II)<sup>15</sup> in the presence of  $\alpha$ -keto aldehydes is the most severe limitation to the practical utility of this catalytic system.

Reactions were carried out by using two procedures. In one (called the "simultaneous" procedure here),  $\alpha$ -keto aldehyde was added slowly to an aqueous solution of GSH containing both gel-immobilized enzymes. In the second (called the "sequential" procedure), immobilized GX-I was used to convert  $\alpha$ -keto aldehyde to the intermediate thioester. The GX-I was then removed, and immobilized GX-II was used in a separate step to hydrolyze the thioester to product. Removal of the GX-II and readdition of GX-I and additional  $\alpha$ -keto aldehyde permitted limited reuse of GSH, howbeit with a decrease in the enantiomeric excess of the product  $\alpha$ -hydroxy acid.

Table I summarizes representative results obtained with these two procedures. These results illustrate several features of our experience with this reaction. First, reactions run with the "simultaneous" procedure give better efficiencies for use of GSH (as reflected in  $\text{TN}_{\text{GSH}}$ ) than those with the "sequential" procedure. Particularly on large scale, however, the  $\text{TN}$ 's for the enzymes (especially GX-II) are lower in the "simultaneous" procedure, probably as a result of deactivation of the enzymes by reaction with  $\alpha$ -keto aldehyde. Since the enzymes are the greatest contributor to the cost of these syntheses, in most cases it is more practical simply to accept GSH as a stoichiometric reagent, use the "sequential" procedure, and recover as much GSH as possible (typically ~50%) for reuse by isolation at the conclusion of the reaction. Second, the

enantiomeric excess in the product is variable. It appears that use of large quantities of GX-II (reflected in low values of  $\text{TN}_{\text{GX-II}}$  in Table I) leads to relatively high values of ee. We conclude from this observation, and from others,<sup>20</sup> that the intermediate  $\alpha$ -hydroxy thiol esters ( $\text{RCHOHCOSG}$ ) epimerize under the reaction conditions: the highest values of ee are observed under conditions in which these species would be expected to have relatively short lifetimes.

### Conclusions

The enzymatic conversion of  $\alpha$ -keto aldehydes to  $\alpha$ -hydroxy acids represents a practical if specialized method for the small-scale (<10 g) preparation of these latter species.<sup>21</sup> The advantage of this method is that the product is obtained in an enantiomerically enriched form. Further, if relatively large quantities of GX-II are used, the enantiomeric excess of the product may be high. The enzymes are commercially available, and the experimental procedure is straightforward. Although most of our work has been carried out with immobilized enzymes, these reactions may also be conducted with soluble enzymes. The stabilities of the soluble enzymes are lower than those of the immobilized enzymes, but the procedure is simplified by omission of the immobilization step and the lower stability is compensated for by avoiding the significant loss in enzymatic activity (50–69% of initial activity) that accompanies the immobilization step. The enzymes appear to accept a useful range of substrates,<sup>5,10,14</sup> although their activity is very low when R is a group that introduces steric bulk close to the keto group.

The deficiencies of this method are also several. First, yields of  $\alpha$ -hydroxy acids are not high, and enzyme costs are such that the reaction is expensive to scale up.<sup>22</sup> Efforts to increase the yield or to produce larger quantities of product seem to be frustrated by spontaneous epimerization of the intermediate  $\text{RCHOHCOSG}$  in the reaction and by deactivation of the glyoxalase enzymes (especially GX-II) by reaction with  $\alpha$ -keto aldehyde (and perhaps also with thioester). Second, only one enantiomer of the  $\alpha$ -hydroxy acids is accessible by this route.

### Experimental Section

**General Procedures.** Glyoxalase I (EC 4.4.1.5, from yeast), glyoxalase II (EC 3.1.2.6, from beef liver), D-lactic dehydrogenase (EC 1.1.1.28), L-lactic dehydrogenase (EC 1.1.1.27), and reduced glutathione were obtained from Sigma Chemical Co. (R)-(+)- $\alpha$ -Methoxy- $\alpha$ -(trifluoromethyl)benzeneacetic acid was from Aldrich. All other materials were reagent grade. pH measurements were made on a Radiometer PH M62 standard pH meter. Enzymatic assays and absorbance measurements were performed on a Gilford 240 spectrophotometer equipped with a constant temperature bath and a strip-chart recorder. A Varian T-60 was used to record 60-MHz <sup>1</sup>H NMR spectra, and a Bruker WH-250 was used for

(20) The use of multiple cycles in the "sequential" reactions leads to a reduced ee. This result can be explained by an  $\alpha$ -hydroxy carboxylate catalyzed epimerization of the  $\alpha$ -hydroxy thiol ester. In addition, phenylglyoxals having electron-withdrawing substituents seem to give mandelic acids with lower enantiomeric excesses than with electron-donating substituents.

(21) Optically active  $\alpha$ -hydroxy acids have been obtained by resolution (Collet, A.; Jacques, J. *Bull. Soc. Chim. Fr.* 1973, 3330–3334) or by the conversion of optically active  $\alpha$ -bromo acids (Levene, P. A.; Mori, T.; Mikeska, L. A. *J. Biol. Chem.* 1927, 75, 337–365) or  $\alpha$ -amino acids (Baker, C. G.; Meister, A. *J. Am. Chem. Soc.* 1951, 73, 1336–1338). In addition, they can be prepared by asymmetric reactions (Meyers, A. I.; Slade, J. *J. Org. Chem.* 1980, 45, 2785–2791 and references contained therein).

(22) Costs: GX-I: \$7.10/100 U; GX-II: \$43.00/100 U; GSH: \$305/mol. The major contributor to the cost of the reactions is the price of GX-II. Attempts to substitute less expensive esterases, such as acylase I, acetyltransferase, carboxylesterase, and *Candida cylindracea* lipase for the GX-II were unsuccessful. With extensive screening it might, however, be possible to develop a better (i.e., less expensive) esterase for these glutathione thiol esters.

(16) Pollak, A.; Blumenfeld, H.; Wax, M.; Baughn, R. L.; Whitesides, G. M. *J. Am. Chem. Soc.* 1980, 102, 6324–6336.

(17) Means, G. E.; Feeney, R. E. "Chemical Modification of Proteins"; Holden-Day: San Francisco, 1971.

(18) Takahashi, K. *J. Biochem.* 1977, 81, 395–402, 403–414.

(19) McLaughlin, J. A.; Pethig, R.; Szent-Györgyi, A. *Proc. Natl. Acad. Sci. U.S.A.* 1980, 77, 949–951.

Table I. Synthesis of Chiral  $\alpha$ -Hydroxy Carboxylic Acids from  $\alpha$ -Keto Aldehydes Using Glyoxalase I (GX-I), Glyoxalase II (GX-II), and Glutathione (GSH)

R	RCOCHO, mmol	GSH, mmol	product (RCHOHCO <sub>2</sub> M) <sup>a</sup>				reaction type <sup>b</sup>	TN <sup>c</sup>		
			M	g	%	ee R, %		GX-I	GX-II	GSH cycles <sup>d</sup>
<i>p</i> -ClPh	278	139	Ca/2	14	26	76 <sup>e</sup>	simul	5 × 10 <sup>7</sup>	1 × 10 <sup>6</sup>	2
Ph	65	7.5	H	7	75	91 <sup>e</sup>		3 × 10 <sup>7</sup>	1 × 10 <sup>5</sup>	9
<i>p</i> -MeOPh	11	2	Ca/2	2	84	94 <sup>e</sup>		4 × 10 <sup>6</sup>	2 × 10 <sup>4</sup>	6
CH <sub>3</sub>	324	7	Zn/2	8	17	97 <sup>f</sup>		4 × 10 <sup>6</sup>	9 × 10 <sup>4</sup>	46
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	28	0.6	Ba/2	4	79	>98 <sup>e</sup>		2 × 10 <sup>6</sup>	5 × 10 <sup>4</sup>	47
	20	25		2	54	>99 <sup>e</sup>	seq	2 × 10 <sup>6</sup>	6 × 10 <sup>5</sup>	0.8
	29	36		2	30	>99 <sup>e</sup>		4 × 10 <sup>6</sup>	4 × 10 <sup>5</sup>	0.8
	67	50		4	30	85 <sup>e,g</sup>		2 × 10 <sup>7</sup>	9 × 10 <sup>5</sup>	1.3
	98	68		12	65	80 <sup>e,g,h</sup>		3 × 10 <sup>7</sup>	1 × 10 <sup>6</sup>	1.4

<sup>a</sup> Yields (in grams and percent based on RCOCHO) are for isolated products. The yield before isolation was 73% for the lactic acid producing system, as measured by enzymatic assay. Suitable assays did not exist for the measurement of other products in solution. <sup>b</sup> Reaction type is discussed in the text. <sup>c</sup> TN = turnover number = moles of RCHOHCO<sub>2</sub>H generated per mole of enzyme or glutathione originally present in the solution. These turnover numbers are not corrected by taking into account enzyme or glutathione present at the conclusion of the reaction. <sup>d</sup> "Cycle" refers to the number of times that the sample in a sequential procedure was used in a sequence of formation of RCHOHCOSG and hydrolysis of this species. <sup>e</sup> Enantiomeric excess determined by NMR analysis, using diastereomeric esters (see the text). <sup>f</sup> Enantiomeric excess determined by enzymatic analysis. <sup>g</sup> Insignificant quantities of active enzymes were recovered upon completion of the reactions except for these cases. An average of 55% GX-I and 35% of GX-II initial activities were recovered at the end of each cycle. <sup>h</sup> Enantiomeric excess was 89% after two cycles.

250-MHz spectra. Distilled water was deionized with a Barnstead ion-exchange column and then redistilled, using a Corning AG-1b type still; this water was used to prepare all solutions. Peristaltic pumps from LKB were used for the addition of substrates and base, while the pH was maintained at a constant value with a Chemtrix pH controller.

Methylglyoxal was steam-distilled under argon before use. (*p*-Chlorophenyl)glyoxal,<sup>23</sup> (*p*-methoxyphenyl)glyoxal,<sup>23</sup> propylglyoxal,<sup>24</sup> *S*-(*p*-bromobenzyl)glutathione,<sup>25</sup> and PAN<sup>16</sup> were synthesized following literature procedures. Glyoxalase I activity was measured in buffered solutions (pH 6.0–8.0) that were 10 mM in MgCl<sub>2</sub>, 3.0 mM in methylglyoxal, and 3.0 mM in GSH at 30 °C. The increase in absorbance at 240 nm ( $\epsilon$  3370 M<sup>-1</sup> cm<sup>-1</sup>)<sup>26</sup> was directly related to enzymic activity. Glyoxalase II activity was determined in buffered solutions (pH 6.0–8.0) containing 0.6 mM *S*-lactoylglutathione at 30 °C, with the decrease in absorbance at 240 nm being measured. The concentrations of methylglyoxal,<sup>26</sup> propylglyoxal,<sup>26</sup> phenylglyoxal,<sup>5</sup> (*p*-methoxyphenyl)glyoxal,<sup>5</sup> (*p*-chlorophenyl)glyoxal,<sup>5</sup> glutathione,<sup>26</sup> and L-lactic acid<sup>26</sup> were measured enzymatically by using literature procedures. D(-)-Chloromandelic acid was prepared by resolution.<sup>21</sup>

**Immobilization of Enzymes.** The glyoxalase enzymes were immobilized following the procedure of Pollak et al.<sup>16</sup> GX-I was immobilized in a 0.3 M Hepes, pH 7.5, solution containing 10 mM MgCl<sub>2</sub>, 5 mM 1,3-dithiopropion-2-ol (DTP), and 0.8 mM *S*-(*p*-bromobenzyl)glutathione.<sup>25</sup> Optimal yields were obtained when a ratio of 0.5 mg of enzyme to 100 mg of PAN-500 (contains 500  $\mu$ mol of active ester groups per g of solid) was used. The average yield of immobilized enzyme was 51%, with 14% of the initial activity present in the wash. GX-II was immobilized in a 0.2 M P<sub>i</sub>, pH 7.0, solution containing 5 mM MgCl<sub>2</sub>, 5 mM DTP, and 8 mM GSH with an enzyme to PAN-500 ratio of 2 mg/100 mg. An average of 31% of the initial activity was present in the gel, with an additional 8% in the wash.

**$\alpha$ -Hydroxypentanoate: "Simultaneous" Procedure.** The pH of 150 mL of a solution containing MOPS (7.5 mmol), MgCl<sub>2</sub> (1.5 mmol), GSH (0.6 mmol), and 1,3-dithiopropion-2-ol (DTP, 0.9 mmol) was adjusted to 6.8 with 2 N NaOH. The solution was

deoxygenated with a stream of argon and soluble GX-I (160 U) and GX-II (100 U) added. Over an 8-day period 69 mL of an aqueous solution of propylglyoxal (28 mmol) was added, together with additional GX-I (180 U) and GX-II (75 U). The pH was maintained at 6.8 with 2 N NaOH by using a pH controller. Upon completion of the reaction, the mixture was acidified to pH 1.5 with concentrated HCl and extracted continuously for 40 h with ether (0.7 L). The ether was concentrated under reduced pressure, and the residue was diluted with H<sub>2</sub>O (40 mL), heated briefly, and neutralized with BaCO<sub>3</sub>. The mixture was boiled and filtered while hot, and the residue was washed with boiling H<sub>2</sub>O (10 mL). The filtrate volume was reduced to 25 mL by gentle boiling, the solution cooled (70 °C), and ethanol (15 mL) added. After cooling to 4 °C, the precipitate was filtered, washed with ether, and dried, yielding 4.1 g of barium  $\alpha$ -hydroxypentanoate (>98% ee R; 79% based on propylglyoxal). The activities of GX-I and GX-II at the end of the reaction were only 6% and 3%, respectively, of the original values.

**$\alpha$ -Hydroxypentanoate: "Sequential" Procedure.** A 100-mL aqueous solution of propylglyoxal (27 mmol) containing GSH (34 mmol), MgCl<sub>2</sub> (1.0 mmol), and DTP (1.0 mmol) was adjusted to pH 7.0 with 10 N NaOH and then deoxygenated with argon. Immobilized GX-I (22 U) was added. After 47 h the GX-I was removed by gentle centrifugation (~3000 rpm) and immobilized GX-II (6.6 U) added to the supernatant. The pH of the suspension was maintained between 6.7 and 7.0 with 1 N NaOH and a pH controller. After 67 h the mixture was centrifuged to remove the GX-II and additional propylglyoxal (31 mmol) and GSH (9 mmol) was added to the supernatant. The pH of the solution was adjusted to 6.9 with 10 N NaOH, the solution deoxygenated, and immobilized GX-I (22 U) added. Fifty hours were required for completion of the reaction, at which time the GX-I was removed and immobilized GX-II (6.5 U) added to the supernatant. Seventy hours later the suspension was centrifuged to remove the GX-II, and to the resulting solution was added propylglyoxal (40 mmol) and GSH (25 mmol). Following adjustment of pH 6.8 with 10 N NaOH, immobilized GX-I (40 U) was added. The suspension was centrifuged after 48 h to isolate the GX-I, and immobilized GX-II (5.4 U) was added. Forty-three hours later an additional portion of GX-II (5.1 U) was added. After 71 h the reaction was complete, the GX-II was removed by centrifugation, and the supernatant was acidified to pH 1.5 with concentrated HCl. The acidified solution was extracted with ether and barium  $\alpha$ -hydroxypentanoate (12.0 g, 32 mmol; 80% ee R; 65% yield based on propylglyoxal) isolated as described above. The average amount of initial activity recovered at the end of each step was 56% for GX-I and 37% for GX-II.

***p*-Chloromandelate.** A "simultaneous" procedure, utilizing immobilized GX-I and II, converted a solution of (*p*-chlorophenyl)glyoxal (279 mmol, dissolved in Me<sub>2</sub>SO) to *p*-chloromandelic acid (3.5 g, 19 mmol) and calcium *p*-chloromandelate

(23) Arnold, R. T.; Fuson, R. C. *J. Am. Chem. Soc.* 1936, 58, 1295–1296.

(24) Royals, E. E.; Robinson, A. G. *J. Am. Chem. Soc.* 1956, 78, 4161–4164. The solution of  $\alpha$ -keto aldehyde obtained by acid hydrolysis of the corresponding acetal was used without purification.

(25) Vince, R.; Daluge, S.; Wadd, W. B. *J. Med. Chem.* 1971, 14, 402–404. *S*-(*p*-Bromobenzyl)glutathione is a competitive inhibitor of glyoxalase I, causing 50% inhibition at a concentration of 9  $\mu$ M with an  $\alpha$ -keto hemithioacetal concentration of 0.51 mM.

(26) "Methods of Enzymatic Analysis", 2nd English ed.; Bergmeyer, H. U., Ed.; Academic Press: New York, 1974. GX-I, pp 469–470; RCOCHO, R = Me or *n*-Pr, pp 1496–1498; GSH, pp 1643–1647; D-lactate, pp 1492–1495; L-lactate, pp 1464–1468.

(11 g, 27 mmol). The final reaction mixture was 14% (v/v) Me<sub>2</sub>SO. Calcium carbonate was used in the preparation of calcium *p*-chloromandelate. The overall yield was 26% (based on (*p*-chlorophenyl)glyoxal) and the product was composed of a 76% ee of the *R* enantiomer. The recovered GX-I and GX-II had 3% and 59%, respectively, of their initial activities.

**Mandelic Acid.** An aqueous suspension of phenylglyoxal (65 mmol in 350 mL of H<sub>2</sub>O, pH 6.3) was converted in a "simultaneous" process with immobilized enzymes to mandelic acid (7.4 g, 49 mmol; 91% ee *R*; 75% based on phenylglyoxal). The recovered enzymes were not active.

***p*-Methoxymandelate.** A "simultaneous" procedure, using soluble GX-I and GX-II, converted 256 mL of an aqueous solution of (*p*-methoxyphenyl)glyoxal (11 mmol) to the corresponding  $\alpha$ -hydroxy acid. Calcium *p*-methoxymandelate (1.9 g, 4.6 mmol; 94% ee *R*; 84% based on (*p*-methoxyphenyl)glyoxal) was isolated with calcium carbonate. No enzymatic activity remained upon completion of the reaction.

**Lactate.** Methylglyoxal solutions are known to contain significant amounts of acid, presumably lactic acid.<sup>4</sup> A steam-distilled solution (450 mL of 2.97 M methylglyoxal, 0.48 M acid) was treated with 16 portions of Dowex-1  $\times$  8/50 (OH<sup>-</sup>) (15-mL portions, 20 min each) to give a solution (380 mL) of 1.62 M methylglyoxal and 8.7 mM acid. A "simultaneous" procedure, using immobilized enzymes, converted 200 mL of this solution (324 mmol methylglyoxal) to lactic acid, which was isolated as the zinc salt (7.9 g, 28 mmol; 97% ee *R*; 17% based on methylglyoxal), using zinc carbonate. No enzymatic activity was recovered.

**Determination of Enantiomeric Compositions.** D- and L-lactate were measured by using enzymatic assays.<sup>26</sup> The same value of enantiomeric excess was found for the lactate present in the reaction mixture before workup and for the zinc lactate isolated as product. This observation indicates that no enantiomeric enrichment occurs during workup. Assays for other materials were based on Mosher's method,<sup>27,28</sup> chemical shifts for

the OCH<sub>3</sub> protons of the diastereomeric forms of RCH(OCOC\*(CF<sub>3</sub>)(OCH<sub>3</sub>)Ph)CO<sub>2</sub>CH<sub>3</sub> differed by ~0.1 ppm, and those for the CO<sub>2</sub>CH<sub>3</sub> group by ~0.03 ppm. At 250 MHz, the latter gave base line separations and were used to determine enantiomeric excess. The absolute configurations of the compounds were assigned by comparison with spectra of compounds derived from resolved or enriched  $\alpha$ -hydroxy methyl esters. The methyl esters used were prepared by using a procedure described below for methyl 2-hydroxypentanoate. Control experiments carried out with the methyl esters of mandelic and *p*-chloromandelic acids showed that the procedure for the determination of the enantiomeric compositions does not result in enantiomeric enrichment.

**Methyl 2-Hydroxypentanoate.** The ester was prepared by placing the barium salt (0.5 g) of 2-hydroxypentanoic acid in MeOH (20 mL), adding acetyl chloride (0.5 mL), and heating the mixture under reflux for 30 min. The mixture was cooled, filtered to remove BaCl<sub>2</sub>, and concentrated under reduced pressure. Ether (10 mL) was added, residual BaCl<sub>2</sub> removed by filtration, and solvent removed under vacuum. The resulting liquid was used directly in determination of enantiomeric excess.

**Acknowledgment.** We thank Dr. Stephen Moore for obtaining the 250-MHz <sup>1</sup>H NMR spectra.

**Registry No.** (*p*-Chlorophenyl)glyoxal, 4998-15-6; phenylglyoxal, 1074-12-0; (*p*-methoxyphenyl)glyoxal, 1076-95-5; methylglyoxal, 78-98-8; propylglyoxal, 7332-93-6; (*R*)-4-chloro- $\alpha$ -hydroxybenzeneacetic acid 1/2 Ca, 79083-97-9; (*R*)- $\alpha$ -hydroxybenzeneacetic acid, 611-71-2; (*R*)-4-methoxy- $\alpha$ -hydroxybenzeneacetic acid 1/2 Ca, 79083-98-0; (*R*)- $\alpha$ -hydroxypropanoic acid 1/2 Zn, 18975-98-9; (*R*)- $\alpha$ -hydroxypentanoic acid 1/2 Ba, 79083-99-1.

(28) The method of Reuben for determining enantiomeric purity of unmodified  $\alpha$ -hydroxy carboxylic acids led to formation of gels (Reuben, *J. Am. Chem. Soc.* 1980, 102, 2232-2237). The resolution of the method is such that it is only useful for mixtures containing significant quantities of both enantiomers. Mixtures of Eu(dcm)<sub>3</sub> (McCreary, M. D.; Lewis, D. W.; Wernick, D. L.; Whitesides, G. M. *J. Am. Chem. Soc.* 1974, 96, 1038-1054) and compounds of the structure RCHO<sub>2</sub>CH<sub>3</sub> gave shifted spectra but no usable splitting of enantiomeric signals.

(27) Dale, J. A.; Dull, D. L.; Mosher, H. S. *J. Org. Chem.* 1969, 34, 2543-2549.

## Isolation of a Diazoalkane Intermediate in the Photoc Bamford-Stevens Reaction

Alex Nickon\* and Pamela St. John Zurer

Department of Chemistry, The Johns Hopkins University, Baltimore, Maryland 21218

Received June 1, 1981

Arylsulfonylhydrazones decompose in alkaline media on thermolysis (Bamford-Stevens reaction) or on UV irradiation. The *thermal* method is known to produce a diazo hydrocarbon, which gives products by carbenic and/or cationic routes according to the reaction conditions. In contrast, mechanistic information on the *photoc* decomposition is lacking. We now report a successful isolation of a diazo hydrocarbon from direct photolysis of a tosylhydrazone sodium salt and present evidence that the diazo compound is an intermediate in the overall *photoc* Bamford-Stevens reaction. Our key substrate was spiro[12.11]tetracosan-13-one (5a), synthesized from cyclododecanone by a three-step sequence involving reductive coupling, epoxidation of the derived olefin, and rearrangement of the oxirane. Photoc Bamford-Stevens reaction of the tosylhydrazone sodium salt of 5a under different alkaline conditions was brought about by UV irradiation through a Vycor filter (i.e.,  $\lambda >$  ca. 220 nm). The products were (*E*)-spiroalkene 11 (75-80%), (*Z*)-spiroalkene 12 (10-14%), and saturated hydrocarbons (10-11%). The alkanes were not individually identified but are expected products of intramolecular C-H insertions by a carbene. By using lower energy light (i.e., Pyrex filter;  $\lambda >$  ca. 270 nm), we isolated instead 13-diazospiro[12.11]tetracosane (14) in 95% yield. This diazo hydrocarbon was characterized spectroscopically, by elemental analysis, and by an independent synthesis. Then, on separate irradiation with Vycor-filtered light, 14 gave a product mixture, which, after correction for a small contribution from cationic pathways, contained *E* olefin 11, *Z* olefin 12, and saturated hydrocarbons in a ratio of 78:12:11, the same as that from the uninterrupted *photoc* Bamford-Stevens reactions. We also found that *thermal* Bamford-Stevens conditions give markedly different product ratios, viz., 90% 11, ca. 0% 12, and 10% alkanes. These differences suggest that the "carbene" (or diazo hydrocarbon) derived thermally and the one derived photically behave differently with respect to the stereoselectivity of the 1,2 H shift that produces olefin.

Thermal decomposition of arylsulfonylhydrazones in alkaline media (the Bamford-Stevens reaction) provides

an important synthetic route to olefins and cyclopropanes.<sup>1,2</sup> At elevated temperatures the anions of the