

35 hexane/EtOAc): TLC analysis (65:53 hexane/EtOAc) $R_f = 0.35$; $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 7.57 (1 H, dd, $J = 14.9, 11.7$ Hz), 5.98 (1 H, d, $J = 12.1$ Hz), 5.81 (1 H, d, $J = 15.3$ Hz), 4.19 (2 H, q, $J = 7.2$ Hz), 3.97 (4 H, br s), 2.56–2.53 (2 H, m), 2.39–2.36 (2 H, m), 1.75–1.71 (4 H, m), 1.28 (3 H, t, $J = 7.2$ Hz); $^{13}\text{C NMR}$ (125 MHz, CDCl_3) δ 167.5 (s), 150.7 (d), 140.0 (d), 121.6 (d), 119.8 (s), 108.3 (s), 64.4 (t), 60.1 (t), 35.9 (t), 35.2 (t), 34.1 (t), 26.2 (t), 14.3 (q). HRMS–EI m/z : calcd for $\text{C}_{14}\text{H}_{20}\text{O}_4$ 252.1361, found 252.1369.

Acknowledgment. Financial support of this work by the National Institutes of Health (GM34927) is gratefully acknowledged. We thank Dr. Celia Kingsbury for obtaining the results given in Table 1 entry 8, the NIH (SIG 1-S10-RR06301) for NMR instrumentation funding, and the NSF (CHE-93000831) for GC-MS instrumentation funding.

JO980841C

Additions and Corrections

Vol. 63, 1998

J. Edgar Anderson*, **Anthony I. Ijeh**, **Christine Storch Daniele Casarini**, and **Lodovico Lunazzi***. Eclipsed Conformation of the Exocyclic N–CH₂ Bond in *N*-Neopentylpiperidines and the Stereodynamic Consequences As Studied by Dynamic NMR Spectroscopy and Molecular Mechanics Calculations.

Page 3312, Table 2. In column 4, numbers **2b** and **2e** should be interchanged. The data in columns 5 and 6 are correct.

JO984086V

S0022-3263(98)04986-X

Published on Web 06/25/1998