

1.5 Hz, Py), 9.26 (1H, d, $^3J = 12.5$ Hz, =CH–), 13.16 (1H, d, $^3J = 12.5$ Hz, NH); ^1H NMR (360 MHz, DMSO- d_6) δ 1.01 (6H, s, 2CH₃), 2.37 (2H, s, CH₂), 2.44 (2H, s, CH₂), 5.22 (2H, br s, NH₂), 7.04 (1H, dd, $^3J = 4.6$ and 7.8 Hz, Py), 7.27 (1H, dd, $^3J = 7.8$, $^4J = 1.4$ Hz, Py), 7.79 (1H, dd, $^3J = 4.8$, $^4J = 1.4$ Hz, Py), 9.02 (1H, d, $^3J = 12.6$ Hz, =CH–), 12.62 (1H, d, $^3J = 12.6$, NH); ^{13}C NMR (90.5 MHz, DMSO) δ 22.9 (CH₃), 30.6, 50.72 (CH₂), 51.0 (CH₂), 109.1, 121.9, 125.4, 134.1, 137.4, 138.8, 147.1 (=CH–), 195.3

(C=O), 198.9 (C=O); ^{15}N NMR (36.5 MHz, CDCl₃) $^1J(^{15}\text{N}, ^1\text{H}) = 89.9$ Hz (NH), $^1J(^{15}\text{N}, ^1\text{H}) = 78.5$ Hz.

Acknowledgment. The authors thank Professor Allan R. Pinhas of the University of Cincinnati for his help in the preparation of this paper.

JO981405T

Additions and Corrections

Vol. 62, 1997

Michael A. Calter, T. Keith Hollis, Larry E. Overman,*

Joseph Ziller, and G. Greg Zipp. First Enantioselective Catalyst for the Rearrangement of Allylic Imidates to Allylic Amides.

Page 1449, Scheme 1. The following correction to Scheme 1 should be made: **b:** R¹ = CH₂OTBDPS, R³ = *n*-Bu, R⁴ = CCl₃, R⁵ = H.

The rearrangement of **1b** is described in Mehmandoust, M.; Petit, Y.; Larchevêque, M. *Tetrahedron Lett.* **1992**, *33*, 4313.

JO984028T

10.1021/jo984028t

Published on Web 02/03/1999