

1257, 1122, 1086, 1228, 961, 883, 836, 793, 739, 723, 699 (cm^{-1});
 ^1H NMR δ 7.72–7.33 (m, 12H), 4.27 (dd, $J = 0.8, 11.6$ Hz, 1H),
3.30 (d, $J = 11.6$ Hz, 1H), 3.24 (s, 1H), 1.81 (s, 3H), 1.26 (d, $J = 0.8$ Hz, 3H), 0.79 (s, 3H); ^{13}C NMR δ 173.1 (s), 134.6, 134.5, 130.9,
130.8, 128.3, 128.2 (t), 133.9, 133.8 (s), 83.2 (s), 81.0 (t), 38.3 (s),
30.2 (q), 28.6 (q), 8.9 (t), –2.0 (q).

Acknowledgment. We thank the Deutschen Akademischen Austauschdienst for a fellowship for Md.A.H. Financial support by the Fonds der Chemischen Industrie is gratefully acknowledged.

JO990478P

Additions and Corrections

Vol. 65, 1998

**Eric D. Soli, Amy S. Manoso, Michael C. Patterson,
Philip DeShong,* David A. Favor, Ralph Hirschmann,
and Amos B. Smith, III.** Azide and Cyanide Displacements via
Hypervalent Silicate Intermediates.

Page 3171. Reference 17 should be a citation to the studies of glycosyl azides from the Györgydeák group at Lajos Kossuth University, Debrecen, Hungary reported in: Györgydeák, Z.; Szilágyi, L.; Paulsen, H. *J. Carbohydr. Chem.* **1993**, *12*, 139–163. The paper cited in ref 17 in the manuscript should have been included in ref 18 instead.

We apologize for having failed to include this citation.

JO9940042

10.021/jo9940042

Published on Web 07/23/1999