

Additions and Corrections

Vol. 64, 1999

Richard E. Taylor* and **Jaroslav Zajicek**. The Conformational Properties of Epothilone.

Page 7227. The observed coupling constant J_{obs} for H6–H7 was 4.8 Hz in CD_2Cl_2 . This value is significantly different than the observed coupling constant J_{obs} for H6–H7 in DMSO/ D_2O (9.5 Hz). This suggests that conformers other than conformer A (such as conformer B) are more prevalent in CD_2Cl_2 .

Page 7228. Table 1 should read as follows.

Table 1. Selected Chemical Shifts, NOE's, and Calculated Distances for Epothilone A (s = Strong, m = Medium)

no.	^1H shift		atomic distance (Å)	
	(ppm, CD_2Cl_2)	NOE	conformer A	conformer B
2a	2.40	2a–7 (m)	4.49	2.25
		2a–21 (m)	2.63	4.43
3	4.18	3–6 (s)	4.75	2.24
		3–7 (m)	6.24	3.54
		3–21 (s)	2.54	2.48
		3–22 (s)	2.48	3.88
6	3.21	6–9a (m)	2.18	4.91
		6–21 (s)	2.25	2.41
		6–24 (s)	4.77	2.27
7	3.73	7–23 (s)	2.55	3.81
		7–24 (s)	2.38	2.69
8	1.70			
9a	1.40			
21	1.37	21–23 (s)	2.39	2.44
22	1.06			
23	1.15			
24	0.99			

JO0040240


10.1021/jo0040240

Published on Web 07/06/2000

Vol. 65, 2000

Hiroyuki Furuta*, Hiromitsu Maeda, and Atsuhiro Osuka*. Stability and Structure of Doubly N-Confused Porphyrins.

Page 4222–4226, Figure 3. Optimized bond distances and angles of **3-b** and bond angles of **5-a** were mistyped. Corrected structures are shown below.



JO004031O

10.1021/jo004031o

Published on Web 07/13/2000