


Chemoselective Deprotection of Cyclic *N,O*-Aminals Using Catalytic Bismuth(III) Bromide in Acetonitrile

Xin Cong,^{†,‡} Fang Hu,[‡] Ke-Gang Liu,[‡]
Qing-Jiang Liao,[†] and Zhu-Jun Yao^{*,‡}

Department of Medicinal Chemistry, China Pharmaceutical University, 24 Tong Jia Xiang Road, Nanjing 210009, China, and State Key Laboratory of Bioorganic and Natural Product Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Road, Shanghai 200032, China

yaoz@mail.sioc.ac.cn


Received February 4, 2005



Cyclic *N,O*-aminals can be chemoselectively and efficiently deprotected using a catalytic amount of bismuth(III) bromide in acetonitrile at room temperature. This selectivity was also achieved in the presence of terminal *O,O*-acetal functionality. The susceptibility of various other groups to cleavage was also investigated. This method has advantages of ease of operation and use of nontoxic and inexpensive reagents in catalytic amounts.

Serine and its derivatives are frequently used in the course of many syntheses as *N,O*-aminal-protected chiroins such as the Garner aldehyde.¹ However, the related cyclic *O,O*-acetals are also commonly used protecting groups for 1,2- or 1,3-diols. To our knowledge, cyclic acetals are usually more resistant to cleavage than the corresponding acyclic acetals. Methods of deprotecting cyclic *O,O*-acetals can often be applied to cleaving cyclic *N,O*-aminals and acyclic acetals as well. Although numerous approaches have been developed for deprotection of cyclic *O,O*-acetals,² few examples³ have been reported where *N,O*-aminals are selectively deprotected in the presence of *O,O*-acetals. Typical conditions for acid-catalyzed deprotection of acetals include DOWEX 50 W (H⁺) resin/methanol,⁴ trifluoroacetic/H₂O,⁵ *p*-TsOH/methanol,⁶ aqueous 60–80% AcOH,⁷ Amberlyst 15/methanol, or Amberlyst 15/acetone/H₂O.⁸ However, many of these involve strong acids, corrosive reagents, and elevated temperatures. Under such conditions, selectivity in cleav-

SCHEME 1



age between cyclic *O,O*-acetals and *N,O*-aminals would be difficult.

Recently, considerable effort has been focused on developing mild, selective methods for acetal deprotection. Several methods have been reported for cleaving acetals and ketals under nearly neutral conditions, wherein mild Lewis acids are often adopted instead of strong acids. For example, PdCl₂(MeCN) in moist acetonitrile or in acetone can deprotect acetals and ketals efficiently. However, these hydrolyses are not consistently reproducible and must be shielded from light.⁹ Selective deprotection of acyclic acetals has been reported using Bi(NO₃)₃·5H₂O in CH₂Cl₂.¹⁰ Several cyclic *O,O*-acetals have been deprotected utilizing catalytic Bi(OTf)₃·4H₂O in refluxing THF/H₂O (4:1).¹¹ Bismuth(III) compounds are attractive because they have suitable acidity yet are nontoxic, easy to handle, and inexpensive.¹²

We report herein that bismuth(III) bromide in acetonitrile is a highly efficient catalytic system for selective deprotection of cyclic *N,O*-aminals (Scheme 1). Since few mild methods exist for the selective deprotection of *N,O*-aminals, we examined the use of this system on a variety of other functionalities. It was found that the method proved to be highly efficient and easy to apply; furthermore, chemoselective deprotection of cyclic *N,O*-aminals was achieved in the presence of cyclic *O,O*-acetals. The reactions proceeded very well in acetonitrile with a catalytic amount of bismuth(III) bromide at room temperature.

Protected cyclic *N,O*-aminals were conveniently prepared from L-Garner's aldehyde¹³ through reduction and protection of the resultant primary alcohols. As shown

(4) (a) Ho, P.-T. *Tetrahedron Lett.* **1978**, *19*, 1623–1626. (b) Devel, L.; Hamon, L.; Becker, H.; Thellend, A.; Vidal-Cros, A. *Carbohydr. Res.* **2003**, *338*, 1591–1602.

(5) (a) Ellison, R. A.; Lukenbach, E. R.; Chiu, C.-W. *Tetrahedron Lett.* **1975**, *16*, 499–502. (b) Zou, B.; Wei, J.-J.; Cai, G.-R.; Ma, D.-W. *Org. Lett.* **2003**, *5*, 3503–3506. (c) Yves, L.; Brian, J. F.; Julian, A.; Francoise, P.; Joshua, R. *J. Org. Chem.* **1986**, *51*, 789–793.

(6) (a) Okamoto, H.; Hara, O.; Makino, K.; Hamada, Y. *J. Org. Chem.* **2002**, *67*, 9210–9215. (b) Akitami, I.; Makoto, U.; Sadao, S. *Tetrahedron Lett.* **1977**, *18*, 3473–3476.

(7) (a) Marvin, L. L.; John, J.-S. *J. Org. Chem.* **1969**, *34*, 3505–3512. (b) Alessandro, D.; Giandomenico, M.; Alberto, M. *J. Org. Chem.* **2002**, *67*, 4475–4486.

(8) (a) Chun, J.; He, L.-L.; Byun, H.-S.; Robert, B. *J. Org. Chem.* **2000**, *65*, 7634–7640. (b) Cappola, G. M. *Synthesis* **1984**, 1021–1022.

(9) (a) Bruce, H. L.; Daniel, P.; Joseph, M.; Hiyoshizo, K. *Tetrahedron Lett.* **1985**, *26*, 705–708. (b) Kadota, I.; Kawada, M.; Muramatsu, Y.; Yamamoto, Y. *Tetrahedron: Asymmetry* **1997**, *8*, 3887–3893.

(10) Kyle, J. E.; Michael, S. P.; Laura, C. W.; Ram, S. M. *J. Org. Chem.* **2000**, *65*, 8399–8401.

(11) Marc, D. C.; Dusan, S.; Sarapa, R.; Laura, C. W.; Ram, S. M. *J. Org. Chem.* **2002**, *67*, 1027–1030.

(12) Nicholas, M. L.; Laura C. W.; Ram, S. M. *Tetrahedron* **2002**, *58*, 8373–8397.

(13) (a) Giuliano, D. M.; Maria, C. D. G.; Francesco, M.; Domenico, M.; Giovanni, Z. *Synthesis* **1995**, 1155–1158. (b) Philip, G.; Jung, M. P. *J. Org. Chem.* **1987**, *52*, 2361–2364. (c) Alexander, M.; Richard, J. K. T.; Robert, J. W.; Norman, L. *Synthesis* **1994**, 31–33.

[†] China Pharmaceutical University.

[‡] Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences.

(1) Coppola, G. M.; Schuster, H. F. *Asymmetric Synthesis—Construction of Chiral Molecules Using Amino Acids*; John Wiley & Sons: New York, 1987.

(2) Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 3rd ed.; John Wiley & Sons: New York, 1999.

(3) Marshall, J. A.; Beaudoin, S. *J. Org. Chem.* **1996**, *61*, 581–586.

TABLE 1. Deprotection of Cyclic *N,O*-Aminals Using BiBr₃ in MeCN

substrates products


Entry	Substrate ^a	BiBr ₃ (mol%)/Time ^b	Product ^a	Yield (%) ^c
1	3 : R ¹ =Cbz; R ² =OMOM	10%/5h	4 : R ¹ =Cbz; R ² =OMOM	97%
2	5 : R ¹ =Cbz; R ² =OTBDMS	10%/8h	6 : R ¹ =Cbz; R ² =OH	85%
3	7 : R ¹ =Cbz; R ² =OAc	10%/4h	8 : R ¹ =Cbz; R ² =OAc	97%
4	9 : R ¹ =Boc; R ² =OMOM	10%/7h	10 : R ¹ =Boc; R ² =OMOM	99%
5	11 : R ¹ =Boc; R ² =OTBDMS	10%/7h	12 : R ¹ =Boc; R ² =OH	85%
6	13 : R ¹ =Boc; R ² =OAc	10%/1h	14 : R ¹ =Boc; R ² =OAc	94%
7	15 : R ¹ =Boc; R ² =OTBDPS	10%/6h	16 : R ¹ =Boc; R ² =OTBDPS	92%
8		20%/7h		90%

^a All substrates and products were fully characterized by spectroscopic methods (see Supporting Information). ^b Reaction was carried out following the general procedure described in this paper. ^c Isolated yields.

in Table 1, deprotection of the cyclic *N,O*-aminals was achieved in excellent yields. Amine protecting groups such as Boc and Cbz were stable under the conditions used. MOM (entries 1 and 4) and TBDPS (entry 7) groups, as well as acetate (entries 3 and 6) and unsaturated ester functionalities (entry 8), were also stable under the reaction conditions. However, the TBDMS groups (entries 2 and 5) were simultaneously cleaved using this protocol.¹³ The reaction rate decreased when the preferred solvent (MeCN) was replaced with acetone or CH₂-Cl₂. In the latter cases, significant quantities of unreacted starting materials were indicated by TLC monitoring. It is noteworthy here that the reaction rates could be accelerated by adding a catalytic amount of H₂O such that reaction times could be reduced to 1 h. However, under these conditions the reaction system becomes weakly acidic.¹⁴

To ascertain the reactivity of *O,O*-acetals under the above conditions, the reaction of cyclic *O,O*-acetal **19** was

SCHEME 2



examined (Scheme 2). Treatment of **19** with 10 mol % BiBr₃ in MeCN at room temperature for 48 h afforded only 16% yield of diol **20**. The results were slightly improved (up to 35% yield) when 20 mol % BiBr₃ was used at room temperature. However, the reactions led

(14) Joginder, S. B.; Jame, V.; Joel, S.; Oljan, R.; Thomas, B. *Tetrahedron Lett.* **2000**, *41*, 6021–6024.

to more complex mixtures if the amount of BiBr₃ was increased to 50 mol % or the reaction temperature was elevated to reflux. These results indicated that selective deprotection of the cyclic *N,O*-aminals using a lower amount of BiBr₃ in MeCN at room temperature might be achieved in the presence of cyclic *O,O*-acetals. Thus, substrate **21** embodying both a cyclic *N,O*-aminal and a cyclic *O,O*-acetal was prepared from alcohol **22**¹⁵ and examined (Scheme 2). Treatment of **21** with 20 mol % BiBr₃ in MeCN for 4 h at room temperature afforded alcohol **22** (69%) and triol **23** (21%). These results indicate that BiBr₃ in MeCN may be a practical choice for selective removal of *N,O*-aminals in the presence of *O,O*-acetals.

In summary, we have demonstrated that the use of catalytic amounts (10–20 mol %) of bismuth(III) bromide in acetonitrile at room temperature provides a mild and effective method for selective deprotection of cyclic *N,O*-aminals. These cyclic *N,O*-aminals can be converted to the corresponding amino alcohols in satisfactory to excellent yields in the presence of MOM, TBDPS, acetate, and cyclic *O,O*-acetal groups. Advantages of this method include the simplicity of procedure and the use of inexpensive, nontoxic reagents in catalytic amounts.

Experimental Section

General Procedure for Deprotection of Cyclic *N,O*-Aminals. To a solution of cyclic *N,O*-aminal **1** (1.0 mmol) in MeCN (10 mL) was added bismuth(III) bromide (45 mg, 0.1 mmol) at room temperature. The reaction mixture was stirred at room temperature, and the progress of the reaction was monitored by TLC. When all starting material had disappeared, the reaction mixture was quenched by adding saturated aqueous NaHCO₃ (5 mL). The layers were separated, and the aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated. The residue was purified by flash column chromatography on silica gel to afford the pure product **2**.

Benzyl (*S*)-1-Hydroxy-3-(methoxymethoxy)propan-2-yl-carbamate (4**).** To a solution of (*S*)-benzyl 4-((methoxymethoxy)methyl)-2,2-dimethyloxazolidine-3-carboxylate (**3**) (56 mg, 0.18 mmol) in MeCN (1.8 mL) was added bismuth(III) bromide (8 mg, 0.018 mmol) at room temperature. The reaction mixture was stirred at room temperature for 5 h. When all starting material had disappeared, the reaction mixture was quenched by adding

saturated aqueous NaHCO₃ (2 mL). The layers were separated, and the aqueous layer was extracted with ethyl acetate (3 × 10 mL). The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated in vacuo to give the crude product, which was purified by flash column chromatography on silica gel (petroleum ether/ethyl acetate = 2:1) to give **4** (47 mg, 97%) as a white wax. [α]_D²²: 1.4 (c 1.18, CH₂Cl₂). IR (neat): 3360, 1945, 2883, 1719, 1561, 1500, 1213, 1147, 1030 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ 2.69 (br, 1H), 3.34 (s, 3H), 3.63–3.89 (m, 5H), 4.61 (s, 2H), 5.10 (brs, 2H), 5.45 (d, *J* = 7.2 Hz, 1H), 7.31–7.37 (m, 5H) ppm. ¹³C NMR (75 MHz, CDCl₃): δ 52.1, 55.4, 63.2, 66.9, 67.9, 96.81, 128.1 (2C), 128.2, 128.5 (2C), 136.3, 156.4 ppm. ESIMS (*m/z*, %): 292.2 (M + Na⁺, 100%). HR-ESI-MS calcd for C₁₃H₁₉NO₅Na (M + Na⁺), 292.1155; found, 292.1151.

***N*-(*R*)-2-Hydroxy-1-((4*R*,5*S*)-2,2-dimethyl-5-((*R*)-2,2-dimethyl-1,3-dioxolan-4-yl)-1,3-dioxolan-4-yl)ethyl)acetamide (**22**)¹⁵ and Triol **23**.** Compound **21** (100 mg, 0.29 mmol) was converted to amino alcohol **22** and triol **23** by the above general procedure using 20 mol % BiBr₃. The crude products were separated by flash column chromatography on silica gel (petroleum ether/ethyl acetate = 4:1) to give pure alcohol **22** (61 mg, 69%) as a colorless oil and triol **23** (16 mg, 21%) as a colorless wax. Data for **22**: [α]_D²⁴: 14.4 (c 1.36, CHCl₃). ¹H NMR (300 MHz, CDCl₃): δ 1.38 (s, 6H), 1.41 (s, 3H), 1.43 (s, 3H), 2.02 (s, 3H), 3.20 (br, 1H), 3.84–3.93 (m, 4H), 4.01–4.06 (m, 1H), 4.08–4.10 (m, 2H), 4.18 (dd, *J* = 8.4 Hz, 1H) ppm. Data for **23**: [α]_D²⁴: 15.6 (c 1.30, CH₃OH). IR (neat): 3318, 2988, 2939, 2887, 1653, 1555, 1374, 1239, 1214, 1081, 1060 cm⁻¹. ¹H NMR (300 MHz, CD₃OD): δ 1.36 (s, 3H), 1.38 (s, 3H), 1.98 (s, 3H), 3.51–3.81 (m, 5H), 3.92 (dd, *J* = 7.2 Hz, 1H), 4.02–4.08 (m, 1H), 4.13 (dd, *J* = 7.5 Hz, 1H), 6.42 (brs, 1H) ppm. ¹³C NMR (75 MHz, CD₃OD): δ 23.2, 27.9, 28.0, 56.6, 62.4, 65.0, 75.1, 80.2, 80.5, 111.3, 173.8 ppm. ESI-MS (*m/z*, %): 264.1 (M + H⁺, 20%), 286.1 (M + Na⁺, 100%). HR-ESI-MS calcd for C₁₁H₂₁NO₆Na (M + Na⁺), 286.1261; found, 286.1260.

Acknowledgment. This work was financially supported by the Major State Basic Research and Development Program (G2000077500), NSFC (20425205 and 20321202), the Chinese Academy of Sciences (KGCX2-SW-209), and Shanghai Municipal Commission of Science and Technology (03DZ19203 and 04DZ14901). The authors thank Dr. Terrence R. Burke, Jr. for his help with manuscript preparation.

Supporting Information Available: Detailed experimental procedures and characterization data for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO050228A

(15) Liu, K.-G.; Yan, S.; Wu, Y.-L.; Yao, Z.-J. *J. Org. Chem.* **2002**, *67*, 6758–6763.