


Vol. 70, 2005

**Miriam Benitez, Gerhard Bringmann, Michael Dreyer, Hermenegildo Garcia,\* Heiko Ihmels,\* Michael Waidelich, and Kathrin Wissel.** Design of a Chiral Mesoporous Silica and Its Application as a Host for Stereoselective Di- $\pi$ -methane Rearrangements.

Page 2318. Scheme 2. One methylene group was missing in the structure of the silylating reagent above the arrow. The correct structure should be as follows:



In addition, compound **6** (*N,N'*-bis-[3-(triethoxysilyl)propylaminocarbonyl]-*trans*-(1*R*,2*R*)-(-)-cyclohexadiamine) in Scheme 2 was previously prepared: Moreau, J. J. E.; Vellutini, L.; Chi Man, M. W.; Bied, C. *J. Am. Chem. Soc.* **2001**, *123*, 1509–1510.

Spectroscopic data of compound **6**: IR (KBr disk)  $\nu$  3330, 1635, and 1600  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR  $\delta$  ( $\text{CDCl}_3$ )  $\delta$  0.6 (t,  $\text{SiCH}_2$ ), 1.2 (t,  $\text{CH}_3$ ), 1.3–2.1 (m,  $\text{CH}_2$ ), 2.8–3.7 (m, NCH and  $\text{NCH}_2$ ), 3.8 (t,  $\text{OCH}_2$ ), 4.9 (m, NH), and 5.3 (NH);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.4 ( $\text{CH}_2$ ), 18.3 ( $\text{CH}_3$ ), 23.5 ( $\text{CH}_2$ ), 24.8 ( $\text{CH}_2$ ), 33.2 ( $\text{CH}_2$ ), 43.0 ( $\text{CH}_2$ ), 54.7 (CH), 58.1 ( $\text{CH}_2$ ), 159.0 (C=O).

JO056026W

10.1021/jo056026w

Published on Web 05/18/2005