


## Structural Effects on the $\beta$ -Scission Reaction of Tertiary Arylcarbinyloxy Radicals. The Role of $\alpha$ -Cyclopropyl and $\alpha$ -Cyclobutyl Groups

Massimo Bietti,\* Giacomo Gente, and Michela Salamone

Dipartimento di Scienze e Tecnologie Chimiche, Università "Tor Vergata", Via della Ricerca Scientifica, 1 I-00133 Rome, Italy

bietti@uniroma2.it


Received May 2, 2005



A product and time-resolved kinetic study on the reactivity of tertiary arylcarbinyloxy radicals bearing  $\alpha$ -cyclopropyl and  $\alpha$ -cyclobutyl groups has been carried out. Both the 1-cyclopropyl-1-phenylethoxy (**1** $\cdot$ ) and  $\alpha,\alpha$ -dicyclopropylphenylmethoxy (**2** $\cdot$ ) radicals undergo  $\beta$ -scission to give cyclopropyl phenyl ketone as the major or exclusive product with rate constants higher than that measured for the cumyloxy radical. It is proposed that in the transition state for  $\beta$ -scission of **1** $\cdot$  and **2** $\cdot$ , formation of the C=O double bond is assisted by overlap with the C–C bonding orbitals of the cyclopropane ring. With tertiary arylcarbinyloxy radicals bearing  $\alpha$ -cyclobutyl groups such as the 1-cyclobutyl-1-phenylethoxy (**4** $\cdot$ ) and 1-cyclobutyl-1-phenylpropoxy (**5** $\cdot$ ) radicals, the fragmentation regioselectivity is essentially governed by the stability of the radical formed by  $\beta$ -scission. Accordingly, **4** $\cdot$  undergoes exclusive C-cyclobutyl bond cleavage to give acetophenone, whereas with **5** $\cdot$ , competition between C-cyclobutyl and C-ethyl bond cleavage, leading to propiophenone and cyclobutylphenyl ketone in a 2:1 ratio, is observed.

Recently, while investigating the *O*-neophyl rearrangement of ring-substituted 1,1-diaryloxy radicals, we observed that among the radicals studied the cyclopropyl[bis(4-methoxyphenyl)]methoxy radical displayed the lowest reactivity, undergoing 1,2-aryl shift with  $k = 2.6 \times 10^5 \text{ s}^{-1}$ .<sup>1</sup> No evidence for the formation of products deriving from C-cyclopropyl  $\beta$ -scission but only of the rearrangement product cyclopropyl 4-methoxyphenyl ketone was obtained. Quite interestingly, the rearrangement rate constant measured for this radical is lower than the rate constant measured previously for C-methyl  $\beta$ -scission of the 4-methoxycumyloxy radical,  $k = 1.0 \times 10^6 \text{ s}^{-1}$  (Scheme 1: An = 4-MeOC<sub>6</sub>H<sub>4</sub>),<sup>2,3</sup> leading to the suggestion that in alkoxy radicals C-cyclopropyl  $\beta$ -scis-

**SCHEME 1**



sion occurs at a significantly lower rate than C-Me  $\beta$ -scission, in line with the slightly higher stability of the methyl radical as compared to the cyclopropyl one (on the basis of the CH<sub>3</sub>–H and *c*-C<sub>3</sub>H<sub>5</sub>–H BDEs: 105.0 and 106.3 kcal mol<sup>-1</sup>, respectively).<sup>4,5</sup>

C–C  $\beta$ -scission is one of the most important reactions of alkoxy radicals, and accordingly, this reaction has

(1) Aureliano Antunes, C. S.; Bietti, M.; Ercolani, G.; Lanzalunga, O.; Salamone, M. *J. Org. Chem.* **2005**, *70*, 3884–3891.


(2) Baciocchi, E.; Bietti, M.; Salamone, M.; Steenken, S. *J. Org. Chem.* **2002**, *67*, 2266–2270.

(3) Unfortunately, it was not possible to study the reactivity of the 1,1-di(4-methoxyphenyl)ethoxy radical as the instability of the precursor 1,1-di(4-methoxyphenyl)ethanol did not allow the synthesis of the corresponding *tert*-butyl peroxide (see ref 1).

(4) Luo, Y.-R. *Handbook of Bond Dissociation Energies in Organic Compounds*; CRC Press LLC: Boca Raton, FL, 2003.

(5) For a critical discussion on the relative stabilities of alkyl radicals, see refs 6 and 7.


## SCHEME 2




been thoroughly investigated<sup>8–16</sup> and has found increasing application in synthetically useful procedures.<sup>17–21</sup> With tertiary alkoxy radicals bearing different alkyl groups it is possible, at least in principle, to obtain three different C–C bond fragmentations (Scheme 2), and consequently, the intramolecular selectivity in the  $\beta$ -scission reactions of alkoxy radicals has been the subject of detailed investigation.

The main conclusion of these studies was that cleavage generally leads to the most stable possible alkyl radical, and it was also shown that other factors such as the stability of the product ketone (in some  $\beta$ -scission reactions leading to the same radical R•)<sup>8a,13b</sup> or, when dealing with cycloalkoxy radicals, the release of ring strain associated to ring opening<sup>13a,c,15,22–24</sup> may play an important role. Along this line, following our continuous interest in the role of structural effects on the  $\beta$ -scission reactions of alkoxy radicals and taking advantage of the fact that arylcarbinoyloxy radicals display an absorption band in the visible region of the spectrum,<sup>2,25</sup> it seemed

## CHART 1



## SCHEME 3



- 1a: R<sub>1</sub> = Me, R<sub>2</sub> = cyp  
 2a: R<sub>1</sub> = R<sub>2</sub> = cyp  
 3a: R<sub>1</sub> = R<sub>2</sub> = Me  
 4a: R<sub>1</sub> = Me, R<sub>2</sub> = cyb  
 5a: R<sub>1</sub> = Et, R<sub>2</sub> = cyb  
 6a: R<sub>1</sub> = R<sub>2</sub> = cyb

particularly interesting to study the reactivity of tertiary arylcarbinoyloxy radicals bearing one or two  $\alpha$ -cyclopropyl groups to test whether the presence of these groups influences the reactivity toward  $\beta$ -scission of these radicals. For this purpose, we have carried out a product and time-resolved kinetic study on the reactivity of the 1-cyclopropyl-1-phenylethoxy and dicyclopropylphenylmethoxy radicals (**1•** and **2•**, respectively) whose structures are shown in Chart 1. The results obtained for these radicals have been compared with those obtained analogously for the cumyloxy radical (**3•**).

The study has been also extended to tertiary arylcarbinoyloxy radicals bearing one or two  $\alpha$ -cyclobutyl groups (radicals **4•–6•** in Chart 1). However, because time-resolved experiments have shown that **4•** displays a reactivity which exceeds the time resolution of the ns laser flash photolysis apparatus employed (see below), with radicals **5•** and **6•**, which can be reasonably expected to display a higher reactivity than **4•**, only product studies have been carried out.

## Results

**Product Studies.** Arylcarbinoyloxy radicals **1•–6•** have been generated photochemically by visible light irradiation of CH<sub>2</sub>Cl<sub>2</sub> solutions containing the parent arylalkanols (**1a–6a**), (diacetoxy)iodobenzene (DIB), and I<sub>2</sub>. It is well established that under these conditions the DIB/I<sub>2</sub> reagent converts alcohols into intermediate hypoidites which are then photolyzed to give alkoxy radicals,<sup>18,19,22,26</sup> precursors of the observed reaction products. With arylalkanols **1a–6a**, the alkoxy radicals so formed are expected to undergo  $\beta$ -scission to give the corresponding aryl ketones (Scheme 3: cyp = cyclopropyl; cyb = cyclobutyl).

Argon-saturated CH<sub>2</sub>Cl<sub>2</sub> solutions containing arylalkanols **1a–6a** (10 mM), DIB (22 mM), and I<sub>2</sub> (10 mM) were irradiated with visible light ( $\lambda_{\text{max}} \approx 480$  nm) at  $T = 20$  °C. The irradiation time was chosen in such a way as

(6) Coote, M. L.; Pross, A.; Radom, L. *Org. Lett.* **2003**, *5*, 4689–4692.

(7) Matsunaga, N.; Rogers, D. W.; Zavitsas, A. A. *J. Org. Chem.* **2003**, *68*, 3158–3172.

(8) (a) Newcomb, M.; Daublain, P.; Horner, J. H. *J. Org. Chem.* **2002**, *67*, 8669–8671. (b) Horner, J. H.; Choi, S.-Y.; Newcomb, M. *Org. Lett.* **2000**, *2*, 3369–3372.

(9) (a) Nakamura, T.; Watanabe, Y.; Suyama, S.; Tezuka, H. *J. Chem. Soc., Perkin Trans. 2* **2002**, 1364–1369. (b) Nakamura, T.; Busfield, W. K.; Jenkins, I. D.; Rizzardo, E.; Thang, S. H.; Suyama, S. *J. Org. Chem.* **2000**, *65*, 16–23.

(10) Wilsey, S.; Dowd, P.; Houk, K. N. *J. Org. Chem.* **1999**, *64*, 8801–8811.

(11) (a) Avila, D. V.; Brown, C. E.; Ingold, K. U.; Luszytk, J. *J. Am. Chem. Soc.* **1993**, *115*, 466–470. (b) Neville, A. G.; Brown, C. E.; Rayner, D. M.; Ingold, K. U.; Luszytk, J. *J. Am. Chem. Soc.* **1989**, *111*, 9269–9270.

(12) (a) Banks, J. T.; Scaiano, J. C. *J. Am. Chem. Soc.* **1993**, *115*, 6409–6413. (b) Baignée, A.; Howard, J. A.; Scaiano, J. C.; Stewart, L. C. *J. Am. Chem. Soc.* **1983**, *105*, 6120–6123. (c) Mendenhall, G. D.; Stewart, L. C.; Scaiano, J. C. *J. Am. Chem. Soc.* **1982**, *104*, 5109–5114.

(13) (a) Walling, C.; Clark, R. T. *J. Am. Chem. Soc.* **1974**, *96*, 4530–4533. (b) Walling, C. *Pure Appl. Chem.* **1967**, *15*, 69–80. (c) Walling, C.; Padwa, A. *J. Am. Chem. Soc.* **1963**, *85*, 1593–1597.

(14) (a) Bacha, J. D.; Kochi, J. K. *J. Org. Chem.* **1965**, *30*, 3272–3278. (b) Kochi, J. K. *J. Am. Chem. Soc.* **1962**, *84*, 1193–1197.

(15) Greene, F. D.; Savitz, M. L.; Osterholtz, F. D.; Lau, H. H.; Smith, W. N.; Zanet, P. M. *J. Org. Chem.* **1963**, *28*, 55–64.

(16) Gray, P.; Williams, A. *Chem. Rev.* **1959**, *59*, 239–328.

(17) Yoshimitsu, T.; Sasaki, S.; Arano, Y.; Nagaoka, H. *J. Org. Chem.* **2004**, *69*, 9262–9268.

(18) Boto, A.; Hernández, R.; Montoya, A.; Suárez, E. *Tetrahedron Lett.* **2004**, *45*, 1559–1563. Francisco, C. G.; Gonzalez, C. C.; Paz, N. R.; Suárez, E. *Org. Lett.* **2003**, *5*, 4171–4173. Alonso-Cruz, C. R.; Kennedy, A. R.; Rodriguez, M. S.; Suárez, E. *Org. Lett.* **2003**, *5*, 3729–3732. Suárez, E.; Rodriguez, M. S. In *Radicals in Organic Synthesis*; Renaud, P.; Sibi, M. P., Eds.; Wiley-VCH: Weinheim, Germany, 2001; Vol. 2, pp 440–454.

(19) Togo, H.; Katohgi, M. *Synlett.* **2001**, *5*, 565–581.

(20) (a) Zhang, W. In *Radicals in Organic Synthesis*; Renaud, P.; Sibi, M. P., Eds.; Wiley-VCH: Weinheim, Germany, 2001; Vol. 2, pp 234–245. (b) Dowd, P.; Zhang, W. *Chem. Rev.* **1993**, *93*, 2091–2115.

(21) Yet, L. *Tetrahedron* **1999**, *55*, 9349–9403.

(22) Aureliano Antunes, C. S.; Bietti, M.; Lanzalunga, O.; Salamone, M. *J. Org. Chem.* **2004**, *69*, 5281–5289.







(23) (a) Beckwith, A. L. J.; Hay, B. P. *J. Am. Chem. Soc.* **1989**, *111*, 2674–2681. (b) Beckwith, A. L. J.; Hay, B. P. *J. Am. Chem. Soc.* **1989**, *111*, 230–234.

(24) Bietti, M.; Lanzalunga, O.; Salamone, M. *J. Org. Chem.* **2005**, *70*, 1417–1422.

(25) (a) Avila, D. V.; Ingold, K. U.; Di Nardo, A. A.; Zerbetto, F.; Zgierski, M. Z.; Luszytk, J. *J. Am. Chem. Soc.* **1995**, *117*, 2711–2718. (b) Avila, D. V.; Luszytk, J.; Ingold, K. U. *J. Am. Chem. Soc.* **1992**, *114*, 6576–6577.

(26) Courtneidge, J. L.; Luszytk, J.; Pagè, D. *Tetrahedron Lett.* **1994**, *35*, 1003–1006.

**TABLE 1.** Product Distributions and Yields Observed after Irradiation of Argon-Saturated CH<sub>2</sub>Cl<sub>2</sub> Solutions Containing Arylalkanols **1a–6a**, DIB, and I<sub>2</sub><sup>a</sup>

substrate	products	yield (%)
<b>1a</b>		45 <sup>b</sup>
<b>2a</b>		24
<b>3a</b>		15
<b>4a</b>		> 99 <sup>c</sup> 65 <sup>d</sup>
<b>5a</b>		51 <sup>d,e</sup>
<b>6a</b>		56 <sup>d</sup>


<sup>a</sup> [DIB] = 22 mM, [I<sub>2</sub>] = 10 mM, [substrate] = 10 mM, irradiation wavelength = 480 nm, irradiation time = 15 min, *T* = 20 °C. <sup>b</sup> Product yields: cyclopropyl phenyl ketone 45%, acetophenone ≤ 0.5%. <sup>c</sup> Irradiation time = 2 min. Under these conditions, complete conversion of the substrate (**4a**) was observed. <sup>d</sup> Irradiation time = 1 min, using only 4 of the 10 15-W lamps available. <sup>e</sup> Product yields: propiophenone 34%, cyclobutyl phenyl ketone 17%.

to avoid complete substrate conversion. After workup of the reaction mixture, the reaction products were identified by GC (comparison with authentic samples) and GC-MS and were quantitatively determined, together with the unreacted substrate by GC, using bibenzyl as internal standard. The product distributions and yields observed after irradiation of argon-saturated CH<sub>2</sub>Cl<sub>2</sub> solutions containing substrates **1a–6a**, DIB, and I<sub>2</sub> are collected in Table 1.

The reactions of α,α-dicyclopropylphenylmethanol (**2a**) and 2-phenyl-2-propanol (**3a**) led to the formation of cyclopropyl phenyl ketone and acetophenone, respectively, as the exclusive reaction product. With 1-cyclopropyl-1-phenylethanol (**1a**), formation of cyclopropyl phenyl ketone as the major product, accompanied by traces of acetophenone, was instead observed, and a lower limit for the product ratio cyclopropyl phenyl ketone/acetophenone ≥ 90 was determined.


The reactions of 1-cyclobutyl-1-phenylethanol (**4a**) and α,α-dicyclobutylphenylmethanol (**6a**) led to the formation of acetophenone and cyclobutyl phenyl ketone, respectively, as the exclusive reaction product. With 1-cyclobutyl-1-phenylpropanol (**5a**), formation of propiophenone and cyclobutyl phenyl ketone in a 2:1 ratio was observed.

**Time-Resolved Kinetic Studies.** Alkoxy radicals **1•**, **2•**, and **4•** have been generated in MeCN by 266-nm laser flash photolysis (LFP) of the parent *tert*-butyl peroxides **1p**, **2p**, and **4p** (equation 1: cyp = cyclopropyl; cyb = cyclobutyl), which in turn have been synthesized by reaction of arylalkanols **1a**, **2a**, and **4a** with *tert*-butyl hydroperoxide in the presence of *p*-toluenesulfonic acid according to a slight modification of a previously de-



**FIGURE 1.** Time-resolved absorption spectra observed after 266-nm LFP of **1p** (10.4 mM) in an argon-saturated MeCN solution at 192 ns (empty circles), 352 ns (filled circles), and 1.3 μs (empty squares) after the 8 ns, 10 mJ laser pulse. Insets: (a) Zoom-in of the 260–600 nm region. (b) First-order decay of radical **1•** monitored at 485 nm. (c) Corresponding first-order buildup of absorption at 240 nm, assigned to the product cyclopropyl phenyl ketone.

scribed procedure (see Supporting Information).<sup>2,27</sup> To obtain information on the effect of solvent polarity on the β-scission reaction, radicals **1•** and **2•** have been also generated in MeCN/H<sub>2</sub>O 1:1 and CF<sub>3</sub>CH<sub>2</sub>OH.



**1p:** R<sub>1</sub> = Me, R<sub>2</sub> = cyp

**2p:** R<sub>1</sub> = R<sub>2</sub> = cyp

**4p:** R<sub>1</sub> = Me, R<sub>2</sub> = cyb


The cumyloxy radical **3•** was instead generated in MeCN and CF<sub>3</sub>CH<sub>2</sub>OH by 266-nm LFP of dicumyl peroxide (5.0 × 10<sup>-3</sup> M) as described previously.<sup>25</sup>

In MeCN, alkoxy radicals **1•** and **2•** display a broad absorption band in the visible region of the spectrum centered around 485 nm, which is unaffected by oxygen, in line with previous observations on the spectral properties and reactivity toward oxygen of the cumyloxy radical **3•**.<sup>2,11a,25,28</sup> The visible absorption band undergoes a first-order decay, leading in both cases to a corresponding strong increase in absorption in the UV region of the spectrum, assigned, on the basis of the product studies described above, to the product cyclopropyl phenyl ketone formed by β-scission in radicals **1•** and **2•**. Quite interestingly, as compared to MeCN, in MeCN/H<sub>2</sub>O 1:1 and CF<sub>3</sub>-CH<sub>2</sub>OH, a red-shift in the position of the visible absorption band maximum of radicals **1•–3•** of ≈ 30 nm is observed.<sup>2</sup>

As an example, Figure 1 displays the time-resolved absorption spectra observed after LFP of an argon-

(27) Hendrickson, W. H.; Nguyen, C. C.; Nguyen, J. T.; Simons, K. T. *Tetrahedron Lett.* **1995**, *36*, 7217–7220.

(28) Actually, the time-resolved absorption spectrum of the α,α-dicyclopropylphenylmethoxy radical (**2•**) recorded in MeCN (see Figure S1 in the Supporting Information) shows a very broad absorption band extending from approximately 550 nm down to the UV region, characterized by maxima at ≈ 490, 420, and 360 nm.



**FIGURE 2.** Time-resolved absorption spectra observed after 266-nm LFP of dicumyl peroxide (5.0 mM) in an argon-saturated  $\text{CF}_3\text{CH}_2\text{OH}$  solution at 48 (filled circles), 120 (empty circles), 258 (filled squares), and 810 ns (empty squares) after the 8 ns, 10 mJ laser pulse. Insets: (a) Zoom-in of the 260–600 nm region. (b) First-order decay of radical  $3^*$  monitored at 515 nm. (c) Corresponding first-order buildup of absorption at 250 nm, assigned to the product acetophenone.

saturated MeCN solution containing peroxide **1p**, showing after 192 ns (empty circles) a broad absorption band centered at 485 nm, which is assigned to the radical  $1^*$ . This species undergoes a first-order decay (inset b) accompanied by a corresponding buildup of optical density at 240 nm (inset c) assigned to cyclopropyl phenyl ketone (an isosbestic point is visible at 290 nm) which, accordingly, is characterized by an absorption maximum at this wavelength. As expected, both the 485-nm and the 240-nm bands were not affected by the presence of oxygen. Also visible is a relatively stable absorption band centered at 280 nm which, on the basis of the comparison with literature data,<sup>2,29</sup> is assigned to the *tert*-butoxy radical, formed according to eq 1.<sup>30</sup>

The time-resolved absorption spectra observed after LFP of an argon-saturated MeCN solution containing peroxide **2p** are instead displayed in Figure S1 (see Supporting Information).

Figure 2 displays the time-resolved absorption spectra observed after LFP of an argon-saturated  $\text{CF}_3\text{CH}_2\text{OH}$  solution containing dicumyl peroxide. The spectrum recorded after 48 ns is characterized by a broad absorption band in the visible region centered at 515 nm and by a band in the UV region between 300 and 350 nm, which are assigned to the cumyloxy radical  $3^*$ .<sup>2,25</sup> As compared to the spectrum of  $3^*$  recorded in MeCN,<sup>2,25</sup> a 30-nm red-shift in the position of the visible absorption band maximum is observed (see later). This species undergoes a first-order decay (inset b) accompanied by a corresponding buildup of optical density at 250 nm (inset c) and 280 nm assigned to acetophenone (an isosbestic point is visible at 300 nm).

With peroxide **4p**, the time-resolved absorption spectrum observed after LFP of an argon-saturated MeCN solution did not show an absorption band in the visible

**TABLE 2.** Visible Absorption Band Maximum Wavelengths, Decay, and Buildup Rate Constants for Alkoxy Radicals ( $1^*$ – $4^*$ ) Measured at  $T = 25^\circ\text{C}$

radical	conditions		$\lambda_{\text{max}}/\text{nm}^a$	$k_t/\text{s}^{-1b}$	$k_f/\text{s}^{-1c}$
$1^*$	MeCN	Ar	485	$3.0 \times 10^6$	$3.2 \times 10^6$
	MeCN	$\text{O}_2$	485	$3.1 \times 10^6$	$3.0 \times 10^6$
	MeCN/ $\text{H}_2\text{O}^d$	Ar	515	$9.3 \times 10^6$	$8.9 \times 10^6$
	$\text{CF}_3\text{CH}_2\text{OH}$	Ar	515	$2.1 \times 10^7$	$2.2 \times 10^7$
$2^*$	MeCN	Ar	490 <sup>e</sup>	$1.3 \times 10^6$	$1.3 \times 10^6$
	MeCN	$\text{O}_2$	490 <sup>e</sup>	$1.1 \times 10^6$	$1.4 \times 10^6$
	MeCN/ $\text{H}_2\text{O}^d$	Ar	490 <sup>e</sup>	$4.6 \times 10^6$	$4.5 \times 10^6$
	$\text{CF}_3\text{CH}_2\text{OH}$	Ar	<sup>f</sup>	$1.6 \times 10^7$	<sup>g</sup>
$3^*$	MeCN	Ar	485	$7.1 \times 10^5$	
	$\text{CF}_3\text{CH}_2\text{OH}$	Ar	515	$6.1 \times 10^6$	
$4^*$	MeCN	Ar			$\geq 8 \times 10^7$ <sup>h</sup>

<sup>a</sup> Visible absorption band maximum of the alkoxy radical. <sup>b</sup> Measured following the decay of the alkoxy radical at the visible absorption band maximum. <sup>c</sup> Measured following the buildup of the product aryl ketone at 240 or 250 nm. <sup>d</sup> MeCN/ $\text{H}_2\text{O}$  1:1 (v/v). <sup>e</sup> See ref 28 and Figure S1 in the Supporting Information. <sup>f</sup>  $\lambda_{\text{max}}$  between 480 and 530 nm. The low intensity of the visible absorption band did not allow a precise determination of the band maximum under these conditions. <sup>g</sup> A fast buildup was observed, but because of an initial bleaching in the kinetic traces, it was not possible to measure the buildup rate constant. <sup>h</sup> Measured at  $T = -10^\circ\text{C}$ . Under these conditions, the radical was too reactive to be studied by ns LFP and, on the basis of the buildup at 250 nm, only a lower limit could be determined.

region, but only the formation, immediately after the laser pulse, of a stable species absorbing around 250 nm assigned on the basis of the results of product studies described above to acetophenone formed after C-cyclobutyl  $\beta$ -scission in  $4^*$ .

The reactivity of alkoxy radicals  $1^*$  and  $2^*$  was measured spectrophotometrically by monitoring the decrease in optical density at their visible absorption maxima and the corresponding buildup in the UV region because of the formation of cyclopropyl phenyl ketone, which were found to follow in all cases first-order kinetics. The corresponding rate constants thus obtained are reported in Table 2. Radical  $4^*$  was instead too reactive to be studied by ns LFP and, on the basis of the buildup of the product acetophenone at 250 nm described above, even at  $T = -10^\circ\text{C}$  only a lower limit for its reactivity could be determined ( $k \geq 8 \times 10^7 \text{ s}^{-1}$ ).

In the same table, the decay rate constants for the cumyloxy radical  $3^*$  measured at 485 and 515 nm in MeCN and  $\text{CF}_3\text{CH}_2\text{OH}$ , respectively, are also included.

## Discussion


The results of product studies collected in Table 1 show that the reaction of 1-cyclopropyl-1-phenylethanol (**1a**) leads to the formation of cyclopropyl phenyl ketone and acetophenone in a  $\geq 90$  ratio. This result clearly indicates that in  $1^*$  cleavage of the C-methyl bond is strongly favored over that of the C-cyclopropyl one, as expected on the basis of the higher stability of the methyl radical as compared to the cyclopropyl one discussed above.<sup>4,5</sup> However, since the difference in BDE between  $\text{CH}_3\text{--H}$  and *c*- $\text{C}_3\text{H}_5\text{--H}$  has been reported to be around 1 kcal  $\text{mol}^{-1}$ ,<sup>4</sup> the large product ratio observed in the reaction of **1a** suggests that the stability of the radical formed after  $\beta$ -scission is not the only factor which governs the fragmentation regioselectivity in  $1^*$ .

(29) Tsentelovich, Y. P.; Kulik, L. V.; Gritsan, N. P.; Yurkovskaya, A. V. *J. Phys. Chem. A* **1998**, *102*, 7975–7980.

(30) The slight increase in absorption observed at 280 nm is attributed to the formation of cyclopropyl phenyl ketone, which, in addition to the 240-nm band described above, is also characterized by a weak absorption band centered at 276 nm.

Quite surprisingly, the results collected in Table 2 show that as compared to **3**<sup>•</sup>, an increase in the rate of  $\beta$ -scission is observed with both **1**<sup>•</sup> and **2**<sup>•</sup>. In particular, taking into account the results of product studies, it appears that both C-methyl bond cleavage in **1**<sup>•</sup> and C-cyclopropyl bond cleavage in **2**<sup>•</sup> occur faster than C-methyl bond cleavage in **3**<sup>•</sup>, whereas C-cyclopropyl bond cleavage is significantly slower in **1**<sup>•</sup> than in **2**<sup>•</sup>. These observations clearly indicate that the presence of an  $\alpha$ -cyclopropyl group plays an important role in the alkoxy radical fragmentation reaction.

In this context, it is well-known that cyclopropyl substituents strongly stabilize carbocations by interaction of the cyclopropyl C–C bonding orbitals with the vacant p orbital of the carbocation.<sup>31</sup> These orbitals, because of angle strain, have an abnormal amount of p character, and thus when the cyclopropyl group adopts a bisected conformation (with respect to the nodal plane of the vacant p orbital),<sup>32</sup> effective overlap may occur. Accordingly, the efficacy of cyclopropane as a neighboring group in solvolytic reactions is well documented.<sup>34</sup>



Along this line, it is reasonable to propose that in the transition state for C-methyl  $\beta$ -scission of **1**<sup>•</sup> and C-cyclopropyl  $\beta$ -scission of **2**<sup>•</sup>, formation of the C=O double bond is assisted by overlap with the C–C bonding orbitals of the cyclopropane ring. An analogous assistance is clearly not possible for C-cyclopropyl  $\beta$ -scission in **1**<sup>•</sup> and for C-methyl  $\beta$ -scission in the cumyloxy radical **3**<sup>•</sup>.

Quite importantly, cyclopropyl participation involving overlap between the C–C bonding orbitals of the cyclopropane ring and developing p orbitals has been also proposed in other reactions, such as for example, the thermolysis of azo compounds<sup>35</sup> and the decarbonylation of tricyclic alkenones.<sup>36,37</sup>

By combining the product ratio determined for the reaction of **1a** (Table 1) with the decay rate constant

(31) See, for example: Olah, G. A.; Prakash Reddy, V.; Surya Prakash, G. K. *Chem. Rev.* **1992**, *92*, 69–95 and references therein.

(32) The preference for the bisected conformation over the perpendicular one in cyclopropyl substituted cations has been confirmed in several studies (see, for example, ref 33).

(33) Childs, R. F.; Faggiani, R.; Lock, C. J. L.; Mahendran, M.; Zweep, S. D. *J. Am. Chem. Soc.* **1986**, *108*, 1692–1693. Buss, V.; Gleiter, R.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1971**, *93*, 3927–3933. Baldwin, J. E.; Fogelsong, W. D. *J. Am. Chem. Soc.* **1968**, *90*, 4303–4310.


(34) See, for example: Roberts, D. D. *J. Org. Chem.* **1999**, *64*, 1341–1343. Lowry, T. H.; Richardson, K. S. *Mechanism and Theory in Organic Chemistry*, 3rd ed.; Harper & Row: New York, 1987, and references therein.

(35) Schmidt, H.; Schweig, A.; Trost, B. M.; Neubold, H. B.; Scudder, P. H. *J. Am. Chem. Soc.* **1974**, *96*, 622–624. Allred, E. L.; Voorhees, K. J. *J. Am. Chem. Soc.* **1973**, *95*, 620–621. Allred, E. L.; Johnson, A. L. *J. Am. Chem. Soc.* **1971**, *93*, 1300–1301.


(36) Battiste, M. A.; Visnick, M. *Tetrahedron Lett.* **1978**, *19*, 4771–4774. Halton, B.; Battiste, M. A.; Rehberg, R.; Deyrup, C. L.; Brennan, M. E. *J. Am. Chem. Soc.* **1967**, *89*, 5964–5965.

(37) Cyclopropyl participation has been also proposed in the ring-opening of tricyclic cyclopropyl ketones, rationalized in terms of the overlap between the scissile cyclopropane C–C bond and the adjacent p orbital (see ref 38).

#### SCHEME 4



#### SCHEME 5



measured for **1**<sup>•</sup> (Table 2), the partial rate constants for C-methyl bond cleavage assisted by the presence of an adjacent cyclopropyl group ( $k_{\text{Me(ass)}} = 3.1 \times 10^6 \text{ s}^{-1}$ ) and for unassisted C-cyclopropyl bond cleavage ( $k_{\text{cyp(unass)}} = \leq 3.4 \times 10^4 \text{ s}^{-1}$ ) can be derived (Scheme 4).<sup>39,40</sup>

The rate constants measured for **2**<sup>•</sup> and **3**<sup>•</sup> (Table 2), corrected for the statistical factor, provide instead the assisted rate constant for C-cyclopropyl bond cleavage ( $k_{\text{cyp(ass)}} = 6.5 \times 10^5 \text{ s}^{-1}$ ) and the unassisted rate constant for C-methyl bond cleavage ( $k_{\text{Me(unass)}} = 3.6 \times 10^5 \text{ s}^{-1}$ ), respectively (Scheme 5).

Since C-methyl cleavage in **1**<sup>•</sup> and C-cyclopropyl cleavage in **2**<sup>•</sup>, as well as C-cyclopropyl cleavage in **1**<sup>•</sup> and C-methyl cleavage in **3**<sup>•</sup>, lead to the formation of the same ketone (cyclopropyl phenyl ketone and acetophenone, respectively), comparison between the pertinent rate constants provides the kinetic effect of the stability of the alkyl radical on the  $\beta$ -scission reaction.

In both the assisted and unassisted reaction, C-methyl cleavage is favored over C-cyclopropyl cleavage, in line with the slightly higher stability of the methyl radical as compared to the cyclopropyl one.<sup>4,5</sup> However, the reactivity ratio decreases on going from the unassisted ( $k_{\text{Me(unass)}}/k_{\text{cyp(unass)}} = \geq 10.6$ ) to the assisted reaction ( $k_{\text{Me(ass)}}/k_{\text{cyp(ass)}} = 4.8$ ), clearly indicating that, as compared to the methyl radical, formation of the cyclopropyl radical benefits to a greater extent from cyclopropyl assistance. The same conclusion arises from the comparison between the assisted and unassisted  $\beta$ -scission reactions leading

(38) Monti, S. A. *J. Org. Chem.* **1970**, *35*, 380–383. Monti, S. A.; Bucheck, D. J.; Shepard, J. C. *J. Org. Chem.* **1969**, *34*, 3080–3084.

(39) The product ratio for **1a** has been determined in  $\text{CH}_2\text{Cl}_2$ , while the decay rate constant for **1**<sup>•</sup> has been measured in MeCN. In this respect, no significant difference in product ratio should be observed on going from  $\text{CH}_2\text{Cl}_2$  to MeCN as the solvent is expected to behave in the same way on the two competing fragmentation reactions of the alkoxy radical.

(40) The decay rate constants for radicals **1**<sup>•</sup>–**3**<sup>•</sup> have been also measured in  $\text{CH}_2\text{Cl}_2$  as  $3.3 \times 10^6$ ,  $1.4 \times 10^6$ , and  $1.3 \times 10^6 \text{ s}^{-1}$ , respectively; values which are slightly higher than the corresponding ones measured in the more polar solvent MeCN (see Table 2). However, as the rate constants for  $\beta$ -scission of alkoxy radicals are known to decrease by decreasing solvent polarity (see, for example, refs 2, 11a, and 29), the observation of an increase in rate on going from MeCN to  $\text{CH}_2\text{Cl}_2$  can be rationalized in terms of the contribution to the decay rate constant of a hydrogen abstraction reaction from the solvent  $\text{CH}_2\text{Cl}_2$ .

to the same radical, where, neglecting differences in stability between the product ketones, cyclopropyl assistance is relatively more important in C-cyclopropyl than in C-methyl cleavage ( $k_{\text{cyp(ass)}}/k_{\text{cyp(unass)}} = \geq 19.1$  and  $(k_{\text{Me(ass)}}/k_{\text{Me(unass)}} = 8.6)$ .

As observed previously for the cumyloxy radical **3**<sup>•</sup>,<sup>2,11a</sup> also the rate constants for  $\beta$ -scission of radicals **1**<sup>•</sup> and **2**<sup>•</sup> increase by increasing solvent polarity. Solvent effects on the rate constant for  $\beta$ -scission of the cumyloxy radical have been studied in detail,<sup>11a</sup> and the increase in rate observed with increasing solvent polarity has been interpreted in terms of the stabilization of the transition state for  $\beta$ -scission through increased solvation of the incipient carbonyl product. Quite interestingly, a fairly good correlation between  $\log k_{\beta}$  and the Dimroth-Reichardt  $E_{\text{T}}^{\text{N}}$  parameter has been reported for the  $\beta$ -scission reactions of the cumyloxy radical **3**<sup>•</sup> in several solvents.<sup>11a</sup> In this context, also the  $k_{\beta}$  value measured for **3**<sup>•</sup> in CF<sub>3</sub>CH<sub>2</sub>OH (characterized by  $E_{\text{T}}^{\text{N}} = 0.898$ )<sup>41</sup> fits very well into the  $\log k_{\beta}$  versus  $E_{\text{T}}^{\text{N}}$  correlation.<sup>42</sup>

Going from MeCN to MeCN/H<sub>2</sub>O 1:1 and CF<sub>3</sub>CH<sub>2</sub>OH results in a red-shift in  $\lambda_{\text{max}}$  (vis) of approximately 30 nm for radicals **1**<sup>•</sup>–**3**<sup>•</sup>.<sup>2</sup> The observation of this effect in H<sub>2</sub>O and CF<sub>3</sub>CH<sub>2</sub>OH but not in other protic solvents such as AcOH, *t*BuOH, and dipolar aprotic ones such as MeCN requires some comment. On the basis of Abraham's  $\alpha_2^{\text{H}}$  parameters,<sup>43</sup> CF<sub>3</sub>CH<sub>2</sub>OH and AcOH are characterized by similar hydrogen bond acidities ( $\alpha_2^{\text{H}} = 0.567$  and 0.550, respectively), values that are significantly higher than that determined for H<sub>2</sub>O ( $\alpha_2^{\text{H}} = 0.353$ ). Clearly, this solvent property alone cannot be invoked to explain the observed solvent effect on  $\lambda_{\text{max}}$  (vis). On the other hand, both H<sub>2</sub>O and CF<sub>3</sub>CH<sub>2</sub>OH are characterized by  $E_{\text{T}}^{\text{N}}$  values that are significantly higher than those measured for MeCN, *t*BuOH, and AcOH,<sup>44</sup> suggesting that the combination of different solute/solvent interaction forces (well represented by this empirical parameter) contributes to this effect.<sup>45</sup>

The reactions of 1-cyclobutyl-1-phenylethanol (**4a**) and  $\alpha,\alpha$ -dicyclobutylphenylmethanol (**6a**) led to the formation of acetophenone and cyclobutyl phenyl ketone, respectively, as the exclusive reaction product. With 1-cyclobutyl-1-phenylpropanol (**5a**), formation of propiophenone and cyclobutyl phenyl ketone in a 2:1 ratio was instead observed. Clearly in **4**<sup>•</sup>, cleavage of the C-cyclobutyl bond is strongly favored over that of the C-methyl one, as expected on the basis of the large difference in stability between the cyclobutyl and the methyl radical (on the

basis of the *c*-C<sub>4</sub>H<sub>7</sub>–H and CH<sub>3</sub>–H BDEs: 96.8 and 105.0 kcal mol<sup>-1</sup>, respectively).<sup>4</sup> Differences in radical stability can also be invoked to explain the fragmentation regioselectivity observed with **5**<sup>•</sup>, as the *c*-C<sub>4</sub>H<sub>7</sub>–H BDE is reported to be lower than the CH<sub>3</sub>CH<sub>2</sub>–H one (96.8 and 100.5 kcal mol<sup>-1</sup>, respectively).<sup>4</sup> An observation that is also supported by kinetic data where, however, even though the high reactivity of **4**<sup>•</sup> did not allow the determination of its fragmentation rate constant but only of a lower limit ( $k_{\beta} \geq 8 \times 10^7 \text{ s}^{-1}$  at  $T = -10 \text{ }^{\circ}\text{C}$ , see Table 2), the comparison of this limit with the rate constant measured for C-ethyl  $\beta$ -scission in the 2-(4-methylphenyl)-2-butoxy radical ( $k_{\beta} = 1.1 \times 10^7 \text{ s}^{-1}$  at  $T = -15 \text{ }^{\circ}\text{C}$ )<sup>24</sup> indicates that in tertiary arylcarbinyloxy radicals cleavage of the C-cyclobutyl bond is favored over that of the C-ethyl one.

In conclusion, by means of product and time-resolved kinetic studies, convincing evidence has been provided for cyclopropyl participation in the  $\beta$ -scission reactions of tertiary arylcarbinyloxy radicals bearing  $\alpha$ -cyclopropyl groups. It is proposed that  $\beta$ -scission proceeds through a transition state where formation of the C=O double bond is assisted by overlap with the C–C bonding orbitals of the cyclopropane ring. The kinetic data clearly show that both C-methyl and C-cyclopropyl bond cleavage are significantly accelerated by cyclopropyl assistance which, moreover, is relatively more important in the latter than in the former  $\beta$ -scission reaction. In particular, C-cyclopropyl bond cleavage, which is expected to be a relatively slow reaction, when assisted by the presence of an additional cyclopropyl group becomes faster than C-methyl cleavage in the cumyloxy radical. With tertiary arylcarbinyloxy radicals bearing  $\alpha$ -cyclobutyl groups, the fragmentation regioselectivity is essentially governed by the stability of the radical formed by  $\beta$ -scission, with cleavage of the C-cyclobutyl bond being favored over that of the C-ethyl one and being the exclusive fragmentation pathway when in competition with C-methyl bond cleavage.

## Experimental Section

**Materials.** Spectroscopic grade MeCN and CF<sub>3</sub>CH<sub>2</sub>OH were used as received. CH<sub>2</sub>Cl<sub>2</sub> was purified prior to use by column chromatography over basic alumina. (Diacetoxy)iodobenzene (DIB), iodine,  $\alpha,\alpha$ -dicyclopropylphenylmethanol (**2a**), and 2-phenylpropan-2-ol (**3a**) were of the highest commercial quality available and were used as received. Dicumyl peroxide was recrystallized twice from methanol.

Details of the synthesis of arylalkanols **1a**, **4a**–**6a**, and of the *tert*-butyl arylalkyl peroxides (**1p**, **2p**, and **4p**) are given in the Supporting Information. The purity of the arylalkanols (**1a**–**6a**) employed in the product studies was always  $\geq 99\%$ .

**Product Analysis.** All the reactions were carried out under an argon atmosphere. Irradiations were performed with visible light (4–10  $\times$  15 W lamps with emission between 400 and 550 nm,  $\lambda_{\text{max}} \approx 480 \text{ nm}$ ). The reactor was a cylindrical flask equipped with a water cooling jacket thermostated at  $T = 20 \text{ }^{\circ}\text{C}$ . Irradiation times were chosen in such a way as to avoid complete substrate consumption. In a typical experiment, a solution of the alcohol (10 mM) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) containing (diacetoxy)iodobenzene (22 mM) and iodine (10 mM) was irradiated for times ranging between 1 and 15 min under Ar bubbling. The reaction mixture was then poured into water and extracted with CH<sub>2</sub>Cl<sub>2</sub> (2  $\times$  10 mL). The combined organic layers were washed with a 10% aqueous thiosulfate solution (2  $\times$  30 mL) and water (2  $\times$  30 mL) and were dried over

(41) Reichardt, C. *Solvents and Solvent Effects in Organic Chemistry*, 3rd ed.; Wiley-VCH: Weinheim, Germany, 2003.

(42) As it has been pointed out that the  $E_{\text{T}}^{\text{N}}$  parameter is mainly related to the solvent anion solvating ability (see ref 41, pp 462–463), the observation of a  $\log k_{\beta}$  versus  $E_{\text{T}}^{\text{N}}$  correlation reasonably reflects the development of negative charge on the oxygen atom of the forming carbonyl compound. We thank a referee for this suggestion.

(43) Abraham, M. H.; Grellier, P. L.; Prior, D. V.; Duce, P. P.; Morris, J. J.; Taylor, P. J. *J. Chem. Soc., Perkin Trans. 2* **1989**, 699–711.

(44) (CH<sub>3</sub>)<sub>2</sub>COH, MeCN, AcOH, CF<sub>3</sub>CH<sub>2</sub>OH, and H<sub>2</sub>O are characterized by the following  $E_{\text{T}}^{\text{N}}$  values: 0.389, 0.460, 0.648, 0.898, and 1.000, respectively (see ref 41).

(45) In this connection, it is important to point out that in water the red-shift in the cumyloxy radical  $\lambda_{\text{max}}$  (vis) is expected to be significantly greater than 30 nm. See, for example, ref 46 for a discussion on the effect of water on the 4-methoxycumyloxy radical  $\lambda_{\text{max}}$  (vis).

(46) Baciocchi, E.; Bietti, M.; Lanzalunga, O.; Steenken, S. *J. Am. Chem. Soc.* **1998**, *120*, 11516–11517.

anhydrous sodium sulfate. Reaction products were identified by GC (comparison with authentic samples) and GC-MS and were quantitatively determined by GC using bibenzyl as internal standard. Excellent mass balances ( $\geq 95\%$ ) were obtained in all experiments. Blank experiments performed in the absence of irradiation, keeping the solution in the dark, showed the formation of negligible amounts of reaction products.

**Laser Flash Photolysis Studies.** Laser flash photolysis experiments were carried out with a laser kinetic spectrometer using the fourth harmonic (266 nm) of a Q-switched Nd:YAG laser. The laser energy was adjusted to  $\leq 10$  mJ/pulse by the use of the appropriate filter. A 3-mL Suprasil quartz cell (10 mm  $\times$  10 mm) was used for all experiments. Argon- or oxygen-saturated MeCN, MeCN/H<sub>2</sub>O 1:1 (v/v), or CF<sub>3</sub>CH<sub>2</sub>OH solutions of peroxides **1p**, **2p**, and dicumyl peroxide (between  $5.0 \times 10^{-3}$  and  $1.0 \times 10^{-2}$  M,  $A_{266} \approx 1.0$ ) were used. All the experiments were carried out at  $T = 25 \pm 0.5$  °C under magnetic stirring. LFP of peroxide **4p** ( $1.0 \times 10^{-2}$  M) was instead carried out in argon-saturated MeCN solution at  $T = -10$  °C. Rate constants were obtained by averaging four to eight values each consisting

of the average of at least three laser shots and were reproducible to within 10%.

**Acknowledgment.** Financial support from the Ministero dell'Istruzione dell'Università e della Ricerca (MIUR) is gratefully acknowledged. We thank Prof. Osvaldo Lanzalunga for assistance in the synthesis and characterization of the peroxides, Prof. Basilio Pispisa and Dr. Lorenzo Stella for the use and assistance in the use of a LFP equipment, and Mr. Luigi Gastaldo (Helios Italquartz srl) for providing us with a photochemical reactor.

**Supporting Information Available:** Details on the synthesis and characterization of the precursor arylalkanols and *tert*-butyl arylalkyl peroxides. Time-resolved absorption spectra observed after LFP of peroxide **2p**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO050883I