


A High-Yield, One-Step Synthesis of *o*-Phenylene Ethynylene Cyclic Trimer via Precipitation-Driven Alkyne Metathesis

Wei Zhang, Scott M. Brombosz,[†] Jose L. Mendoza,[‡] and Jeffrey S. Moore*

Roger Adams Laboratory, Departments of Chemistry and Materials Science & Engineering, University of Illinois at Urbana–Champaign, Urbana, Illinois 61801

jsmoore@uiuc.edu


Received August 23, 2005



A shape-persistent, conjugated *o*-phenylene ethynylene cyclic trimer was prepared in one step from tetrasubstituted benzene monomer **4** in 86% isolated yield through precipitation-driven alkyne metathesis. The template-free, selective generation of the molecular triangle **5** is a thermodynamically favored process and under equilibrium control. A novel tetrameric macrocycle **7** was generated via scrambling dynamic metathesis between tricycle **5** and hexacycle **6** using this dynamic covalent chemistry.

There has been growing interest in shape-persistent, arylene ethynylene macrocycles as building blocks for molecular materials and supramolecular chemistry.¹ These macrocycles have rigid, noncollapsible hydrocarbon backbones and are able to organize in solution, the molten phase, and solid state (noncovalent nanotubes, discotic liquid crystals, guest–host complexes, etc.). Among these structures, the conjugated cyclic trimer comprising three *o*-phenylene ethynylene units (**1**) has attracted great interest.² Macrocycle **1** and its derivatives are structural units of *graphyne*.^{2a} The distance from the center of the 12-membered dehydroannulene to the center

of the carbon–carbon triple bonds is about 1.2 Å, which provides a suitable geometry for π -alkylene/transition-metal bonding interactions.^{2c} The metal can be either in or out of the plane of the macrocycle. The interesting structural and conducting properties of the transition-metal complexes derived from **1** have been extensively explored.^{2b–d}



1

Although there have been a series of phenylene ethynylene trimeric macrocycles reported to date, their preparation is mainly restricted to cross-coupling approaches, and usually only low yields are obtained.^{3,4} The primary disadvantage of the coupling approach is that the product distribution is kinetically determined. Since these irreversible reactions do not have the ability to correct undesired bond formations, oligomer growth that overshoots the length of target macrocycles cannot contribute to the yield of the desired product.

On the other hand, given the successful examples of dynamic covalent chemistry in organic synthesis,⁵ we envisioned a one-step approach to arylene ethynylene macrocycles from monomers could be accomplished via reversible alkyne metathesis. By utilizing molybdenum catalyst that is both functional group tolerant and active near room temperature,⁶ we have accomplished one-step, gram-scale synthesis of hexameric phenylene ethynylene macrocycles.⁷ Macrocycle formation is a thermodynamically favored process under equilibrium control.⁸

Recently, Vollhardt et al. synthesized trimeric phenylene ethynylene macrocycles through alkyne metathesis catalyzed by Schrock's catalyst.⁹ Although the un-

(3) For reviews on synthesis of dehydrobenzoannulenes, see: (a) Marsden, J. A.; Palmer, G. J.; Haley, M. M. *Eur. J. Org. Chem.* **2003**, 2355–2369. (b) Youngs, W. J.; Tessier, C. A.; Bradshaw, J. D. *Chem. Rev.* **1999**, *99*, 3153–3180. (c) Haley, M. M. *Synlett* **1998**, 557–565 and ref 1f.

(4) For recent examples, see: (a) Li, Y.; Zhang, J.; Wang, W.; Miao, Q.; She, X.; Pan, X. *J. Org. Chem.* **2005**, *70*, 3285–3287. (b) Iyoda, M.; Sirinintasak, S.; Nishiyama, Y.; Vorasingha, A.; Sultana, F.; Nakao, K.; Kuwatani, Y.; Matsuyama, H.; Yoshida, M.; Miyake, Y. *Synthesis* **2004**, 1527–1531. (c) Nishinaga, T.; Miyata, Y.; Nodera, N.; Komatsu, K. *Tetrahedron* **2004**, *60*, 3375–3382. (d) Iyoda, M.; Vorasingha, A.; Kuwatani, Y.; Yoshida, M. *Tetrahedron Lett.* **1998**, *39*, 4701–4704. (e) Solooki, D.; Ferrara, J. D.; Malaba, D.; Bradshaw, J. D.; Tessier, C. A.; Youngs, W. J. *Inorg. Synth.* **1997**, *31*, 122–128.

(5) Rowan, S. J.; Cantrill, S. J.; Cousins, G. R. L.; Sanders, J. K. M.; Stoddart, J. F. *Angew. Chem., Int. Ed.* **2002**, *41*, 898–952.

(6) (a) Zhang, W.; Kraft, S.; Moore, J. S. *Chem. Commun.* **2003**, 832–833. (b) Zhang, W.; Kraft, S.; Moore, J. S. *J. Am. Chem. Soc.* **2004**, *126*, 329–335.

(7) Zhang, W.; Moore, J. S. *J. Am. Chem. Soc.* **2004**, *126*, 12796. Bunz previously reported the synthesis of hexameric macrocycles via alkyne metathesis. Under their conditions, which required high temperature (150 °C), the desired products were obtained in only 0.5–6% yields. Ge, P.-H.; Fu, W.; Herrmann, W. A.; Herdtweck, E.; Campana, C.; Adams, R. D.; Bunz, U. H. F. *Angew. Chem., Int. Ed.* **2000**, *39*, 3607–3610.

(8) Zhang, W.; Moore, J. S. *J. Am. Chem. Soc.* **2005**, *127*, 11863–11870.


[†] Summer intern from Illinois Wesleyan University.

[‡] Summer intern from Instituto Tecnológico y de Estudios Superiores de Monterrey Campus Monterrey.

(1) For reviews, see: (a) Höger, S. *Angew. Chem., Int. Ed.* **2005**, *44*, 3806–3808. (b) Höger, S. *Chem. Eur. J.* **2004**, *10*, 1320–1329. (c) Zhao, D.; Moore, J. S. *Chem. Commun.* **2003**, 807–818. (d) Yamaguchi, Y.; Yoshida, Z. *Chem. Eur. J.* **2003**, *9*, 5430–5440. (e) Bunz, U. H. F.; Rubin, Y.; Tobe, Y. *Chem. Soc. Rev.* **1999**, *28*, 107–119. (f) Haley, M. M.; Pak, J. J.; Brand, S. C. *Top. Curr. Chem.* **1999**, *201*, 81–130. (g) Moore, J. S. *Acc. Chem. Res.* **1997**, *30*, 402–413. (h) Young, J. K.; Moore, J. S. In *Modern Acetylene Chemistry*; Stang, P. J., Diederich, F., Eds.; VCH: Weinheim, 1995; Chapter 12.

(2) (a) Eickmeier, C.; Junga, H.; Matzger, A. J.; Scherhag, F.; Shim, M.; Vollhardt, K. P. C. *Angew. Chem., Int. Ed.* **1997**, *36*, 2103–2108. (b) Ferrara, J. D.; Tessier-Youngs, C.; Youngs, W. J. *J. Am. Chem. Soc.* **1985**, *107*, 6719–6721. (c) Youngs, W. J.; Kinder, J. D.; Bradshaw, J. D.; Tessier, C. A. *Organometallics* **1993**, *12*, 2406–2407. (d) Ferrara, J. D.; Djebli, A.; Tessier-Youngs, C.; Youngs, W. J. *J. Am. Chem. Soc.* **1988**, *110*, 647–649.

substituted cyclic product **1** can be obtained in 54% yield at 80 °C, methoxy substituted analogue **2** was only obtained in 28% yield even after 140 h, with 20% catalyst loading. To further improve the synthesis of trimeric phenylene ethynylene macrocycles and also to explore the scope of our newly developed synthetic protocol, we now report the efficient, one-step preparation of an alkoxy substituted *o*-phenylene ethynylene cyclic trimer using the precipitation-driven alkyne metathesis.⁷ Under equilibrium control, the thermodynamically favored cyclic trimer **5** was obtained in high yield.



Sonogashira cross-coupling¹⁰ of 1,2-diethynyl-4,5-dihexyloxybenzene (**3**)¹¹ and 4-benzoyl-4'-bromobiphenyl gave the desired tetrasubstituted benzene monomer **4** (Scheme 1). The metathesis reaction of **4** was performed in CCl₄ catalyzed by a molybdenum alkylidyne catalyst⁶ (10 mol % loading) generated in situ from the corresponding molybdenum triamide and *p*-nitrophenol. The reaction mixture was stirred for 22 h at 30 °C, during which time the precipitation of the byproduct diarylacetylene drove the reaction to completion. After removal of the precipitate, the filtrate was concentrated and the macrocyclic product was purified by column chromatography, affording triangular cycle **5** as a yellow solid with 86% isolated yield.

The obtained cyclic trimer **5** was fully characterized by ¹H NMR, ¹³C NMR, FAB-MS, GPC, and elemental analysis, which unambiguously demonstrate the desired cyclic trimer. The ¹H NMR spectrum of **5** revealed the presence of only one type of aromatic proton (δ 6.72) and the ¹³C NMR spectrum showed a unique signal corresponding to the acetylene carbon (δ 91.8), which are consistent with *D*_{3h} symmetry for the triangular structure. FAB mass spectrometry showed the molecular ion at m/z = 900.6269 (calcd 900.6268), supporting the structure of cyclic trimer **5**. GPC trace of macrocycle **5** (Figure 1) showed the monodisperse product distribution (PD = 1.00) and a M_n of 1090, further supporting the generation of the target structure. Finally, the elemental analysis result is also consistent with the expected composition of **5**.¹²

SCHEME 1. Synthesis of *o*-Phenylene Ethynylene Cyclic Trimer **5**





FIGURE 1. GPC trace of purified cyclic trimer **5** (THF, 25 °C).

The reversibility of macrocycle **5** formation was demonstrated by a scrambling experiment, in which a 1:2 mixture of two different phenylene ethynylene macrocycles (**5** and **6**¹³) was subjected to metathesis (eq 1). The reaction was conducted in CCl₄ at 30 °C for 22 h followed by FD-MS analysis (Figure 2). A tetrameric macrocycle, presumably structure **7** (m/z = 1200) having two *o*-disubstituted units from **5** (m/z = 900) and two *m*-disubstituted units from **6** (m/z = 1800) was observed, demonstrating the reversibility of macrocycle formation.

This template-free,^{14,15} selective generation of triangular cycle **5** via precipitation-driven alkyne metathesis is thus confirmed to be a thermodynamically controlled process.⁸ The initial growth of the oligomers is driven by the poor solubility of the diarylacetylene byproduct. Once the chains are sufficiently long, the oligomers can transform into corresponding cyclic oligomers ($n \geq 3$) through intramolecular metathesis with nearly complete removal of end groups. At this point, open-chain and macrocyclic intermediates can equilibrate leading predominantly to the thermodynamically more stable trimeric macrocycles.

In conclusion, the cyclic trimer **5** has been successfully synthesized by using precipitation-driven alkyne metathesis. The selective generation of cyclic trimer is a thermodynamically favored process and under equilibrium control. The successful synthesis of **5** as well as generation of a tetracycle via scrambling metathesis of tricycle **5** and hexacycle **6** further demonstrates the general scope of the one-step protocol for arylene ethynylene macrocycle preparation and the importance of dynamic covalent chemistry in constructing thermodynamically favored molecular architectures in high yields.





FIGURE 2. FD-MS of crude product from scrambling metathesis of macrocycle **5** and **6**.



Experimental Section

Macrocycle (5). In an argon-filled glovebox, a solution of molybdenum triamide⁶ (7.5 mg, 0.011 mmol) and *p*-nitrophenol (4.7 mg, 0.034 mmol) in CCl_4 (1.0 mL) was added to a solution of monomer **4** (94.3 mg, 0.11 mmol) in CCl_4 (2.1 mL). The flask was sealed and removed from the glovebox. The resulting mixture was stirred for 22 h at 30 °C. After removal of the precipitate by vacuum filtration, the filtrate was concentrated in vacuo and the macrocyclic product was purified by column

chromatography (*n*-hexane/ CH_2Cl_2 , 3/2, v/v) to give macrocycle **5** as a yellow solid (28.9 mg, 86%): mp 125–126 °C; ^1H NMR (CDCl_3 , 400 MHz) δ 6.72 (s, 6H), 3.95 (t, $J = 7.6$ Hz, 12H), 1.80 (m, 12H), 1.30–1.50 (m, 36H), 0.90 (t, $J = 6.4$ Hz, 18H); ^{13}C NMR (CDCl_3 , 100 MHz) δ 149.0, 119.6, 115.5, 91.8, 68.9, 31.5, 29.0, 25.6, 22.5, 14.0; MS (FAB) m/z 900.7 (66), 816.6 (15), 731.5 (7), 307.1 (20), 154.1 (100); HR-MS ($\text{C}_{60}\text{H}_{84}\text{O}_6$) calcd 900.6268, found 900.6269; TLC $R_f = 0.34$ (*n*-hexane/ CH_2Cl_2 , 1/1, v/v); GPC

(9) Miljanica, O. Š.; Vollhardt, K. P. C.; Whitener, G. D. *Synlett* **2003**, 29–34.

(10) (a) Sonogashira, K.; Tohda, Y.; Hagihara, N. *Tetrahedron Lett.* **1975**, 4467–4470. (b) Sonogashira, K.; Yatake, T.; Tohda, Y.; Takahashi, S.; Hagihara, N. *J. Chem. Soc., Chem. Commun.* **1977**, 291–292. (c) Sonogashira, K. In *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., Ed.; Wiley-Interscience: New York, 2002; pp 493–529.

(11) (a) Zhou, Q.; Carroll, P. J.; Swager, T. M. *J. Org. Chem.* **1994**, 59, 1294–1301. (b) Bowles, D. M.; Palmer, G. J.; Landis, C. A.; Scott, J. L.; Anthony, J. E. *Tetrahedron* **2001**, 57, 3753–3760.

(12) See the Experimental Section for detailed characterization data.

(13) Macrocycle **6** was synthesized from the corresponding *m*-dialkoxybenzene monomer via precipitation-driven alkyne metathesis (see the Supporting Information for details). On MALDI-MS of the metathesis product, only hexameric ($m/z = 1800$) and pentameric ($m/z = 1500$) macrocycles were observed. Such a macrocycle mixture was subjected to the subsequent scrambling metathesis experiment without further separation.

(14) Höger applied a template-cyclization to synthesize phenylene ethynylene macrocycles in excellent yields. However, the covalently attached templates require the introduction of specific functional groups into the cyclization precursors, and extra synthetic steps are needed for attachment and cleavage. See: (a) Höger, S. *J. Polym. Sci. Part A: Polym. Chem.* **1999**, 37, 2685–2698. (b) Höger, S.; Meckenstock, A. D. *Chem. Eur. J.* **1999**, 5, 1686–1691. (c) Höger, S.; Enkelmann, V. *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 2713–2716.

(15) Sanders utilized noncovalent template approach in synthesis of large shape-persistent rings. Cyclic porphyrin oligomers were obtained in 50–70% yield. (a) Anderson, H. L.; Sanders, J. K. M. *Angew. Chem., Int. Ed. Engl.* **1990**, 29, 1400–1403. (b) McCallien, D. W. J.; Sanders, J. K. M. *J. Am. Chem. Soc.* **1995**, 117, 6611–6612.

1090 (M_n), 1.00 (M_w/M_n). Anal. Calcd for $C_{60}H_{54}O_4$ (900.6): C, 79.96; H, 9.39. Found: C, 79.87; H, 9.44.

Acknowledgment. This material is based upon work supported by the National Science Foundation under Grant No. 0345254 and the U.S. Department of Energy, Division of Materials Sciences, under Award No. DEFG02-91ER45439, through the Frederick Seitz

Materials Research Laboratory at the University of Illinois at Urbana–Champaign.

Supporting Information Available: Characterization data for compound **3**, **4**, and precursors to **6**; experimental details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO0517803