


## On the Influence of the C2–O2 and C3–O3 Bonds in 4,6-*O*-Benzylidene-Directed $\beta$ -Mannopyranosylation and $\alpha$ -Glucopyranosylation

David Crich\* and Olga Vinogradova

Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, Illinois 60607-7061

dcrich@uic.edu

Received July 7, 2006




The synthesis of 4,6-*O*-benzylidene-protected 2-deoxyarabino-, 3-deoxyarabino-, and 3-deoxyribothio-glycosides is described and their glycosylation reactions, with activation by either 1-benzenesulfinyl piperidine/trifluoromethanesulfonic anhydride or diphenyl sulfoxide/trifluoromethanesulfonic anhydride, studied. In contrast to the corresponding 4,6-*O*-benzylidene-protected glucosyl and mannosyl donors, which are  $\alpha$ - and  $\beta$ -selective, respectively, poor diastereoselectivity is observed in all cases. The reasons for this poor selectivity are discussed in terms of the interaction between the C2–O2 and C3–O3 bonds in the glucosyl and mannosyl donors and of the influence of this interaction on the ease of formation of the intermediate glycosyl oxacarbenium ions.

### Introduction

One of the more intriguing aspects of the 4,6-*O*-benzylidene-directed  $\beta$ -mannopyranosylation reactions developed in this laboratory<sup>1–6</sup> is the change in selectivity observed with the corresponding glucopyranosyl and galactopyranosyl donors, which are  $\alpha$ -selective.<sup>4,6–9</sup> This contrast in selectivity, originally seen with donors **1** and **2**,<sup>7</sup> on activation with trifluoromethanesulfonic anhydride in the presence of a hindered non-nucleophilic base such as 2,4,6-tri-*tert*-butylpyrimidine (TTBP),<sup>10</sup> and subsequently observed with donors **3** and **4**, on activation with 1-benzenesulfinyl piperidine (BSP) and triflic anhydride,<sup>4</sup> is reflected in the reduced  $\beta$ -selectivity seen on going from the

3,4-*O*-carbonate-protected 6-deoxymanno (rhamno) donor **5** to the corresponding gluco donor **6**.<sup>11</sup>




This difference in selectivity between the 4,6-*O*-benzylidene-protected manno- and glucopyranose series has been examined computationally by Nukuda and Whitfield and their co-workers, who viewed the problem in terms of the conformations of the intermediate oxacarbenium ions and of the protonated glycosides resulting from methanol attack on the oxacarbenium ions.<sup>12</sup>

- (1) Crich, D.; Sun, S. *J. Org. Chem.* **1997**, *62*, 1198–1199.  
 (2) Crich, D.; Sun, S. *Tetrahedron* **1998**, *54*, 8321–8348.  
 (3) Crich, D.; Sun, S. *J. Am. Chem. Soc.* **1998**, *120*, 435–436.  
 (4) Crich, D.; Smith, M. *J. Am. Chem. Soc.* **2001**, *123*, 9015–9020.  
 (5) Crich, D.; Smith, M. *J. Am. Chem. Soc.* **2002**, *124*, 8867–8869.  
 (6) Crich, D.; de la Mora, M.; Vinod, A. U. *J. Org. Chem.* **2003**, *68*, 8142–8148.  
 (7) Crich, D.; Cai, W. *J. Org. Chem.* **1999**, *64*, 4926–4930.  
 (8) Bousquet, E.; Khitri, M.; Lay, L.; Nicotra, F.; Panza, L.; Russo, G. *Carbohydr. Res.* **1998**, *311*, 171–181.  
 (9) Kim, K. S.; Kang, S. S.; Seo, Y. S.; Kim, H. J.; Jeong, K.-S. *Synlett* **2003**, 1311–1314.  
 (10) Crich, D.; Smith, M.; Yao, Q.; Picione, J. *Synthesis* **2001**, 323–326.

- (11) Crich, D.; Jayalath, P. *J. Org. Chem.* **2005**, *70*, 7252–7259.

## SCHEME 1. Glycosylation Mechanism



However, it was recognized that the analysis was based on thermodynamic considerations and might not be relevant to kinetically determined stereoselectivity. It is well-known that the anomeric effect is higher in mannopyranosides than in glucopyranosides, leading to a greater thermodynamic preference for the  $\alpha$ -anomers in the mannose series,<sup>13</sup> and it is evident that the glycosylation stereochemistries obtained with donors **1–4** at reaction temperatures from  $-78$  to  $-60$  °C are kinetic in origin. Accordingly, while the geometries of the 4,6-*O*-benzylidene-protected oxocarbenium ions computed by the Nukuda and Whitfield groups are taken into account in the discussion below, the rationale for glycosylation stereochemistry presented by these authors is not further considered. Differences in reactivity of mannopyranosyl and glucopyranosyl oxocarbenium ions have also been discussed recently by Lucero and Woerpel,<sup>14</sup> in the context of a broader investigation into the synthesis of C-glycosides and the influence of substituents on the stereoselectivity of oxocarbenium ion trapping.<sup>15</sup> In the absence of a conformationally restricting group, such as the benzylidene acetal, these authors considered the tetra-*O*-benzylmannopyranosyl cation to exist predominantly in the  $^3H_4$  conformer but to react via the less populated  $^4H_3$  form, in a typical Curtin–Hammett kinetic scheme to afford the  $\alpha$ -C-glycosides in high yield. The  $\alpha$ -selective C-glycosylation of the tetra-*O*-benzylglucopyranosyl cation was also considered to take place through the  $^4H_3$  conformer.<sup>14</sup> With the benzylidene acetal in place, and the  $^3H_4$  conformer of the oxocarbenium ion excluded from consideration, the problem is simplified and only the  $^4H_3$  conformer need be considered along with the somewhat related  $B_{2,5}$  and  $^4E$  conformers preferred by the benzylidene-protected manno- and glucopyranosyl cations according to the calculations of Nukuda, Whitfield, and their co-workers.<sup>12</sup>

We currently view the mechanism of the 4,6-*O*-benzylidene-directed glycosylations, which are preformed, operationally by preactivation of the donor before addition of the acceptor, as proceeding through the intermediacy of a covalent glycosyl triflate **7**, which has been demonstrated spectroscopically to have the  $\alpha$ -configuration in both the manno and the gluco series.<sup>7,16</sup> This intermediate, which is formed rapidly and which is stable for many hours at low temperature in  $CD_2Cl_2$ , acts as a reservoir for a transient contact ion pair **8** (CIP) that is the true actual glycosylating species in the  $\beta$ -mannosylation reaction.<sup>17</sup> This




FIGURE 1. Destabilizing interaction in the contact ion pair, illustrated for the 4,6-*O*-benzylidene-protected glycosyl triflates.

contact ion pair is in turn in equilibrium with a solvent separated ion pair (SSIP) **9**, which we presume to be the source of the  $\alpha$ -glycosides (Scheme 1). The function of the benzylidene acetal is to shift the series of equilibria between **7** and the two ions as far as possible toward the covalent triflate **7**, thereby minimizing the population of the SSIP **9** and, at least in the mannose series, the formation of  $\alpha$ -glycosides. The benzylidene acetal exerts its influence through a combination of torsional factors that come into play as the covalent triflate chair conformer flattens to the oxocarbenium ion in **8**<sup>18,19</sup> and by maintaining the C5–C6 bond in the most electron-withdrawing *tg* conformer.<sup>20</sup> On this basis, the  $\alpha$ -selective glycosylations arise through the SSIP **9** and the problem is reduced to one of understanding the reasons for its greater population in the gluco series. There remains the alternative possibility that the  $\alpha$ -glucosides are formed via the intermediacy of a second contact ion pair, with the triflate counterion shielding the  $\beta$ -face of the oxocarbenium ion, which would necessarily be in equilibrium with a covalent  $\beta$ -glycosyl triflate, much as envisaged by Lemieux in his halide ion promoted  $\alpha$ -glycosylations.<sup>21</sup> While we cannot exclude this mechanism conclusively, we consider it unlikely on the grounds that our NMR experiments provided no evidence for the existence of a  $\beta$ -glycosyl triflate.<sup>7</sup>

We considered that the shift toward the SSIP **9** implied by the change from  $\beta$ - to  $\alpha$ -selectivity observed on going from the manno to the gluco series might be due to greater repulsion between the O-2 substituent and the triflate counterion in the CIP **8** in the gluco series, which would have the effect of facilitating SSIP formation (Figure 1).

Alternatively, we were intrigued by the notion that the O2–C2–C3–O3 torsional interaction plays a significant role in the observed differing selectivities between the manno and gluco series. Effectively, as the mannosyl triflate **10** collapses to the ion pair, in either the  $^4H_3$  chair **11** or the  $B_{2,5}$  conformer **12** preferred by Nukuda and Whitfield,<sup>12</sup> the O2–C2–C3–O3 torsion angle decreases (Table 1), whereas when the gluco triflate **13** is transformed into either the  $^4H_3$  chair **14** or the  $^4E$  conformer **15** favored by the computations,<sup>12</sup> the corresponding torsion angle is increased (Table 2).

Examination of the change in the O2–C2–C1–O5 torsion angle with formation of the oxocarbenium ion is also instructive. In mannose (Table 1) there is an increase in this torsion angle on going from the covalent triflate **10** to the  $^4H_3$  oxocarbenium ion **11** and maximization of this torsion angle in the  $B_{2,5}$  conformer **12** of the oxocarbenium ion. On the other hand, in glucose the O2–C2–C1–O5 torsion angle, which begins at

(12) Nukada, T.; Berces, A.; Whitfield, D. M. *Carbohydr. Res.* **2002**, *337*, 765–774.

(13) Lemieux, R. U. In *Molecular Rearrangements, Part 2*; De Mayo, P., Ed.; Wiley: New York, 1964; pp 709–769.

(14) Lucero, C. G.; Woerpel, K. A. *J. Org. Chem.* **2006**, *71*, 2641–2647.

(15) Smith, D. M.; Woerpel, K. A. *Org. Biomol. Chem.* **2006**, 1195–1201.

(16) Crich, D.; Sun, S. *J. Am. Chem. Soc.* **1997**, *119*, 11217–11223.


(17) Crich, D.; Chandrasekera, N. S. *Angew. Chem., Int. Ed.* **2004**, *43*, 5386–5389.

(18) Fraser-Reid, B.; Wu, Z. C.; Andrews, W.; Skowronski, E. *J. Am. Chem. Soc.* **1991**, *113*, 1434–1435.


(19) Andrews, C. W.; Rodebaugh, R.; Fraser-Reid, B. *J. Org. Chem.* **1996**, *61*, 5280–5289.

(20) Jensen, H. H.; Nordstrom, M.; Bols, M. *J. Am. Chem. Soc.* **2004**, *126*, 9205–9213.

(21) Lemieux, R. U.; Hendriks, K. B.; Stick, R. V.; James, K. *J. Am. Chem. Soc.* **1975**, *97*, 4056–4062.

**TABLE 1.** Torsional Angle Values for Mannosyl Triflate (**10**) and the  $^4H_3$  (**11**) and  $B_{2,5}$  (**12**) Conformers of the Mannosyl Cation<sup>22</sup>


	O2–C2–C3–O3 torsion angle (deg)	O2–C2–C1–O5 torsion angle (deg)
<b>10</b>	60	60
<b>11</b>	45	135
$\Delta$ ( <b>10</b> $\rightarrow$ <b>11</b> )	-15	+75
<b>12</b>	60	180
$\Delta$ ( <b>10</b> $\rightarrow$ <b>12</b> )	0	+120

**TABLE 2.** Torsional Angle Values for Glucosyl Triflate (**13**) and the  $^4H_3$  (**14**) and  $^4E$  (**15**) Conformers of the Glucosyl Cation


	O2–C2–C3–O3 torsion angle (deg)	O2–C2–C1–O5 torsion angle (deg)
<b>13</b>	60	180
<b>14</b>	75	105
$\Delta$ ( <b>13</b> $\rightarrow$ <b>14</b> )	+15	-75
<b>15</b>	90	120
$\Delta$ ( <b>13</b> $\rightarrow$ <b>15</b> )	+30	-60

the maximum of 180° in the covalent triflate **13**, undergoes a significant reduction regardless of whether the oxacarbenium ion exists in the  $^4H_3$  or  $^4E$  conformations **14** and **15**, respectively (Table 2). By analogy with Bols' rationale for the benzylidene effect,<sup>20</sup> it might be inferred that the higher O2–C2–C1–O5 torsion angle in the mannose oxacarbenium ions **12** and **13** maximizes the electron-withdrawing effect of the C2–O2 bond, as compared to that in the glucosyl oxacarbenium ions **14** and **15**. This in turn would contribute toward a destabilization of the mannosyl oxacarbenium ions with respect to their glucosyl counterparts, which translates into a longer lived glucosyl oxacarbenium ion and a concomitant loss of selectivity.

The effect of the O2–O3 interaction on the rate of acid-catalyzed hydrolyses of methyl glycopyranosides was one of several factors considered by Edward,<sup>23</sup> Feather and Harris,<sup>24</sup> and others<sup>25</sup> in some of the classical studies in the area, but has subsequently been overlooked. With a view to probing these effects we have investigated the glycosylation stereoselectivity of a set of 4,6-*O*-benzylidene protected 2- and 3-deoxythioglycosides and describe here the results of our investigations.

## Results


**Preparation of Donors.** The synthesis of the 2-deoxy donor **19** was straightforward and followed the protocol outlined in Scheme 2 starting from triacetyl glucal.

(22) Torsion angles were determined with ring conformations fixed according to the parameters provided by Bérces and co-workers (<http://www.sao.nrc.ca/ibs/6ring.html>). Bérces, A.; Whitfield, D. M.; Nukuda, T. *Tetrahedron* **2001**, *57*, 477–491.


(23) Edward, J. T. *Chem. Ind.* **1955**, 1102–1104.

(24) Feather, M. S.; Harris, J. F. *J. Org. Chem.* **1965**, *30*, 153–157.

(25) Overend, W. G.; Rees, C. W.; Sequeira, J. S. *J. Chem. Soc.* **1962**, 3429–3440.

**SCHEME 2.** Synthesis of the 2-Deoxy Donor **19**<sup>a</sup>

<sup>a</sup> Reagents and conditions: (a) HCl, toluene, 0 °C; (b) PhSH, (*i*-Pr)<sub>2</sub>EtN, toluene, rt, 47% over two steps; (c) NaOMe, MeOH; (d) PhCH(OMe)<sub>2</sub>, *p*-TsOH, DMF, 50 °C, 60% over two steps; (e) NaH, BnBr, THF, reflux, 73%.

**SCHEME 3.** Preparation of the 3-Deoxy Ribo Donor **25**<sup>a</sup>

<sup>a</sup> Reagents and conditions: (a) IR-120 (H<sup>+</sup>), H<sub>2</sub>O, 80 °C; (b) Ac<sub>2</sub>O, pyridine, DMAP, rt, 85% for two steps; (c) Bu<sub>3</sub>SnH, AIBN, toluene, reflux, 78%; (d) PhSH, BF<sub>3</sub>·Et<sub>2</sub>O, DCM, 0 °C, 97%,  $\alpha/\beta = 1:4$ ; (e) NaOMe, MeOH; (f) PhCH(OMe)<sub>2</sub>, *p*-TsOH, DMF, 50 °C, 96% for two steps; (g) NaH, BnBr, THF, reflux, 93%.

Obvious routes to the 3-deoxy donors beginning with the Barton deoxygenation of 1,2,5,6-di-*O*-acetoneglucopyranose<sup>26</sup> and proceeding through 3-deoxy- $\beta$ -D-ribo-hexopyranose and 1,2,4,6-tetra-*O*-acetyl-3-deoxy- $\beta$ -D-ribo-hexopyranose were not satisfactory owing to formation of furanose isomers in significant amounts.<sup>27</sup> Eventually, a synthesis was devised in which 1,2,5,6-di-*O*-acetoneglucopyranose was converted to the known 3-deoxy-3-iodoalofuranose derivative **20**,<sup>28,29</sup> which on deprotection in acidic media and subsequent acetylation gave mainly the desired pyranose form. Radical deiodination then afforded **21**<sup>27</sup> in good yield. Treatment with thiophenol and boron trifluoride afforded a 4:1 separable mixture of the anomeric thioglycosides **22** and **23**, of which only the major  $\beta$ -isomer was processed through the subsequent routine steps of the synthesis to give the 3-deoxy ribo donor **25** (Scheme 3).

Adapting a classical scheme for the formation of  $\beta$ -mannopyranosides from the corresponding  $\beta$ -glucopyranosides,<sup>30–35</sup> Dess–Martin oxidation<sup>36</sup> of **24** gave the corresponding ketone, which was stereoselectively reduced with L-Selectride to give

(26) Barton, D. H. R.; McCombie, S. W. *J. Chem. Soc., Perkin Trans. I* **1975**, 1574–1585.

(27) Withers, S. G.; Percival, M. D.; Street, I. P. *Carbohydr. Res.* **1989**, *187*, 43–66.

(28) Russell, R. N.; Weigel, T. M.; Han, O.; Liu, H.-w. *Carbohydr. Res.* **1990**, *201*, 95–114.


(29) Classon, B.; Liu, Z.; Samuelsson, B. *J. Org. Chem.* **1988**, *53*, 6126–6130.

(30) Theander, O. *Acta Chem. Scand.* **1958**, *12*, 1883–1885.

(31) Barresi, F.; Hindsgaul, O. In *Modern Methods in Carbohydrate Synthesis*; Khan, S. H., O'Neill, R. A., Eds.; Harwood Academic Publishers: Amsterdam, 1996; p 251–276.

(32) Pozsgay, V. In *Carbohydrates in Chemistry and Biology*; Ernst, B., Hart, G. W., Sinay, P., Eds.; Wiley-VCH: Weinheim, 2000; Vol. 1, p 319–343.

(33) Ito, Y.; Ohnishi, Y. In *Glycoscience: Chemistry and Chemical Biology*; Fraser-Reid, B., Kuniaki, T., Thiem, J., Eds.; Springer-Verlag: Berlin, 2001; Vol. 2, p 1589–1619.


SCHEME 4. Preparation of the 3-Deoxy Arabino Donor 27<sup>a</sup>

<sup>a</sup> Reagents and conditions: (a) Dess–Martin periodinane, DCM, rt; (b) L-Selectride, THF,  $-78\text{ }^{\circ}\text{C}$ , 81% for two steps; (c) NaH, BnBr, THF, reflux, 94%.

the 3-deoxyarabino product **26**. Benzylation then afforded the desired donor **27** (Scheme 4).

**Investigation of Key Intermediates.** Exploratory experiments conducted in  $\text{CD}_2\text{Cl}_2$  by variable-temperature  $^1\text{H}$  NMR spectroscopy revealed the 2-deoxy donor **19** to be fully activated by the BSP/ $\text{TF}_2\text{O}$  combination within minutes at  $-60\text{ }^{\circ}\text{C}$ . However, only minor signals were observed in the region anticipated<sup>16</sup> ( $\delta$  6.0–6.3) for the anomeric signal of the putative glycosyl triflate intermediate **28**. On warming to  $-30\text{ }^{\circ}\text{C}$ , one major compound, the glycol **29**,<sup>37</sup> was formed, as determined from key resonances at  $\delta$  6.38 (dd,  $J = 6.0, 1.2\text{ Hz}$ ) and  $\delta$  4.86 (dd,  $J = 6.0, 2.0\text{ Hz}$ ) indicative of the olefinic hydrogens, along with other unidentified byproducts. The result suggested the triflate **28** to be too unstable for the glycosylations in the 2-deoxy series. This behavior appears to be typical of the 2-deoxy series as a parallel experiment with donor **30**<sup>38</sup> also failed to afford an observable glycosyl triflate but provided instead tribenzyl glucal **31** on warming to  $-40\text{ }^{\circ}\text{C}$ . Activation of thioglycoside **19** with 2 equiv of diphenyl sulfoxide<sup>39,40</sup> and 1.1 equiv of triflic anhydride was also rapid and provided a putative intermediate glycosyl sulfonium ion **32** of the type observed by Gin in his studies on the activation of anomeric hemiacetals with diphenyl sulfoxide and triflic anhydride.<sup>41</sup> On warming, decomposition of this intermediate was complete by  $-50\text{ }^{\circ}\text{C}$ . In the 3-deoxy series, the chemistry reverted to the more typical pattern observed previously in our laboratory<sup>4,7,11,16,42</sup> with clean activation of both **25** and **27** by the BSP/ $\text{TF}_2\text{O}$  couple at  $-60\text{ }^{\circ}\text{C}$  in  $\text{CD}_2\text{Cl}_2$  and the formation of the putative glycosyl triflates **33** and **34**, respectively, neither of which underwent decomposition until the temperature was raised to  $-10\text{ }^{\circ}\text{C}$ .

**Glycosylation Reactions.** In view of the variable-temperature NMR experiments, couplings to the 2-deoxy donor **19** were conducted at  $-60\text{ }^{\circ}\text{C}$  with preactivation by a combination of 2 equiv of diphenyl sulfoxide and 1.1 equiv of triflic anhydride in the presence of TTBP, before addition of the acceptor alcohol.



Satisfactory yields were obtained in this manner but selectivities were poor, approaching a maximum of 4:1  $\beta/\alpha$  with a primary alcohol (Table 3). As determined by NMR spectroscopy of the crude reaction mixtures, comparable results were obtained when diphenyl sulfoxide was replaced by BSP, but isolated yields were reduced owing to difficulties in purification and are hence not reported.

TABLE 3. Glycosylation of the 2-Deoxy Donor 19

Acceptor	Product	$\alpha/\beta$ ratio <sup>a</sup>	Yield, %
		1:1.5	75
		1:1.7	75
		1:4	61
		1:1.9	30

<sup>a</sup> Determined by  $^1\text{H}$  NMR spectroscopy on the reaction mixture.

Glycosylation reactions with the 3-deoxy donors **25** and **27** were conducted with preactivation by the BSP/ $\text{TF}_2\text{O}$  couple in the presence of TTBP before addition of the acceptor and are thought to proceed via the intermediacy of the glycosyl triflates **33** and **34**, respectively, on the basis of the variable-temperature experiments. The results of these experiments, all of which proceeded with good yield but modest selectivity, are reported in Tables 4 and 5.

While the assignment of stereochemistry in the 2-deoxy series (Table 3) and in the 3-deoxy ribo series (Table 4) was straightforward based on the  $^3J_{\text{H}_1, \text{H}_2}$  coupling constants, that in the 3-deoxy arabino series relies on the magnitude of the  $^1J_{\text{C}_1, \text{H}_1}$  coupling constants.<sup>43,44</sup>

(34) Lichtenhaler, F. W.; Lergenmüller, M.; Peters, S.; Varga, Z. *Tetrahedron: Asymmetry* **2003**, *14*, 727–736.

(35) Ekborg, G.; Lindberg, B.; Lonngrén, J. *Acta Chem. Scand. B* **1972**, *26*, 3287–3292.

(36) Dess, P. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155–4156.

(37) Roush, W. R.; Sebesta, D. P.; Bennett, C. E. *Tetrahedron* **1997**, *53*, 8825–8836.

(38) Crich, D.; Ritchie, T. J. *J. Chem. Soc., Perkin Trans. 1* **1990**, 945–954.







(39) Codée, J. D. C.; van den Bos, L. J.; Litjens, R. E. J. N.; Overkleef, H. S.; van Boeckel, C. A. A.; van Boom, J. H.; van der Marel, G. A. *Tetrahedron* **2004**, *60*, 1057–1064.

(40) Codée, J. D. C.; Litjens, R. E. J. N.; Van den Bos, L. J.; Overkleef, H. S.; Van der Marel, G. *Chem. Soc. Rev.* **2005**, *34*, 769–782.

(41) Garcia, B. A.; Gin, D. Y. *J. Am. Chem. Soc.* **2000**, *122*, 4269–4279.

(42) Crich, D.; Hutton, T. K.; Banerjee, A.; Jayalath, P.; Picione, J. *Tetrahedron: Asymmetry* **2005**, *16*, 105–119.


TABLE 4. Glycosylation of the 3-Deoxy Donor 25

Acceptor	Product	$\alpha/\beta$ ratio <sup>a</sup>	Yield, %
		1.2:1	95
		4.7:1	76
		1:1.8	92

<sup>a</sup> Determined by <sup>1</sup>H NMR spectroscopy on the reaction mixture.

## Discussion

The lack of stereoselectivity observed with the 2-deoxy donor **19** (Table 3) is not surprising in view of the well-known difficulties in the stereoselective formation of 2-deoxyglycopyranosidic linkages, in general,<sup>45–47</sup> and is somewhat anticipated on the basis of the variable-temperature experiments and the manifest instability of the intermediates. These problems obviously arise from the greater lability of the 2-deoxy systems, which is the result of the absence of the electron-withdrawing C2–O2 bond and the greater facility of oxocarbenium ion formation that this affords.<sup>24,25,48–50</sup> Accordingly, no further account is taken here of the results presented in Table 3. We do note, however, the reduced  $\alpha$ -selectivity seen with the donor **19** as compared to the high  $\alpha$ -selectivity seen by Kahne and co-workers with a series of 2,6-dideoxy glycosyl sulfoxides in the course of their syntheses of ciclamycin:<sup>51,52</sup> evidently, the 4,6-*O*-benzylidene acetal is not without influence over the course of these reactions. In a similar vein, we note the  $\beta$ -selective glycosylations reported by Hashimoto on activation of the 2-deoxyglycosyl phosphite **49**, and related phosphates, with trimethylsilyl triflate at  $-94$  °C, which were considered to take place via intermediate glycosyl triflates, something that cannot be excluded on the basis of the experiments presented here.<sup>53</sup>



The results obtained with the 3-deoxy donors **25** and **27** (Tables 4 and 5) are more germane to the question in hand.

(43) Bock, K.; Pedersen, C. *J. Chem. Soc., Perkin Trans. 2* **1974**, 293–297.

(44) Duus, J. O.; Gotfredsen, C. H.; Bock, K. *Chem. Rev.* **2000**, *100*, 4589–4614.

(45) Thiem, J.; Klaffke, W. *Topics Curr. Chem.* **1990**, *154*, 285–332.







(46) Marzabadi, C. H.; Franck, R. W. *Tetrahedron* **2000**, *56*, 8385–8417.

(47) Veyrières, A. In *Carbohydrates in Chemistry and Biology*; Ernst, B., Hart, G. W., Sinay, P., Eds.; Wiley-VCH: Weinheim, 2000; Vol. 13, pp 367–405.

(48) Green, L. G.; Ley, S., V. In *Carbohydrates in Chemistry and Biology*; Ernst, B., Hart, G. W., Sinay, P., Eds.; Wiley-VCH: Weinheim, 2000; Vol. 1, pp 427–448.

(49) Zhang, Z.; Ollmann, I. R.; Ye, X.-S.; Wischnat, R.; Baasov, T.; Wong, C.-H. *J. Am. Chem. Soc.* **1999**, *121*, 734–753.

TABLE 5. Glycosylation of the 3-Deoxy Donor 27

Acceptor	Product	$\alpha/\beta$ ratio <sup>a</sup>	Yield, %
		1.9:1	85
		1.3:1	80
		3.6:1	83

<sup>a</sup> Determined by <sup>1</sup>H NMR spectroscopy on the reaction mixture.

Various 3-deoxyglycopyranosyl donors have been described in the literature, but with rare exceptions,<sup>54</sup> all carried ester protecting groups on O2 to direct the glycosylation stereochemistry<sup>55–58</sup> and are not relevant to the present discussion. Interestingly, although the coupling was carried out under very different conditions to the ones employed here, the benzyl-protected 3-deoxyglucosyl chloride **50** gave an approximately 1:1 mixture of two anomers on coupling to methyl 2,3,6-tri-*O*-benzyl- $\beta$ -D-glucopyranoside by the Lemieux chloride ion-catalyzed protocol.<sup>21,54</sup> Classical work on the acid-catalyzed hydrolysis of simple glycopyranosides revealed that removal of the C3–O3 bond has only a relatively minor accelerating effect,<sup>23–25,59</sup> as might also be inferred from more recent work on the disarming effect of ester protecting groups and their location on the pyranoside ring.

In the 3-deoxy ribo series (Table 4) there is a significant preference toward the formation of the  $\beta$ -glycosides, away from the  $\alpha$ -selectivity previously observed with the glucosyl donors **2** and **4**. This does not accord with the hypothesis of Figure 1 in which repulsion between the 2-*O*-benzyl group and the triflate counterion is responsible for a general loosening of the ion pair in the glucose series. Possibly, the move away from  $\alpha$ -selectivity with donor **25** is a reflection of a minor change in conformation of the oxocarbenium ion in the resulting CIP which facilitates  $\beta$ -face attack. As noted below, conformation effects have been shown to play a major role in the stereoselectivity of these glycosylation reactions.

In the 3-deoxyarabino series (Table 5) there is a strong shift away from the high  $\beta$ -selectivity typically observed with donors **1** and **3**. Obviously the absence of the electron-withdrawing C3–

(50) Namchuk, M. N.; McCarter, J. D.; Becalski, A.; Andrews, T.; Withers, S. G. *J. Am. Chem. Soc.* **2000**, *122*, 1270–1277.

(51) Raghavan, S.; Kahne, D. *J. Am. Chem. Soc.* **1993**, *115*, 1580–1581.

(52) Gildersleeve, J.; Smith, A.; Sakurai, D.; Raghavan, S.; Kahne, D. *J. Am. Chem. Soc.* **1999**, *121*, 6176–6182.

(53) Hashimoto, S.; Sano, A.; Sakamoto, H.; Nakajima, M.; Yanagiya, Y.; Ikegami, S. *Synlett* **1995**, 1271–1273.

(54) Bock, K.; Pedersen, H. *Acta Chem. Scand.* **1987**, *B41*, 617–628.

(55) van Dorst, J. A. L. M.; van Heusden, C. J.; Tikkanen, J. M.; Kamerling, J. P.; Vliegthart, J. F. G. *Carbohydr. Res.* **1997**, *297*, 209–227.


(56) Oscarson, S.; Svahnberg, P. *Carbohydr. Res.* **1998**, *309*, 207–212.

(57) Oscarson, S.; Tedebark, U. *Carbohydr. Res.* **1995**, *278*, 271–287.

(58) Kováč, P.; Edgar, K. J. *Carbohydr. Res.* **1990**, *201*, 79–83.

(59) Richards, G. N. *Chem. Ind.* **1955**, 228–228.

O3 bond cannot be totally neglected here, but it seems apparent that the loss of the compression of the O2–C2–C3–O3 torsion angle seen in the mannose series (Table 1) on going from the covalent triflate to the oxacarbenium ion leads to a higher population of CIP, and consequently of the SSIP, with an overall reduction in  $\beta$ -stereoselectivity. In support of this argument, we recall the mannosyl donor **51** and its rhamnosyl counterparts **52** and **53**, which adopt a half-chair conformation in which the O2–C2–C3–O3 torsion angle is 25–30°, as opposed to the 60° in **1** and **3**, and which are highly  $\alpha$ -selective in homogeneous glycosylation reactions:<sup>60–62</sup> subtle conformational effects play a major role in these glycosylation reactions, even to the point of overriding powerful electronic effects.



Finally, we return to the possibility of the formation of the  $\alpha$ -glucosides via the intermediacy of covalent  $\beta$ -glucosyl triflate and an associated transient  $\beta$ -face contact ion pair. While this mechanism cannot be ruled out as a contributing factor to the overall  $\alpha$ -selectivity seen in the glucose series,<sup>4,6–9</sup> the poor selectivity seen with the 3-deoxy donors **25** and **27**, irrespective of the configuration at C2, suggests that it is not a major pathway.

## Experimental Section

**General Coupling Protocol for S-Phenyl 3-O-Benzyl-4,6-O-benzylidene-2-deoxy- $\beta$ -D-arabino-thiohexopyranoside (19) Using Ph<sub>2</sub>SO (Method A).** To a stirred solution of **19** (100 mg, 0.23 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (6 mL) containing Ph<sub>2</sub>SO (93 mg, 0.46 mmol), TTBP (114 mg, 0.46 mmol), and activated 3 Å powdered molecular sieves was added Tf<sub>2</sub>O (42.6  $\mu$ L, 0.25 mmol) at –60 °C under argon atmosphere. After 5 min, a solution of acceptor (0.28 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.4 mL) was added, and the reaction mixture was stirred for an additional 2 min at –60 °C, then warmed to room temperature, filtered, washed (saturated aqueous NaHCO<sub>3</sub>, brine), and dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was removed under reduced pressure. The crude was filtered through a pad of silica gel (with ethyl acetate as an eluent) and chromatographed using SiO<sub>2</sub>.

**General Coupling Protocol for S-Phenyl 2-O-Benzyl-4,6-O-benzylidene-3-deoxy- $\beta$ -D-ribo-thiohexopyranoside (25) and S-Phenyl 2-O-Benzyl-4,6-O-benzylidene-3-deoxy- $\beta$ -D-arabino-thiohexopyranoside (27) Using BSP (Method B).** To a stirred solution of **25** or **27** (50 mg, 0.12 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL) containing BSP (24 mg, 0.12 mmol), TTBP (57 mg, 0.23 mmol), and activated 3 Å powdered molecular sieves was added Tf<sub>2</sub>O (21.3  $\mu$ L, 0.13 mmol) at –60 °C under argon atmosphere. After 5 min, a solution of acceptor (0.14 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1.2 mL) was added, and the reaction mixture was stirred for an additional 2 min at –60 °C, then warmed to room temperature, filtered, washed (saturated aqueous NaHCO<sub>3</sub>, brine), and dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was removed under reduced pressure. The crude was filtered through a pad of silica gel (with ethyl acetate as an eluent) and chromatographed using SiO<sub>2</sub>.

**Methyl 4-O-(3'-O-Benzyl-4',6'-O-benzylidene-2'-deoxy- $\alpha$ -D-arabino-hexopyranosyl)-2,3-O-isopropylidene- $\alpha$ -L-rhamnopyra-**

**noside (39 $\alpha$ ) and Methyl 4-O-(3'-O-Benzyl-4',6'-O-benzylidene-2'-deoxy- $\beta$ -D-arabino-hexopyranosyl)-2,3-O-isopropylidene- $\alpha$ -L-rhamnopyranoside (39 $\beta$ ).** Prepared by method A and purified by means of radial chromatography, eluting consecutive plates with hexanes to 1:9 diethyl ether/hexanes, and then with hexanes to 1:19 ethyl acetate/hexanes. Combined yield 94 mg (75%, 1:1.5  $\alpha/\beta$ ). **39 $\alpha$** : [ $\alpha$ ]<sub>D</sub><sup>24</sup> +50.0 (c 0.53, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.52–7.57 (m, 2H), 7.27–7.43 (m, 8H), 5.64 (s, 1H), 5.00 (d, J = 3.5 Hz, 1H), 4.87 (d, J = 11.9 Hz, 1H), 4.86 (s, 1H), 4.70 (d, J = 11.9 Hz, 1H), 4.29 (dd, J = 10.0, 5.0 Hz, 1H), 4.07–4.15 (m, 3H), 4.03 (ddd, J = 11.2, 9.1, 5.1 Hz, 1H), 3.75 (t, J = 10.1 Hz, 1H), 3.71 (t, J = 9.4 Hz, 1H), 3.65 (dd, J = 10.0, 6.3 Hz, 1H), 3.37 (s, 3H), 3.31–3.36 (m, 1H), 2.27 (dd, J = 13.2, 5.1 Hz, 1H), 1.82 (ddd, J = 13.3, 11.2, 3.9 Hz, 1H), 1.52 (s, 3H), 1.34 (s, 3H), 1.26 (d, J = 6.4 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  138.7, 137.7, 128.8, 128.4, 128.2, 127.7, 127.6, 126.1, 109.1, 101.3, 98.6, 98.0, 84.0, 80.3, 76.9, 75.9, 73.0, 72.8, 69.1, 64.8, 63.0, 54.8, 36.6, 28.2, 26.4, 17.5; ESIHRMS calcd for C<sub>30</sub>H<sub>38</sub>NaO<sub>9</sub> [M + Na]<sup>+</sup> 565.2414, found 565.2409. **39 $\beta$** : white solid; mp 108–111 °C; [ $\alpha$ ]<sub>D</sub><sup>24</sup> –40.3 (c 0.31, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.49–7.54 (m, 2H), 7.27–7.42 (m, 8H), 5.61 (s, 1H), 5.04 (dd, J = 9.9, 2.0 Hz, 1H), 4.86 (s, 1H), 4.81 (d, J = 12.1 Hz, 1H), 4.72 (d, J = 12.3 Hz, 1H), 4.28 (dd, J = 10.5, 5.0 Hz, 1H), 4.13–4.18 (m, 1H), 4.09 (d, J = 5.5 Hz, 1H), 3.83 (t, J = 10.3 Hz, 1H), 3.77 (ddd, J = 11.2, 9.0, 5.1 Hz, 1H), 3.58–3.70 (m, 3H), 3.37 (s, 3H), 3.30–3.36 (m, 1H), 2.36 (ddd, J = 12.9, 5.1, 2.1 Hz, 1H), 1.65 (ddd, J = 12.8, 11.2, 9.9 Hz, 1H), 1.51 (s, 3H), 1.35 (s, 3H), 1.28 (d, J = 5.9 Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  138.5, 137.6, 128.9, 128.4, 128.2, 127.7, 127.6, 126.1, 109.3, 101.3, 97.9, 83.2, 78.5, 77.8, 76.1, 74.6, 72.4, 69.0, 66.6, 64.1, 54.8, 37.8, 27.9, 26.4, 17.5; ESIHRMS calcd for C<sub>30</sub>H<sub>38</sub>NaO<sub>9</sub> [M + Na]<sup>+</sup> 565.2414, found 565.2411.

**3-O-(3'-O-Benzyl-4',6'-O-benzylidene-2'-deoxy- $\alpha$ -D-arabino-hexopyranosyl)-1,2:5,6-di-O-isopropylidene- $\alpha$ -D-glucofuranose (40 $\alpha$ ) and 3-O-(3'-O-Benzyl-4',6'-O-benzylidene-2'-deoxy- $\beta$ -D-arabino-hexopyranosyl)-1,2:5,6-di-O-isopropylidene- $\alpha$ -D-glucofuranose (40 $\beta$ ).** Prepared by method A and purified by means of radial chromatography, eluting consecutive plates with hexanes to 1:9 ethyl acetate/hexanes and then with 7:13 or 1:4 diethyl ether: hexanes for **40 $\alpha$**  and **40 $\beta$** , respectively. **40 $\alpha$**  yield 39 mg (29%), **40 $\beta$**  yield 62 mg (46%). **40 $\alpha$** : [ $\alpha$ ]<sub>D</sub><sup>24</sup> +38.4 (c 0.25, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.50–7.57 (m, 2H), 7.28–7.45 (m, 8H), 5.89 (d, J = 3.7 Hz, 1H), 5.66 (s, 1H), 5.21 (d, J = 3.7 Hz, 1H), 4.87 (d, J = 11.7 Hz, 1H), 4.69 (d, J = 12.5 Hz, 1H), 4.56 (d, J = 3.7 Hz, 1H), 4.33 (dd, J = 9.5, 4.4 Hz, 1H), 4.24 (d, J = 2.9 Hz, 1H), 4.09–4.18 (m, 2H), 3.96–4.09 (m, 3H), 3.78–3.90 (m, 2H), 3.73 (t, J = 8.8 Hz, 1H), 2.29 (dd, J = 13.2, 5.1 Hz, 1H), 1.78–1.86 (m, 1H), 1.50 (s, 3H), 1.41 (s, 3H), 1.33 (s, 3H), 1.32 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  138.6, 137.4, 129.0, 128.4, 128.3, 127.6, 126.0, 112.1, 109.3, 105.3, 101.3, 99.2, 84.1, 83.8, 81.4, 80.2, 73.1, 72.6, 72.4, 69.0, 67.9, 63.9, 36.1, 26.9, 26.3, 25.4; ESIHRMS calcd for C<sub>32</sub>H<sub>40</sub>NaO<sub>10</sub> [M + Na]<sup>+</sup> 607.2519, found 607.2524. **40 $\beta$** : [ $\alpha$ ]<sub>D</sub><sup>24</sup> –15.8 (c 0.36, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.47–7.54 (m, 2H), 7.27–7.44 (m, 8H), 5.92 (d, J = 3.7 Hz, 1H), 5.61 (s, 1H), 4.83 (d, J = 12.1 Hz, 1H), 4.67–4.73 (m, 2H), 4.49 (d, J = 3.9 Hz, 1H), 4.28–4.38 (m, 4H), 4.00–4.07 (m, 2H), 3.82 (t, J = 10.3 Hz, 1H), 3.67–3.77 (m, 2H), 3.34 (td, J = 13.9, 5.0 Hz, 1H), 2.26 (ddd, J = 13.2, 4.8, 2.2 Hz, 1H), 1.69 (ddd, J = 13.2, 10.2 Hz, 1H), 1.50 (s, 3H), 1.44 (s, 3H), 1.38 (s, 3H), 1.32 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  138.4, 137.4, 129.0, 128.4, 128.3, 127.7, 126.0, 111.9, 108.5, 105.1, 101.3, 98.1, 82.84, 82.76, 80.3, 79.4, 74.2, 73.4, 72.6, 68.8, 66.8, 66.0, 37.5, 26.8, 26.6, 26.3, 25.5; ESIHRMS calcd for C<sub>32</sub>H<sub>40</sub>NaO<sub>10</sub> [M + Na]<sup>+</sup> 607.2519, found 607.2524.

**6-O-(3'-O-Benzyl-4',6'-O-benzylidene-2'-deoxy- $\alpha$ -D-arabino-hexopyranosyl)-1,2:3,4-di-O-isopropylidene- $\alpha$ -D-galactopyranose (41 $\alpha$ ) and 6-O-(3'-O-Benzyl-4',6'-O-benzylidene-2'-deoxy- $\beta$ -D-arabino-hexopyranosyl)-1,2:3,4-di-O-isopropylidene- $\alpha$ -D-galactopyranose (41 $\beta$ ).** Prepared by method A and purified by

(60) Crich, D.; Cai, W.; Dai, Z. *J. Org. Chem.* **2000**, *65*, 1291–1297.

(61) Crich, D.; Vinod, A. U.; Picione, J.; Wink, D. *J. ARKIVOC* **2005**, (vi), 339–344.

(62) Crich, D.; Vinod, A. U.; Picione, J. *J. Org. Chem.* **2003**, *68*, 8453–8458.

means of radial chromatography, eluting consecutive plates with hexanes to 3:17 ethyl acetate/hexanes and then with 1:3 to 3:7 diethyl ether/hexanes. Combined yield 82 mg (61%, 1:4  $\alpha/\beta$ ). **41 $\alpha$** :  $[\alpha]^{24}_D +16.9$  (c 0.14, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.48–7.54 (m, 2H), 7.23–7.41 (m, 8H), 5.62 (s, 1H), 5.53 (d,  $J = 5.3$  Hz, 1H), 4.97 (d,  $J = 3.5$  Hz, 1H), 4.85 (d,  $J = 12.0$  Hz, 1H), 4.68 (d,  $J = 12.0$  Hz, 1H), 4.63 (dd,  $J = 7.9, 2.3$  Hz, 1H), 4.32 (dd,  $J = 5.0, 2.3$  Hz, 1H), 4.23–4.29 (m, 2H), 4.05 (ddd,  $J = 11.1, 9.1, 5.3$  Hz, 1H), 3.97 (td,  $J = 6.6, 1.8$  Hz, 1H), 3.87 (td,  $J = 9.9, 4.7$  Hz, 1H), 3.64–3.79 (m, 4H), 2.32 (dd,  $J = 13.2, 5.3$  Hz, 1H), 1.80 (ddd,  $J = 13.4, 11.1, 3.8$  Hz, 1H), 1.55 (s, 3H), 1.45 (s, 3H), 1.35 (s, 3H), 1.34 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  138.8, 137.6, 128.9, 128.3, 128.2, 127.6, 127.5, 126.1, 109.4, 108.6, 101.3, 98.1, 96.4, 83.9, 73.0, 71.0, 70.6, 69.1, 65.9, 65.8, 63.1, 36.5, 26.2, 26.0, 25.0, 24.6; FABHRMS calcd for C<sub>32</sub>H<sub>40</sub>NaO<sub>10</sub> [M + Na]<sup>+</sup> 607.2519, found 607.2515. **41 $\beta$** :  $[\alpha]^{24}_D -51.4$  (c 0.21, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.48–7.53 (m, 2H), 7.26–7.42 (m, 8H), 5.59 (s, 1H), 5.55 (d,  $J = 5.0$  Hz, 1H), 4.79 (d,  $J = 12.3$  Hz, 1H), 4.71 (d,  $J = 12.1$  Hz, 1H), 4.66 (dd,  $J = 9.8, 2.1$  Hz, 1H), 4.60 (dd,  $J = 8.0, 2.5$  Hz, 1H), 4.29–4.34 (m, 2H), 4.21 (dd,  $J = 7.9, 1.8$  Hz, 1H), 3.97–4.05 (m, 2H), 3.82 (t,  $J = 10.3$  Hz, 1H), 3.64–3.77 (m, 3H), 3.35 (td,  $J = 9.6, 5.0$  Hz, 1H), 2.44 (ddd,  $J = 13.0, 5.0, 2.1$  Hz, 1H), 1.69–1.78 (m, 1H), 1.54 (s, 3H), 1.45 (s, 3H), 1.34 (s, 3H), 1.31 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  138.7, 137.7, 129.1, 128.6, 128.4, 127.9, 127.8, 126.3, 109.6, 108.9, 101.5, 101.3, 96.5, 83.2, 74.6, 72.5, 71.7, 70.9, 70.6, 69.4, 69.1, 68.0, 66.8, 37.7, 26.3, 26.2, 25.2, 24.6; ESIHRMS calcd for C<sub>32</sub>H<sub>40</sub>NaO<sub>10</sub> [M + Na]<sup>+</sup> 607.2519, found 607.2521.

**1-Adamantyl 3-O-Benzyl-4,6-O-benzylidene-2'-deoxy- $\alpha$ -D-arabino-hexopyranoside (42 $\alpha$ ) and 1-Adamantyl 3-O-Benzyl-4,6-O-benzylidene-2'-deoxy- $\beta$ -D-arabino-hexopyranoside (42 $\beta$ )**. Prepared by method A and purified by means of radial chromatography, eluting consecutive plates with hexanes to 1:39 ethyl acetate/hexanes and then with 1:9 diethyl ether/hexanes. Combined yield 33 mg (30%, 1:1.9  $\alpha/\beta$ ). **42 $\alpha$** : white solid; mp 160–162 °C;  $[\alpha]^{24}_D +54.7$  (c 0.086, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.50–7.55 (m, 2H), 7.22–7.42 (m, 8H), 5.63 (s, 1H), 5.35 (d,  $J = 3.3$  Hz, 1H), 4.87 (d,  $J = 11.7$  Hz, 1H), 4.69 (d,  $J = 11.7$  Hz, 1H), 4.22 (dd,  $J = 10.3, 5.0$  Hz, 1H), 4.01–4.15 (m, 2H), 3.73 (t,  $J = 10.4$  Hz, 1H), 3.67 (t,  $J = 9.3$  Hz, 1H), 2.08–2.18 (m, 4H), 1.73–1.86 (m, 7H), 1.56–1.68 (m, 6H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  139.0, 137.8, 128.8, 128.3, 128.2, 127.6, 127.5, 126.0, 101.2, 91.2, 84.4, 74.2, 73.4, 73.0, 69.2, 62.7, 42.5, 38.1, 36.3, 30.6; ESIHRMS calcd for C<sub>30</sub>H<sub>36</sub>NaO<sub>5</sub> [M + Na]<sup>+</sup> 499.2461, found 499.2472. **42 $\beta$** : slightly yellow solid; mp 86–88 °C;  $[\alpha]^{24}_D -11.5$  (c 0.19, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.47–7.55 (m, 2H), 7.24–7.43 (m, 8H), 5.59 (s, 1H), 4.89 (dd,  $J = 9.9, 2.3$  Hz, 1H), 4.80 (d,  $J = 12.0$  Hz, 1H), 4.72 (d,  $J = 12.0$  Hz, 1H), 4.27 (dd,  $J = 10.5, 5.0$  Hz, 1H), 3.83 (t,  $J = 10.4$  Hz, 1H), 3.70–3.79 (m, 1H), 3.67 (t,  $J = 8.9$  Hz, 1H), 3.35 (td,  $J = 14.0, 5.0$  Hz, 1H), 2.11–2.22 (m, 4H), 1.71–1.88 (m, 7H), 1.56–1.70 (m, 6H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  138.6, 137.6, 128.9, 128.4, 128.2, 127.63, 127.56, 126.1, 101.3, 93.1, 82.9, 75.0, 74.9, 72.3, 69.1, 66.5, 42.5, 39.1, 36.2, 30.6; ESIHRMS calcd for C<sub>30</sub>H<sub>36</sub>NaO<sub>5</sub> [M + Na]<sup>+</sup> 499.2461, found 499.2452.

**Methyl 4-O-(2'-O-Benzyl-4',6'-O-benzylidene-3'-deoxy- $\alpha$ -D-ribo-hexopyranosyl)-2,3-O-isopropylidene- $\alpha$ -L-rhamnopyranoside (43 $\alpha$ ) and Methyl 4-O-(2'-O-Benzyl-4',6'-O-benzylidene-3'-deoxy- $\beta$ -D-ribo-hexopyranosyl)-2,3-O-isopropylidene- $\alpha$ -L-rhamnopyranoside (43 $\beta$ )**. Prepared by method B. Purified by means of radial chromatography, eluting with hexanes to 3:17 diethyl ether/hexanes to give **43 $\alpha$**  (33 mg, 53%) and **43 $\beta$**  (26 mg, 42%). **43 $\alpha$** : white solid; mp 138–142 °C;  $[\alpha]^{22}_D +31.1$  (c 0.074, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.48–7.52 (m, 2H) 7.28–7.40 (m, 8H), 5.50 (s, 1H), 4.93 (d,  $J = 3.3$  Hz, 1H), 4.86 (s, 1H), 4.67 (d,  $J = 12.1$  Hz, 1H), 4.56 (d,  $J = 12.3$  Hz, 1H), 4.21–4.28 (m, 2H), 4.12 (d,  $J = 5.5$  Hz, 1H), 4.06 (td,  $J = 9.8, 5.0$  Hz, 1H), 3.75–3.82 (m, 1H), 3.57–3.67 (m, 2H), 3.49 (ddd,  $J = 11.6, 9.5, 4.0$  Hz, 1H), 3.39 (dd,  $J = 10.1, 7.5$  Hz, 1H), 3.36 (s, 3H), 2.31

(ddd,  $J = 11.0, 4.3, 4.1$  Hz, 1H), 2.04 (ddd,  $J_1 = J_2 = J_3 = 11.7$  Hz, 1H), 1.52 (s, 3H), 1.37 (d,  $J = 6.2$  Hz, 3H), 1.35 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  137.8, 137.5, 129.1, 128.5, 128.4, 127.9, 127.7, 126.3, 109.1, 101.8, 97.9, 81.1, 76.1, 74.3, 71.1, 69.5, 64.9, 64.3, 54.7, 29.81, 29.75, 28.2, 26.5, 17.3; FABHRMS calcd for C<sub>30</sub>H<sub>38</sub>NaO<sub>9</sub> [M + Na]<sup>+</sup> 565.2414, found 565.2390. **43 $\beta$** :  $[\alpha]^{22}_D -43.9$  (c 0.098, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.45–7.49 (m, 2H), 7.27–7.39 (m, 8H), 5.49 (s, 1H), 5.03 (d,  $J = 7.5$  Hz, 1H), 4.87 (s, 1H), 4.82 (d,  $J = 12.1$  Hz, 1H), 4.65 (d,  $J = 12.1$  Hz, 1H), 4.23–4.29 (m, 2H), 4.11 (d,  $J = 5.7$  Hz, 1H), 3.63–3.76 (m, 3H), 3.51 (ddd,  $J = 11.7, 8.9, 4.3$  Hz, 1H), 3.31–3.43 (m, 5H), 2.40 (dt,  $J = 12.0, 4.7$  Hz, 1H), 1.79 (ddd,  $J_1 = J_2 = J_3 = 11.7$  Hz, 1H), 1.53 (s, 3H), 1.36 (s, 3H), 1.30 (d,  $J = 5.9$  Hz, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  138.6, 137.4, 129.1, 128.4, 127.7, 127.6, 126.2, 109.3, 103.1, 101.7, 98.1, 78.2, 77.9, 77.6, 76.3, 76.0, 75.5, 72.8, 70.0, 69.3, 64.2, 55.0, 34.9, 27.9, 26.4, 17.7; FABHRMS calcd for C<sub>30</sub>H<sub>38</sub>NaO<sub>9</sub> [M + Na]<sup>+</sup> 565.2414, found 565.2398.

**3-O-(2'-O-Benzyl-4',6'-O-benzylidene-3'-deoxy- $\alpha$ -D-ribo-hexopyranosyl)-1,2:5,6-di-O-isopropylidene- $\alpha$ -D-glucopyranose (44 $\alpha$ ) and 3-O-(2'-O-Benzyl-4',6'-O-benzylidene-3'-deoxy- $\beta$ -D-ribo-hexopyranosyl)-1,2:5,6-di-O-isopropylidene- $\alpha$ -D-glucopyranose (44 $\beta$ )**. Prepared by method B. Purified by means of radial chromatography, eluting with hexanes to 1:4 diethyl ether/hexanes, followed by normal-phase HPLC, eluting with hexanes to 3:2 ethyl acetate/hexanes. Combined yield 51 mg (76%, 4.7:1  $\alpha/\beta$ ). **44 $\alpha$** :  $[\alpha]^{22}_D +38.7$  (c 0.20, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.45–7.52 (m, 2H), 7.27–7.41 (m, 8H), 5.96 (d,  $J = 3.5$  Hz, 1H), 5.52 (s, 1H), 5.21 (d,  $J = 3.2$  Hz, 1H), 4.60–4.67 (m, 3H), 4.53 (dt,  $J = 8.6, 5.4$  Hz, 1H), 4.27–4.34 (m, 2H), 4.10 (dd,  $J = 8.5, 2.6$  Hz, 1H), 4.06 (d,  $J = 5.6$  Hz, 2H), 3.75–3.83 (m, 1H), 3.70 (t,  $J = 10.2$  Hz, 1H), 3.61 (ddd,  $J = 11.7, 4.2, 3.9$  Hz, 1H), 3.49–3.57 (m, 1H), 2.29 (ddd,  $J = 11.3, 4.2, 4.1$  Hz, 1H), 2.04 (ddd,  $J_1 = J_2 = J_3 = 11.7$  Hz, 1H), 1.51 (s, 3H), 1.43 (s, 3H), 1.33 (s, 3H), 1.28 (s, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  137.9, 137.2, 129.2, 128.5, 128.4, 127.8, 127.4, 126.1, 112.0, 109.2, 105.2, 101.7, 97.5, 84.1, 81.4, 80.1, 73.9, 72.1, 70.9, 69.3, 67.3, 65.0, 29.8, 27.1, 26.9, 26.4, 25.4; FABHRMS calcd for C<sub>32</sub>H<sub>40</sub>NaO<sub>10</sub> [M + Na]<sup>+</sup> 607.2519, found 607.2524. **44 $\beta$** :  $[\alpha]^{22}_D -50.0$  (c 0.042, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$ : 7.45–7.49 (m, 2H), 7.28–7.40 (m, 8H), 5.84 (d,  $J = 3.7$  Hz, 1H), 5.49 (s, 1H), 4.66 (d,  $J = 11.7$  Hz, 1H), 4.63 (d,  $J = 12.1$  Hz, 1H), 4.54–4.58 (m, 2H), 4.33–4.39 (m, 3H), 4.29 (dd,  $J = 10.5, 5.0$  Hz, 1H), 4.04–4.09 (m, 2H), 3.72 (t,  $J = 10.3$  Hz, 1H), 3.54 (ddd,  $J = 16.2, 4.6, 4.5$  Hz, 1H), 3.31–3.43 (m, 2H), 2.47 (dt,  $J = 12.0, 4.7$  Hz, 1H), 1.75 (ddd,  $J_1 = J_2 = J_3 = 11.9$  Hz, 1H), 1.50 (s, 3H), 1.45 (s, 3H), 1.38 (s, 3H), 1.28 (s, 3H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$ : 137.9, 137.2, 129.2, 128.5, 128.4, 127.9, 127.5, 126.1, 111.9, 108.6, 105.1, 103.1, 101.6, 82.6, 80.5, 80.2, 75.7, 75.3, 73.4, 72.6, 70.4, 69.0, 66.0, 34.6, 26.7, 26.6, 26.2, 25.6; FABHRMS calcd for C<sub>32</sub>H<sub>40</sub>NaO<sub>10</sub> [M + Na]<sup>+</sup> 607.2519, found 607.2509.

**1-Adamantyl 2-O-Benzyl-4,6-O-benzylidene-3-deoxy- $\alpha$ -D-ribo-hexopyranoside (45 $\alpha$ ) and 1-Adamantyl 2-O-Benzyl-4,6-O-benzylidene-3-deoxy- $\beta$ -D-ribo-hexopyranoside (45 $\beta$ )**. Prepared by method B. Purified by means of radial chromatography, eluting consecutive plates with hexanes to 1:19 diethyl ether/hexanes, and then with hexanes to 1:39 ethyl acetate/hexanes. Combined yield 50.4 mg (92%, 1:1.8  $\alpha/\beta$ ). **45 $\alpha$** : white solid; mp 119–122 °C;  $[\alpha]^{22}_D +43.6$  (c 0.078, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.45–7.53 (m, 2H), 7.27–7.43 (m, 8H), 5.50 (s, 1H), 5.28 (d,  $J = 3.5$  Hz, 1H), 4.62 (d,  $J = 12.3$  Hz, 1H), 4.58 (d,  $J = 12.3$  Hz, 1H), 4.20 (dd,  $J = 10.2, 4.7$  Hz, 1H), 4.00 (td,  $J = 9.9, 4.8$  Hz, 1H), 3.63 (t,  $J = 10.4$  Hz, 1H), 3.53–3.59 (m, 1H), 3.49 (ddd,  $J = 11.4, 9.4, 4.4$  Hz, 1H), 2.08–2.26 (m, 5H), 1.82–1.96 (m, 6H), 1.59–1.70 (m, 6H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  138.3, 137.5, 129.0, 128.5, 128.3, 127.7, 127.6, 126.2, 101.7, 89.7, 74.5, 74.0, 70.6, 69.6, 63.7, 42.6, 36.3, 30.7, 30.1; FABHRMS calcd for C<sub>30</sub>H<sub>36</sub>NaO<sub>5</sub> [M + Na]<sup>+</sup> 499.2461, found 499.2447. **45 $\beta$** :  $[\alpha]^{22}_D -19.2$  (c 0.052, CHCl<sub>3</sub>); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  7.43–7.51 (m, 2H), 7.27–7.41 (m, 8H), 5.47 (s, 1H), 4.85 (d,  $J = 12.1$  Hz, 1H),

4.78 (d,  $J = 7.5$  Hz, 1H), 4.68 (d,  $J = 12.1$  Hz, 1H), 4.26 (dd,  $J = 10.5, 5.0$  Hz, 1H), 3.74 (t,  $J = 10.3$  Hz, 1H), 3.51 (ddd,  $J = 11.7, 8.9, 4.3$  Hz, 1H), 3.35–3.44 (m, 2H), 2.40 (dt,  $J = 12.0, 4.6$  Hz, 1H), 2.15–2.22 (m, 3H), 1.90–1.97 (m, 3H), 1.81–1.88 (m, 3H), 1.76 (ddd,  $J_1 = J_2 = J_3 = 11.7$  Hz, 1H), 1.60–1.71 (m, 6H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  138.6, 137.4, 129.1, 128.3, 127.8, 127.6, 126.2, 101.6, 98.2, 76.2, 75.43, 75.37, 73.0, 70.0, 69.3, 42.8, 36.3, 35.3, 30.7; FABHRMS calcd for  $\text{C}_{30}\text{H}_{36}\text{NaO}_5$  [ $\text{M} + \text{Na}$ ] $^+$  499.2461, found 499.2447.

**Methyl 4-*O*-(2'-*O*-Benzyl-4',6'-*O*-benzylidene-3'-deoxy- $\alpha$ -D-arabino-hexopyranosyl)-2,3-*O*-isopropylidene- $\alpha$ -L-rhamnopyranoside (46 $\alpha$ ) and Methyl 4-*O*-(2'-*O*-Benzyl-4',6'-*O*-benzylidene-3'-deoxy- $\beta$ -D-arabino-hexopyranosyl)-2,3-*O*-isopropylidene- $\alpha$ -L-rhamnopyranoside (46 $\beta$ ).** Prepared by method B. Purified by means of radial chromatography, eluting with hexanes to 3:17 diethyl ether/hexanes. Combined yield 53.0 mg (85%, 1.9:1  $\alpha/\beta$ ). Analytical samples of both isomers were obtained by reversed-phase HPLC separation, eluting with 1:1 water/acetonitrile to acetonitrile. **46 $\alpha$** : white solid; mp 152–155 °C;  $[\alpha]_D^{25} +54.6$  ( $c$  0.108,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.48–7.54 (m, 2H), 7.27–7.41 (m, 8H), 5.59 (s, 1H), 4.84 (s, 1H), 4.79 (s, 1H), 4.67 (d,  $J = 12.3$  Hz, 1H), 4.56 (d,  $J = 12.3$  Hz, 1H), 4.23 (dd,  $J = 10.1, 4.8$  Hz, 1H), 4.06–4.15 (m, 3H), 3.98–4.06 (m, 1H), 3.78 (t,  $J = 9.9$  Hz, 1H), 3.65–3.69 (m, 1H), 3.54–3.63 (m, 1H), 3.33–3.41 (m, 4H), 2.23 (dt,  $J = 13.1, 3.3$  Hz, 1H), 1.92–2.01 (m, 1H), 1.52 (s, 3H), 1.34 (s, 3H), 1.13 (d,  $J = 6.4$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  137.8, 137.7, 129.0, 128.5, 128.3, 127.9, 127.8, 126.2, 109.2, 102.0, 98.5, 98.0 ( $^1J_{\text{CH}} = 166.5$ ), 80.3, 76.9, 76.0, 74.6, 74.2, 71.3, 69.4, 65.1, 64.8, 54.8, 29.3, 28.2, 26.5, 17.3; ESIHRMS calcd for  $\text{C}_{30}\text{H}_{39}\text{O}_9$  [ $\text{M} + \text{H}$ ] $^+$  543.25886, found 543.25986. **46 $\beta$** : white solid; mp 158–162 °C;  $[\alpha]_D^{25} -67.1$  ( $c$  0.082,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.46–7.51 (m, 2H), 7.26–7.42 (m, 8H), 5.56 (s, 1H), 5.10 (d,  $J = 0.9$  Hz, 1H), 4.88 (s, 1H), 4.84 (d,  $J = 12.5$  Hz, 1H), 4.68 (d,  $J = 12.5$  Hz, 1H), 4.25 (dd,  $J = 10.5, 5.0$  Hz, 1H), 4.14–4.18 (m, 1H), 4.10 (d,  $J = 5.5$  Hz, 1H), 4.04 (ddd,  $J = 11.8, 9.2, 4.5$  Hz, 1H), 3.91 (t,  $J = 10.4$  Hz, 1H), 3.80–3.82 (m, 1H), 3.63–3.73 (m, 2H), 3.45 (td,  $J = 9.7, 4.9$  Hz, 1H), 3.40 (s, 3H), 2.31 (dt,  $J = 13.1, 3.9$  Hz, 1H), 1.72–1.79 (m, 1H), 1.54 (s, 3H), 1.33–1.37 (m, 6H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  138.7, 137.6, 129.0, 128.32, 128.28, 127.7, 127.6, 126.1, 109.3, 101.9, 100.9 ( $^1J_{\text{CH}} = 160.0$ ), 97.9, 78.5, 77.7, 76.1, 74.6, 74.1, 73.3, 71.2, 69.1, 64.3, 54.9, 33.9, 27.9, 26.5, 17.7; FABHRMS calcd for  $\text{C}_{30}\text{H}_{38}\text{NaO}_9$  [ $\text{M} + \text{Na}$ ] $^+$  565.2414, found 565.2408.

**3-*O*-(2'-*O*-Benzyl-4',6'-*O*-benzylidene-3'-deoxy- $\alpha$ -D-arabino-hexopyranosyl)-1,2:5,6-di-*O*-isopropylidene- $\alpha$ -D-glucofuranose (47 $\alpha$ ) and 3-*O*-(2'-*O*-Benzyl-4',6'-*O*-benzylidene-3'-deoxy- $\beta$ -D-arabino-hexopyranosyl)-1,2:5,6-di-*O*-isopropylidene- $\alpha$ -D-glucofuranose (47 $\beta$ ).** Prepared by method B. Purification by means of radial chromatography, eluting with hexanes to 1:3 diethyl ether/hexanes, gave crude products. Further purification of crude **47 $\alpha$**  and **47 $\beta$**  by means of normal-phase HPLC, eluting with hexanes to 1:1 ethyl acetate/hexanes and hexanes to 3:2 ethyl acetate/hexanes respectively, afforded pure products (53.5 mg, combined 80% yield, 1.3:1  $\alpha/\beta$ ). **47 $\alpha$** : white solid; mp 145–148 °C;  $[\alpha]_D^{25} +37.6$  ( $c$  0.492,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.48–7.52 (m, 2H), 7.28–7.40 (m, 8H), 5.88 (d,  $J = 3.5$  Hz, 1H), 5.61 (s, 1H), 5.15 (s, 1H), 4.64 (d,  $J = 12.3$  Hz, 1H), 4.57–4.62 (m, 2H), 4.34 (d,  $J =$

$2.4$  Hz, 1H), 4.28–4.31 (m, 1H), 4.00–4.15 (m, 5H), 3.81–3.89 (m, 2H), 3.71–3.75 (m, 1H), 2.24 (dt,  $J = 13.0, 3.3$  Hz, 1H), 1.96 (td,  $J = 12.5, 3.0$  Hz, 1H), 1.51 (s, 3H), 1.42 (s, 3H), 1.33 (s, 6H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  137.7, 137.5, 129.1, 128.5, 128.4, 127.9, 127.6, 126.1, 112.1, 109.4, 105.3, 101.9, 98.1 ( $^1J_{\text{CH}} = 172.0$ ), 84.1, 81.5, 79.9, 74.5, 74.1, 72.5, 71.3, 69.3, 67.9, 66.0, 29.4, 26.9, 26.3, 25.5; EIHRMS calcd for  $\text{C}_{32}\text{H}_{40}\text{O}_{10}$  [ $\text{M}$ ] $^+$  584.2622, found 584.2659. **47 $\beta$** :  $[\alpha]_D^{25} -30.3$  ( $c$  0.330,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.44–7.51 (m, 2H), 7.27–7.40 (m, 8H), 5.92 (d,  $J = 3.8$  Hz, 1H), 5.56 (s, 1H), 4.72 (d,  $J = 12.6$  Hz, 1H), 4.69 (s, 1H), 4.64 (d,  $J = 12.6$  Hz, 1H), 4.40–4.47 (m, 2H), 4.33–4.38 (m, 2H), 4.28 (dd,  $J = 10.4, 4.8$  Hz, 1H), 4.15–4.20 (m, 1H), 4.02–4.11 (m, 2H), 3.87 (t,  $J = 10.4$  Hz, 1H), 3.71–3.75 (m, 1H), 3.46 (td,  $J = 9.6, 5.0$  Hz, 1H), 2.33 (dt,  $J = 13.2, 4.2$  Hz, 1H), 1.68–1.78 (m, 1H), 1.51 (s, 3H), 1.45 (s, 3H), 1.36 (s, 3H), 1.32 (s, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  138.0, 137.4, 129.1, 128.43, 128.36, 127.8, 126.1, 112.0, 108.6, 105.1, 101.8, 100.2 ( $^1J_{\text{CH}} = 155.4$ ), 82.7, 80.5, 80.2, 73.8, 73.6, 73.3, 72.6, 71.0, 69.0, 66.1, 32.8, 26.8, 26.6, 26.3, 25.5; EIHRMS calcd for  $\text{C}_{32}\text{H}_{40}\text{O}_{10}$  [ $\text{M}$ ] $^+$  584.2622, found 584.2602.

**1-Adamantyl 2-*O*-Benzyl-4,6-*O*-benzylidene-3-deoxy- $\alpha$ -D-arabino-hexopyranoside (48 $\alpha$ ) and 1-Adamantyl 2-*O*-Benzyl-4,6-*O*-benzylidene-3-deoxy- $\beta$ -D-arabino-hexopyranoside (48 $\beta$ ).** Prepared by method B. Purification by means of radial chromatography, eluting with hexanes to 1:19 ethyl acetate/hexanes, gave crude products. Further purification of crude **48 $\alpha$**  and **48 $\beta$**  by means of normal-phase HPLC, eluting with hexanes to 3:2 ethyl acetate/hexanes, afforded pure **48 $\alpha$**  (35.4 mg, 65%) and **48 $\beta$**  (9.6 mg, 18%). **48 $\alpha$** : white solid; mp 110–114 °C;  $[\alpha]_D^{25} +61.9$  ( $c$  0.690,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.49–7.53 (m, 2H), 7.28–7.41 (m, 8H), 5.59 (s, 1H), 5.15 (s, 1H), 4.67 (d,  $J = 12.1$  Hz, 1H), 4.59 (d,  $J = 12.1$  Hz, 1H), 4.19 (dd,  $J = 10.3, 4.8$  Hz, 1H), 3.96–4.10 (m, 2H), 3.78 (t,  $J = 10.2$  Hz, 1H), 3.54–3.57 (m, 1H), 2.21 (dt,  $J = 12.8, 3.5$  Hz, 1H), 2.11–2.17 (m, 3H), 2.06 (td,  $J = 12.2, 2.9$  Hz, 1H), 1.74–1.83 (m, 6H), 1.57–1.68 (m, 6H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  138.1, 137.8, 129.0, 128.5, 128.3, 127.8, 127.7, 126.2, 101.9, 91.0 ( $^1J_{\text{CH}} = 166.5$ ), 76.7, 74.6, 74.5, 71.2, 69.5, 64.9, 42.5, 36.3, 30.6, 29.2; FABHRMS calcd for  $\text{C}_{30}\text{H}_{36}\text{NaO}_5$  [ $\text{M} + \text{Na}$ ] $^+$  499.2461, found 499.2492. **48 $\beta$** :  $[\alpha]_D^{25} -33.2$  ( $c$  0.196,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.41–7.50 (m, 4H), 7.25–7.40 (m, 6H), 5.55 (s, 1H), 4.88 (d,  $J = 13.1$  Hz, 1H), 4.85 (d,  $J = 0.9$  Hz, 1H), 4.79 (d,  $J = 12.9$  Hz, 1H), 4.24 (dd,  $J = 10.4, 4.8$  Hz, 1H), 4.02 (ddd,  $J = 11.8, 9.2, 4.4$  Hz, 1H), 3.88 (t,  $J = 10.4$  Hz, 1H), 3.57–3.62 (m, 1H), 3.45 (td,  $J = 9.6, 5.0$  Hz, 1H), 2.30 (ddd,  $J = 13.0, 3.8, 3.7$  Hz, 1H), 2.14–2.20 (m, 3H), 1.58–1.91 (m, 13H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  138.8, 137.7, 129.0, 128.3, 128.2, 127.9, 127.4, 126.2, 101.9, 95.3 ( $^1J_{\text{CH}} = 152.6$ ), 75.4, 74.9, 74.0, 73.0, 70.9, 69.3, 42.4, 36.3, 34.1, 30.6; FABHRMS calcd for  $\text{C}_{30}\text{H}_{36}\text{NaO}_5$  [ $\text{M} + \text{Na}$ ] $^+$  499.2461, found 499.2413.

**Acknowledgment.** We thank Dr. Tom Hutton for helpful discussions and the NIH (GM 62160) for support of this work.

**Supporting Information Available:** Details of the preparation of all donors and copies of NMR spectra for all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO061417B