


## The Synthesis of a Geminally Perfluoro-*tert*-butylated $\beta$ -Amino Acid and its Protected Forms as a Potential Pharmacokinetic Modulator and Reporter for Peptide-Based Pharmaceuticals

Zhong-Xing Jiang and Y. Bruce Yu\*

Department of Pharmaceutics and Pharmaceutical Chemistry,  
College of Pharmacy, University of Utah, Salt Lake City, Utah  
84112

bruce.yu@utah.edu

Received August 4, 2006



To modulate and report the pharmacokinetics of peptide-based pharmaceuticals, a novel geminally perfluoro-*tert*-butylated  $\beta$ -amino acid ( $\beta$ Fa) and its Fmoc- and Boc-protected forms were designed and synthesized.  $\beta$ Fa was incorporated into a model tripeptide via standard solid-phase chemistry. Both the amino acid (free and protected) and the tripeptide show a sharp singlet  $^{19}\text{F}$  NMR signal. Reversed-phase chromatography and 1-octanol/water partition measurements demonstrate that  $\beta$ Fa is extremely hydrophobic.

We are interested in the design and synthesis of fluorinated amino acids as modulators and reporters of peptide pharmacokinetics. The potential benefits brought by fluorinated amino acids include prolonged in vivo half-life,<sup>1</sup> enhanced membrane permeability,<sup>2</sup> and noninvasive detection via  $^{19}\text{F}$  magnetic resonance spectroscopy.<sup>3</sup>

A generic, highly fluorinated amino acid that can dramatically increase the lipophilicity of a peptide and give a sharp singlet  $^{19}\text{F}$  NMR signal is highly desirable for pharmacokinetic modulation and reporting purposes. To this end, we designed a geminally perfluoro-*tert*-butylated  $\beta$ -amino acid ( $\beta$ Fa, **1**), as<sup>1</sup>

(1) Hsieh, K.-H.; Needleman, P.; Marshall, G. R. *J. Med. Chem.* **1987**, *30*, 1097–1100.

(2) (a) Ortial, S.; Durand, G.; Poeggeler, B.; Polidori, A.; Pappolla, M. A.; Böker, J.; Hardeland, R.; Pucci, B. *J. Med. Chem.* **2006**, *49*, 2812–2820. (b) Perino, S.; Contino-Pépin, C.; Jasseron, S.; Rapp, M.; Maurizis, J.-C.; Pucci, B. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 1111–1114. (c) Park, B. K.; Kitteringham, N. R.; O'Neill, P. M. *Ann. Rev. Pharmacol. Toxicol.* **2001**, *41*, 443–470. (d) Gerebtzoff, G.; Li-Blatter, X.; Fischer, H.; Frenzler, A.; Seelig, A. *ChemBioChem* **2004**, *5*, 676–684.

(3) Wolf, W.; Presant, C. A.; Waluch, V. *Adv. Drug Delivery Rev.* **2000**, *41*, 55–74.




FIGURE 1. Structures of target molecules.




FIGURE 2. Structures of two model tripeptides, formyl-Gly- $\beta$ Fa-Gly-amide (**4**) and formyl-Gly-L-Trp-Gly-amide (**5**).


shown in Figure 1. We also need  $\beta$ Fa in its Fmoc- and Boc-protected forms (**2** and **3**, respectively) for solid-phase peptide synthesis. The fluorine atoms are introduced into the amino acid through two symmetrically positioned perfluoro-*tert*-butyl groups. Such a spherically symmetric arrangement of the 18 fluorine atoms in  $\beta$ Fa ensures that they have an identical chemical environment and avoids  $^{19}\text{F}$ – $^{19}\text{F}$  or  $^{19}\text{F}$ – $^1\text{H}$  coupling. As a result, we anticipate this amino acid to give a single  $^{19}\text{F}$  NMR signal.  $\beta$ Fa is achiral and needs no side-chain protection, significantly simplifying the synthesis of both the free amino acid and its protected forms. Considering that the electron-withdrawing capacity of perfluoro-*tert*-butyl groups can potentially weaken the basicity of the amino group and hence its reactivity during solid-phase peptide synthesis, a  $\beta$ -amino acid is adopted to ensure sufficient separation between the amino group and the two perfluoro-*tert*-butyl groups. This also allows better steric accommodation of the bulky perfluoro-*tert*-butyl groups.

To demonstrate the feasibility of  $\beta$ Fa to be incorporated into peptides via solid-phase synthesis, we designed the following tripeptide, formyl-Gly- $\beta$ Fa-Gly-amide **4**. This peptide serves dual purposes. First, it demonstrates that protected  $\beta$ Fa can indeed be incorporated into peptides via solid-phase peptide synthesis; second, it demonstrates that  $\beta$ Fa can indeed increase the hydrophobicity of a peptide.  $\beta$ Fa is positioned in the middle of the tripeptide to demonstrate that  $\beta$ Fa can be placed in any position of a peptide, not just the N- and C-termini. The N- and C-termini of the tripeptide are formylated and amidated, respectively. These modifications abolish terminal charges of the tripeptide so that they will not interfere with hydrophobicity and 1-octanol/water partition measurements. Note that the disentanglement of hydrophobic interactions from electrostatic interactions in peptides/proteins is far from trivial.<sup>4</sup> Two glycines, which have no side chains, flank the central  $\beta$ Fa. The purpose is to abolish nearest-neighbor interactions that can interfere with hydrophobicity measurements.<sup>5</sup> Hence, this tripeptide provides a “clean” model system for hydrophobicity and


(4) (a) Yu, Y.; Monera, O. D.; Hodges, R. S.; Privalov, P. L. *J. Mol. Biol.* **1996**, *255*, 367–372. (b) Yu, Y.; Monera, O. D.; Hodges, R. S.; Privalov, P. L. *Biophys. Chem.* **1996**, *59*, 299–314.

(5) Kovacs, J. M.; Mant, C. T.; Hodges, R. S. *Biopolymers* **2006**, *84*, 283–297.


## SCHEME 1. Synthesis of 1



## SCHEME 2. Synthesis of 2



## SCHEME 3. Synthesis of 3



1-octanol/water partition measurements. A reference tripeptide, **5**, contains tryptophan (Trp) in place of  $\beta$ Fa. Trp is the most hydrophobic among the 20 natural amino acids<sup>5</sup> and serves as an excellent reference point for hydrophobicity measurements. Structures of the two tripeptides (**4** and **5**) are shown in Figure 2.

The commercially available pentaerythritol (**6**) provides an ideal starting material for the synthesis of compound **1**. Our synthesis commenced with the selective protection of pentaerythritol (Scheme 1). Protecting two of the four hydroxyl groups in pentaerythritol **6** with *p*-methoxybenzaldehyde gave the diol **7** with a good yield.<sup>6</sup> As the acidity of the hydroxyl group in perfluoro-*tert*-butanol is enhanced by the three electron-withdrawing  $-\text{CF}_3$  groups, perfluoro-*tert*-butanol is a good substrate for the Mitsunobu reaction to form perfluoro-*tert*-butyl ethers.<sup>7</sup> Thus, the Mitsunobu reaction was employed to introduce two perfluoro-*tert*-butyl groups into compound **7** in just one step to give fluorinated ether **8** with a 98% yield after the reaction mixture was stirred at 45 °C for 30 h. Such a high yield was achieved by carrying out the reaction in a sealed vessel and in the presence of 4 Å molecular sieves. Neither FC-72 ( $\text{C}_6\text{F}_{14}$ ) nor HFE-7100 (a mixture of *n*- $\text{C}_4\text{F}_9\text{OCH}_3$  and *i*- $\text{C}_4\text{F}_9\text{OCH}_3$ ) could extract compound **8** from the acetonitrile/water (95%/5%) solution of the reaction mixture. Instead, standard flash chromatography was used to purify ether **8** with a 98% yield. The *p*-methoxybenzylidene acetal protecting group was cleaved off of compound **8** by powdered aluminum chloride in the presence of anisole to give 1,3-diol **9** with quantitative yield. Treatment of 1,3-diol **9** with thionyl chloride gave a cyclic sulfite intermediate which was then oxidized by in situ-generated ruthenium tetroxide to give the cyclic sulfate **10** with an 84% yield in two steps. Ring opening of the cyclic sulfate **10** with sodium azide, followed by hydrolysis of the resulting sulfonic acid, gave the azido compound **11** with excellent yield which was then subjected to Jones oxidation to give the carboxylic acid **12** with an 85% yield. Hydrogenation of the azido group in compound **12** yielded the fluorinated amino acid **1** with

excellent yield. This completes the synthesis of the free amino acid, as depicted in Scheme 1.

To obtain the Fmoc-protected form of the amino acid, the amino group of compound **1** reacted with 9-fluorenylmethoxycarbonyl chloride (FmocCl) to give compound **2** with a 96% yield on a 13.3-gram scale, as depicted in Scheme 2.

To obtain the Boc-protected form of the amino acid, the azido group of compound **12** was reduced to the amino group, which then reacted with di-*tert*-butyl dicarbonate ( $\text{Boc}_2\text{O}$ ) to give compound **3** with a 96% yield on a 1.3-gram scale, as depicted in Scheme 3.

The two tripeptides (**4** and **5**) were synthesized using standard Fmoc chemistry on a solid support.<sup>8</sup> Both the carboxyl group and the amino group of  $\beta$ Fa showed good reactivity during solid-phase reaction. Hence,  $\beta$ Fa can be incorporated into any position of a target peptide through its carboxyl group, amino group, or both. Tripeptide **4** was purified by normal-phase HPLC, while tripeptide **5** was purified by reversed-phase HPLC due to their different solubilities in water and methanol. The molecular mass and the purity of each tripeptide were verified by mass spectrometry and analytical HPLC, respectively.

The  $^{19}\text{F}$  NMR spectra of the free amino acid **1** and the tripeptide **4** are shown in Figure 3. As designed, all 18 fluorine atoms give a sharp singlet  $^{19}\text{F}$  signal (line width  $\sim 0.01$  ppm) in both the free amino acid **1** (left) and the tripeptide **4** (right). This is also the case for the two protected forms (compounds **2** and **3**) of this amino acid (see Supporting Information).

The hydrophobicity of  $\beta$ Fa was evaluated in the context of tripeptide **4**, with tripeptide **5** serving as a reference point. To this end, we used the reversed-phase chromatography method developed by Hodges and co-workers, who determined the relative hydrophobicity order of 23 amino acids.<sup>5</sup> Figure 4 shows the chromatogram of the co-injection of the two tripeptides. Clearly, tripeptide **4** is much more retentive than tripeptide **5** in reversed-phase chromatography, proving that  $\beta$ Fa is much more hydrophobic than Trp, the most hydrophobic natural amino acid.

The 1-octanol/water partition coefficients ( $P_{\text{oct}}$ ) of the two tripeptides were evaluated.  $P_{\text{oct}}$  is a standard physicochemical parameter in assessing membrane permeability of peptides and

(6) Aguilera, B.; Romero-Ramirez, L.; Abad-Rodriguez, J.; Corrales, G.; Nieto-Sampedro, M.; Fernandez-Mayoralas, A. *J. Med. Chem.* **1998**, *41*, 4599–4606.

(7) Sebesta, D. P.; O'Rourke, S. S.; Pieken, W. A. *J. Org. Chem.* **1996**, *61*, 361–362.

(8) Chan, W. C.; White, P. D. *Fmoc Solid Phase Peptide Synthesis: A Practical Approach*; Oxford University Press: New York, 2000; pp 1–75.




FIGURE 3. Chemical shift (ppm) of  $^{19}\text{F}$  NMR in free amino acid **1** (left) and tripeptide **4** (right) (376 MHz,  $\text{CD}_3\text{OD}$ ,  $\text{C}_6\text{F}_6$  as internal standard).




FIGURE 4. Retention behavior of tripeptides **4** and **5** in reversed-phase HPLC.

other drugs.<sup>9</sup> For **5** (formyl-Gly-L-Trp-Gly-amide),  $P_{\text{Oct}} = 1/9.5$ , as determined by analytical HPLC at 280 nm (Trp signal) of the 1-octanol and aqueous phases. Therefore, in spite of the hydrophobicity of Trp, **5** still prefers water to 1-octanol. As for **4** (formyl-Gly- $\beta\text{Fa}$ -Gly-amide), after equilibration, its existence in water can be detected by neither analytical HPLC nor  $^{19}\text{F}$  NMR, while its existence in 1-octanol can be readily detected by both analytical HPLC and  $^{19}\text{F}$  NMR (see Supporting Information). Based on this result, we conclude that, for **4**,  $P_{\text{Oct}} \gg 10$ . Hence,  $P_{\text{Oct}}(\mathbf{4})$  is over 100 times larger than  $P_{\text{Oct}}(\mathbf{5})$ .

Results of the hydrophobicity measurement and the 1-octanol/water partition coefficient measurement are entirely consistent with each other, both showing that, in the context of a peptide,  $\beta\text{Fa}$  is very hydrophobic and strongly prefers the organic over the aqueous phase. All of these bode well for its potential use as a membrane permeability enhancer for peptide-based pharmaceuticals.

In conclusion, a novel, highly fluorinated  $\beta$ -amino acid ( $\beta\text{Fa}$ ) has been designed and synthesized. The Fmoc- and Boc-protected forms of this amino acid have also been synthesized with high yield. The syntheses were highly efficient with an overall

yield of ca. 65% at multigram synthesis scale.  $\beta\text{Fa}$  can be incorporated into peptides using standard solid-phase synthesis. As designed, all 18 fluorine atoms give a sharp singlet  $^{19}\text{F}$  NMR signal in both the free amino acid and the tripeptide context, auguring well for  $^{19}\text{F}$ -MRS monitoring.  $\beta\text{Fa}$  is much more hydrophobic than Trp, the most hydrophobic natural amino acid, and results in a much stronger preference of organic over aqueous phases.

### Experimental Section

**2-(4-Methoxyphenyl)-5,5-bis(2,2,2-trifluoro-1,1-bis-trifluoromethylethoxymethyl)-[1,3]-dioxane 8.** To a stirred mixture of compound **7** (24.4 g, 96.1 mmol), triphenylphosphine (75.5 g, 288.5 mmol), and 4 Å molecular sieves (20.0 g) in tetrahydrofuran (500 mL) at 0 °C was added dropwise diethylazodicarboxylate (50.2 g, 288.2 mmol). After the addition, the reaction mixture was allowed to warm to room temperature and was stirred for an additional 20 min. Then, perfluoro-*tert*-butanol (68.0 g, 288.0 mmol) was added in one portion, and the resulting mixture was stirred at 45 °C for 30 h in a sealed vessel. The mixture was evaporated to dryness, and the residue was dissolved in ethyl ether (600 mL). After filtered through a pad of Celite, the filtrate was washed with brine (300 mL), dried over sodium sulfate, and concentrated in vacuo. The residue was purified by flash column chromatography on silica gel (*n*-hexane/ethyl acetate = 20/1) to give **8** as a solid (65.1 g, 98%): mp 89–90 °C;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  3.78 (s, 1H), 3.80 (s, 4H), 3.89 (s, 2H), 4.13 (s, 1H), 4.16 (s, 1H), 4.49 (s, 2H), 5.39 (s, 1H), 6.88–6.92 (m, 2H), 7.33–7.37 (m, 2H);  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -73.42 (s);  $^{13}\text{C}$  NMR (100.7 MHz,  $\text{CDCl}_3$ )  $\delta$  39.2, 55.3, 66.2, 67.7, 68.7, 102.4, 113.8, 120.1 (q,  $J = 292.6$  Hz), 120.3 (q,  $J = 292.6$  Hz), 127.3, 129.9, 160.3; MS (CI)  $m/z$  691 ( $\text{M}^+ + 1$ , 100), 690 ( $\text{M}^+$ , 17), 583 (22); HRMS (CI) calcd for  $\text{C}_{21}\text{H}_{17}\text{F}_{18}\text{O}_5$ , 691.0787; found, 691.0792.

**2-Azidomethyl-3-(2,2,2-trifluoro-1,1-bis-trifluoromethylethoxy)-2-(2,2,2-trifluoro-1,1-bis-trifluoromethylethoxymethyl)propan-1-ol 11.** Sodium azide (4.4 g, 66.9 mmol) was added to a stirred solution of compound **10** (21.2 g, 33.4 mmol) in dimethylformaldehyde (120 mL). The reaction mixture was stirred at 60 °C for 4 h. The solvent was removed under vacuo, and the residue was dissolved in tetrahydrofuran (120 mL). Sulfuric acid (0.87 mL) and water (0.32 mL) were added to the stirred tetrahydrofuran solution, and the resulting mixture was stirred at room temperature for an additional 1 h. After removing the solvent, the residue was redissolved in dichloromethane (200 mL) and extracted with perfluorohexane (100 mL  $\times$  4). The combined extraction was washed with dichloromethane (10 mL) and concentrated under vacuo to give the pure azide **11** as a clear oil (19.3 g, 97%):  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  3.47 (s, 2H), 3.63 (s, 2H), 4.02 (s, 4H);  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -73.21 (s);  $^{13}\text{C}$  NMR (100.7

(9) (a) Abbruscato, T.; Williams, S.; Misicka, A.; Lipkowski, A. W.; Hryby, V. J.; Davis, T. P. *J. Pharmacol. Exp. Ther.* **1996**, *276*, 1049–1057. (b) Gentry, C. L.; Eggleton, R. D.; Gillespie, T.; Abbruscato, T. J.; Bechowski, H. B.; Hruby, V. J.; Davis, T. P. *Peptides* **1999**, *20*, 1229–1238.

MHz, CDCl<sub>3</sub>) δ 45.8, 49.8, 60.2, 66.8, 79.6 (m), 120.2 (q, *J* = 293.3 Hz); MS (CI) *m/z* 598 (M<sup>+</sup> + 1, 72), 570 (100); HRMS (CI) calcd for C<sub>13</sub>H<sub>10</sub>F<sub>18</sub>N<sub>3</sub>O<sub>3</sub>, 598.0435; found, 598.0418.

**2-Aminomethyl-3-(2,2,2-trifluoro-1,1-bis-trifluoromethylethoxy)-2-(2,2,2-trifluoro-1,1-bis-trifluoromethylethoxymethyl)propionic Acid 1.** A mixture of palladium on carbon (2.5 g) in methanol (200 mL) was degassed for 2 min and stirred under a hydrogen atmosphere for 30 min. A solution of acid **12** (10.7 g, 17.5 mmol) in methanol (10 mL) was then added, and the resulting mixture was stirred at room temperature under a hydrogen atmosphere for an additional 30 h. After solvent removal, the resulting residue was purified by flash column chromatography on silica gel (methanol/dichloromethane = 1/10) to give the amino acid **1** as a solid (10.1 g, 98%): mp 182–184 °C; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 2.99 (s, 2H), 4.22 (d, *J* = 9.2 Hz, 2H), 4.49 (d, *J* = 8.4 Hz, 2H); <sup>19</sup>F NMR (376 MHz, CD<sub>3</sub>OD) δ -71.00 (s); <sup>13</sup>C NMR (100.7 MHz, CD<sub>3</sub>OD) δ 41.7, 52.1, 69.9, 80.9 (m), 121.7 (q, *J* = 292.6 Hz), 175.0; MS (CI) *m/z* 586 (M<sup>+</sup> + 1, 100); HRMS (CI) calcd for C<sub>13</sub>H<sub>10</sub>F<sub>18</sub>NO<sub>4</sub>, 586.0322; found, 586.0285.

**2-[(9H-Fluoren-9-ylmethoxycarbonylamino)methyl]-3-(2,2,2-trifluoro-1,1-bis-trifluoromethylethoxy)-2-(2,2,2-trifluoro-1,1-bis-trifluoromethylethoxymethyl)propionic Acid 2.** To a stirred solution of amino acid **1** (10.1 g, 17.2 mmol) in tetrahydrofuran (100 mL) and water (100 mL) was added sodium carbonate (4.6 g, 42.9 mmol). After all of the sodium carbonate was dissolved, the resulting mixture was cooled to 0 °C, and 9-fluorenylmethyl chloroformate (6.7 g, 25.9 mol) was added in three portions. The resulting reaction mixture was stirred at room temperature overnight. The solvent was removed under vacuo, and the residue was purified by flash column chromatography on silica gel (*n*-hexane/ethyl acetate = 5/1) to give the acid **2** as a white solid (13.3 g, 96%): mp 104–105 °C; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 3.45 (s, 2H), 4.16 (m), 4.27 (m, 4H), 4.46 (d, *J* = 8.8 Hz, 2H), 7.26 (t, *J* = 7.2 Hz, 2H), 7.35 (t, *J* = 7.2 Hz, 2H), 7.60 (d, *J* = 7.6 Hz, 2H), 7.74 (d, *J* = 7.2 Hz, 2H); <sup>19</sup>F NMR (376 MHz, CD<sub>3</sub>OD) δ -71.00 (s); <sup>13</sup>C NMR (100.7 MHz, CD<sub>3</sub>OD) δ 36.9, 42.8, 53.7, 68.2, 69.0,

80.9 (m), 120.9, 121.7 (q, *J* = 292.6 Hz), 126.2, 128.1, 128.8, 142.6, 145.2, 158.8, 174.4; MS (CI) *m/z* 808 (M<sup>+</sup> + 1, 100); HRMS (CI) calcd for C<sub>28</sub>H<sub>20</sub>F<sub>18</sub>NO<sub>6</sub>, 808.1003; found, 808.1010.

**3-tert-Butoxycarbonylamino-2,2-bis(2,2,2-trifluoro-1,1-bis-trifluoromethylethoxymethyl)propionic Acid 3.** A mixture of palladium on carbon (200 mg) in methanol (20 mL) was degassed for 2 min and stirred under a hydrogen atmosphere for 30 min. A solution of acid **12** (1.2 g, 2.0 mmol) and di-*tert*-butyl dicarbonate (872 mg, 4.0 mmol) in methanol (5 mL) was then added, and the resulting mixture was stirred at room temperature under an atmosphere of hydrogen gas for 30 h. After solvent removal, the resulting residue was purified by flash column chromatography on silica gel (*n*-hexane/ethyl acetate = 5/1) to give the amino acid **3** as a solid (1.32 g, 96%): mp 128–130 °C; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 1.36 (s, 9H), 3.31 (s, 2H), 4.17 (d, *J* = 8.0 Hz, 2H), 4.39 (d, *J* = 8.0 Hz, 2H); <sup>19</sup>F NMR (376 MHz, CD<sub>3</sub>OD) δ -71.01 (s); <sup>13</sup>C NMR (100.7 MHz, CD<sub>3</sub>OD) δ 28.8, 42.8, 54.1, 69.6, 80.7, 81.1 (m), 121.9 (q, *J* = 293.3 Hz), 158.1, 177.2; MS (CI) *m/z* 686 (M<sup>+</sup> + 1, 10), 644 (100); HRMS (CI) calcd for C<sub>18</sub>H<sub>18</sub>F<sub>18</sub>NO<sub>6</sub>, 686.0847; found, 686.0815.

**Acknowledgment.** This work was supported by grants from the NIH (EB002880 and EB004416) and the Sidney Kimmel Foundation for Cancer Research. Y.B.Y. is a Kimmel scholar.

**Supporting Information Available:** Experimental procedures and product characterization for compounds **7**, **9**, **10**, and **12**, synthesis and partition procedures for **4** and **5**, copies of <sup>1</sup>H, <sup>19</sup>F, and <sup>13</sup>C NMR spectra for compounds **8**, **9**, **10**, **11**, **12**, **1**, **2**, and **3**, copies of HRMS spectra for compounds **1**, **2**, and **3**, copies of <sup>1</sup>H NMR and HPLC spectra for compounds **4** and **5**, copy of <sup>19</sup>F NMR spectra for compound **4**, and copies of HPLC spectra of partition test for compounds **4** and **5**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO0616308