


Short and Efficient Total Synthesis of Luotonin A and 22-Hydroxyacuminatine Using A Common Cascade Strategy

Hai-Bin Zhou,[†] Guan-Sai Liu,^{†,‡} and Zhu-Jun Yao^{*,†,‡}

State Key Laboratory of Bioorganic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Road, Shanghai 200032, China, and Department of Chemistry, University of Science and Technology of China, Hefei, Anhui 230026, China

yaoz@mail.sioc.ac.cn

Received April 21, 2007



Total syntheses of 22-hydroxyacuminatine and luotonin A were achieved in direct fashion and high overall yields (16% in eight steps and 47% in five steps, respectively). A mild cascade methodology was successfully applied in both syntheses as a common strategy. The new approach presents the advantages of short routes, high overall yields, use of stable intermediates, and ease of operations.

The potent antitumor activities and clinical applications of camptothecin-family alkaloids have attracted intense interest worldwide. Their parent representative analogue, camptothecin (CPT, **1**), serves as an excellent lead drug in anticancer research because of its potent antitumor activity (Figure 1).¹ The primary cellular target of CPT and many CPT derivatives is the covalent binary complex formed between DNA and topoisomerase I during DNA relaxation, and the stabilization of this complex by CPT is believed to lead to cell death.² Many imaginative syntheses of camptothecin and its analogues (including clinically used anticancer drugs topotecan (**2a**)³ and irinotecan (**2b**)⁴ and




FIGURE 1. Structures of camptothecin (**1**), 22-hydroxyacuminatine (**3**), luotonin A, (**4**), and two representative drugs, **2a** and **2b**.

some drug candidates⁵ as well) have emerged from numerous research groups over the past decades.⁶ 22-Hydroxyacuminatine (**3**) is another quinoline alkaloid isolated from Chinese medicinal plant (*Camptotheca acuminata*) (in a low isolated yield of 0.00006%) in 1989, showing significant cytotoxic activity against the murine leukemia P-388 cells (ED₅₀ 1.32 μg/mL) and KB (ED₅₀ 0.61 μg/mL) in vitro.⁷ As the only natural alkaloid containing a benz[6,7]indolizino[1,2-*b*]quinolin-11(13*H*)-one core, 22-hydroxyacuminatine attracted lots of attention. Three total syntheses of 22-hydroxyacuminatine have been achieved.⁸ Luotonin A (**4**) is one more CPT-family alkaloid isolated from another Chinese medicinal plant (*Peganum nigellastrum*) in 1997. Luotonin A also presents potent cytotoxicity against P-388 cells (IC₅₀ 1.8 μg/mL).⁹ Although it lacks the lactone functionality of CPT, luotonin A was recently demonstrated to stabilize the binary complex between human DNA and topoisomerase I and mediate topoisomerase I-dependent cytotoxicity in intact cells (IC₅₀ 5.7–12.6 μM).¹⁰ Several syntheses of luotonin A also have been reported.¹¹ Most importantly, Hecht and co-workers found that 14-azacamptothecin, a hybrid of luotonin A and CPT, is a water-soluble potent topoisomerase I poison.¹² Thus, further development of practical methods and general strategies for acquiring CPT-family alkaloids, including 22-

(4) (a) Negoro, S.; Fukuoka, M.; Masuda, N.; Takada, M.; Kusunoki, Y.; Matsui, K.; Takifuji, N.; Kudoh, S.; Niitani, H.; Taguchi, T. *J. Natl. Cancer Inst.* **1991**, *83*, 1164–1168. (b) Kawato, Y.; Aonuma, M.; Hirota, Y.; Kuga, H.; Sato, K. *Cancer Res.* **1991**, *51*, 4187–4191.

(5) (a) Cragg, G. M.; Newman, D. J. *J. Nat. Prod.* **2004**, *67*, 232–244. (b) Butler, M. S. *Nat. Prod. Rep.* **2005**, *22*, 162–195.

(6) For reviews on camptothecin and its derivatives, see: (a) Du, W. *Tetrahedron* **2003**, *59*, 8649–8687. (b) Wall, M. E.; Wani, M. C. In *The Alkaloids*; Cordell, G. A., Ed.; Academic Press: San Diego, 1998; Vol. 50, Chapter 13, pp 509–520. (c) Hutchinson, C. R. *Tetrahedron* **1981**, *37*, 1047–1065. For a related effort on cascade protocol for camptothecin, see: (d) Fortunak, J. M. D.; Mastrocola, A. R.; Mellinger, M.; Sisti, N. J.; Wood, J. L.; Zhuang, Z.-P. *Tetrahedron Lett.* **1996**, *37*, 5679–5682.

(7) Lin, L. Z.; Cordell, G. A. *Phytochemistry* **1989**, *28*, 1295–1297.

(8) (a) Ma, Z.; Lee, D. Y. W. *Tetrahedron Lett.* **2004**, *45*, 6721–6723. (b) Xiao, X.; Antony, S.; Pommier, Y.; Cushman, M. *J. Med. Chem.* **2006**, *49*, 1408–1412. (c) Babjak, M.; Kanazawa, A.; Anderson, R. J.; Greene, A. E. *Org. Biomol. Chem.* **2006**, *4*, 407–409.

(9) Ma, Z. Z.; Hano, Y.; Nomura, T.; Chen, Y.-J. *Heterocycles* **1997**, *46*, 541–546.

(10) Cagir, A.; Jones, S. H.; Gao, R.; Eisenhauer, B. M.; Hecht, S. M. *J. Am. Chem. Soc.* **2003**, *125*, 13628–13629.

[†] State Key Laboratory of Bioorganic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences.

[‡] Department of Chemistry, University of Science and Technology of China.

(1) Wall, M. E.; Wani, M. C.; Cook, C. E.; Palmer, K. H.; McPhail, A. T.; Sim, G. A. *J. Am. Chem. Soc.* **1966**, *88*, 3888–3890.

(2) (a) Hsiang, Y. H.; Hertzberg, R.; Hecht, S. M.; Liu, L. F. *J. Biol. Chem.* **1985**, *260*, 14873–14878. (b) Kohn, K. W.; Pommier, Y. *Ann. N.Y. Acad. Sci.* **2000**, *922*, 11–26. (c) Staker, B. L.; Hjerrild, K.; Feese, M. D.; Behnke, C. A.; Burgin, A. B., Jr.; Stewart, L. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 15387–15392.

(3) Kingsbury, W. D.; Boehm, J. C.; Jakas, D. R.; Holden, K. G.; Hecht, S. M.; Gallagher, G.; Caranfa, M. J.; McCabe, F. L.; Faucette, L. F.; Johnson, R. K.; Hertzberg, R. P. *J. Med. Chem.* **1991**, *34*, 98–107.





FIGURE 2. Retrosynthesis of 22-hydroxyacuminatine (**3**) and luotonin A (**4**) using a common cascade strategy.

hydroxyacuminatine (**3**), luotonin A (**4**), and other new CPT derivatives, is of great value.

More recently, we reported a mild cascade methodology to construct variously substituted indolizino[1,2-*b*]quinolin-9(11*H*)-ones, the tetracyclic A/B/C/D-ring core of CPT-family alkaloids.^{6d,13} This highly efficient cascade reaction triggered by bis(triphenyl)oxodiphosphonium trifluoromethanesulfonate under mild conditions has been successfully applied in total synthesis of camptothecin (**1**).¹³ In order to examine the power of this methodology, two additional naturally occurring alkaloids, 22-hydroxyacuminatine (**3**) and luotonin A (**4**) were selected for its further applications. Both of them are notable with structural similarities to camptothecin (**1**) in identical A–C rings. Herein, we report our results of total syntheses of 22-hydroxyacuminatine (**3**) and luotonin A (**4**) using the above-mentioned cascade reaction as a common strategy. These two targets were successfully achieved in very short steps and high overall yields. Figure 2 illustrates their general retrosynthetic analysis. With the consideration of employing our cascade reaction conditions, chemically stable amides **5** and **6** have to be prepared at first, respectively.

Synthesis of 22-hydroxyacuminatine (**3**) started from the commercially available 2,6-dicyanotoluene **9** (Scheme 1).

SCHEME 1. Eight-Step Formal Total Synthesis of 22-Hydroxyacuminatine (**3**)



Condensation of 2,6-dicyanotoluene **9** with diethyl oxalate in the presence of potassium *tert*-butoxide, followed by a basic treatment, yielded 5-cyano-1-oxo-1,2-dihydroisoquinoline-3-carboxylic acid (**10**) in 63% yield.¹⁴ Treatment of **10** with oxalyl chloride followed by reaction with excess methanol afforded ester **11** (88%). *N*-Propargylation of **11** was carried out with propargyl bromide, K₂CO₃, LiBr, and a catalytic amount of tetrabutylammonium bromide in toluene, giving a new pyridone **8** in 91% yield.¹⁵ Basic hydrolysis of methyl ester **8** afforded acid **7** in 98% yield. Conversion of acid **7** to its corresponding acid chloride followed by treatment with aniline afforded amide **5** (70% yield in two steps). Amide precursor **5** was then treated with bis(triphenyl)oxodiphosphonium trifluoromethanesulfonate (prepared in situ) at room temperature for half an hour.¹³ The known advanced intermediate **12**^{8b} was finally obtained (91% yield), containing the whole skeleton of 22-hydroxyacuminatine (**3**). Conversion of **12** to 22-hydroxyacuminatine (**3**) has been previously carried out by Cushman and co-workers using a two-step DIBAL-H reduction procedure (51% yield).^{8b} Thus, using our mild cascade reaction, we accomplished a straightforward synthesis of the known key intermediate **12** (six steps and 31% yield) starting from commercially available 2,6-dicyanotoluene **9**. This represents a formal total synthesis of 22-hydroxyacuminatine (**3**) in eight steps and 16% overall yield.

Employing a similar strategy, total synthesis of luotonin A (**4**) was accomplished in five steps (Scheme 2). Reaction of the commercially available anthranilamide **13** with diethyl oxalate yielded a quinazolinone derivative **14** in 81% yield.¹⁶ Ethyl ester **14** was then hydrolyzed with LiOH to give acid **15**.¹⁷ Conversion

(11) Synthesis of luotonin A: (a) Wang, H.; Ganesan, A. *Tetrahedron Lett.* **1998**, *39*, 9097–9098. (b) Ma, Z. Z.; Hano, Y.; Nomura, T.; Chen, Y.-J. *Heterocycles* **1999**, *51*, 1593–1596. (c) Kelly, T. R.; Chamberland, S.; Silva, R. A. *Tetrahedron Lett.* **1999**, *40*, 2723–2724. (d) Molina, P.; Tarraga, A.; Gonzalez-Tejedor, A. *Synthesis* **2000**, *11*, 1523–1525. (e) Toyota, M.; Komori, C.; Ihara, M. *Heterocycles* **2002**, *56*, 101–104. (f) Toyota, M.; Komori, C.; Ihara, M. *ARKIVOC* **2003**, 15–23. (g) Dallavalle, S.; Merlini, L. *Tetrahedron Lett.* **2002**, *43*, 1835–1838. (h) Yadav, J. S.; Reddy, B. V. S. *Tetrahedron Lett.* **2002**, *43*, 1905–1908. (i) Osborne, D.; Stevenson, P. J. *Tetrahedron Lett.* **2002**, *43*, 5469–5470. (j) Lee, E. S.; Park, J. G.; Jahng, Y. *Tetrahedron Lett.* **2003**, *44*, 1883–1886. (k) Harayama, T.; Morikami, Y.; Shigeta, Y.; Abe, H.; Takeuchi, Y. *Synlett* **2003**, 6, 847–848. (l) Harayama, T.; Hori, A.; Serban, G.; Morikami, Y.; Matsumoto, T.; Abe, H.; Takeuchi, Y. *Tetrahedron* **2004**, *60*, 10645–10650. (m) Mhaske, S. B.; Argade, N. P. *J. Org. Chem.* **2004**, *69*, 4563–4566. (n) Chavan, S. P.; Sivappa, R. *Tetrahedron* **2004**, *60*, 9931–9936. (o) Twin, H.; Batey, R. A. *Org. Lett.* **2004**, *6*, 4913–4916. (p) Ma, Z. Z.; Hano, Y.; Nomura, T. *Heterocycles* **2005**, *65*, 2203–2219. (q) Bowman, W. R.; Cloonan, M. O.; Fletcher, A. J.; Stein, T. *Org. Biomol. Chem.* **2005**, *3*, 1460–1467. (r) Tangirala, R.; Antony, S.; Agama, K.; Pommier, Y.; Curran, D. P. *Synlett* **2005**, *18*, 2843–2846. (s) Servais, A.; Azzouz, M.; Lopes, D.; Courillon, C.; Malacria, M. *Angew. Chem., Int. Ed.* **2007**, *46*, 576–579.

(12) (a) Cheng, K.; Rahier, N. J.; Eisenhauer, B. M.; Thomas, S. J.; Gao, R.; Hecht, S. M. *J. Am. Chem. Soc.* **2005**, *127*, 838–839. (b) Elban, M. A.; Sun, W.; Eisenhauer, B. M.; Gao, R.; Hecht, S. M. *Org. Lett.* **2006**, *8*, 3513–3516.

(13) Zhou, H.-B.; Liu, G.-S.; Yao, Z.-J. *Org. Lett.* **2007**, *9*, 2003–2006.


(14) Wenkert, E.; Johnston, D. B. R.; Dave, K. G. *J. Org. Chem.* **1964**, *29*, 2534–2542.

(15) (a) Ricoh, I.; Halvorsen, K.; Dubrule, C.; Lattes, A. J. *J. Org. Chem.* **1994**, *59*, 415. (b) Liu, H.; Ko, S.-B.; Josien, H.; Curran, D. P. *Tetrahedron Lett.* **1995**, *36*, 8917.

(16) Joshi, V.; Chaudhari, R. P. *Indian J. Chem.* **1987**, *26B*, 602–604.

(17) Rastogi, R.; Sharma, S. *Indian J. Chem.* **1982**, *21B*, 744–746.

SCHEME 2. Five-Step Total Synthesis of Luotonin A



of **15** to its corresponding acid chloride, followed by coupling with aniline, afforded the amide **16** in 67% yield (from **14**). *N*-Alkylation of amide **16** with propargyl bromide provided the precursor amide **6** in 87% yield.¹⁵ Finally, the cascade annulation was carried out by simple treatment of **6** with bis(triphenyl)oxodiphosphonium trifluoromethanesulfonate at room temperature for 1 h, affording luotonin A (**4**) in 99% yield. This five-step synthesis of luotonin A represents an efficient overall yield of 47% from the commercially available **13**. To best of our knowledge, this is the shortest synthesis route for luotonin A. Compared with a similar strategy previously used by Batey and Twin,¹¹ our approach presents the advantages of a shorter route, higher overall yield (especially for the key cascade annulation step), and the use of the stable amide intermediate.

In conclusion, highly efficient and concise total syntheses of 22-hydroxyacuminatine and luotonin A were accomplished in direct fashions (eight steps and five steps, respectively) and high overall yields (16% and 47%, respectively). A mild and efficient cascade reaction was successfully employed as the common annulation strategy. Further applications of this useful and highly efficient cascade annulation protocol to construction of new CPT derivatives and other structurally related bioactive molecules are underway in our laboratory, and these results will be reported in due course.

Experimental Section

7-Cyano-12*H*-5,11a-diazadibenzo[*b,h*]fluoren-11-one (**12**).^{8b}

To a solution of triphenylphosphine oxide (0.77 g, 2.75 mmol) in dry CH₂Cl₂ (20 mL) was added trifluoromethanesulfonic anhydride (0.23 mL, 1.38 mmol) slowly at 0 °C. After the mixture was stirred at 0 °C for 10 min, amide **5** (0.30 g, 0.92 mmol) was then added at the same temperature. The reaction was allowed to warm to room temperature. After 0.5 h, the reaction mixture was quenched by addition of 10% aqueous NaHCO₃ solution. The aqueous layer was extracted with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄, filtered, and concentrated. The crude product was purified by flash chromatography on silica gel to afford **12**^{8b} (0.26 g, 91%). ¹H NMR (CDCl₃, 300 MHz): δ 8.77 (1H, d, *J* = 8.1 Hz), 8.40 (1H, s), 8.28 (1H, d, *J* = 8.4 Hz), 8.09 (1H, d, *J* = 7.5 Hz), 8.02 (1H, s), 7.94 (1H, d, *J* = 7.8 Hz), 7.85 (1H, t, *J* = 7.8 Hz), 7.67 (1H, t, *J* = 7.5 Hz), 7.63 (1H, t, *J* = 8.0 Hz), 5.41 (2H, s). HRMS (MALDI, *m/z*) calcd for C₂₀H₁₁N₃O (M+H)⁺: 310.0980; found, 310.0975. IR (KBr): 2229, 1664, 1637, 1591, 1550, 1473, 1394, 1377, 1296, 1253, 1205, 868, 816, 760, 705, 599 cm⁻¹. Anal. calcd for C₂₀H₁₁N₃O: C, 77.66; H, 3.58; N, 13.58; found: C, 77.65; H, 3.58; N, 13.57.

Luotonin A (4).^{9,11} Compound **4** was prepared by a procedure similar to that for compound **12**, and it was isolated in 99% yield by column chromatography on silica gel (CH₂Cl₂/CH₃OH, 150:1 to 80:1) as a white solid; mp: 278–280 °C dec. ¹H NMR (CDCl₃, 500 MHz): δ 8.46 (1H, d, *J* = 8.5 Hz), 8.43–8.40 (2H, m), 8.11 (1H, d, *J* = 8.1 Hz), 7.92 (1H, d, *J* = 8.1 Hz), 7.86 (1H, d, *J* = 8.2 Hz), 7.82 (1H, d, *J* = 8.2 Hz), 7.66 (1H, t, *J* = 7.5 Hz), 7.57 (1H, t, *J* = 7.5 Hz), 5.32 (2H, s). ¹³C NMR (CDCl₃, 125 MHz): δ 160.6, 152.6, 151.2, 149.5, 134.6, 131.5, 130.72, 130.68, 129.5, 128.8 (2C), 128.5, 127.9, 127.4, 126.4, 121.3, 47.3. IR (KBr): 1678, 1630, 1606, 1572, 1467, 1360, 1326, 1237, 769, 690, 604 cm⁻¹. HRMS (ESI, *m/z*) calcd for C₁₈H₁₂N₃O (M + H)⁺: 286.0975; found, 286.0990. Anal. calcd for C₁₈H₁₁N₃O: C, 75.78; H, 3.89; N, 14.73; found: C, 75.80; H, 4.10; N, 14.64.

Acknowledgment. This project is financially supported by National Natural Science Foundation (20425205, 20621062), Chinese Academy of Sciences (KGCX2-SW-213, KJCX2-YW-H08), and the Shanghai Municipal Committee of Science and Technology (06QH14016, 04DZ14901).

Supporting Information Available: Experimental details and characterization of new compounds, copies of NMR spectra for new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO070837D