


Mild Conditions for Pd-Catalyzed Carboamination of *N*-Protected Hex-4-enylamines and 1-, 3-, and 4-Substituted Pent-4-enylamines. Scope, Limitations, and Mechanism of Pyrrolidine Formation

Myra Beaudoin Bertrand,[†] Joshua D. Neukom, and John P. Wolfe*

Department of Chemistry, University of Michigan, 930 N. University Avenue,
Ann Arbor, Michigan, 48109-1055

jpwolfe@umich.edu

Received July 23, 2008



The use of the weak base Cs_2CO_3 in Pd-catalyzed carboamination reactions of *N*-protected γ -aminoalkenes with aryl bromides leads to greatly increased tolerance of functional groups and alkene substitution. Substrates derived from (*E*)- or (*Z*)-hex-4-enylamines are stereospecifically converted to 2,1'-disubstituted pyrrolidine products that result from suprafacial addition of the nitrogen atom and the aryl group across the alkene. Transformations of 4-substituted pent-4-enylamine derivatives proceed in high yield to afford 2,2-disubstituted products, and *cis*-2,5- or *trans*-2,3-disubstituted pyrrolidines are generated in good yield with excellent diastereoselectivity from *N*-protected pent-4-enylamines bearing substituents at C1 or C3. The reactions tolerate a broad array of functional groups, including esters, nitro groups, and enolizable ketones. The scope and limitations of these transformations are described in detail, along with models that account for the observed product stereochemistry. In addition, deuterium labeling experiments, which indicate these reactions proceed via *syn*-aminopalladation of intermediate palladium(aryl)(amido)complexes regardless of degree of alkene substitution or reaction conditions, are also discussed.

Introduction

Substituted pyrrolidines are common motifs displayed in a wide array of interesting biologically active compounds, including natural products and pharmaceutically relevant molecules.¹ Due to the significance of pyrrolidine derivatives, a large number of methods have been developed for their synthesis.^{1a,2} However, few existing methods effect closure of the pyrrolidine ring with concomitant formation of a C–C bond and a stereocenter

adjacent to the ring.³ In addition, the invention of new methods for pyrrolidine synthesis that can be used to prepare many different analogs from a common precursor has great potential utility in medicinal chemistry applications.

Over the past several years, our group has developed and explored a new method for the synthesis of substituted pyrro-


[†] Current address: Discovery Chemistry, Bristol-Myers Squibb Company, Route 206 and Province Line Road, Princeton, NJ, 08543.

(1) For recent reviews, see: (a) Bellina, F.; Rossi, R. *Tetrahedron* **2006**, *62*, 7213–7256. (b) Hackling, A. E.; Stark, H. *Chem. BioChem* **2002**, *3*, 946–961. (c) Lewis, J. R. *Nat. Prod. Rep.* **2001**, *18*, 95–128.

(2) For recent reviews, see: (a) Coldham, I.; Hufton, R. *Chem. Rev.* **2005**, *105*, 2765–2810. (b) Pyne, S. G.; Davis, A. S.; Gates, N. J.; Hartley, J. P.; Lindsay, K. B.; Machan, T.; Tang, M. *Synlett* **2004**, 2670–2680. (c) Felpin, F.-X.; Lebreton, J. *Eur. J. Org. Chem.* **2003**, 3693–3712. (d) Pichon, M.; Figadere, B. *Tetrahedron: Asymmetry* **1996**, *7*, 927–964.

(3) For examples of Cu-catalyzed intramolecular carboamination of *N*-(arylsulfonyl)-2-allylanilines and related derivatives, see: (a) Sherman, E. S.; Fuller, P. H.; Kasi, D.; Chemler, S. R. *J. Org. Chem.* **2007**, *72*, 3896–3905. (b) Sherman, E. S.; Chemler, S. R.; Tan, T. B.; Gerlits, O. *Org. Lett.* **2004**, *6*, 1573–1575. For Pd(II)-catalyzed alkoxyacylation of alkenes bearing tethered nitrogen nucleophiles, see: (c) Harayama, H.; Abe, A.; Sakado, T.; Kimura, M.; Fugami, K.; Tanaka, S.; Tamaru, Y. *J. Org. Chem.* **1997**, *62*, 2113–2122. For carboamination reactions between alkenes and *N*-allylsulfonamides, see: (d) Scarborough, C. C.; Stahl, S. S. *Org. Lett.* **2006**, *8*, 3251–3254. For carboamination of vinylcyclopropanes, see: (e) Larock, R. C.; Yum, E. K. *Synlett* **1990**, 529–530. For 1,1-carboamination of alkenes, see: (f) Harris, G. D., Jr.; Herr, R. J.; Weinreb, S. M. *J. Org. Chem.* **1992**, *57*, 2528–2530. (g) Harris, G. D., Jr.; Herr, R. J.; Weinreb, S. M. *J. Org. Chem.* **1993**, *58*, 5452–5464. (h) Larock, R. C.; Yang, H.; Weinreb, S. M.; Herr, R. J. *J. Org. Chem.* **1994**, *59*, 4172–4178.

lidines via Pd-catalyzed carboamination reactions of *N*-substituted γ -aminoalkenes with aryl/alkenyl halides (eq 1).^{4–6} These transformations effect the construction of the pyrrolidine ring with simultaneous formation of one C–N bond, one C–C bond, and up to two stereocenters from simple precursors.³ In addition, this method can easily be used to generate many different analogs from a single alkene substrate, as a large number of aryl, heteroaryl, and alkenyl halides are readily available.



Despite the utility of these transformations, the original reaction conditions employed the strong base NaOtBu as a reagent, which led to several significant limitations. For example, many common functional groups such as esters, nitro groups, or enolizable ketones were not tolerated under the strongly basic conditions. In addition, the scope of this method was usually limited to substrates bearing terminal alkenes, as almost all attempted transformations of disubstituted alkenes either resulted in competing base-mediated substrate decomposition or formation of complex mixtures of regioisomers.^{7,8}

We recently reported the discovery of mild conditions for Pd-catalyzed carboamination reactions of *N*-protected γ -aminoalkenes, in which the strong base NaOtBu is replaced with the weak base Cs₂CO₃. These conditions significantly expand the scope of the carboamination method by allowing the use of functionalized aryl halide coupling partners.⁹ In this Article we describe our full studies on the development, scope, and limitations of Pd-catalyzed carboamination reactions conducted under these new conditions. In addition to illustrating improved functional group tolerance, these studies demonstrate that transformations of disubstituted alkenes can now be achieved when an appropriate catalyst system is employed. We also present experiments that provide additional evidence for our previously described mechanistic hypothesis,⁵ which involves alkene *syn*-aminopalladation as a key step in the catalytic cycle. These experiments suggest the pyrrolidine products are formed through a common mechanism regardless of the degree of alkene substitution or the nature of the base (NaOtBu vs Cs₂CO₃).

(4) For reviews, see: (a) Wolfe, J. P. *Eur. J. Org. Chem.* **2007**, 571–582. (b) Wolfe, J. P. *Synlett* **2008**, accepted for publication.

(5) (a) Ney, J. E.; Wolfe, J. P. *Angew. Chem., Int. Ed.* **2004**, *43*, 3605–3608. (b) Lira, R.; Wolfe, J. P. *J. Am. Chem. Soc.* **2004**, *126*, 13906–13907. (c) Bertrand, M. B.; Wolfe, J. P. *Tetrahedron* **2005**, *61*, 6447–6459. (d) Ney, J. E.; Wolfe, J. P. *J. Am. Chem. Soc.* **2005**, *127*, 8644–8651. (e) Yang, Q.; Ney, J. E.; Wolfe, J. P. *Org. Lett.* **2005**, *7*, 2575–2578. (f) Ney, J. E.; Hay, M. B.; Yang, Q.; Wolfe, J. P. *Adv. Synth. Catal.* **2005**, *347*, 1614–1620. (g) Bertrand, M. B.; Wolfe, J. P. *Org. Lett.* **2006**, *8*, 2353–2356.

(6) For closely related approaches to the synthesis of piperazines, imidazolidin-2-ones, and isoxazolidines via Pd-catalyzed alkene carboamination reactions, see: (a) Nakhla, J. S.; Wolfe, J. P. *Org. Lett.* **2007**, *9*, 3279–3282. (b) Fritz, J. A.; Nakhla, J. S.; Wolfe, J. P. *Org. Lett.* **2006**, *8*, 2531–2534. (c) Fritz, J. A.; Wolfe, J. P. *Tetrahedron* **2008**, *64*, 6838–6852. (d) Peng, J.; Jiang, D.; Lin, W.; Chen, Y. *Org. Biomol. Chem.* **2007**, *5*, 1391–1396. (e) Peng, J.; Lin, W.; Yuan, S.; Chen, Y. *J. Org. Chem.* **2007**, *72*, 3145–3148. (f) Dongol, K. G.; Tay, B. Y. *Tetrahedron Lett.* **2006**, *47*, 927–930.

(7) Secondary carbamates decompose to afford reactive isocyanates when heated in the presence of NaOtBu. See: Tom, N. J.; Simon, W. M.; Frost, H. N.; Ewing, M. *Tetrahedron Lett.* **2004**, *45*, 905–906.

(8) The only substrates of this type that have been effectively employed in Pd-catalyzed carboamination reactions are *N*-protected cyclopent-2-enylethylamine derivatives. See refs 5a, 5c, and 5d.

(9) A portion of these studies have been previously communicated. See: (a) Bertrand, M. B.; Leathen, M. L.; Wolfe, J. P. *Org. Lett.* **2007**, *9*, 457–460.

TABLE 1. Summary of Reaction Optimization^a

entry	base	Pd	solvent	yield (%)
1	NaOtBu	Pd ₂ (dba) ₃	toluene	81
2	Cs ₂ CO ₃	Pd ₂ (dba) ₃	toluene	38
3	Cs ₂ CO ₃	Pd(OAc) ₂	toluene	63
4	Cs ₂ CO ₃	Pd(OAc) ₂	dioxane	82 ^b


^a Conditions: 1.0 equiv substrate, 1.2 equiv ArBr, 2.3 equiv base, 1 mol % Pd₂(dba)₃ (2 mol % Pd) or 2 mol % Pd(OAc)₂, 2 mol % Dpe-phos (with Pd₂(dba)₃) or 4 mol % Dpe-phos (with Pd(OAc)₂), solvent (0.25 M), 105 °C, 17–27 h (reaction times are unoptimized).
^b Reaction was conducted at 100 °C.

Results

Optimization Studies. To develop mild conditions for Pd-catalyzed carboamination reactions we initially examined the coupling of a simple substrate, *N*-Boc-pent-4-enylamine (**1**), with an electron-neutral aryl bromide (4-bromo-*tert*-butylbenzene) in the presence of a catalyst composed of Pd₂(dba)₃/Dpe-phos.¹⁰ As shown in Table 1, this transformation afforded **2** in 81% yield using our original, strongly basic, conditions (NaOtBu, toluene, 105 °C). A number of organic and inorganic bases were screened under otherwise similar reaction conditions, and disappointing results were obtained. In general, use of organic bases (e.g., Et₃N) failed to provide any of the desired product, and most inorganic bases were also ineffective, although small amounts of **2** were occasionally observed.¹¹ The most promising results were obtained with Cs₂CO₃, but the yield of **2** was still modest (38%, entry 2).¹²

To further optimize conditions with Cs₂CO₃ as base, we examined the effect of other reaction parameters. We quickly discovered that use of Pd(OAc)₂ in place of Pd₂(dba)₃ provided significantly improved yields (63%, entry 3). After conducting reactions in several different solvents, we arrived at much improved conditions in which **2** was obtained in 82% yield when dioxane was used as solvent (entry 4).¹³

(10) Dpe-phos = bis(2-diphenylphosphino)phenyl ether. Xantphos = 9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene. Nixantphos = 4,6-bis(diphenylphosphino)phenoxazine. BINAP = 2,2'-diphenylphosphino-1,1'-binaphthyl. Dppe = 1,2-bis(diphenylphosphino)ethane.



(11) In most instances the use of other bases led to very slow consumption of starting material and/or competing Heck arylation of the alkene moiety.

(12) The use of Cs₂CO₃ and other weak bases in Pd-catalyzed *N*-arylation reactions of amines with aryl halides has been reported. For reviews, see: (a) Muci, A. R.; Buchwald, S. L. *Top. Curr. Chem.* **2002**, *219*, 131–209. (b) Hartwig, J. F. In *Modern Arene Chemistry*; Astruc, D., Ed.; Wiley-VCH, Weinheim, 2002; pp 107–168. (c) Schlummer, B.; Scholz, U. *Adv. Synth. Catal.* **2004**, *346*, 1599–1626.

TABLE 2. Palladium-Catalyzed Carboamination of *N*-Protected γ -Aminoalkenes with Functionalized Aryl Bromides^a

Entry	Amine	Aryl bromide	Product	Rxn. Time (h)	Yield (%) ^b
1				15 11	75 (77) ^{c,e}
2				27 26	82 (81) ^{c,e}
3				20	78 ^d
4				18	76 ^d
5				28	71
6				18 15	79 (72) ^{c,e}
7				18	76 ^{d,f}
8				16 18	88 (17) ^c
9				18	88 ^d

^a Conditions: 1.0 equiv amine, 1.2 equiv ArBr, 2.3 equiv Cs₂CO₃, 2 mol % Pd(OAc)₂, 4 mol % Dpe-phos, dioxane (0.2–0.25 M), 100–105 °C. ^b Yield refers to average isolated yield obtained in two or more experiments. ^c NaOtBu was used in place of Cs₂CO₃ and toluene was used as solvent. ^d Reaction was conducted at 85 °C in DME solvent. ^e Pd₂(dba)₃ was used in place of Pd(OAc)₂. ^f Dppe was used in place of Dpe-phos.

Studies on Scope and Limitations. Reactions of Terminal Alkene Substrates. Having successfully optimized conditions for the coupling of **1** with 4-bromo-*tert*-butylbenzene, we proceeded to explore the utility of these conditions for transformations of different substrate combinations. As shown in Tables 2 and 3, a number of *N*-protected pent-4-enylamine derivatives are effectively converted to 2-benzylpyrrolidines in good yield. Although the functional group tolerance of Pd-catalyzed carboamination reactions is modest when NaOtBu is used as base, a broad variety of functional groups are tolerated with the new, mild reaction conditions. As shown below, aldehydes, enolizable ketones, nitro groups, methyl esters, and alkyl acetates can be incorporated into the pyrrolidine products. In addition, the carboamination reactions of electron-rich, electron-neutral, and heterocyclic aryl bromides proceed with good chemical yields. However, use of dppe¹⁰ as ligand was necessary to suppress competing N-arylation when 4-bromonitrobenzene was used as a coupling partner (Table 2, entry 7).

(13) In some cases use of DME as solvent provided comparable results to those obtained with dioxane (see Table 2, entries 3, 4, 7, and 9).

TABLE 3. Palladium-Catalyzed Carboamination of Substituted *N*-Protected γ -Aminoalkenes with Functionalized Aryl Bromides^a

Entry	Amine	Aryl bromide	Product	dr	Rxn. Time (h)	Yield (%) ^b
1				12:1	20	80
2				15:1	16	76
3				14:1	18	73
4				>20:1	16	75
5				>20:1	18	74
6				>20:1	23 18	71 (62) ^{c,d}
7				>20:1	20	73


^a Conditions: 1.0 equiv amine, 1.2 equiv ArBr, 2.3 equiv Cs₂CO₃, 2 mol % Pd(OAc)₂, 4 mol % Dpe-phos, dioxane (0.2–0.25 M), 100 °C. ^b Yield refers to average isolated yield obtained in two or more experiments. ^c NaOtBu was used in place of Cs₂CO₃ and toluene was used as solvent. ^d Pd₂(dba)₃ was used in place of Pd(OAc)₂.

Reactions of unfunctionalized aryl bromides with *N*-Boc- or *N*-acyl-pent-4-enylamine (**1** or **3**) provided similar yields regardless of whether NaOtBu or Cs₂CO₃ was used as base (Table 2, entries 1, 2, and 6). However, the new reaction conditions provided excellent results for the carboamination of Cbz-protected substrate **4** with 2-bromonaphthalene (88% yield of **11**, entry 8), whereas use of NaOtBu for this transformation provided only 17% yield of **11**. Cleavage of the Cbz-group from the substrate was problematic when the reaction was conducted with the stronger base.

The mild conditions are also effective for stereoselective reactions of *N*-Boc- and *N*-Cbz-protected pent-4-enylamine derivatives bearing substituents adjacent to the nitrogen atom and/or adjacent to the alkene. For example, substrates **13** and **14**, which are substituted at the allylic position, were converted to *trans*-2,3-disubstituted pyrrolidines **18**–**20** with 12–15:1

diastereoselectivity (Table 3, entries 1–3).¹⁴ The synthesis of *cis*-2,5-disubstituted pyrrolidine products **21** and **22** was achieved in good yields and >20:1 diastereoselectivity through carboamination of substrates **15** and **16**, which are substituted adjacent to the nitrogen atom (entries 4 and 5). The coupling of **17** with bromobenzene provided **23**, which is an intermediate in the synthesis of the natural product preussin,^{5g} in 71% yield with >20:1 dr (Table 3, entry 6). Similarly good results were obtained for the coupling of **17** with methyl-4-bromobenzoate (Table 3, entry 7).

Reactions of Hex-4-enylamine Derivatives. Palladium-catalyzed carboamination reactions of substrates bearing internal alkenes are of considerable potential utility, as these transformations would create two new contiguous stereocenters in one step. However, efforts to convert these substrates to substituted pyrrolidines under conditions in which NaOtBu was used as a base were largely unsuccessful. Pd-catalyzed reactions of aryl bromides with hex-4-enylamines bearing *N*-aryl substituents provided complex mixtures of products,^{5a} and transformations involving *N*-Boc-protected substrates proceeded in low yield. For example, treatment of (*E*)-alkene **25** with 2-bromonaphthalene and NaOtBu in the presence of the Pd(OAc)₂/Dpe-phos catalyst afforded only 26% yield of **26** (eq 2).^{15,16}



To improve the efficiency of reactions involving acyclic internal alkene substrates, we investigated the cyclization of **25** using Cs₂CO₃ as base and dioxane as solvent. After considerable screening, we were gratified to find that use of Nixantphos¹⁰ as ligand led to the conversion of **25** to pyrrolidine product **26**, which results from suprafacial addition across the alkene, in 59% yield with >20:1 dr (Table 4, entry 1). Although Nixantphos provided the best results for the coupling of **25** with 2-bromonaphthalene, subsequent experiments indicated that use of (±)-BINAP as ligand provided the best results for most systems examined. Side products resulting from competing *N*-arylation of the substrate were usually formed in lesser amounts with BINAP than Nixantphos.

Yields in the transformations of (*E*)-alkene **25** were generally modest, although diastereoselectivities were uniformly high (>20:1 dr). Relatively long reaction times were required for these transformations (27–44 h), and in some instances (entries 2 and 3) the formation of ca. 10–20% of regioisomeric side products (**31**, **33**)¹⁷ was also observed. Electron-poor aryl bromides could be employed as coupling partners, but use of electron-rich aryl bromides, such as 4-bromoanisole, led to low conversions (<50%) and formation of large amounts of side-products resulting from competing Heck arylation of the alkene. Reactions of Cbz-protected (*E*)-alkene substrate **27** proceeded in modest yield, but with excellent (≥20:1) regioselectivity and diastereoselectivity (entries 4 and 5).

(14) See Supporting Information for details on the assignment of product stereochemistry.

(15) Bertrand, M. B.; Wolfe, J. P. Unpublished results.

(16) Use of Nixantphos as the ligand for this transformation resulted in the formation of **26** in 22% yield with >20:1 dr.

TABLE 4. Carboamination of *N*-Protected Hex-4-enylamines^a

Entry	Amine	Aryl bromide	Product	Rxn. Time (h)	Yield (%) ^b
1	25	2-bromonaphthalene	26	23	59
2	25	4-bromonitrobenzene	30 + 31 (11:1) ^c	36	50
3	25	4-bromobenzoic acid methyl ester	32 + 33 (5:1) ^d	44	44 ^e
4	27	4-bromobenzoic acid methyl ester	34	42	43 ^{e,f}
5	27	2-bromonaphthalene	35	30	43 ^{e,f,g}
6	28	2-bromonaphthalene	36	27	55 ^e
7	28	4-bromobenzoic acid methyl ester	37	21	55 ^e
8	28	4-bromoacetophenone	38	31	60 ^{e,h}
9	28	2-bromopyridine	39	53	62 ^e
10	29	4-bromobenzoic acid methyl ester	40	22	49 ^e
11	29	4-bromoacetophenone	41	53	44 ^{e,g}

^a Conditions: 1.0 equiv amine, 1.2 equiv ArBr, 2.3 equiv Cs₂CO₃, 5 mol % Pd(OAc)₂, 7.5 mol % Nixantphos, dioxane (0.25 M), 100 °C.


^b Yield refers to average isolated yield obtained in two or more experiments. All reactions proceeded with >20:1 diastereoselectivity.

^c Ar = 3-nitrophenyl. ^d Ar = 4-carbomethoxyphenyl. ^e (±)-BINAP used in place of Nixantphos. ^f A trace amount (ca. 1–5%) of a regioisomer analogous to **31** and **33** was also obtained. ^g This reaction was conducted with 2.0 equiv ArBr, 10 mol % Pd(OAc)₂, and 15 mol % (±)-BINAP. ^h This reaction proceeded to 94% conversion.


Palladium-catalyzed carboamination reactions of (*Z*)-alkene substrates **28** and **29** also afforded products resulting from suprafacial addition to the alkene with >20:1 dr and moderate chemical yield (Table 4, entries 6–11). However, several significant differences in reactivity between (*E*)- and (*Z*)-alkene substrates were observed. The reactions of **28** and **29** were usually faster than transformations of (*E*)-alkene substrates **25** and **27**. In addition, the regioselectivity in reactions of **28** and **29** with *m*- and *p*-substituted aryl bromides was uniformly high (>20:1). For example, the coupling of **25** with methyl 4-bromobenzoate required 44 h and afforded a 5:1 mixture of regioisomers **32** and **33** (entry 3), whereas the analogous reaction

of **28** was complete in 21 h, and afforded a single pyrrolidine product (entry 7, **37**).

A final difference in the reactivity of (*E*)- versus (*Z*)-alkenes was observed when we attempted to couple **25** and **28** with 2-bromochlorobenzene (eqs 3 and 4). Surprisingly, the reaction of this aryl halide with (*Z*)-alkene **28** failed to provide the expected 2-(*sec*-phenethyl)pyrrolidine product, but instead afforded regioisomer **42** in 70% yield (eq 3).¹⁸ In contrast, the analogous reaction of *N*-protected (*E*)-alkene substrate **25** with 2-bromochlorobenzene gave a complex mixture of products. This mixture contained less than 10% of **42**, and little or none of the expected 2-(*sec*-phenethyl)pyrrolidine derivative (eq 4).



Attempts to couple cyclopentene-derived substrate **43** with 4-bromobiphenyl under the optimized mild reaction conditions also led to surprising results.¹⁹ As shown in eq 5, this transformation afforded less than 5% of the expected product **45**. Instead, benzocyclobutene **44** was generated in 72% yield and >20:1 dr.²⁰ In contrast, when the strong base NaOtBu was employed for this transformation the expected *N*-Boc-6-aryl octahydrocyclopenta[*b*]pyrrole **45** was formed in 51% yield with >20:1 dr (eq 6).



Reactions of 4-Substituted Pent-4-enylamine Derivatives.

N-Protected γ -aminoalkenes bearing substituents at the 4-position are another class of compounds that are not suitable substrates for Pd-catalyzed carboamination reactions conducted under strongly basic conditions. For example, the conversion

(17) We have been unable to separate these regioisomers from the major product. The structures of **31** and **33** have been assigned on the basis of analysis of EI-MS fragmentation patterns and the observation of a triplet (assigned as the CH₃ signal from the 2-ethyl group) at ~0.9 ppm in the ¹H NMR spectrum of the mixture. Similar products have been observed in analogous reactions that generate *N*-aryl pyrrolidines. See ref 5a.

(18) This product contained ca. 8% of an unidentified impurity.

(19) These results have been previously communicated. See: Bertrand, M. B.; Wolfe, J. P. *Org. Lett.* **2007**, *9*, 3073–3075.

(20) This transformation is effective with a number of different aryl bromides. See ref 19.

TABLE 5. Palladium-Catalyzed Carboamination of 4-Substituted Pent-4-enylamine Derivatives^a

Entry	Amine	Aryl bromide	Product	Rxn. Time (h)	Yield (%) ^b
1	46	2-bromonaphthalene	47	19	73
2	46	2-bromopyridine	52	18	75 ^c
3	46	2-bromo-4-methoxyphenyl	53	36	66
4	48	4- <i>t</i> -butylbromobenzene	54	19	71
5	48	4-methoxycarbonylbromobenzene	55	20	61
6	49	4-bromobiphenyl	56	25	83
7	49	4-bromobenzoyl methyl ester	57	42	69 ^d
8	50	2,4-dibromochlorobenzene	58	20	75
9	51	4-bromobenzotrifluoride	59	7	72
10	51	4-bromo-2-methoxynaphthalene	60	7	78


^a Conditions: 1.0 equiv amine, 1.2 equiv ArBr, 2.3 equiv Cs₂CO₃, 2 mol % Pd(OAc)₂, 4 mol % Nixantphos, dioxane (0.25 M), 100 °C.

^b Yield refers to average isolated yield obtained in two or more experiments. Products **56–60** with two stereocenters were obtained with >20:1 dr. ^c Reaction was conducted with 4 mol % Pd(OAc)₂ and 8 mol % Nixantphos. ^d Reaction was conducted with 5 mol % Pd(OAc)₂ and 10 mol % Nixantphos and proceeded to 80% conversion.

of **46** to **47** proceeded in only 31% yield when NaOtBu was employed as base (eq 7). Fortunately, use of the mild reaction conditions described above (Cs₂CO₃, dioxane) with catalysts supported by Nixantphos^{10,21} provided excellent results with these substrates. These new conditions facilitated the preparation of **47** in 73% yield (Table 5, entry 1).

As shown in Table 5, various aryl bromides were effectively coupled with **46** and the Cbz-protected analog **48**, including electron-poor, electron-rich and heteroaryl derivatives (entries



(21) Use of Dpe-phos as the ligand for this reaction afforded **47** in 43% isolated yield.



1–5). The 4-substituted pent-4-enylamine derivatives are more reactive than the *N*-protected hex-4-enylamines, and in most cases carboaminations of **46** and **48** can be effectively achieved using only 2 mol % Pd, compared to the higher catalyst loadings (5–10 mol % Pd) that are required for (*E*)- or (*Z*)-alkene substrates **25** and **27–29**.

These conditions are also effective for the coupling of methylene cyclopentane and cyclohexane derivatives **49–51** with several different aryl bromides. Reactions of these substrates provided fused bicyclic products **56–60** in good yield with >20:1 diastereoselectivity (Table 5, entries 6–10).

Attempted Reactions of Trisubstituted Alkene Derivatives. To further test the degree of alkene substitution tolerated in Pd-catalyzed carboamination reactions, we briefly examined reactions of trisubstituted alkenes **61** and **62** (eqs 8 and 9). Unfortunately, neither of these substrates was converted to the desired pyrrolidine product. Reactions of **61** provided mainly products resulting from Heck arylation of the alkene, whereas no reaction was observed between substrate **62** and 4-bromotert-butylbenzene at temperatures up to 175 °C with several different catalyst systems.




Deuterium Labeling Experiments. In our previous investigations on Pd-catalyzed carboamination reactions that employ NaOtBu as base, we determined that transformations of cyclopent-2-enylethylamine-derived substrates (e.g., **43**) afford products resulting from suprafacial addition to the alkene (eq 6).^{5a,c,d,22} This proved to be a key piece of data that suggested the carboaminations proceed via a mechanism involving a rare *syn*-aminopalladation process (see below). However, prior to the studies described in this Article, we had not elucidated the stereochemistry of alkene addition in reactions using the weak base Cs₂CO₃,²³ nor had alkene addition stereochemistry been examined with terminal alkene substrates. Moreover, although reactions of terminal alkenes provide pyrrolidine products regardless of whether NaOtBu or Cs₂CO₃ is used as base,

(22) For related studies on the mechanism of Pd-catalyzed carboetherification reactions of γ -hydroxyalkenes that generate tetrahydrofuran products, see: Hay, M. B.; Wolfe, J. P. *J. Am. Chem. Soc.* **2005**, *127*, 16468–16476.


(23) The isolation of 5- and 6-aryl *N*-Boc-octahydrocyclopenta[*b*]pyrrole side products in the conversion of **43** to benzocyclobutene derivatives when certain aryl halides were employed was consistent with this mechanism. However, the formation of these side products was only observed in two cases. See ref 19.

SCHEME 1



transformations of cyclopent-2-enylethylamine derivative **43** provide much different products depending on the nature of the base (eqs 5 and 6). Thus, two significant questions about the mechanism of Pd-catalyzed carboamination reactions remained unresolved: (a) Do reactions of internal alkenes, terminal alkenes, and cycloalkenes all proceed via a similar mechanism? (b) Do reactions that employ Cs₂CO₃ as base occur through a pathway that is mechanistically analogous to reactions in which NaOtBu is used?

The results shown above in Table 4 and eq 2 indicate that transformations of acyclic 1,2-disubstituted alkene substrates occur with suprafacial addition using either NaOtBu or Cs₂CO₃ as base, and are stereospecific. To address questions about the mechanism of carboamination reactions of terminal alkenes, we prepared a simple terminal alkene substrate **63** bearing a deuterium label at C5. As shown below (eq 10), treatment of **63** with methyl 2-bromobenzoate²⁴ in the presence of Cs₂CO₃ and the Pd(OAc)₂/Dpe-phos catalyst afforded **64** in 72% yield as a single stereoisomer that results from suprafacial addition to the alkene. No side products resulting from loss or migration of deuterium were observed.





To determine if the nature of the base had an effect on the stereochemistry of addition, the reaction of **63** with methyl 2-bromobenzoate was also conducted using NaOtBu as base. In addition to the expected product **64**, this transformation provided three additional products **65a–c** that result from scrambling of the alkoxide groups on the carbamate and ester appendages (Scheme 1). Fortunately, treatment of this mixture with LiAlH₄ afforded a single product stereoisomer **66** that derives from suprafacial alkene addition in the carboamination reaction.

Conversion of Pyrrolidine Products to Tetrahydropyrrolo-isoquinolin-5-ones. In addition to significantly expanding the scope of pyrrolidine-forming reactions, the mild reaction conditions described above can be used in a high-yielding three-

(24) This aryl bromide was chosen to facilitate elucidation of product stereochemistry. This was achieved by conversion of **64** to a tetrahydroisoquinoline derivative through treatment with TFA followed by Na₂CO₃. See Supporting Information for complete details.


step reaction sequence for the conversion of *N*-Boc-protected γ -aminoalkenes to substituted tetrahydropyrroloisoquinolin-5-ones. As shown below (eqs 11 and 12), *N*-Boc pyrrolidine derivatives **8**, **20**, and **57**, which were prepared from **1**, **14**, and **49** (Table 2, entry 5; Table 3, entry 3; Table 5, entry 7), were transformed to **67–69** in good to excellent yields via treatment with TFA followed by Na_2CO_3 . These heterocyclic scaffolds are displayed in several biologically active natural products and pharmaceutical leads.²⁵



Discussion

Synthetic Scope and Limitations. Palladium-catalyzed carboamination reactions of aryl bromides with γ -aminoalkenes bearing *N*-acyl, *N*-Boc, or *N*-Cbz protecting groups are mild and efficient means for the stereoselective synthesis of functionalized pyrrolidines. Although the scope of these transformations is somewhat limited when the strong base NaOtBu is employed as a key reagent,^{5c} the new reaction conditions described in this Article,⁹ which replace NaOtBu with Cs_2CO_3 , are effective with a much broader range of substrates. A number of functional groups are tolerated under these conditions, including methyl esters, alkyl acetates, enolizable ketones, and nitro groups. In addition, the reactions can also be conducted with aryl bromides that are electron-neutral, electron-poor, or heteroaromatic. Electron-rich aryl bromides can be effectively coupled with pent-4-enylamine derived substrates, although reactions of these electrophiles with hex-4-enylamine derivatives are slow and generally fail to proceed beyond 50% conversion even with relatively high catalyst loadings (10 mol % Pd). Aryl bromides bearing substituents in the *meta*- or *para*-positions provided good results with all *N*-protected γ -aminoalkenes examined in these studies, and *ortho*-substituted derivatives can be coupled with several different *N*-protected pent-4-enylamines. In addition, the use of methyl 2-bromobenzoate as the aryl halide electrophile leads to the formation of 2-benzylpyrrolidine derivatives that can be converted to substituted tetrahydropyrroloisoquinolin-5-ones in good yield through a two step sequence of Boc-deprotection followed by intramolecular amide bond formation (eqs 11 and 12). However, carboamination reactions of *ortho*-substituted aryl bromides with hex-4-enylamine derivatives were generally not effective, except for the conversion of (*Z*)-alkene **28** and 2-bromochlorobenzene to 2-(1-

SCHEME 2. Catalytic Cycle



phenylethyl)pyrrolidine **42** (eq 3). Efforts to couple alkenyl bromide electrophiles were also unsuccessful, although alkenyl bromides can be used in carboamination reactions of terminal alkene substrates when NaOtBu is employed as base.^{5c}


The mild reaction conditions described in this Article also broaden the scope of Pd-catalyzed carboaminations with respect to the degree of alkene substitution that is tolerated. Transformations that use NaOtBu as base are generally limited to terminal alkenes, but moderate yields can be obtained with the cycloalkene substrate *N*-Boc-cyclopent-2-enylethylamine (**43**). In contrast, the use of Cs_2CO_3 allows for coupling reactions of substrates bearing either pendant terminal alkenes or disubstituted alkenes. However, substrate **43** is converted to benzocyclobutene derivatives under these conditions, rather than to substituted pyrrolidines.¹⁹ In addition, reactions of very sterically encumbered trisubstituted alkenes failed to generate pyrrolidine products. It appears likely that the development or discovery of new catalyst systems will be required in order to effect carboamination of these compounds. When Cs_2CO_3 is employed as base, the relative reactivity of different alkenes decreases in the order of: terminal alkenes > 1,1-disubstituted alkenes > (*Z*)-1,2-disubstituted alkenes > (*E*)-1,2-disubstituted alkenes \gg trisubstituted alkenes. This order of reactivity mirrors that of many metal-catalyzed transformations that involve alkene insertion processes.²⁶

Reaction Mechanism and Origin of Regioisomers. The pyrrolidine-forming carboamination reactions between *N*-protected γ -aminoalkenes and aryl bromides appear to be mechanistically similar to other Pd-catalyzed carboetherification and carboamination reactions that afford oxygen^{22,27} or nitrogen⁵ heterocycles. A simplified catalytic cycle for the pyrrolidine-forming reactions is shown below (Scheme 2). The transformations are presumably initiated by oxidative addition of the aryl bromide to a Pd(0) catalyst that is generated in situ from the combination of $\text{Pd}(\text{OAc})_2$ and phosphine ligand. The resulting $\text{L}_n\text{Pd}(\text{Ar})(\text{Br})$ intermediate **70** is likely converted to palladium(aryl)(amido) complex **71** through reaction with the amine substrate and base.¹² Intramolecular *syn*-aminopalladation of **71** can provide **72**,^{5,28,29} which is transformed to the observed product via C–C bond-forming reductive elimination.³⁰

(25) (a) Yang, C. W.; Chuang, T. H.; Wu, P. L.; Huang, W. H.; Lee, S. J. *Biochem. Biophys. Res. Commun.* **2007**, *354*, 942–948. (b) Chuang, T. H.; Lee, S. J.; Yang, C. W.; Wu, P. L. *Org. Biomol. Chem.* **2006**, *5*, 860–867. (c) Aboul-Ela, M. A.; El-Lakany, A. M.; Hammoda, H. M. *Pharmazie* **2004**, *59*, 894–896. (d) Zuccotto, F.; Zvelebil, M.; Brun, R.; Chowdhury, S. F.; Di Lucrezia, R.; Leal, I.; Maes, L.; Ruiz-Perez, L. M.; Gonzalez Pacanowska, D.; Gilbert, I. H. *Eur. J. Med. Chem.* **2001**, *36*, 395–405.

(26) (a) Heck, R. F. *Org. React.* **1982**, *27*, 345–390. (b) Brase, S.; deMeijere, A. In *Metal-Catalyzed Cross-Coupling Reactions*; Diederich, F., Stang, P. J., Eds.; Wiley VCH: Weinheim, Germany, 1998; Chapter 3. (c) Link, J. T. *Org. React.* **2002**, *60*, 157–534.


(27) (a) Wolfe, J. P.; Rossi, M. A. *J. Am. Chem. Soc.* **2004**, *126*, 1620–1621. (b) Hay, M. B.; Hardin, A. R.; Wolfe, J. P. *J. Org. Chem.* **2005**, *70*, 3099–3107.

SCHEME 3. Formation of **31** and **33**

Two key pieces of evidence support this mechanistic hypothesis, along with the notion that the mechanism for pyrrolidine formation is not dependent on the nature of the base.³¹ First of all, the transformations proceed with suprafacial addition across both terminal and internal alkenes, as illustrated by the examples shown in Table 4, eqs 2, 6, and 10, and Scheme 1. The suprafacial addition is observed when either NaOtBu or Cs₂CO₃ is employed as base, and is consistent with the conversion of **71** to **72** via *syn*-aminopalladation.

The formation of 2-ethylpyrrolidine derivatives **31** and **33** (Table 4, entries 2 and 3) and 2-(1-phenylethyl)pyrrolidine **42** (eq 3) is also consistent with the mechanism shown above, and cannot satisfactorily be explained by alternative mechanisms that involve carbopalladation rather than aminopalladation.³² As shown above (Scheme 3), products **31** and **33** are most likely generated through *syn*- β -hydride elimination from intermediate **72** (via transition state **73**) to give **74**. Reinsertion of the alkene into the Pd-H bond of **74** with the opposite regioselectivity would yield **75**, which can undergo a second series of β -hydride elimination/reinsertion processes to provide **77**. Reductive elimination from **77** could then afford isomers arylated at the 3-position, whereas additional β -hydride elimination/reinsertion steps could ultimately generate isomers arylated at the 4- or 5-position.

The formation of 3-arylpiperidine side products is commonly observed in Pd-catalyzed carboamination reactions of γ -(*N*-arylamino) alkenes bearing relatively electron-rich *N*-aryl groups (e.g., Ph or PMP).^{5a} In contrast, the formation of mixtures of regioisomers is usually not observed in reactions of *N*-Boc-, *N*-acyl-, or *N*-Cbz-protected substrates, except for couplings of (*E*)-disubstituted alkenes **25** and **27** with electron-poor aryl

SCHEME 4. Formation of **42**

bromides (Table 4, entries 2–3). The absence of these side-products in most reactions of *N*-Boc and *N*-acyl-protected substrates may be due to decreased rates of β -hydride elimination from intermediate **72** relative to the analogous *N*-aryl derivative. This may be due to stabilization of **72** through chelation of the metal to the carbonyl of the amide or carbamate,³³ or through inductive effects induced by the amide/carbamate protecting groups. These electron-withdrawing groups likely destabilize the developing positive charge on C2 in the transition state for β -hydride elimination (**73**).³⁴ The fact that a similar regioisomer was not generated when the electron-neutral 2-bromonaphthalene was coupled with **25** may be due to the decreased electrophilicity of the arylpalladium complex (**72**, Ar = 2-naphthyl), which could slow the rate of β -hydride elimination. However, the reasons for formation of only trace amounts (ca. 1–5%) of regioisomers in transformations of Cbz-protected substrate **27** are unclear.

The pathway by which **42** is generated is likely similar to the mechanism for the formation of **31** and **33**, but with initial β -elimination of a hydride from the methyl group of **72a** (Scheme 4) rather than from C2 (Scheme 3). The formation of regioisomers analogous to **42** was not observed with *p*- or *m*-substituted aryl bromides. This suggests the steric bulk of 2-bromochlorobenzene may increase the rate of β -hydride elimination from the less hindered methyl group relative to the more hindered C2 position, or slow the rate of reductive elimination from **72a** or **77** relative to **79**.³⁵

Interestingly, the formation of regioisomers similar to **31** and **33** was not observed in reactions of *N*-Boc-protected (*Z*)-alkene substrate **28**. This may be due to severe developing allylic strain interactions in the transition state for β -hydride elimination, from steric destabilization of the resulting (Ar)Pd(H)(alkene) complex, or both. As shown in Scheme 5, alkene insertion from **71** would yield **72**, which could undergo β -hydride elimination from transition state **73** to provide **74**. This intermediate could then be converted to the observed regioisomers as described above. A similar sequence of transformations from (*Z*)-alkene-derived intermediate **71a** could lead to the analogous intermediate **74a**. However, the eclipsing interaction between the methyl group and the carbamate protecting group in transition state **73a** would result in developing allylic strain similar to that in an *S*-*cis*-

(28) For examples of *syn*-alkene insertion into Pt-N bonds, see: (a) Cowan, R. L.; Troglor, W. C. *J. Am. Chem. Soc.* **1989**, *111*, 4750–4761. (b) Cowan, R. L.; Troglor, W. C. *Organometallics* **1987**, *6*, 2451–2453. For examples of *syn*-alkene insertion into Ni-N bonds, see: (c) VanderLende, D. D.; Abboud, K. A.; Boncella, J. M. *Inorg. Chem.* **1995**, *34*, 5319–5326. For examples of *syn*-alkene insertion into Ir-N bonds, see: (d) Casalnuovo, A. L.; Calabrese, J. C.; Milstein, D. *J. Am. Chem. Soc.* **1988**, *110*, 6738–6744. For examples of *syn*-alkene insertion into Rh-N bonds, see: (e) Zhao, P.; Krug, C.; Hartwig, J. F. *J. Am. Chem. Soc.* **2005**, *127*, 12066–12073. For examples of *syn*-insertion of dimethyl acetylenedicarboxylate into Pd-N bonds, see: (f) Villanueva, L. A.; Abboud, K. A.; Boncella, J. M. *Organometallics* **1992**, *11*, 2963–2965. For a review on alkene aminopalladation (*syn*- and *anti*-), see: (g) Minatti, A.; Muniz, K. *Chem. Soc. Rev.* **2007**, *36*, 1142–1152.

(29) For other Pd-catalyzed reactions that likely proceed via alkene *syn*-aminopalladation, see: (a) Helaja, J.; Gottlich, R. *J. Chem. Soc., Chem. Commun.* **2002**, 720–721. (b) Tsutsui, H.; Narasaka, K. *Chem. Lett.* **1999**, 45–46. (c) Brice, J. L.; Harang, J. E.; Timokhin, V. I.; Anastasi, N. R.; Stahl, S. S. *J. Am. Chem. Soc.* **2005**, *127*, 2868–2869.

(30) Milstein, D.; Stille, J. K. *J. Am. Chem. Soc.* **1979**, *101*, 4981–4991.

(31) The formation of benzocyclobutene **44** from the Pd-catalyzed reaction of **43** with 4-bromobiphenyl when Cs₂CO₃ is used as a base appears to arise from a divergence in mechanism prior to formation of intermediate palladium amido complex **71**. For further discussion, see ref 19.


(32) For a detailed discussion of all mechanistic possibilities in Pd-catalyzed reactions of unsaturated amines or alcohols that generate heterocyclic products, see ref 4b.

(33) (a) Zhang, L.; Zetterberg, K. *Organometallics* **1991**, *10*, 3806–3813. (b) Lee, C.-W.; Oh, K. S.; Kim, K. S.; Ahn, K. H. *Org. Lett.* **2000**, *2*, 1213–1216. (c) Oestreich, M.; Dennison, P. R.; Kodanko, J. J.; Overman, L. E. *Angew. Chem., Int. Ed.* **2001**, *40*, 1439–1442. (d) Clique, B.; Fabritius, C.-H.; Couturier, C.; Monteiro, N.; Balme, G. *Chem. Commun.* **2003**, 272–273.

(34) Mueller, J. A.; Sigman, M. S. *J. Am. Chem. Soc.* **2003**, *125*, 7005–7013.

(35) It is also possible that the *o*-chloro substituent facilitates β -hydride elimination by effecting dissociation of one arm of the chelating phosphine ligand. For a discussion of the effect of *o*-halo groups on ligand substitution processes, see: Kuniyasu, H.; Yamashita, F.; Terao, J.; Kambe, N. *Angew. Chem., Int. Ed.* **2007**, *46*, 5929–5933.

SCHEME 5




locked-(*Z,Z*)-1,4-diene. This presumably raises the energy of the transition state to a sufficient degree such that competing β -elimination of the C2-hydrogen atom is minimized. Further evidence for the high energy of transition state 73a comes from the formation of regioisomer 42 in the reaction of (*Z*)-alkene 28 with 2-bromochlorobenzene (eq 3). This product results from β -hydride elimination of 72a from the less electronically favorable methyl-position.³⁶

Ligand Effects. Three ligands that were surveyed in Pd-catalyzed carboamination reactions that use Cs₂CO₃ as a base were found to provide good results with the majority of substrate combinations described in this paper. In most cases, Dpe-phos provided satisfactory yields in transformations of 1- or 3-substituted pent-4-enylamine derivatives. However, more sterically encumbered 4-substituted substrates were converted to pyrrolidine products most efficiently when Nixantphos was employed as ligand. Finally, highest yields in carboaminations of hex-4-enylamine derivatives were usually obtained with (\pm)-BINAP.


The precise origin of the observed ligand effects is not clear, but is likely influenced by the effect of ligand bite angle on the relative rates of carboamination vs competing Heck arylation or N-arylation of the substrate. In general, use of Nixantphos led to greater amounts of N-arylated side products than Dpe-phos or BINAP, whereas the latter two ligands led to greater amounts of products resulting from Heck arylation. The relatively wide bite angle³⁷ of Nixantphos (>108°)³⁸ compared to Dpe-phos (104°) and BINAP (93°) may accelerate the aminopalladation step (71 to 72) as well as competing C–N bond-forming reductive elimination from 71,³⁹ the change in relative rates appears to depend on substrate structure. However, the electronic properties of the ligand are also likely of importance, as the related wide-bite angle ligand Xantphos (108°)³⁷ was inferior to both Nixantphos and Dpe-phos, and provided large amounts of N-arylated side products.

Stereochemistry. The stereochemistry of pyrrolidine products generated in Pd-catalyzed carboamination reactions of *N*-protected γ -aminoalkenes appears to be determined in the *syn*-

SCHEME 6. Stereochemistry of 18–20



SCHEME 7. Stereochemistry of 21 and 22



aminopalladation step of the catalytic cycle illustrated above (Scheme 2, 71 to 72). The *syn*-aminopalladation ultimately leads to suprafacial addition of the nitrogen atom and the aromatic group across the substrate C–C double bond. Moreover, it appears likely that the alkene insertion proceeds through a highly organized cyclic transition state, in which the alkene π -system is eclipsed with the Pd–N bond,⁴⁰ and substituents on the tether between the amine and the alkene are oriented to minimize nonbonding interactions. For example, transformations of substrates 13–14 bearing allylic substituents provide *trans*-2,3-disubstituted pyrrolidine products 18–20 (Table 3, entries 1–3). As shown above (Scheme 6), *syn*-aminopalladation via 80, in which the C3-methyl group is oriented in a pseudoequatorial position, would provide the observed major diastereomers.

The conversion of substrates 15 and 16 to *cis*-2,5-disubstituted pyrrolidines 21 and 22 can be rationalized through a similar transition state model. As shown in Scheme 7, *syn*-aminopalladation via transition state 82 would generate 83, which can be converted to the observed products through reductive elimination. Cyclization through 82, in which the C1-phenyl group is oriented in a pseudoaxial position appears to be favored over transition state 84, which suffers from A^(1,3)-strain between the C1-phenyl group and the nitrogen protecting group.^{41,42}

Summary and Conclusion

In conclusion, the mild reaction conditions described in this Article greatly expand the scope of Pd-catalyzed alkene carboamination chemistry. The combined use of the weak base

(36) The transition state for β -hydride elimination from the methyl group would contain a developing partial positive charge on a primary carbon atom.

(37) (a) Ozawa, F.; Kubo, A.; Matsumoto, Y.; Hayashi, T.; Nishioka, E.; Yanagi, K.; Moriguchi, K. *Organometallics* **1993**, *12*, 4188–4196. (b) van Haaren, R. J.; Goubitz, K.; Fraanje, J.; van Strijdonck, G. P. F.; Overing, H.; Coussens, B.; Reek, J. N. H.; Kamer, P. C. J.; van Leeuwen, P. W. N. M. *Inorg. Chem.* **2001**, *40*, 3363–3372.

(38) The bite angle of flexible ligands is dependent on coordination environment. No X-ray structures of (Nixantphos)PdX₂ complexes have been reported, and thus we cannot provide a precise value for the Nixantphos bite angle for comparison to the bite angles of Xantphos, Dpe-Phos, and BINAP, which were obtained from X-ray structures of their PdX₂ complexes. However, the metal-independent calculated natural bite angle of Nixantphos is 114°, whereas that of Xantphos is 111°. Thus, we feel it is reasonable to estimate that the bite angle of (Nixantphos)PdX₂ should exceed that of the analogous Xantphos complex.

(39) Xantphos has previously been demonstrated to provide good results in Pd-catalyzed N-arylation reactions of amides and carbamates. See: Yin, J.; Buchwald, S. L. *J. Am. Chem. Soc.* **2002**, *124*, 6043–6048.

(40) Related alkene carbopalladation processes that occur in intramolecular Heck reactions are believed to proceed through similar transition states. Other possible transition states in which the alkene is perpendicular to the M–C bond appear to be significantly higher in energy. See: Overman, L. E. *Pure. Appl. Chem.* **1994**, *66*, 1423–1431, and references therein.

(41) For further discussion of A^(1,3)-strain in transformations involving *N*-substituted amines, see: (a) Hoffmann, R. W. *Chem. Rev.* **1989**, *89*, 1841–1860. (b) Hart, D. J. *J. Am. Chem. Soc.* **1980**, *102*, 397–398. (c) Williams, R. M.; Sinclair, P. J.; Zhai, D.; Chen, D. *J. Am. Chem. Soc.* **1988**, *110*, 1547–1557. (d) Kano, S.; Yokomatsu, T.; Iwasawa, H.; Shibuya, S. *Heterocycles* **1987**, *26*, 2805–2809.

(42) The conversion of 17 to 23 and 24 also occurs through a transition state in which the C-1 substituent (C₉H₁₉) is pseudoaxial. For further discussion of this transformation in the context of a total synthesis of (+)-preussin, see ref 5g.

Cs₂CO₃, dioxane solvent, Pd(OAc)₂ precatalyst, and the appropriate phosphine ligand allows for the stereoselective synthesis of substituted pyrrolidines from *N*-protected γ -aminoalkenes with good tolerance of functional groups. The pyrrolidine products are formed in moderate to excellent yields, with diastereoselectivities ranging from 12 to >20:1. The mild conditions also greatly increase tolerance of alkene substitution, and for the first time substrates bearing acyclic disubstituted alkenes can be successfully employed in these reactions.

The observed selectivity for suprafacial addition with both internal and terminal alkene substrates provides additional evidence in support of our proposed reaction mechanism, which involves intramolecular *syn*-aminopalladation as a key step. In addition, the results described above suggest this mechanism of pyrrolidine-formation is operational regardless of the degree of alkene substitution, and with either Cs₂CO₃ or NaOtBu as the stoichiometric base.

Future studies will be directed toward the development/discovery of highly active catalysts that can transform tri- and tetrasubstituted alkene derivatives, along with additional applications of these transformation to the synthesis of biologically active molecules.

Experimental Section

General Procedure for Pd-Catalyzed Carboamination Reactions of Aryl Bromides. A flame-dried Schlenk tube equipped with a magnetic stirbar was cooled under a stream of nitrogen and charged with the aryl bromide (1.2 equiv), Pd(OAc)₂ (2 mol %), Dpe-phos, dppe, (\pm)-BINAP, or Nixantphos (4 mol %) and Cs₂CO₃ (2.3 equiv). The tube was purged with nitrogen and a solution of the *N*-protected amine substrate (1.0 equiv) in dioxane (5 mL/mmol substrate) was then added. The resulting mixture was heated to 100 °C with stirring until the starting material had been consumed as

determined by GC analysis. The reaction mixture was cooled to room temperature and saturated aqueous NH₄Cl (1 mL) and ethyl acetate (1 mL) were added. The layers were separated, the aqueous layer was extracted with ethyl acetate (3 \times 5 mL), and the combined organic layers were dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo. The crude product was then purified by flash chromatography on silica gel.

2-(4-*tert*-Butylbenzyl)pyrrolidine-1-carboxylic Acid *tert*-Butyl Ester (2). The general procedure was employed for the reaction of 4-*tert*-butyl bromobenzene (52 μ L, 0.30 mmol) with **1** (47 mg, 0.25 mmol) using Dpe-phos as ligand. This procedure afforded 66 mg (83%) of the title compound as a pale yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 7.38–7.26 (m, 2 H), 7.22–7.07 (m, 2 H), 4.12–3.84 (m, 1 H), 3.49–3.23 (m, 2 H), 3.23–2.96 (m, 1 H), 2.60–2.42 (m, 1 H), 1.92–1.67 (m, 4 H), 1.52 (s, 9 H), 1.32 (s, 9 H); ¹³C NMR (100 MHz, CDCl₃) δ 154.5, 148.9, 136.1, 129.1, 125.2, 79.0, 58.8, 46.4, 40.0, 34.3, 31.4, 29.7, 28.6, 22.7; IR (film) 1695 cm⁻¹. Anal. calcd for C₂₀H₃₁NO₂: C, 75.67; H, 9.84; N, 4.41. Found: C, 75.46; H, 9.88; N, 4.38.

Acknowledgment. The authors thank the NIH-NIGMS (GM 071650) for financial support of this work. M.B.B. was supported by a Bristol-Myers Squibb Award in Synthetic Organic Chemistry. Additional funding for these studies was provided by the Camille and Henry Dreyfus Foundation (New Faculty Award, Camille Dreyfus Teacher Scholar Award), Research Corporation (Innovation Award), Eli Lilly, Amgen, GlaxoSmithKline, and 3M.

Supporting Information Available: Experimental procedures, spectroscopic data, descriptions of stereochemical assignments, and copies of ¹H and ¹³C NMR spectra for all new compounds reported. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO801631V