


## Total Synthesis of 20-Norsalvinorin A. 1. Preparation of a Key Intermediate

Ylva E. Bergman,<sup>†</sup> Roger Mulder,<sup>‡</sup> and Patrick Perlmutter<sup>\*†</sup>

School of Chemistry, Monash University, PO Box 23,  
Victoria 3800, Australia, and CSIRO Molecular and Health  
Technologies, Bag 10 Clayton South, Victoria 3169, Australia

patrick.perlmutter@sci.monash.edu.au

Received December 14, 2008



The key tricyclic intermediate **3a**, for the total synthesis of the C<sub>20</sub>-nor analogue of salvinorin A, was prepared in seven steps from 3-furaldehyde. Key steps involved a highly regio- and diastereoselective Lewis acid assisted Diels–Alder reaction followed by base-promoted epimerization and a completely stereoselective conjugate reduction.

Salvinorin A (**1**) is a highly potent and selective  $\kappa$ -opioid receptor (KOR) agonist, both in vivo and in vitro, with no significant affinity for any other opioid receptor subtype.<sup>1</sup> It was originally isolated in 1982 from the perennial herb *Salvia divinorum* by Ortega<sup>2</sup> and independently by Valdés.<sup>3</sup> Two total syntheses of salvinorin A have been reported. One by Evans<sup>4</sup> in 29 steps via a transannular Michael reaction cascade and the other by Hagiwara<sup>5</sup> in 20 steps from enantiomerically pure Wieland–Miescher ketone. In addition, partial syntheses have been disclosed by Hügel<sup>6</sup> and Forsyth.<sup>7</sup>

Because of the remarkable affinity for the KOR, the functional groups of salvinorin A have been extensively modified, and subsequent SAR studies have begun to clarify the pharmaco-





FIGURE 1. Summary of the outcomes of SAR studies on salvinorin A (**1**).

### SCHEME 1. Retrosynthetic Analysis of 20-Norsalvinorin Analogue Including the Target Intermediate **3a**



phore. The rapidly expanding library of salvinorin A analogues includes modifications at the C<sub>1</sub>, C<sub>2</sub>, C<sub>4</sub>, C<sub>17</sub>, and C<sub>8</sub> position (Figure 1).<sup>8</sup> To date no studies have evaluated the importance, if any, of the C<sub>20</sub>-methyl group. From its location on the salvinorin skeleton modification of C<sub>20</sub> is clearly very difficult. Hence we decided to prepare the C<sub>20</sub>-nor analogue of salvinorin A through total synthesis. Herein we describe a short, expeditious, synthesis of key intermediate **3a** (Scheme 1).

From the retrosynthesis shown in Scheme 1, the key step is an *endo-syn* selective intermolecular Diels–Alder addition of a semicyclic dihydropyran-based diene (**4a**). It is envisaged that the C<sub>18</sub>-methoxycarbonyl group will be introduced via Pd-catalyzed carbonylation of the enoltriflate of **3a**.<sup>9</sup> A sequence of conjugate reduction of the unsaturated ester followed by allylic oxidation and Evans<sup>4</sup> conjugate reduction of the resulting unsaturated lactone would provide the 20-norsalvinorin A skeleton. Demethylation followed by acetylation completes the synthesis of **2**.

<sup>†</sup> Monash University.

<sup>‡</sup> CSIRO Molecular and Health Technologies.

(1) Roth, B. L.; Baner, K.; Westkaemper, R.; Siebert, D.; Rice, K. C.; Steinberg, S.; Ernsberger, P.; Rothman, R. B. *Proc. Nat. Acad. Sci. U.S.A.* **2002**, *99*, 11934–11939.

(2) Ortega, A.; Blount, J. F. *J. Chem. Soc., Perkin Trans. 1* **1982**, 2505–2508.

(3) Valdés, I.; Leander, J.; Butler, W. M.; Hatfield, G. M.; Paul, A. G.; Koreeda, M. *J. Org. Chem.* **1984**, *49*, 4716–4720.

(4) Scheerer, J. R.; Lawrence, J. F.; Wang, G. C.; Evans, D. A. *J. Am. Chem. Soc.* **2007**, *129*, 8968–8969.

(5) Nozawa, M.; Suka, Y.; Hoshi, T.; Suzuki, T.; Hagiwara, H. *Org. Lett.* **2008**, *10*, 1365–1368.


(6) Lingham, A. R.; Hügel, H. M.; Rook, T. J. *Aust. J. Chem.* **2006**, *59*, 340–348.

(7) Burns, A. C.; Forsyth, C. J. *Org. Lett.* **2008**, *10*, 97–100.


(8) Prisinzano, T. E.; Rothman, R. B. *Chem. Rev.* **2008**, *108*, 1732–1743.

(9) Hagiwara, H.; Hamano, K.; Nozawa, M.; Hoshi, T.; Suzuki, T.; Kido, F. *J. Org. Chem.* **2005**, *70*, 2250–2255.

**SCHEME 2. Stannic Chloride Mediated Diels–Alder Reactions of Diene **4b** with 2-Methoxy-5-methyl-1,4-benzoquinone (**6**)**



**SCHEME 3. Titanium-tetrachloride Mediated Diels–Alder Reactions of Diene **4a** with 2-Methoxy-5-methyl-1,4-benzoquinone (**6**)**




Model studies of the Diels–Alder reaction were carried out with a phenyl-substituted diene **4b** (prepared in three steps<sup>10</sup> from benzaldehyde) with 2-methoxy-5-methyl-1,4-benzoquinone<sup>11</sup> (**6**). By analogy with the work of Carreño, it was anticipated that a mixture of *endo-anti* and *endo-syn* adducts would be obtained.<sup>12</sup> However, unlike Carreño's dienophiles, which reacted completely after 16 h at room temperature, no reaction was observed between **4b** and **6** after 3 days at  $80\text{ }^\circ\text{C}$ . In order to increase the reactivity of **4b**, we turned to precomplexation of the methoxyquinone with metal salts.


Thus, employing the method of Reusch,<sup>13</sup> **4b** was added to a precomplexed mixture of **6** and  $\text{SnCl}_4$  at low temperature. Reaction was found to commence upon warming to approximately  $-45\text{ }^\circ\text{C}$ . After 4 h at this temperature, the reaction was quenched, and an inseparable mixture of two products was obtained. However, allowing the reaction mixture to warm to  $0\text{ }^\circ\text{C}$  produced the desired adduct **7b** in 51% yield (Scheme 2). Apparently at this higher temperature the second product either is converted to **7b** via a retro-Diels–Alder pathway or simply decomposes under the conditions.

Diene **4a**, prepared according to Snapper's protocol in three steps from 3-furaldehyde,<sup>10</sup> was then added to **6** under conditions similar to those just described for **4b** (Scheme 3). Only decomposition was observed when the reaction was warmed to  $0\text{ }^\circ\text{C}$ . At lower temperatures very poor conversion was obtained with a highest yield of only  $\sim 20\%$ . Switching to the use of  $\text{TiCl}_4$  gave decomposition at  $-45\text{ }^\circ\text{C}$ . Running the reaction at

**SCHEME 4. Epimerization of Adduct **7a** To Adjust the Stereochemistry at  $\text{C}_{10}$**



**SCHEME 5. Attempted Conjugate Reduction of Epimerized Adduct **8b****



$-78\text{ }^\circ\text{C}$  in dichloromethane gave a 1:1 mixture of two adducts. Changing the solvent to toluene gave no reaction at  $-78\text{ }^\circ\text{C}$ . However, at  $-45\text{ }^\circ\text{C}$  a 1:1 ratio of adducts was again obtained, providing the desired *endo-syn* product **7a** in 37% isolated yield.

Although yields were modest (37% and 51% for **7a** and **7b**, respectively), the major product in each case was formed with full regiocontrol with the three newly created stereogenic centers (at  $\text{C}_5$ ,  $\text{C}_9$ , and  $\text{C}_{10}$ ) in the same orientation as the natural product. Significantly, the relative stereochemistry between  $\text{C}_5$  and  $\text{C}_{12}$  in **7a** is also correct for the natural product. Interestingly a small amount ( $\sim 10\%$ ) of **8a** (vide infra) was also present in the product. Thus, some epimerization of the adduct had occurred after cycloaddition.

With **7a** and **7b** in hand, it remained for us to adjust the relative stereochemistry at  $\text{C}_{10}$  and introduce the correct stereochemistry at  $\text{C}_2$  (Scheme 4). Thus treatment of **7a** with DBU at room temperature in dichloromethane for 1 h gave smooth conversion to a single product, **8a** (adduct **7b** behaved similarly, cleanly epimerizing to **8b**).

A variety of conditions were explored in order to effect reduction of **8a** and **8b**. All attempts to achieve this transformation with copper-<sup>14</sup> and zinc-based<sup>15</sup> reagents led to mixtures of 1,2-reduction products. In the case of L-Selectride<sup>16</sup> smooth conversion to a single 1,2-reduction product, **9b**, was observed (Scheme 5).

Finally, alkaline dithionite was employed as the reducing system.<sup>17</sup> Unlike previous reductions, this led to exclusive conjugate reduction installing the correct relative stereochemistry at  $\text{C}_2$  without loss of OMe. Similar results were obtained for both **8a** and **8b** (Scheme 6). A crystal structure for **3b** was obtained that confirmed our assignments of relative stereochemistry (see Supporting Information).

(10) Cheung, A. K.; Murelli, R.; Snapper, M. L. *J. Org. Chem.* **2004**, *69*, 5712–5719.

(11) Takizawa, Y.; Iwasa, Y.; Suzuki, T.; Miyaji, H. *Chem. Lett.* **1993**, 1863–1864.

(12) Carreño, C. M.; Urbano, A.; Di Vitta, C. *J. Org. Chem.* **1998**, *63*, 8320–8330.


(13) Tou, J. S.; Reusch, W. *J. Org. Chem.* **1980**, *45*, 5012–5014.

(14) Rainka, M. P.; Aye, Y.; Buchwald, S. L. *Proc. Nat. Acad. Sci. U.S.A.* **2004**, *101*, 5821–5823.

(15) Marchand, A. P.; Reddy, M. *Synthesis* **1991**, 198–200.

(16) Comins, D. L.; Dehghani, A. *Tetrahedron Lett.* **1992**, *33*, 6299–6302.

(17) de Vries, J. G.; Kellogg, R. M. *J. Org. Chem.* **1980**, *45*, 4126–4129.

**SCHEME 6. Successful Conjugate Reduction of Epimerized Adducts 8a and 8b**


With a short route to intermediate **3a** now developed, studies directed toward the synthesis of **2** and evaluation of its KOR selectivity are underway in our laboratories and will be published in due course.

**Experimental Section**

**rac-(2S,6aR,10aS,10bR)-9-Methoxy-6a-methyl-2-phenyl-6,6a,10a,10b-tetrahydro-1H-benzof[isochromene-7,10(2H,4H)-dione (7b).** To a stirring solution of 2-methoxy-5-methyl-1,4-benzoquinone (**6**) (2.50 g, 16.43 mmol) in dry  $\text{CH}_2\text{Cl}_2$  (120 mL) under an atmosphere of argon was added a 1.0 M solution of  $\text{SnCl}_4$  in  $\text{CH}_2\text{Cl}_2$  (13.1 mL, 13.1 mmol), dropwise at  $-78^\circ\text{C}$ . Stirring was continued at this temperature for 1 h, after which a solution of diene **4b** (1.84 g, 9.86 mmol) in  $\text{CH}_2\text{Cl}_2$  (30 mL) was added slowly. The reaction was stirred at  $-78^\circ\text{C}$  for 15 min and then at  $0^\circ\text{C}$  for 5 h. Quenching of the reaction was done at  $-78^\circ\text{C}$  by the addition of brine (20 mL), the mixture was warmed to room temperature, and the layers were separated. The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  ( $2 \times 30$  mL), and the combined organic layers were washed with brine (70 mL), dried over  $\text{MgSO}_4$ , filtered, and concentrated in vacuo to yield a brown residue. This residue was subsequently stirred with aqueous sodium bisulfite ( $\text{NaHSO}_3$ ) solution for 1 h and then extracted with  $\text{CH}_2\text{Cl}_2$ . The combined organic layers were washed and dried in the same manner as above, and the resulting brown foam was purified by flash chromatography (2:3 EtOAc/hexanes) to afford **7b** (1.70 g, 51%) as a pale orange solid. Mp  $144\text{--}146^\circ\text{C}$ .  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.23–7.29 (5H, m), 5.80 (1H, s), 5.61 (1H, d,  $J = 1.4$  Hz), 4.39 (1H, dd,  $J = 11.0, 2.0$  Hz), 4.31 (1H, d,  $J = 12.4$  Hz), 4.11 (1H, dq,  $J = 12.5, 2.8, 1.4$  Hz), 3.71 (3H, s), 3.17 (1H, d,  $J = 7.2$  Hz), 2.89 (1H, m), 2.81 (1H, ddt,  $J = 17.9, 4.7, 1.6$  Hz), 1.95 (1H, q,  $J = 23.6, 12.6$  Hz), 1.81 (1H, dq,  $J = 17.9, 5.6, 2.3$  Hz), 1.58 (1H, ddd,  $J = 12.5, 4.2, 2.0$  Hz), 1.34 (3H, s).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  200.8, 195.7, 162.3, 141.9, 133.5, 128.5, 127.7, 126.1, 119.4, 109.6, 80.7, 72.7, 56.6, 55.7, 47.7, 38.5, 36.9, 31.8, 26.9. IR (neat):  $\nu$  3063w, 3030w, 2937m, 2844m, 1708m, 1662s, 1606s  $\text{cm}^{-1}$ . ESI-MS:  $m/z$  361.2  $[\text{M} + \text{Na}]^+$ . ESI-HRMS calcd for  $[\text{M} + \text{Na}]^+$  ( $\text{C}_{21}\text{H}_{22}\text{NaO}_4$ ):  $m/z$  361.1416, found 361.1405. Anal. Calcd for  $\text{C}_{21}\text{H}_{22}\text{O}_4$ : C 74.54, H 6.55. Found: C 74.15, H 6.52.

**rac-(2S,6aR,10aS,10bR)-2-(Furan-3-yl)-9-methoxy-6a-methyl-6,6a,10a,10b-tetrahydro-1H-benzof[isochromene-7,10(2H,4H)-dione (7a).** To a stirring solution of 2-methoxy-5-methyl-1,4-benzoquinone (**6**) (860 mg, 5.65 mmol) in dry toluene (45 mL) under an atmosphere of argon was added a 1.0 M solution of  $\text{TiCl}_4$  in  $\text{CH}_2\text{Cl}_2$  (4.52 mL, 4.52 mmol) dropwise at  $-78^\circ\text{C}$ . Stirring was continued at this temperature for 1 h, after which a solution of diene **4a** (598 mg, 3.39 mmol) in toluene (5 mL) was added slowly. The reaction was stirred at  $-78^\circ\text{C}$  for 15 min and then at  $-45^\circ\text{C}$  for 5 h. Quenching of the reaction was done at  $-78^\circ\text{C}$  by the addition of brine (10 mL), the mixture was warmed to room temperature, and the layers were separated. The aqueous layer was extracted with  $\text{Et}_2\text{O}$  ( $2 \times 15$  mL), and the combined organic layers were washed with brine (30 mL), dried over  $\text{MgSO}_4$ , filtered, and concentrated in vacuo. The crude residue was purified by flash chromatography (2:3 EtOAc/hexanes) to afford **7a** (411 mg, 37%) as a yellow gum.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.33 (1H, t,  $J =$

1.6 Hz), 7.32 (1H, m), 6.32 (1H, m), 5.83 (1H, s), 5.58 (1H, m), 4.39 (1H, dd,  $J = 11.0, 1.7$  Hz), 4.24 (1H, d,  $J = 12.5$  Hz), 4.08 (1H, dm,  $J = 12.5$ ), 3.76 (3H, s), 3.16 (1H, d,  $J = 6.7$  Hz), 2.74–2.86 (2H, m), 2.04 (1H, q,  $J = 23.7, 12.4$  Hz), 1.80 (1H, dq,  $J = 17.8, 5.6, 2.8$  Hz), 1.57 (1H, ddd,  $J = 12.4$  Hz, 4.3, 2.0 Hz), 1.33 (3H, s).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  200.7, 195.6, 162.4, 143.2, 139.1, 133.3, 126.6, 119.4, 109.5, 108.7, 73.4, 72.5, 56.6, 55.6, 47.8, 37.1, 36.3, 31.9, 26.5. IR (neat):  $\nu$  3065w, 2945m, 2840m, 1711m, 1661s, 1609s  $\text{cm}^{-1}$ . ESI-MS:  $m/z$  351.2  $[\text{M} + \text{Na}]^+$ . ESI-HRMS Calcd for  $[\text{M} + \text{Na}]^+$  ( $\text{C}_{19}\text{H}_{20}\text{NaO}_5$ ):  $m/z$  351.1208, found 351.1205.

**rac-(2S,6aR,10aR,10bR)-2-(Furan-3-yl)-9-methoxy-6a-methyl-6,6a,10a,10b-tetrahydro-1H-benzof[isochromene-7,10(2H,4H)-dione (8a).** To a stirring solution of **7a** (63 mg, 0.19 mmol) in dry  $\text{CH}_2\text{Cl}_2$  (3 mL) under an atmosphere of argon was added DBU (2.8  $\mu\text{L}$ , 0.019 mmol), and the resulting solution was stirred for 1 h at room temperature. The solvent was evaporated, and the brown residue was immediately purified on a short plug of silica gel (2:3 EtOAc/hexanes) to yield **8a** (55 mg, 87%) as a yellow gum.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.37 (1H, m), 7.35 (1H, t,  $J = 1.8$  Hz), 6.37 (1H, m), 5.81 (1H, s), 5.62 (1H, m), 4.63 (1H, dd,  $J = 11.3, 1.6$  Hz), 4.25 (1H, d,  $J = 12.5$  Hz), 4.16 (1H, dm,  $J = 12.4$  Hz), 3.79 (3H, s), 2.95 (1H, m), 2.78 (1H, d,  $J = 9.8$  Hz), 2.57 (1H, ddd,  $J = 12.7, 4.6, 2.0$  Hz), 2.51 (1H, dm,  $J = 20.0$  Hz), 2.28 (1H, ddt,  $J = 18.5, 5.7, 2.0$  Hz), 1.30 (1H, dt,  $J = 12.7, 11.4$  Hz), 1.14 (3H, s).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  202.4, 194.7, 163.1, 143.2, 139.1, 132.8, 126.7, 119.3, 109.1, 108.7, 72.4, 72.4, 56.5, 56.4, 48.2, 39.8, 33.1, 32.4, 21.1. IR (neat):  $\nu$  2962m, 2844m, 1711m, 1661s, 1605s  $\text{cm}^{-1}$ . ESI-MS:  $m/z$  351.2  $[\text{M} + \text{Na}]^+$  ( $\text{C}_{19}\text{H}_{20}\text{NaO}_5$ ):  $m/z$  351.1208, found 351.1215.

**rac-(2S,6aR,9S,10aR,10bR)-2-(Furan-3-yl)-9-methoxy-6a-methyl-6,6a,8,9,10a,10b-hexahydro-1H-benzof[isochromene-7,10(2H,4H)-dione (3a).** To a stirring solution of **8a** (45 mg, 0.14 mmol) in toluene (4.5 mL) was added Adogen 464 (20  $\mu\text{L}$ ) followed by a solution containing  $\text{NaHCO}_3$  (184 mg, 42.7 mmol) and  $\text{Na}_2\text{S}_2\text{O}_4$  (tech,  $\sim 85\%$ ) (280 mg, 26.7 mmol) in water (6 mL). The biphasic system was stirred vigorously at room temperature under an atmosphere of nitrogen for 7 h, after which a second portion of  $\text{Na}_2\text{S}_2\text{O}_4$  (tech,  $\sim 85\%$ ) (56 mg, 0.274 mmol) was added. Upon stirring for another 7 h, the layers were separated, and the aqueous layer was extracted with  $\text{Et}_2\text{O}$  ( $2 \times 10$  mL). The combined organic layers were washed with brine (10 mL), dried over  $\text{MgSO}_4$ , filtered, and concentrated in vacuo. The resulting yellow residue was purified by flash chromatography (2:3 EtOAc/hexanes) to yield **3a** (33 mg, 73%) as a white solid. Mp  $142\text{--}145^\circ\text{C}$ .  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.37 (1H, m), 7.35 (1H, t,  $J = 2.1$  Hz), 6.37 (1H, m), 5.56 (1H, m), 4.60 (1H, dd,  $J = 11.2, 1.8$  Hz), 4.25 (1H, d,  $J = 12.6$  Hz), 4.15 (1H, dm,  $J = 12.6$  Hz), 4.04 (1H, ddd,  $J = 12.0, 7.6, 0.9$  Hz), 3.47 (3H, s), 3.09 (1H, dd,  $J = 14.8, 7.6$  Hz), 3.02 (1H, m), 2.95 (1H, dd,  $J = 14.8, 12.0$  Hz), 2.58 (1H, dm,  $J = 18.4$  Hz), 2.44 (1H, d,  $J = 9.9$  Hz), 2.24 (1H, ddd,  $J = 12.5, 4.8, 2.0$  Hz), 2.05 (1H, ddt,  $J = 18.4, 5.7, 2.0$  Hz), 1.19 (1H, q,  $J = 24.0, 11.5$  Hz), 1.03 (3H, s).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  207.7, 205.8, 143.2, 139.1, 132.8, 126.7, 118.9, 108.7, 80.9, 72.4, 72.3, 58.5, 52.9, 49.2, 44.8, 39.5, 33.0, 31.3, 19.1. IR (neat):  $\nu$  2930m, 2907m, 1717s, 1684m, 1654w, 1636w, 1559m  $\text{cm}^{-1}$ .

**Acknowledgment.** Y.B. and P.P. thank the ARC Centre for Green Chemistry for partial funding of this research.

**Supporting Information Available:** General experimental, experimental procedures, and characterization data for compounds **3b**, **4a**, **4b**, **5a**, **5b**, **8b**, and **9b** and  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra for all new compounds, as well as the crystal structure of **3b** in CIF format. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO802623N