


## Efficient Synthesis of Six Tri- to Hexasaccharide Fragments of *Shigella flexneri* Serotypes 3a and/or X O-Antigen, Including a Study on Acceptors Containing *N*-Trichloroacetylglucosamine versus *N*-Acetylglucosamine

Julien Boutet,<sup>†,‡</sup> Catherine Guerreiro, and Laurence A. Mulard\*

Institut Pasteur, Unité de Chimie des Biomolécules (URA CNRS 2128), 28 rue du Dr Roux, F-75015 Paris, France

laurence.mulard@pasteur.fr

Received September 26, 2008



Six tri- to hexasaccharide fragments of the  $\{2\}$ -[ $\alpha$ -D-Glcp-(1 $\rightarrow$ 3)]- $\alpha$ -L-Rhap-(1 $\rightarrow$ 2)- $\alpha$ -L-Rhap-(1 $\rightarrow$ 3)-[Ac $\rightarrow$ 2]- $\alpha$ -L-Rhap-(1 $\rightarrow$ 3)- $\beta$ -D-GlcpNAc-(1 $\rightarrow$ )]<sub>n</sub> polymer ([**(E)AB**<sub>Ac</sub>**CD**]<sub>n</sub>) were synthesized as their propyl glycosides. All targets share the **(E)AB** sequence. Following a thorough investigation on the use of *N*-trichloroacetylglucosamine- versus *N*-acetylglucosamine-containing tri- and tetrasaccharide acceptors, the successful strategy was based on an efficient combination of the trichloroacetimidate chemistry, a trichloroacetyl used as permanent *N*-protection, and an allyl aglycon as temporary and/or permanent anomeric protection of selected building blocks. Use of an **EAB** intermediate orthogonally protected at 2<sub>A</sub> provided both the trisaccharide target and acceptor **12**, the condensation of which with a chain terminator **D** followed by full deprotection, gave tetrasaccharide **D(E)AB**. Alternatively, stepwise glycosylation of **12** with a **D** donor compatible with a selective deblocking at position 3<sub>D</sub> and a 2-*O*-acetyl **C** donor following exposure of OH-3<sub>D</sub> led to a pentasaccharide, which was partially and fully deprotected into free <sub>Ac</sub>**CD(E)AB** and **CD(E)AB**, respectively. Furthermore, chain elongation of the common **D(E)AB** acceptor with a 2<sub>B</sub>-*O*-levulinoyl rhamnobiore donor **BC** and subsequent partial or total deprotection of the resulting hexasaccharide provided **B<sub>Ac</sub>CD(E)AB** and **BCD(E)AB**, respectively. All of the synthesized oligosaccharides are parts of the O-antigen of *Shigella flexneri* 3a, a prevalent serotype. Moreover, the non-*O*-acetylated fragments are also parts of the *S. flexneri* serotype X O-antigen.

### Introduction

Shigellosis, also termed bacillary dysentery, is an invasive infection of the human colon often associated with blood and mucus in the stools.<sup>1</sup> It is one among the many forms of enteric

infections, which altogether rank third among all causes of disease burden worldwide.<sup>2</sup> Shigellosis is a disease of the most impoverished areas and a major health concern particularly in the pediatric population between 1 and 5 years. Humans are the only reservoir of this highly contagious infection associated with increased antibiotic resistance.<sup>1,3,4</sup> The World Health Organization (WHO) has set it as one of its priorities for the

<sup>†</sup> Université Paris Descartes, 12 rue de l'École de Médecine, F-75006 Paris, France.

<sup>‡</sup> Present address: Glycom A/S, DTU, Building 201, DK-2800 KGs, Lyngby, Denmark.

(1) Niyogi, S. K. *J. Microbiol.* **2005**, *43*, 133–143.




FIGURE 1. Repeating units of the O-Ags of *S. flexneri* serotypes Y (I), X (II), 3a (III), and 2a (IV).

development of a vaccine, and a number of candidates have undergone clinical trials. However, as an additional complexity,<sup>4</sup> the large number of pathogens involved seriously hampers the development of an efficacious vaccine against shigellosis, and there is yet no broadly distributed vaccine against this disease.<sup>5</sup> *Shigella flexneri*, one of the four *Shigella* species, is the most frequently isolated causative bacteria worldwide. It is endemic in developing countries and prevails in children less than 5 years old.<sup>5,6</sup> *S. flexneri* is divided into 15 serotypes, differentiated on the basis of the carbohydrate repeating unit of the O-antigen (O-Ag), which is the specific polysaccharide part of the bacterial lipopolysaccharide (LPS),<sup>7</sup> a key player in bacterial virulence and resistance to innate immunity.<sup>8</sup> As a number of different *S. flexneri* serotypes, including serotype 3a, are isolated from patients, there is an inherent need for a multivalent vaccine in order to provide a broad protection.<sup>5,6</sup> Investigations in the field have shown that an initial infection protects against subsequent exposure to homologous serotype.<sup>9,10</sup> This suggests that *S. flexneri* O-Ags are crucial targets used by the host protective adaptive immunity. Interestingly, all *S. flexneri* but serotype 6 share a linear tetrasaccharide backbone, made of three  $\alpha$ -linked L-rhamnosyl residues (A, B, C) and a 2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl residue (D). This tetrasaccharide ABCD (I) is the basic repeating unit (RU) of serotype Y (Figure 1). Additional serotype-specificity is associated with the presence of branched  $\alpha$ -D-glucopyranosyl (E) and O-acetyl decorations.<sup>7</sup> Noteworthy, the LPS glycosylation affects *S. flexneri* 5a O-Ag conformation, resulting in a shortened length.<sup>8,11,12</sup> Impact on bacterial virulence was hypothesized.<sup>8</sup> To our knowledge, the influence of the O-acetylation remains yet unknown. Considering the close resemblance of the O-Ag RUs of the various serotypes, *S. flexneri* was selected as an attractive model to investigate

the key role played by bacterial O-Ag decorations in virulence, resistance to innate immunity and immunodominant epitope composition. Having initiated our study on *S. flexneri* 5a<sup>11,13</sup> and *S. flexneri* 2a,<sup>12,14</sup> we focused more recently on additional serotype-specific glycosylation and O-acetylation patterns.<sup>15,16</sup> Along this line, we report here the synthesis of fragments of *S. flexneri* serotype X and 3a O-Ags, the repeating units of which are the branched pentasaccharides (E)ABCD (II) and (E)AB<sub>Ac</sub>CD (III),<sup>7</sup> respectively (Figure 1). These well-defined synthetic fragments of the native polymer will serve to probe O-Ag structure and antibody recognition as previously reported for serotypes Y,<sup>17</sup> 5a,<sup>11</sup> and 2a.<sup>12</sup> Interestingly, the *S. flexneri* X O-Ag is the non-O-acetylated form of the *S. flexneri* 3a O-Ag.

## Results and Discussion

The syntheses of all glycosylated di- to pentasaccharide fragments of *S. flexneri* 3a and *S. flexneri* X O-Ags having residue A, C, or D, at their reducing end have been described previously.<sup>15,16</sup> We report here the synthesis of the tri-, tetra-, and pentasaccharides EAB (1), D(E)AB (2), <sub>Ac</sub>CD(E)AB (3), and CD(E)AB (4). These compounds complete the panel of frame-shifted glycosylated di- to pentasaccharide fragments of *S. flexneri* 3a O-Ag needed for a detailed investigation of the recognition specificity of *S. flexneri* 3a LPS by protective monoclonal antibodies. In addition, as we are aware of a possible migration of the acetyl group from position 2 of rhamnose C to the *cis*-vicinal hydroxyl in pentasaccharide 3, we also prepared the 2<sub>C</sub>-O-acetylated and non-O-acetylated hexasaccharides B<sub>Ac</sub>CD(E)AB (5), and BCD(E)ABC (6), respectively. Moreover, we propose an optimized protocol for the conversion of a trichloroacetamide moiety vicinal to an ester into the corresponding acetamido alcohol product. Taking advantage of this procedure, two routes to 5 and 6 were studied. The first one involves a 2<sub>D</sub>-acetamido D(E)AB acceptor, and the second one uses a 2<sub>D</sub>-trichloroacetamido D(E)AB acceptor. A comparative study was first made on the more readily available 2<sub>D</sub>-acetamido D(E)A and 2<sub>D</sub>-trichloroacetamido D(E)A acceptors, respectively.

As for other compounds we synthesized in the *S. flexneri* 3a and X series,<sup>15,16</sup> the target products 1–6 were obtained as propyl glycosides. Relying on our previous experience in the synthesis of *S. flexneri* 2a and *S. flexneri* 5a oligosaccharides,<sup>18,19</sup> all glycosylation steps used the efficient trichloroacetimidate

(2) Girard, M. P.; Steele, D.; Chaignat, C. L.; Kiény, M. P. *Vaccine* **2006**, *24*, 2732–2750.

(3) von Seidlein, L.; Kim, D. R.; Ali, M.; Lee, H.; Wang, X.; Thiem, V. D.; Canh do, G.; Chaicumpa, W.; Agtini, M. D.; Hossain, A.; Bhutta, Z. A.; Mason, C.; Sethabutr, O.; Talukder, K.; Nair, G. B.; Deen, J. L.; Kotloff, K.; Clemens, J. *PLoS Med.* **2006**, *3*, e353.

(4) Phalipon, A.; Mulard, L. A.; Sansonetti, P. J. *Microbes Infect.* **2008**, *10*, 1057–1062.

(5) Levine, M. M.; Kotloff, K. L.; Barry, E. M.; Pasetti, M. F.; Sztein, M. B. *Nat. Rev. Microbiol.* **2007**, *5*, 540–553.

(6) Kotloff, K. L.; Winickoff, J. P.; Ivanoff, B.; Clemens, J. D.; Swerdlow, D. L.; Sansonetti, P. J.; Adak, G. K.; Levine, M. M. *Bull. W. H. O.* **1999**, *77*, 651–666.

(7) Lindberg, A. A.; Karnell, A.; Weintraub, A. *Rev. Infect. Dis.* **1991**, *13*, S279–S284.

(8) West, N. P.; Sansonetti, P.; Mounier, J.; Exley, R. M.; Parsot, C.; Guadagnini, S.; Prevost, M. C.; Prochnicka-Chalufour, A.; Delepiere, M.; Tanguy, M.; Tang, C. M. *Science* **2005**, *307*, 1313–1317.

(9) Ferreccio, C.; Prado, V.; Ojeda, A.; Cayazo, M.; Abrego, P.; Guers, L.; Levine, M. M. *Am. J. Epidemiol.* **1991**, *134*, 614–627.

(10) Formal, S. B.; Oaks, E. V.; Olsen, R. E.; Wingfield-Eggleston, M.; Snoy, P. J.; Cogan, J. P. *J. Infect. Dis.* **1991**, *164*, 533–537.

(11) Clément, M. J.; Imbert, A.; Phalipon, A.; Perez, S.; Simenel, C.; Mulard, L. A.; Delepiere, M. *J. Biol. Chem.* **2003**, *278*, 47928–47936.

(12) Vulliez-Le Normand, B.; Saul, F. A.; Phalipon, A.; Bélot, F.; Guerreiro, C.; Mulard, L. A.; Bentley, G. A. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 9976–9981.

(13) Mulard, L. A.; Ughetto-Monfrin, J. J. *Carbohydr. Chem.* **1999**, *18*, 721–753.


(14) Costachel, C.; Sansonetti, P. J.; Mulard, L. A. *J. Carbohydr. Chem.* **2000**, *19*, 1131–1150.

(15) Boutet, J.; Guerreiro, C.; Mulard, L. A. *Tetrahedron* **2008**, *64*, 10558–10572.

(16) Boutet, J.; Mulard, L. A. *Eur. J. Org. Chem.* **2008**, 5526–5542.

(17) Vyas, N. K.; Vyas, M. N.; Chervenak, M. C.; Johnson, M. A.; Pinto, B. M.; Bundle, D. R.; Quijcho, F. A. *Biochemistry* **2002**, *41*, 13575–13586.

(18) Mulard, L. A.; Ughetto-Monfrin, J. J. *Carbohydr. Chem.* **2000**, *19*, 193–220.

SCHEME 1. Synthesis of Propyl Glycosides **1** and **2**

(TCA) chemistry.<sup>20</sup> The allyl group, often used in complex oligosaccharide synthesis,<sup>21,22</sup> was selected for temporary protection at the anomeric position of the various building blocks. Interestingly, in addition to being orthogonal to a number of protecting groups, an allyl ether is easily reduced to a propyl ether upon Pd/C catalyzed hydrogenation. We took advantage of this property to block the reducing end of the target oligosaccharides in a form mimicking the anomery found in the O-Ag.

**Synthesis of Trisaccharide EAB (1) and Tetrasaccharide D(E)AB(2).** We first undertook the preparation of trisaccharide **1** and tetrasaccharide **2**, which did not require chain elongation at the **D** residue and did not comprise the 2-*O*-acetyl **C** residue. As depicted in Scheme 1, the allyl rhamnoside **7**,<sup>16,23</sup> featuring permanent benzyl groups at O-3 and O-4, was glycosylated with the known disaccharide trichloroacetimidate **8**,<sup>24</sup> bearing a benzoyl participating group on position 2<sub>A</sub>. The condensation was run in toluene in the presence of catalytic TfOH to give

the 2<sub>A</sub>-*O*-benzoyl trisaccharide **10** in 93% yield. On the basis of previous observations, made when working on the corresponding **EA** methyl glycoside **13**, which suggested the partial masking at OH-2<sub>A</sub> by the vicinal 2,3,4,6-tetra-*O*-benzyl-D-glucopyranosyl residue,<sup>13</sup> a steric hindrance surrounding OH-2<sub>A</sub> was anticipated. Accordingly, debenzoylation of trisaccharide **10** into alcohol **12** (98%) was achieved using a large excess of sodium methoxide in a 1:2 refluxing mixture of dichloromethane and methanol. Moreover, the recently reported levulinate analogue **9**<sup>16</sup> was investigated as an alternative glycosyl donor. To our satisfaction, condensation of trichloroacetimidate **9** with acceptor **7** led to the fully protected 2<sub>A</sub>-*O*-levulinoyl trisaccharide **11** in 92% yield when the reaction was run in toluene. This solvent was preferred to dichloromethane, in which the reaction led to a lower isolated yield of levulinate **11** (82%). The NMR analysis of **11** showed a <sup>1</sup>J<sub>ClA,H1A</sub> of 173.4 Hz, which indicated an α-AB linkage<sup>25</sup> and ascertained that the levulinic ester had played its role of participating group. Interestingly, treatment of the fully protected **11** in refluxing methanolic sodium methoxide gave alcohol **12** in a satisfactory 94% yield, confirming the potential of donor **9** as an alternative to donor **8**. As shown previously, this is of special interest when orthogonality to acetate is required.<sup>16</sup> Pd/C mediated hydrogenolysis and concomitant allyl reduction of alcohol **12** next provided the linear trisaccharide **1** (77%).

(19) Bélot, F.; Wright, K.; Costachel, C.; Phalipon, A.; Mulard, L. A. *J. Org. Chem.* **2004**, *69*, 1060–1074.

(20) Schmidt, R. R.; Kinzy, W. *Adv. Carbohydr. Chem. Biochem.* **1994**, *50*, 21–123.

(21) Auzanneau, F. I.; Forooghian, F.; Pinto, B. M. *Carbohydr. Res.* **1996**, *291*, 21–41.

(22) Bélot, F.; Guerreiro, C.; Baleux, F.; Mulard, L. A. *Chem. Eur. J.* **2005**, *11*, 1625–1635.

(23) Westerduin, P.; de Haan, P. E.; Dees, M. J.; van Boom, J. H. *Carbohydr. Res.* **1988**, *180*, 195–205.

(24) Mulard, L. A.; Clément, M. J.; Segat-Dioury, F.; Delepierre, M. *Tetrahedron* **2002**, *58*, 2593–2604.

(25) Bock, K.; Pedersen, C. M. *Acta Chem. Scand., Ser. B* **1975**, *29*, 258–264.





## SCHEME 2. Retrosynthetic Analysis of Propyl Glycosides 3–6




TABLE 1. Study on the Condensation of Acceptor 12 and Donor 18

entry	TMSOTf (equiv)	temp (°C)	donor <b>18</b> (equiv)	solvent	yield (%)
1	0.3	-40	1.4	$\text{CH}_2\text{Cl}_2$	35
2	0.3	-40	1.4	toluene	56
3	0.4	-40	1.5	toluene	77
4	0.4	-20	1.5	toluene	70
5	0.3	-40	1.5	toluene	75
6	0.3	-40	1.7	toluene	95

equilibrium, based on NMR data ( $\delta(\text{H-2}_C) = 4.88$  ppm, bs) and ( $\delta(\text{H-3}_C) = 4.93$  ppm, dd), respectively. RP-HPLC analysis also suggested the presence of a third regioisomer. This result confirmed our previous observations on the  $\text{Ac}_2\text{CD(E)A}$  fragment,<sup>15</sup> although exceptions do exist.<sup>34</sup> Subsequent *O*-deacetylation of the propyl glycoside **3**, followed by RP-HPLC purification, gave the free pentasaccharide **4** (63%).

The synthesis of hexasaccharides **5** and **6** was envisioned in the context of a more ambitious goal, that is, the development of a synthetic strategy that would open the way to a variety of larger fragments of *S. flexneri* 3a O-Ag. For that reason, different routes to a suitable  $\text{D(E)AB}$  acceptor were investigated. We first focused on the construction of a fully protected  $\text{D(E)AB}$


intermediate. As an alternative to the pathway described using the **D** donor **18** and the  $\text{EAB}$  trisaccharide acceptor **12** (Scheme 3), the condensation of a  $\text{D(E)A}$  trichloroacetimidate donor having a levulinoyl protecting group at position  $3_D$  **30** and the rhamnoside acceptor **7** was investigated as a potential inroad to a more convergent strategy (Scheme 4). Indeed the  $3_D$ -*O*-levulinoyl protecting pattern was selected in anticipation to any requirement for orthogonality to the  $2_C$ -acetate. Trichloroacetimidate **30** was obtained in three steps from alcohol **27**.<sup>15</sup> Thus, treatment of trisaccharide **27** with levulinic acid in the presence of DCC and DMAP gave the fully protected **28** (90%). Next, allyl glycoside **28** was converted to hemiacetal **29** (80%) following a two-step selective deallylation procedure involving (i) isomerization of the allyl ether into the corresponding prop-1-enyl ether using a cationic iridium complex<sup>35</sup> and (ii) subsequent iodine-mediated hydrolysis.<sup>36</sup> Finally, the  $\text{D(E)A}$  trisaccharide donor **30** was obtained as an anomeric mixture in 85% yield by reacting hemiacetal **29** with trichloroacetimidate in the presence of catalytic DBU. Running the condensation of the latter with the allyl rhamnoside **7** in toluene containing a catalytic amount of TMSOTf gave mostly unreacted **7** and hemiacetal **29**. When toluene was changed for dichloromethane,

(35) Oltvoort, J. J.; van Boeckel, C. A. A.; der Koning, J. H.; van Boom, J. H. *Synthesis* **1981**, 305–308.


(36) Nashed, M. A.; Anderson, L. *Chem. Commun.* **1982**, 1274–1276.

(34) Das, S. K.; Roy, N. *Carbohydr. Res.* **1995**, *271*, 177–183.

## SCHEME 3. Synthesis of Propyl Glycosides 3 and 4



## SCHEME 4. Study on the Synthesis of D(E)AB According to a D(E)A + B Protocol





the acceptor **7** was totally consumed. Two major compounds **31** and **32**, both corresponding to condensation products, as seen by mass spectrometry analysis, were isolated in a 1:1 ratio (88%). Careful NMR analysis confirmed that **31**, the faster eluting product, was the expected tetrasaccharide having an  $\alpha$ -AB linkage ( $^1J_{\text{C1A,H1A}} = 171.3$  Hz). Interestingly, compound **32** was the stereoisomer having a  $\beta$ -AB linkage ( $^1J_{\text{C1A,H1A}} = 155.1$  Hz). Although formation of the latter was anticipated in the absence of a participating group at position  $2_A$  of trichloroacetimidate **30**, its isolation in such a high yield prevented any further investigation of this strategy. The **D** + **EAB** route was therefore adopted.

*N*-Acetyl- $\beta$ -D-glucosamine is present in numerous oligosaccharides of natural origin. However, the acetamido moiety can hardly be used for anchimeric assistance in complex oligosaccharide synthesis. For that reason, a number of *N*-protecting groups serving as participating groups, or on the contrary

preventing anchimeric assistance, were developed.<sup>37</sup> Whether these protecting groups should be regarded as permanent or temporary acetamido masking patterns is often defined on a case by case basis. For example, in the synthesis of a number of oligosaccharides representative of *S. flexneri* 2a O-Ag, with pentasaccharide **IV** (Figure 1), a regioisomer of **II**, as a repeating unit, chain elongation at OH-3<sub>D</sub> was performed using acceptors containing *N*-acetylglucosamine residues.<sup>22</sup> However, others encountered difficulties when attempting to glycosylate poorly reactive hydroxyl groups as part of *N*-acetylglucosamine-containing acceptors.<sup>21,31</sup> Aiming at reducing the number of deprotection steps that could potentially interfere with the 2<sub>C</sub>-acetate at the late stage of the synthesis, the next step was thus to investigate the best protecting group pattern of the amino group of glucosamine **D** when involved in building blocks

(37) Bongat, A. F.; Demchenko, A. V. *Carbohydr. Res.* **2007**, *342*, 374–406.

SCHEME 5. Conversion of the 2<sub>D</sub>-Trichloroacetamide **33** into the 2<sub>D</sub>-Acetamide **36**CHART 1. Study on the Conversion of **33** into **36**<sup>a</sup>

<sup>a</sup> (A) The initial concentration of **33** in MeOH was 0.4 M (5 equiv of MeONa). (B) The CH<sub>2</sub>Cl<sub>2</sub>/MeOH ratio (v/v) was constant at 8.

required for chain elongation at position 3<sub>D</sub>, such as **D(E)AB**. To broaden the analysis, the study was initiated on the more readily available **D(E)A**.

Finding appropriate conditions to ensure efficient conversion of the *N*-trichloroacetyl moiety into the corresponding acetamide was our first goal. Most commonly used methodologies include basic trichloroacetyl removal and subsequent *N*-acetylation of the resulting amine,<sup>24,38,39</sup> direct reduction under neutral conditions such as Bu<sub>3</sub>SnH-mediated radical hydrodechlorination,<sup>22,26</sup> or often at the final stage of the synthesis, nonspecific palladium-mediated reductive hydrodechlorination.<sup>19,39</sup> Avoiding the tin methodology, we focused on finding chemoselective conditions applicable to nonpolar compounds by taking advantage of former observations. Indeed, we have reported the high-yielding transesterification of the triacetate **33**<sup>15</sup> into triol **34**<sup>15</sup> by use of a methanolic solution of sodium methoxide under controlled conditions. However, we also reported *O*-deacetylation and a concomitant vicinal *N*-trichloroacetyl conversion into the corresponding amine in the presence of additional Et<sub>3</sub>N<sup>14</sup> or dichloromethane.<sup>15,24</sup> We now report on a detailed investigation of the latter conditions resulting in the efficient conversion of triacetate **33** into acetamidotriol **36**<sup>15</sup> (Scheme 5).

When running the deprotection step in a mixture of dichloromethane and methanol, two closely migrating compounds were isolated following subsequent *N*-acetylation of the aminotriol intermediate **35**. On the basis of NMR and mass spectrometry data, the more polar one was the expected acetamido triol **36** (HRMS (ESI<sup>+</sup>) for C<sub>58</sub>H<sub>69</sub>NO<sub>15</sub> ([M + H]<sup>+</sup>, 1020.4745) found *m/z* 1020.4739,  $\delta_{\text{CO}} = 173.3$  ppm,  $\delta_{\text{Me}} =$

23.6 ppm),<sup>15</sup> whereas the first migrating compound was identified as the methyl carbamate **37** (HRMS (ESI<sup>+</sup>) for C<sub>58</sub>H<sub>69</sub>NO<sub>16</sub> ([M + Na]<sup>+</sup>, 1058.4514) found *m/z* 1058.4514,  $\delta_{\text{CO}} = 159.4$  ppm,  $\delta_{\text{OMe}} = 52.9$  ppm), the formation of which is tentatively explained via the addition of methanol onto isocyanate **38**.<sup>40</sup> Varying the reaction conditions in terms of dichloromethane/methanol ratio and amount of sodium methoxide highlighted the impact of these two parameters on the consumption of the starting triacetate **33** (Chart 1). Under the best conditions (6 equiv of sodium methoxide in a 8:1 (v/v) mixture of dichloromethane/MeOH), the transformation was completed within 2 h, providing acetamido **36** and carbamate **37** in 90% and 5% yield post *N*-acetylation, respectively. Satisfactorily, treatment of the 3<sub>D</sub>-acetate **39**<sup>15</sup> under the same conditions gave the known 2<sub>D</sub>-acetamido acceptor **40**<sup>15</sup> in 90% yield.<sup>15</sup>

This new route to acceptor **40** involving the isopropylidene donor **18** was compared to the one reported earlier using the triacetate donor **14**.<sup>15</sup> The corresponding overall yields of acetamide **40** from allyl glycoside **41**, 81% via isopropylidene **39** and 70% via triacetate **33**, respectively, were clearly in favor of the methodology just disclosed when working below the millimolar scale (Scheme 6). When working on 2.1 g of acceptor **41**<sup>24</sup> (2.6 mmol), the yield of trisaccharide **40** via route 1 was 86% and involved one purification step only.


The new conditions were successfully applied to the synthesis of the tetrasaccharide acceptor **D(E)AB**, resulting in an improved yield of acetamide **21** from intermediate **19**, 83% versus 77%. Alternatively, selective removal of the 3<sub>D</sub>-acetate in tetrasaccharide **19** by use of potassium carbonate in methanol gave acceptor **20** in 92% yield (Scheme 7).

(38) Sherman, A. A.; Yudina, O. N.; Mironov, Y. V.; Sukhova, E. V.; Shashkov, A. S.; Menshov, V. M.; Nifantiev, N. E. *Carbohydr. Res.* **2001**, *336*, 13–46.


(39) Donohoe, T. J.; Logan, J. G.; Laffan, D. D. *Org. Lett.* **2003**, *5*, 4995–4998.

(40) Nishikawa, T.; Urabe, D.; Tomita, M.; Tsujimoto, T.; Iwabuchi, T.; Isobe, M. *Org. Lett.* **2006**, *8*, 3263–3265.

SCHEME 6. Access to Acceptor 40 by Condensation of 41 with Donor 14 (Route 1) and Donor 18 (Route 2)




SCHEME 7. Conversion of Fully Protected 19 into Acceptor 20 or 21



Having all four different acceptors **20**, **21**, **27**, and **40** in hands, the next step was to synthesize the rhamnose donor **23**. This was easily achieved from the known allyl glycoside **42**.<sup>16</sup> Indeed, selective cleavage of the allyl aglycon gave hemiacetal **43** (93%), which was converted into trichloroacetimidate **23** (84%), when treated with trichloroacetonitrile in the presence of a catalytic amount of DBU (Scheme 8).

We next focused on the condensation of donor **23** with the 3<sub>D</sub>-acetamido acceptor **21** and the subsequent transformation of the condensation product. Taking into account the fact that TfOH-mediated glycosylation of acceptor **21** with trichloroacetimidate **22** in toluene succeeded in providing **25** (Scheme 3), while conventional TMSOTf-mediated glycosylation of acceptor **40** with the same donor was somewhat unsuccessful,<sup>15</sup> we chose to first investigate the use of both catalysts in the condensation of trisaccharide **40** and donor **23**. Running the reaction in the presence of 0.3 equiv of TMSOTf, at  $-20^\circ\text{C}$ , in toluene or dichloromethane, resulted at best in 25% yield of the pentasaccharide **44**, thus confirming our previous observations. Degradation


SCHEME 8. Synthesis of Donor 23



tion of the donor was observed, whereas unreacted **40** was recovered. Turning to TfOH, this time used in larger amounts (0.9 equiv) was more satisfactory (Scheme 9, route a). Indeed, running the reaction in toluene at  $0^\circ\text{C}$  resulted in 74% of isolated pentasaccharide **44**. Since the yield of **44** was not higher when running the condensation between acceptor **40** and donor **23** at  $70^\circ\text{C}$ ,<sup>22</sup> the TMSOTf-mediated condensation of the same donor **23** with the tetrasaccharide acceptor **21** was run at  $0^\circ\text{C}$ , providing hexasaccharide **45** in 65% yield (Scheme 9, route b). However, careful NMR analysis of the condensation products **44** and **45** suggested, in both cases, the presence of small amounts of a coeluting contaminant (5–10%). Attempts to remove this unidentified side product after cleavage of either the levuniloyl ester or the isopropylidene acetal failed (not described). We thus turned to the use of the corresponding 2<sub>D</sub>-trichloroacetamide acceptors **27** and **20**.

When applying the reaction conditions found successful for the condensation of acceptor **27** and the chain-terminator **BC** donor **46**<sup>15</sup> to the glycosylation of **27** and donor **23**, we could confirm the stability of the isopropylidene acetal at  $-78^\circ\text{C}$ . However, the yield of the condensation product **48** (63%) was lower than expected. Indeed, although the reaction was run for a longer period (1 h instead of 15 min), some starting acceptor remained (16%), whereas the donor was degraded (Table 2, entry 1). Interestingly, when the condensation was run on a 2 g




SCHEME 9. Condensation of Acceptors **40** (D(E)A) (Route a) and **21** (D(E)AB) (Route b) with the BC Donor **23**TABLE 2. Study on the Condensation of Acceptor **27** and Donor **23**

entry	temp (°C)	solvent	time	acceptor <b>27</b> (%)	<b>48</b> (%)	<b>49</b> (%)
1	-78	toluene	1 h	16	63	no <sup>a</sup>
2	-78	CH <sub>2</sub> Cl <sub>2</sub>	1 h	30	42	no
3	-40	toluene	30 min	no	66	22
4	-20	toluene	30 min	no	66	13
5	-40	toluene	10 min	no	75	no


<sup>a</sup> no = not observed.

scale of acceptor **27**, the silylated analogue **47** was isolated (11%) in addition to pentasaccharide **48** (52%). Adding more donor or increasing the amount of TMSOTf had no impact. Changing toluene for dichloromethane worsened the outcome of the glycosylation (Table 2, entry 2), since 30% of unreacted acceptor was recovered in addition to **48** (42%). The temperature of the reaction was investigated next in order to overcome this unexpected outcome. Increasing the reaction temperature to -40 °C was highly satisfactory, since neither the remaining acceptor **27** nor the silylated **47** were apparent. In these conditions (Table 2, entry 3), the overall condensation yield was 88% (Scheme 10, route a). However, the poor stability at this temperature of the 4<sub>D</sub>,6<sub>D</sub>-*O*-isopropylidene resulted in the isolation of two products, the expected **48** (<sup>1</sup>J<sub>C1C,H1C</sub> = 167.6 Hz, 66%) together with the corresponding diol **49** (22%, not described), resulting from isopropylidene loss post condensation, as ascertained by mass spectrometry analysis (HRMS (ESI<sup>+</sup>) for C<sub>98</sub>H<sub>112</sub>Cl<sub>3</sub>NO<sub>26</sub> ([M + Na]<sup>+</sup> 1846.643) *m/z* 1846.6403) and reacetalation. Additional increase of the temperature had no positive effect (Table 2, entry 4). Running the condensation in toluene, at -40 °C, for a shorter period of time (10 min), finally provided pentasaccharide **48** in 75% yield (Table 2, entry 5). Under these conditions, the diol **49** was not isolated (Scheme 10, route b).

Having demonstrated the crucial importance of temperature on the outcome of the condensation between the trisaccharide acceptor **27** and the disaccharide donor **23**, we turned to the preparation of hexasaccharide **50** (Scheme 11). Taking advantage of the study made for obtaining pentasaccharide **48**, the condensation of the trichloroacetimidate **23** and the tetrasac-

SCHEME 10. Condensation of Acceptor **27** and Donor **23**

charide acceptor **20** was run in toluene, at -40 °C, for 10 min in the presence of a catalytic amount of TMSOTf. Under those conditions, the yield of the condensation was 80%, matching our expectations. However, two products were isolated, the fully protected **50** (59%) and a more polar product (21%). Mass spectrometry and NMR analysis of the latter indicated the formation of diol **51**, corresponding to the 4<sub>D</sub>,6<sub>D</sub>-*O*-isopropylidene loss. Treatment of **51** with 2-methoxypropene and CSA in DMF confirmed this assumption, since the fully protected **50** was isolated in 93% yield. Since diol **51** was a suitable intermediate in the synthesis targets **5** and **6**, preventing acetal loss was not attempted. To our satisfaction, running the condensation on larger amounts of acceptor **20** (1.1 mmol instead of 0.2 mmol) under the same conditions but for longer time (45 min) to ensure total consumption of the acceptor, provided **50** and **51** in a 3:1 ratio and 80% total yield. Subsequent partial and total deprotection of the condensation product **50** provided hexasaccharides **5** and **6**, respectively. Indeed, TFA-mediated hydrolysis of the 4<sub>D</sub>,6<sub>D</sub>-*O*-isopropylidene of **50** gave diol **51** (89%). The following selective removal of the levuniloyl protecting group in hexasaccharide **51**, using aqueous hydrazine in pyridine/acetic acid, gave triol **52** (84%). Alternatively, reacting **51** with sodium methoxide in refluxing methanol gave tetraol **53** (92%). Conversion of the allyl glycosides **52** and **53** into the propyl glycosides **5** and **6**, respectively, took advantage of the neutral conditions used previously for the synthesis of related *S. flexneri* 3a oligosaccharides.<sup>15,16</sup> In practice, treatment of ethanolic solutions of triol **52** and tetraol **53** under a hydrogen atmosphere (50 bar), in the presence of Pd/C catalyst for 10 days each, allowed concomitant benzyl removal, allyl reduction into propyl, and trichloroacetamide conversion into the required acetamide. The hexasaccharide targets **5** and **6** were finally isolated in 64% and 82% yield,

SCHEME 11. Synthesis of Hexasaccharides **5** and **6** from Acceptor **20** (D(E)AB) and Donor **23** (BC)

respectively. Overall, the efficiency of the **20** + **23** glycosylation step along with this one-step conversion fully supported the selection of trichloroacetamide **20** as the acceptor in the synthesis of hexasaccharides **5** and **6**.

### Conclusion

We have described the synthesis of a linear trisaccharide (**1**) and five branched tetra-, penta-, or hexasaccharides (**2–6**) representative of fragments of *S. flexneri* 3a and/or X O-Ags. All the oligosaccharide targets share the **EAB** sequence, whereby residue **B** is blocked at the anomeric position with a propyl aglycon. They were synthesized from a common **EAB** allyl glycoside **12**, obtained in 18 synthetic steps. A one-step deprotection and concomitant allyl to propyl conversion of alcohol **12** gave the trisaccharide target **1**. Alternatively, glycosylation of acceptor **12** with **D** donors **14** (4 steps) and **18** (5 steps) gave tetrasaccharides **15** and **19**, respectively. On one hand, conventional deprotection of the triacetate **15** gave the tetrasaccharide target **2** (3 steps from intermediate **12**). On the other hand, the  $4_D, 6_D$ -*O*-isopropylidene tetrasaccharide **19** served as key intermediate to either the *N*-trichloroacetyl acceptor **20** or the *N*-acetyl analogue **21**. Noteworthy, the one-step basic conversion of the fully protected **19** into acceptor **21** (83%) relied on in-house conditions, the optimization of which is also disclosed here. Acceptors containing *N*-acetylglucosamine have some limitations, potentially preventing their use in complex oligosaccharide synthesis. In the course of this work, two of such acceptors, trisaccharide **40** and tetrasaccharide **21**, were compared to their *N*-trichloroacetyl counterparts **27** and **20**, respectively. Independently of the acceptor used, our work pointed out (i) the lower stability of the  $4_D, 6_D$ -*O*-isopropylidene acetal when located on the trichloroacetamide derivatives **27** and **20** than when located on the acetamido analogues **40** and **21**, (ii) the crucial impact of temperature on the glycosylation outcome, and (iii) the key input of the condensation duration in controlling the compromise between glycosylation yield and isopropylidene loss. Optimizing both parameters resulted in good yields of glycosylation in all cases. Nevertheless, the presence of tiny amounts of unidentified contaminants associated with the use of the acetamido acceptors **40** and **21** encouraged the use of the *N*-trichloroacetyl derivatives **27** and **20**, despite their higher propensity to acetal loss. Finally, chain elongation of

acceptor **20** with donor **23** provided the fully protected hexasaccharide **50**. Conversion of the latter into targets **5** and **6** took advantage of the feasible concomitant removal and/or transformation of the protecting groups, including benzyl hydrogenolysis, allyl reduction, and reductive trichloroacetamide conversion to acetamide. Therefore, partial deprotection of the allyl glycoside **50** gave monoacetate **5** (3 steps), whereas full deprotection provided hexasaccharide **6** (3 steps). Data reported herein provide the chemical tools for further synthesis of larger fragments of *S. flexneri* serotypes 3a and/or X O-Ags.

### Experimental Section

**Allyl (2,3,4,6-Tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl)-(1 $\rightarrow$ 3)-(2-*O*-benzoyl-4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (**10**).** TfOH (16  $\mu$ L, 186  $\mu$ mol, 0.2 equiv) was added to a solution of acceptor **7**<sup>16,23</sup> (463 mg, 1.2 mmol, 1.3 equiv) and trichloroacetimidate **8**<sup>24</sup> (953 mg, 0.93 mmol) in toluene (10 mL) containing 4 $\text{\AA}$  MS (500 mg), stirred at  $-30^\circ\text{C}$ . The reaction mixture was stirred for 1 h while slowly coming back to rt. TLC ( $\text{CH}_2\text{Cl}_2/\text{EtOAc}$ , 98:2) showed the complete disappearance of the acceptor and the presence of a major less polar product. Et<sub>3</sub>N (300  $\mu$ L) was added. The mixture was filtered and concentrated to dryness. Chromatography of the residue (Chex/EtOAc, 100:0  $\rightarrow$  80:20) gave the allyl glycoside **10** (1.08 g, 93%) as a colorless syrup. Trisaccharide **10** had  $R_f = 0.5$  (Chex/EtOAc, 8:2), <sup>1</sup>H NMR ( $\text{CDCl}_3$ )  $\delta$  8.09–8.07 (m, 2H, CH<sub>Bz</sub>), 7.59 (m, 1H, CH<sub>Bz</sub>), 7.47–7.08 (m, 37H, CH<sub>Ph</sub>), 5.89 (m, 1H, CH=), 5.84 (dd, 1H,  $J_{1,2} = 2.1$  Hz, H-2<sub>A</sub>), 5.37 (d, 1H,  $J_{1,2} = 3.4$  Hz, H-1<sub>E</sub>), 5.28 (m, 1H,  $J_{\text{trans}} = 17.3$  Hz, =CH<sub>2</sub>), 5.20 (m<sub>overlapped</sub>, 1H, =CH<sub>2</sub>), 5.19 (m<sub>overlapped</sub>, 1H, H-1<sub>A</sub>), 5.00 (d, 1H,  $J = 10.2$  Hz, H<sub>Bn</sub>), 4.96 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.86 (d, 1H,  $J = 10.9$  Hz, H<sub>Bn</sub>), 4.84 (d, 1H,  $J = 10.9$  Hz, H<sub>Bn</sub>), 4.78 (d, 1H,  $J_{1,2} = 1.8$  Hz, H-1<sub>B</sub>), 4.75 (d, 1H,  $J = 10.9$  Hz, H<sub>Bn</sub>), 4.72 (s, 2H, H<sub>Bn</sub>), 4.67–4.63 (m, 3H, H<sub>Bn</sub>), 4.60 (d, 1H,  $J = 12.1$  Hz, H<sub>Bn</sub>), 4.47–4.44 (m, 2H, H<sub>Bn</sub>), 4.39 (dd, 1H,  $J_{2,3} = 3.1$  Hz,  $J_{3,4} = 9.8$  Hz, H-3<sub>A</sub>), 4.36 (d, 1H,  $J = 12.0$  Hz, H<sub>Bn</sub>), 4.15 (m, 1H, H<sub>All</sub>), 4.08–4.03 (m, 3H, H-3<sub>E</sub>, H-5<sub>E</sub>, H-2<sub>B</sub>), 3.97–3.90 (m, 3H, H<sub>All</sub>, H-5<sub>A</sub>, H-3<sub>B</sub>), 3.79 (dd, 1H,  $J_{4,5} = 9.4$  Hz,  $J_{3,4} = 9.7$  Hz, H-4<sub>E</sub>), 3.76–3.69 (m, 3H, H-5<sub>B</sub>, H-6<sub>aE</sub>, H-4<sub>A</sub>), 3.62 (dd, 1H,  $J_{2,3} = 9.7$  Hz, H-2<sub>E</sub>), 3.59 (m, 1H,  $J_{5,6} = 1.7$  Hz, H-6<sub>bE</sub>), 3.56 (pt, 1H,  $J_{4,5} = J_{3,4} = 9.4$  Hz, H-4<sub>B</sub>), 1.43 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>A</sub>), 1.35 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>B</sub>); <sup>13</sup>C NMR ( $\text{CDCl}_3$ )  $\delta$  166.0 (C<sub>Bz</sub>), 139.1–138.0 (C<sub>Ph</sub>), 134.2 (CH=), 133.5, 130.4 (3C, CH<sub>Bz</sub>), 129.2–127.7 (CH<sub>Ph</sub>, CH<sub>Bz</sub>), 117.6 (=CH<sub>2</sub>), 99.6 (C-1<sub>A</sub>), <sup>1</sup>J<sub>CH</sub> = 172.0 Hz), 98.2 (C-1<sub>B</sub>), <sup>1</sup>J<sub>CH</sub> = 169.5 Hz), 93.1 (C-1<sub>E</sub>), <sup>1</sup>J<sub>CH</sub> = 169.2 Hz), 82.4 (C-3<sub>E</sub>), 80.9 (C-4<sub>B</sub>), 80.3 (C-4<sub>A</sub>), 80.1 (C-3<sub>B</sub>), 79.3

(C-2<sub>E</sub>), 77.8 (C-4<sub>E</sub>), 76.7, 76.0, 75.9, 75.3, (4C, C<sub>Bn</sub>), 75.1 (C-2<sub>B</sub>), 73.8 (C<sub>Bn</sub>), 72.9 (C-3<sub>A</sub>), 72.6, 72.5 (2C, C<sub>Bn</sub>), 70.6 (C-5<sub>E</sub>), 68.8 (2C, C-2<sub>A</sub>, C-5<sub>B</sub>\*), 68.6 (C-6<sub>E</sub>), 68.4 (C-5<sub>A</sub>\*), 68.0 (C<sub>All</sub>), 18.5, 18.4 (2C, C-6<sub>A</sub>, C-6<sub>B</sub>); HRMS (ESI<sup>+</sup>) for C<sub>75</sub>H<sub>82</sub>O<sub>15</sub> ([M + Na]<sup>+</sup>, 1269.5552) found *m/z* 1269.5563.

**Allyl (2,3,4,6-Tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl)-(1 $\rightarrow$ 3)-(4-*O*-benzyl-2-*O*-levulinoyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (11).** TMSOTf (210  $\mu$ L, 1.2 mmol, 0.3 equiv) was added to a solution of acceptor **7** (1.5 g, 3.9 mmol) and trichloroacetimidate **9**<sup>16</sup> (5.1 g, 5.1 mmol, 1.3 equiv) in toluene (100 mL) containing 4 $\text{\AA}$  MS (3.4 g), stirred at  $-78\text{ }^\circ\text{C}$ . The reaction mixture was stirred for 15 min while slowly coming back to rt. TLC (Tol/EtOAc, 85:15) showed the complete disappearance of the acceptor and the presence of a major less polar product. Et<sub>3</sub>N (1 mL) was added. The mixture was filtered, and concentrated to dryness. Chromatography of the residue (Tol/EtOAc, 95:5  $\rightarrow$  85:15) gave **11** (4.5 g, 92%) as a colorless syrup. Trisaccharide **11** had *R<sub>f</sub>* = 0.5 (Chex/EtOAc, 85:15); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.45–7.12 (m, 35H, CH<sub>Ph</sub>), 5.87 (m, 1H, CH=), 5.61 (dd, 1H, *J*<sub>1,2</sub> = 2.1 Hz, H-2<sub>A</sub>), 5.32 (d, 1H, *J*<sub>1,2</sub> = 3.4 Hz, H-1<sub>E</sub>), 5.31 (m, 1H, *J*<sub>trans</sub> = 17.2 Hz, =CH<sub>2</sub>), 5.23 (m, 1H, *J*<sub>cis</sub> = 10.3 Hz, =CH<sub>2</sub>), 5.06 (d<sub>po</sub>, 1H, *J* = 10.5 Hz, H<sub>Bn</sub>), 5.05 (d<sub>po</sub>, 1H, H-1<sub>A</sub>), 5.00–4.89 (m, 5H, H<sub>Bn</sub>), 4.82 (d<sub>po</sub>, 1H, *J* = 12.2 Hz, H<sub>Bn</sub>), 4.81 (bs<sub>o</sub>, 1H, H-1<sub>B</sub>), 4.74–4.64 (m, 5H, H<sub>Bn</sub>), 4.53 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.38 (d, 1H, *J* = 12.0 Hz, H<sub>Bn</sub>), 4.26 (dd, 1H, *J*<sub>2,3</sub> = 3.2 Hz, *J*<sub>3,4</sub> = 9.6 Hz, H-3<sub>A</sub>), 4.20–4.09 (m, 3H, H<sub>All</sub>, H-3<sub>E</sub>, H-5<sub>E</sub>), 4.03 (dd, 1H, *J*<sub>1,2</sub> = 1.9 Hz, *J*<sub>2,3</sub> = 2.9 Hz, H-2<sub>B</sub>), 3.97 (m, 1H, H<sub>All</sub>), 3.96–3.90 (m, 2H, H-5<sub>A</sub>, H-3<sub>B</sub>), 3.83 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.5 Hz, H-4<sub>E</sub>), 3.75 (dq, 1H, *J*<sub>4,5</sub> = 9.4 Hz, H-5<sub>B</sub>), 3.69 (dd<sub>po</sub>, 1H, *J*<sub>5,6a</sub> = 2.7 Hz, *J*<sub>6a,6b</sub> = 10.9 Hz, H-6<sub>A</sub>E), 3.68 (dd<sub>po</sub>, 1H, *J*<sub>2,3</sub> = 9.6 Hz, H-2<sub>E</sub>), 3.61 (pt<sub>po</sub>, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.4 Hz, H-4<sub>A</sub>), 3.60 (dd<sub>po</sub>, 1H, H-6<sub>B</sub>E), 3.53 (pt, 1H, *J*<sub>3,4</sub> = 9.4 Hz, H-4<sub>B</sub>), 2.55 (m, 4H, H<sub>Lev</sub>), 2.11 (s, 3H, CH<sub>3Lev</sub>), 1.42 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>A</sub>), 1.37 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>B</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  206.2 (C<sub>Lev</sub>), 171.8 (C<sub>Lev</sub>), 138.7–137.7 (C<sub>Ph</sub>), 133.9 (CH=), 129.1–127.6 (CH<sub>Ph</sub>), 117.2 (=CH<sub>2</sub>), 99.2 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 173.4 Hz), 97.9 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.8 Hz), 92.9 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 168.9 Hz), 82.1 (C-3<sub>E</sub>), 80.5 (C-4<sub>B</sub>), 79.9 (C-4<sub>A</sub>), 79.6 (C-3<sub>B</sub>), 79.5 (C-2<sub>E</sub>), 77.8 (C-4<sub>E</sub>), 76.2, 75.6, 75.5 (3C, C<sub>Bn</sub>), 75.1 (C-2<sub>B</sub>), 75.0, 73.4, 72.8 (3C, C<sub>Bn</sub>), 72.3 (C-3<sub>A</sub>), 72.2 (C<sub>Bn</sub>), 70.3 (C-5<sub>E</sub>), 68.4 (C-5<sub>A</sub>), 68.3 (C-6<sub>E</sub>), 68.1 (C-5<sub>B</sub>), 68.0 (C-2<sub>A</sub>), 67.7 (C<sub>All</sub>), 38.0 (CH<sub>2Lev</sub>), 29.7 (CH<sub>3Lev</sub>), 28.2 (CH<sub>2Lev</sub>), 18.1, 18.0 (2C, C-6<sub>A</sub>, C-6<sub>B</sub>); HRMS (ESI<sup>+</sup>) for C<sub>75</sub>H<sub>84</sub>O<sub>16</sub> ([M + Na]<sup>+</sup>, 1263.5657) found *m/z* 1263.5437, ([M + NH<sub>4</sub>]<sup>+</sup>, 1258.6104) found *m/z* 1258.5887.

**Allyl (2,3,4,6-Tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl)-(1 $\rightarrow$ 3)-(4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (12).** **Route a.** Methanolic MeONa (0.5 M, 10 mL, 5 mmol, 6.6 equiv) was added to a solution of benzoyleated **10** (935 mg, 750  $\mu$ mol) in a 1:1 MeOH/CH<sub>2</sub>Cl<sub>2</sub> (v/v) mixture (12 mL), and the mixture was refluxed for 3 h. TLC (Chex/EtOAc, 7:3) showed the complete disappearance of **10** and the presence of a single more polar product. The mixture was neutralized by addition of Dowex X8-200 ion-exchange resin (H<sup>+</sup>) and filtered. Volatiles were evaporated. The resulting syrup was chromatographed (Chex/EtOAc, 100:0  $\rightarrow$  7:3) to give **12** (845 mg, 98%) as a colorless oil.

**Route b.** Methanolic MeONa (0.5 M, 12.2 mL, 6.1 mmol, 1.7 equiv) was added to a solution of fully protected **11** (4.5 g, 3.6 mmol) in MeOH (200 mL), and the mixture was refluxed for 1 h. TLC (Chex/EtOAc, 7:3) showed the complete disappearance of the starting material and the presence of a single more polar product. The mixture was neutralized by addition of Dowex X8-200 ion-exchange resin (H<sup>+</sup>) and filtered. Volatiles were evaporated. The resulting syrup was chromatographed (Tol/EtOAc, 98:2  $\rightarrow$  9:1) to give **12** (3.9 g, 94%). Trisaccharide **12** had *R<sub>f</sub>* = 0.6 (Chex/EtOAc, 7:3); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.47–7.19 (m, 35H, CH<sub>Ph</sub>), 5.93 (m, 1H, CH=), 5.32 (m, 1H, *J*<sub>trans</sub> = 17.2 Hz, =CH<sub>2</sub>), 5.25 (m, 1H, *J*<sub>cis</sub> = 10.3 Hz, =CH<sub>2</sub>), 5.22 (bs, 1H, H-1<sub>A</sub>), 5.05–4.99 (m, 3H, H<sub>Bn</sub>), 4.97 (d<sub>po</sub>, 1H, *J*<sub>1,2</sub> = 3.5 Hz, H-1<sub>E</sub>), 4.91–4.88 (m, 2H, H<sub>Bn</sub>), 4.84 (d, 1H, *J*<sub>1,2</sub> = 1.6 Hz, H-1<sub>B</sub>), 4.83–4.67 (m, 6H, H<sub>Bn</sub>), 4.60 (d, 1H,

*J* = 12.0 Hz, H<sub>Bn</sub>), 4.54 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.36 (d, 1H, *J* = 12.0 Hz, H<sub>Bn</sub>), 4.19 (m, 1H, H<sub>All</sub>), 4.14–4.09 (m, 4H, H-2<sub>A</sub>, H-2<sub>B</sub>, H-3<sub>A</sub>, H-3<sub>E</sub>), 4.04–3.99 (m, 2H, H<sub>All</sub>, H-5<sub>E</sub>), 3.98 (dd, 1H, *J*<sub>2,3</sub> = 2.9 Hz, *J*<sub>3,4</sub> = 9.5 Hz, H-3<sub>B</sub>), 3.92 (dq, 1H, H-5<sub>A</sub>), 3.82 (dd, 1H, *J*<sub>3,4</sub> = 9.4 Hz, *J*<sub>4,5</sub> = 9.8 Hz, H-4<sub>E</sub>), 3.75 (dq, 1H, *J*<sub>4,5</sub> = 9.4 Hz, H-5<sub>B</sub>), 3.66 (dd, 1H, *J*<sub>1,2</sub> = 3.6 Hz, *J*<sub>2,3</sub> = 9.6 Hz, H-2<sub>E</sub>), 3.75 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.3 Hz, H-4<sub>A</sub>), 3.55–3.49 (m, 2H, H-6<sub>A</sub>E, H-4<sub>B</sub>), 3.45 (dd, 1H, *J*<sub>5,6a</sub> = 1.8 Hz, *J*<sub>6a,6b</sub> = 10.8 Hz, H-6<sub>B</sub>E), 1.43 (d, 3H, *J*<sub>5,6</sub> = 6.3 Hz, H-6<sub>A</sub>), 1.39 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>B</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  138.7–137.6 (C<sub>Ph</sub>), 153.9 (CH=), 129.1–127.0 (CH<sub>Ph</sub>), 117.2 (=CH<sub>2</sub>), 101.0 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.3 Hz), 98.1 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.3 Hz), 94.0 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 166.7 Hz), 82.4 (C-3<sub>E</sub>), 80.6 (C-4<sub>B</sub>), 80.0 (C-3<sub>B</sub>), 79.3 (C-4<sub>A</sub>), 79.0 (C-2<sub>E</sub>), 77.8 (C-4<sub>E</sub>), 76.5 (C-3<sub>A</sub>), 75.6, 75.6, 75.5 (3C, C<sub>Bn</sub>), 75.0 (C-2<sub>B</sub>), 74.9, 74.4, 73.4, 72.5 (4C, C<sub>Bn</sub>), 70.7 (C-5<sub>E</sub>), 68.1 (C-5<sub>B</sub>), 68.0 (C-6<sub>E</sub>), 67.7 (C-5<sub>A</sub>), 67.7 (C<sub>All</sub>), 67.4 (C-2<sub>A</sub>), 18.0, 17.9 (2C, C-6<sub>A</sub>, C-6<sub>B</sub>); HRMS (ESI<sup>+</sup>) for C<sub>70</sub>H<sub>78</sub>O<sub>14</sub> ([M + Na]<sup>+</sup>, 1165.5289) found *m/z* 1165.5283, ([M + NH<sub>4</sub>]<sup>+</sup>, 1160.5735) found *m/z* 1160.5674.

**Propyl  $\alpha$ -D-Glucopyranosyl-(1 $\rightarrow$ 3)- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\alpha$ -L-rhamnopyranoside (1).** Alcohol **12** (260 mg, 209  $\mu$ mol) was dissolved in EtOH (10 mL) containing 1 M aq HCl (155  $\mu$ L) and treated with 10% Pd/C catalyst (250 mg), and the suspension was stirred at rt for 1 night, under a hydrogen atmosphere. TLC (iPrOH/H<sub>2</sub>O/NH<sub>3</sub>, 7:1:2) showed the conversion of **12** into a major polar product. The suspension was filtered on Acrodisc LC 25 mm, and the filtrate was concentrated. HPLC purification of the residue (0.01 M aq TFA/CH<sub>3</sub>CN, 100:0  $\rightarrow$  60:40 over 20 min, 5 mL/min, 215 nm, C-18 Kromasil column), followed by freeze-drying, gave target **1** (83 mg, 77%) as a white powder. Trisaccharide **1** had *R<sub>f</sub>* = 0.5 (iPrOH/H<sub>2</sub>O/NH<sub>3</sub>, 4:1:2); <sup>1</sup>H NMR (D<sub>2</sub>O)  $\delta$  5.09 (d, 1H, *J*<sub>1,2</sub> = 3.8 Hz, H-1<sub>E</sub>), 4.98 (d, 1H, *J*<sub>1,2</sub> = 1.7 Hz, H-1<sub>A</sub>), 4.88 (d, 1H, *J*<sub>1,2</sub> = 1.2 Hz, H-1<sub>B</sub>), 4.25 (dd, 1H, *J*<sub>2,3</sub> = 2.7 Hz, H-2<sub>A</sub>), 3.94 (ddd, 1H, *J*<sub>5,6a</sub> = 2.5 Hz, *J*<sub>5,6b</sub> = 4.1 Hz, *J*<sub>4,5</sub> = 10.1 Hz, H-5<sub>E</sub>), 3.90 (dd, 1H, *J*<sub>2,3</sub> = 3.3 Hz, H-2<sub>B</sub>), 3.94 (dd<sub>po</sub>, 1H, *J*<sub>3,4</sub> = 9.6 Hz, H-3<sub>A</sub>), 3.83 (dd, 1H, *J*<sub>3,4</sub> = 9.6 Hz, H-3<sub>B</sub>), 3.79–3.72 (m, 4H, H-3<sub>E</sub>, H-6<sub>A</sub>E, H-6<sub>B</sub>E, H-5<sub>A</sub>), 3.69 (dq, 1H, *J*<sub>4,5</sub> = 9.6 Hz, H-5<sub>B</sub>), 3.63 (dt, 1H, *J* = 7.0 Hz, *J* = 9.8 Hz, H<sub>Pr</sub>), 3.56 (dd<sub>po</sub>, 1H, *J*<sub>2,3</sub> = 9.9 Hz, H-2<sub>E</sub>), 3.54 (pt<sub>po</sub>, *J*<sub>4,5</sub> = 9.5 Hz, H-4<sub>A</sub>), 3.52 (dt, 1H, *J* = 6.3 Hz, H<sub>Pr</sub>), 3.44 (pt, 2H, H-4<sub>E</sub>, H-4<sub>B</sub>), 1.64–1.57 (m, 2H, CH<sub>2</sub>), 1.27 (d, 6H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>A</sub>, H-6<sub>B</sub>), 0.90 (t, 3H, *J* = 7.4 Hz, CH<sub>3</sub>); <sup>13</sup>C NMR (D<sub>2</sub>O)  $\delta$  102.4 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 169.9 Hz), 99.0 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.5 Hz), 95.7 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 168.8 Hz), 79.5 (C-2<sub>B</sub>), 75.6 (C-3<sub>A</sub>), 73.3 (C-3<sub>E</sub>), 72.6 (C-4<sub>E</sub>\*), 72.1 (C-2<sub>E</sub>), 71.8 (C-5<sub>E</sub>), 70.7 (C-3<sub>B</sub>), 70.5 (C-4<sub>A</sub>), 70.2 (C<sub>Pr</sub>), 69.9 (C-4<sub>B</sub>\*), 69.7 (C-5<sub>A</sub>), 69.1 (C-5<sub>B</sub>), 67.1 (C-2<sub>A</sub>), 60.8 (C-6<sub>E</sub>), 22.3 (CH<sub>2</sub>), 17.2, 17.0 (2C, C-6<sub>A</sub>, C-6<sub>B</sub>), 10.3 (CH<sub>3</sub>). HRMS (ESI<sup>+</sup>) for C<sub>21</sub>H<sub>38</sub>NO<sub>14</sub> ([M + Na]<sup>+</sup>, 537.2159) found *m/z* 537.2147.

**Allyl (3,4,6-Tri-*O*-acetyl-2-deoxy-2-trichloroacetamido- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-(4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (15).** TfOH (4  $\mu$ L, 45  $\mu$ mol, 0.3 equiv) was added to a solution of acceptor **12** (163 mg, 143  $\mu$ mol) and trichloroacetimidate **14**<sup>26,27</sup> (110 mg, 185  $\mu$ mol, 1.3 equiv) in toluene (5 mL) containing 4 $\text{\AA}$  MS (300 mg), stirred at  $-40\text{ }^\circ\text{C}$ . The reaction mixture was stirred for 1.5 h at this temperature, then for 1 h at rt. More donor **14** (20 mg, 34  $\mu$ mol, 0.23 equiv) was added, and the reaction mixture was stirred at rt overnight. TLC (Chex/EtOAc, 7:3) showed the complete disappearance of **12** and the presence of a major more polar product. Et<sub>3</sub>N was added. The mixture was filtered and concentrated to dryness. Chromatography of the residue (Chex/EtOAc, 100:0  $\rightarrow$  70:30) gave the allyl glycoside **15** (196 mg, 87%). Tetrasaccharide **15** had *R<sub>f</sub>* = 0.3 (Chex/EtOAc, 7:3). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.42–7.05 (m, 36H, NH, CH<sub>Ph</sub>), 5.87 (m, 1H, CH=), 5.25 (m<sub>overlapped</sub>, 1H, =CH<sub>2</sub>), 5.22 (d<sub>overlapped</sub>, 1H, H-1<sub>E</sub>), 5.18 (m, 1H, *J*<sub>cis</sub> = 10.4 Hz, =CH<sub>2</sub>), 5.08 (bs, 1H, H-1<sub>A</sub>), 5.02 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.7 Hz, H-4<sub>D</sub>), 4.93 (d, 1H, *J* = 10.8 Hz, H<sub>Bn</sub>), 4.87 (d, 1H, *J*<sub>1,2</sub> = 8.5 Hz, H-1<sub>D</sub>), 4.82 (dd, *J*<sub>2,3</sub> = 10.4 Hz, H-3<sub>D</sub>), 4.76 (bs<sub>overlapped</sub>, 1H, H-1<sub>B</sub>), 4.79–4.54 (m, 11H, H<sub>Bn</sub>), 4.46 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.30 (d, 1H, *J* = 12.0



Hz, H<sub>Bn</sub>), 4.21 (ddd, 1H, 1H,  $J_{2,NH} = 8.7$  Hz, H-2<sub>D</sub>), 4.16–4.07 (m, 5H, H-2<sub>A</sub>, H-3<sub>A</sub>, H-3<sub>E</sub>, H<sub>All</sub>, H-5<sub>E</sub>), 3.95–3.87 (m, 5H, H<sub>All</sub>, H-2<sub>B</sub>, H-2<sub>E</sub>, H-6<sub>Ad</sub>, H-3<sub>B</sub>), 3.85–3.79 (m, 3H, H-4<sub>E</sub>, H-5<sub>A</sub>\*, H-6<sub>Bd</sub>), 3.71 (dq, 1H,  $J_{4,5} = 9.4$  Hz, H-5<sub>B</sub>\*), 3.47 (pt, 2H,  $J_{3,4} = J_{4,5} = 9.3$  Hz, H-4<sub>A</sub>, H-4<sub>B</sub>), 3.44–3.39 (m, 2H, H-6<sub>Ad</sub>, H-6<sub>Bd</sub>), 2.93 (ddd, 1H,  $J_{5,6a} = 2.7$  Hz,  $J_{5,6b} = 3.4$  Hz, H-5<sub>D</sub>), 2.03, 2.00, 1.95 (3s, 9H, H<sub>Ac</sub>), 1.37 (d, 3H,  $J_{5,6} = 6.3$  Hz, H-6<sub>A</sub>\*), 1.33 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>B</sub>\*); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 171.0, 170.8, 169.5 (3C, C<sub>Ac</sub>), 162.4 (C<sub>NtCA</sub>), 138.9–137.9 (C<sub>Ph</sub>), 134.2 (CH=), 128.7–127.7 (CH<sub>Ph</sub>), 117.5 (=CH<sub>2</sub>), 101.4 (C-1<sub>A</sub>,  $^1J_{CH} = 175.8$  Hz), 101.0 (C-1<sub>D</sub>,  $^1J_{CH} = 161.1$  Hz), 98.2 (C-1<sub>B</sub>,  $^1J_{CH} = 171.0$  Hz), 95.0 (C-1<sub>E</sub>,  $^1J_{CH} = 166.9$  Hz), 93.1 (CCl<sub>3</sub>), 83.8 (C-3<sub>E</sub>), 81.0, 80.2 (2C, C-4<sub>B</sub>, C-4<sub>A</sub>), 79.3, 79.9 (2C, C-2<sub>E</sub>, C-3<sub>B</sub>), 78.9 (C-4<sub>E</sub>), 76.5 (C<sub>Bn</sub>), 76.0 (C-2<sub>B</sub>), 75.7, 75.6, 75.3 (3C, C<sub>Bn</sub>), 75.0 (C-3<sub>A</sub>), 74.5 (C-2<sub>A</sub>), 74.2, 73.8 (2C, C<sub>Bn</sub>), 73.6 (C-3<sub>D</sub>), 72.2 (C-5<sub>D</sub>), 72.1 (C<sub>Bn</sub>), 70.3 (C-5<sub>E</sub>), 68.9, 68.2 (2C, C-5<sub>A</sub>, C-5<sub>B</sub>), 68.1 (2C, C-4<sub>D</sub>, C-6<sub>E</sub>), 68.0 (C<sub>All</sub>), 61.8 (C-6<sub>D</sub>), 56.1 (C-2<sub>D</sub>), 21.0, 20.9 (3C, C<sub>Ac</sub>), 18.4, 18.2 (2C, C-6<sub>B</sub>, C-6<sub>A</sub>); HRMS (ESI<sup>+</sup>) for C<sub>84</sub>H<sub>94</sub>Cl<sub>3</sub>NO<sub>22</sub> ([M + Na]<sup>+</sup>, 1598.5234) found *m/z* 1598.5170.

**Allyl (2-Deoxy-2-trichloroacetamido-β-D-glucopyranosyl)-(1→2)-[2,3,4,6-tetra-O-benzyl-α-D-glucopyranosyl-(1→3)]-(4-O-benzyl-α-L-rhamnopyranosyl)-(1→2)-3,4-di-O-benzyl-α-L-rhamnopyranoside (16).** Methanolic MeONa (0.5 M, 1.4 mmol, 0.75 mmol, 6 equiv) was added to a solution of triacetate **15** (185 mg, 117 μmol) in MeOH (5 mL), and the mixture was stirred at rt for 3 h. TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 95:5) showed the presence of a new more polar product. The mixture was neutralized by addition of Dowex X8-200 ion-exchange resin (H<sup>+</sup>) and filtered. Volatiles were evaporated. The resulting syrup was chromatographed (Chex/EtOAc, 100:0 → 70:30) to give triol **16** (135 mg, 79%) as a white foam. Tetrasaccharide **16** had *R<sub>f</sub>* = 0.58 (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 9:1). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.52 (d, 1H,  $J_{2,NH} = 6.0$  Hz, NH), 7.43–7.05 (m, 35H, CH<sub>Ph</sub>), 5.87 (m, 1H, CH=), 5.37 (bs, 1H, H-1<sub>A</sub>), 5.26 (m, 1H,  $J_{trans} = 17.3$  Hz, =CH<sub>2</sub>), 5.19 (m, 1H,  $J_{cis} = 10.4$  Hz, =CH<sub>2</sub>), 5.15 (d, 1H,  $J_{1,2} = 3.6$  Hz, H-1<sub>E</sub>), 5.10 (d, 1H,  $J = 11.1$  Hz, H<sub>Bn</sub>), 5.05 (d, 1H,  $J = 11.1$  Hz, H<sub>Bn</sub>), 5.00 (d, 1H,  $J = 12.2$  Hz, H<sub>Bn</sub>), 4.92 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.90 (d, 1H,  $J = 12.3$  Hz, H<sub>Bn</sub>), 4.79 (d, 1H,  $J = 10.9$  Hz, H<sub>Bn</sub>), 4.73–4.59 (m, 5H, H<sub>Bn</sub>), 4.76 (b<sub>overlapped</sub>, 1H, H-1<sub>B</sub>), 4.54 (d<sub>overlapped</sub>, 1H,  $J_{1,2} = 8.3$  Hz, H-1<sub>D</sub>), 4.53 (d<sub>overlapped</sub>, 1H,  $J = 10.2$  Hz, H<sub>Bn</sub>), 4.49 (d, 1H,  $J = 10.9$  Hz, H<sub>Bn</sub>), 4.35 (d, 1H,  $J = 11.9$  Hz, H<sub>Bn</sub>), 4.17 (dd, 1H,  $J_{1,2} = 1.9$  Hz, H-2<sub>B</sub>), 4.15–4.07 (m, 5H, H<sub>All</sub>, H-3<sub>E</sub>, H-2<sub>A</sub>, H-3<sub>A</sub>, H-5<sub>E</sub>), 3.93 (m<sub>po</sub>, 1H, H<sub>All</sub>), 3.91 (dd<sub>po</sub>, 1H,  $J_{2,3} = 9.2$  Hz,  $J_{3,4} = 9.5$  Hz, H-3<sub>B</sub>), 3.83 (dd<sub>po</sub>, 1H,  $J_{3,4} = 2.9$  Hz,  $J_{4,5} = 9.5$  Hz, H-4<sub>E</sub>), 3.81 (dd, 1H,  $J_{2,3} = 9.8$  Hz, H-2<sub>E</sub>), 3.76–3.67 (m, 3H, H-2<sub>D</sub>, H-5<sub>A</sub>, H-5<sub>B</sub>), 3.59 (dd, 1H,  $J_{6a,6} = 12.2$  Hz,  $J_{5,6a} = 2.9$  Hz, H-6<sub>Ad</sub>), 3.51–3.44 (m, 4H, H-4<sub>A</sub>, H-4<sub>B</sub>, H-6<sub>Ad</sub>, H-6<sub>Bd</sub>), 3.11 (dd, 1H, H-4<sub>D</sub>), 3.05 (m, 1H,  $J_{4,5} = 9.8$  Hz, H-5<sub>D</sub>), 2.88 (dd, 1H,  $J_{5,6} = 7.7$  Hz, H-6<sub>Bd</sub>), 2.37 (dd, 1H,  $J_{2,3} = 10.2$  Hz,  $J_{3,4} = 8.3$  Hz, H-3<sub>D</sub>), 1.44 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>A</sub>\*), 1.36 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>B</sub>\*); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 164.9 (C<sub>NtCA</sub>), 139.0–137.6 (C<sub>Ph</sub>), 137.6 (CH=), 129.5–128.1 (CH<sub>Ph</sub>), 111.5 (=CH<sub>2</sub>), 101.2 (C-1<sub>D</sub>), 100.4 (C-1<sub>A</sub>), 98.5 (C-1<sub>B</sub>), 94.6 (C-1<sub>E</sub>), 92.9 (CCl<sub>3</sub>), 83.6 (C-3<sub>E</sub>), 80.9 (C-3<sub>B</sub>), 80.8 (C-4<sub>B</sub>), 80.1 (C-4<sub>A</sub>), 79.5 (C-2<sub>E</sub>), 79.0 (C-4<sub>E</sub>), 76.7 (C<sub>Bn</sub>), 76.7 (C-3<sub>D</sub>), 76.0 (C-5<sub>D</sub>), 75.6, 75.4, 75.3 (4C, C<sub>Bn</sub>), 74.5 (C-2<sub>A</sub>), 74.3 (C-3<sub>A</sub>), 73.9, 73.3 (2C, C<sub>Bn</sub>), 73.0 (C-4<sub>D</sub>), 71.7 (C-2<sub>B</sub>), 70.4 (C-5<sub>E</sub>), 69.0, 68.7 (C-5<sub>A</sub>, C-5<sub>B</sub>), 68.1 (C-6<sub>E</sub>), 68.0 (C<sub>All</sub>), 62.8 (C-6<sub>D</sub>), 58.3 (C-2<sub>D</sub>), 18.3, 18.2 (2C, C-6<sub>B</sub>, C-6<sub>A</sub>). HRMS (ESI<sup>+</sup>) for C<sub>78</sub>H<sub>88</sub>Cl<sub>3</sub>NO<sub>19</sub> ([M + Na]<sup>+</sup>, 1472.4913) found *m/z* 1472.4921.

**Propyl (2-Acetamido-2-deoxy-β-D-glucopyranosyl)-(1→2)-[α-D-glucopyranosyl-(1→3)]-α-L-rhamnopyranosyl-(1→2)-α-L-rhamnopyranoside (2).** **Route a.** Triol **16** (108 mg, 75 μmol) was dissolved in EtOH (10 mL) containing 1 M aq HCl (110 μL) and treated with 10% Pd/C catalyst (108 mg), and the suspension was stirred at rt for 1 night, under a hydrogen atmosphere. TLC (iPrOH/H<sub>2</sub>O/NH<sub>3</sub>, 7:1:2) showed the conversion of **12** into a more polar product. The suspension was filtered on a pad of Celite, and the filtrate was concentrated under vacuum. The residue was dissolved in EtOH (10 mL) containing Et<sub>3</sub>N (100 μL) and treated with 10%

Pd/C catalyst (100 mg), and the suspension was stirred at rt for 1 night, under a hydrogen atmosphere. TLC (iPrOH/H<sub>2</sub>O/NH<sub>3</sub>, 7:1:2) showed reaction completion. The suspension was filtered on a pad of Celite, and the filtrate was concentrated under vacuum. RP-HPLC purification of the residue (0.01 M aq TFA/CH<sub>3</sub>CN, 100:0 → 60:40 over 20 min, 5 mL min<sup>-1</sup>, 215 nm, C-18 Kromasil column), followed by freeze-drying, gave target **2** (36 mg, 68%) as a white powder.

**Route b.** A 2% TfoH solution in toluene (100 μL, 23 μmol, 0.35 equiv) was added to a solution of acceptor **12** (82 mg, 66 μmol) and trichloroacetimidate **14** (82 mg, 135 μmol, 2.0 equiv) in toluene (2 mL) containing 4 Å MS (200 mg), stirred at –30 °C. The reaction mixture was stirred for 1 h at this temperature and then for 1 h at rt. More donor **14** (10 mg, 17 μmol, 0.25 equiv) was added, and the reaction mixture was stirred for 3 h while slowly coming back to rt. Et<sub>3</sub>N was added. The mixture was filtered and concentrated to dryness. Chromatography of the residue (Chex/EtOAc, 100:0 → 70:30) gave contaminated **15** (90 mg). The whole material was solubilized in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) and 0.5 M methanolic sodium methoxide was added (500 μL). The reaction mixture was stirred for 3 h at rt. TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 9:1) showed the presence of a more polar product that reacted with ninhydrin. The mixture was neutralized by addition of Dowex X8-200 ion-exchange resin (H<sup>+</sup>) and filtered. Volatiles were evaporated. The residue was taken in ethanol (2 mL), and acetic anhydride (1 mL) was added. After 2 h at rt, volatiles were evaporated and coevaporated with toluene. The crude material was chromatographed (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 95:5) to give the acetamidotriol **17** (54 mg, 56% from **12**). The latter was dissolved in EtOH (5 mL) containing 1 M aq HCl (110 μL) and treated with 10% Pd/C catalyst (50 mg), and the suspension was stirred at rt for 1 night, under a hydrogen atmosphere. TLC (iPrOH/H<sub>2</sub>O/NH<sub>3</sub>, 7:1:2) showed the conversion of **17** into a more polar product. The suspension was filtered on a pad of Celite, and the filtrate was concentrated under vacuum. The residue was purified as above to give target **2** as a white powder (19 mg, 40% from **12**). Tetrasaccharide **2** had *R<sub>f</sub>* = 0.32 (iPrOH/H<sub>2</sub>O/NH<sub>3</sub>, 4:1:2); HPLC (215 nm): *t<sub>R</sub>* = 14.2 min (Kromasil 5 μm C18 100 Å 4.6 × 250 mm analytical column, using a 0–35% linear gradient over 20 min of CH<sub>3</sub>CN in 0.01 M aq TFA at 1 mL min<sup>-1</sup> flow rate); <sup>1</sup>H NMR (D<sub>2</sub>O) δ 5.15 (d, 1H,  $J_{1,2} = 3.7$  Hz, H-1<sub>E</sub>), 5.07 (d, 1H,  $J_{1,2} = 1.8$  Hz, H-1<sub>A</sub>), 4.86 (d, 1H,  $J_{1,2} = 1.3$  Hz, H-1<sub>B</sub>), 4.79 (d, 1H,  $J_{1,2} = 8.5$  Hz, H-1<sub>D</sub>), 4.40 (dd, 1H,  $J_{2,3} = 2.3$  Hz, H-2<sub>A</sub>), 4.02 (ddd, 1H,  $J_{4,5} = 10.1$  Hz,  $J_{5,6a} = 2.5$  Hz,  $J_{5,6b} = 4.4$  Hz, H-5<sub>E</sub>), 3.92–3.85 (m, 3H, H-2<sub>B</sub>, H-6<sub>Ad</sub>, H-3<sub>A</sub>), 3.84–3.66 (m, 9H, H-3<sub>B</sub>, H-6<sub>Ad</sub>, H-3<sub>E</sub>, H-6<sub>Bd</sub>, H-6<sub>Bd</sub>, H-5<sub>A</sub>, H-2<sub>D</sub>, H-5<sub>B</sub>, H-2<sub>E</sub>), 3.61 (dt, 1H,  $J = 6.9$  Hz,  $J = 9.8$  Hz, H<sub>Pr</sub>), 3.50–3.39 (m, 6H, H-4<sub>E</sub>, H<sub>Pr</sub>, H-3<sub>D</sub>, H-4<sub>B</sub>, H-5<sub>D</sub>, H-4<sub>D</sub>), 3.34 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.7$  Hz, H-4<sub>A</sub>), 2.07 (s, 3H, H<sub>NAc</sub>), 1.61–1.54 (m, 2H, CH<sub>2</sub>), 1.26 (d, 3H,  $J_{5,6} = 6.3$  Hz, H-6<sub>A</sub>\*), 1.12 (d, 3H,  $J_{5,6} = 6.3$  Hz, H-6<sub>B</sub>\*), 0.80 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub>). <sup>13</sup>C NMR (D<sub>2</sub>O) δ 174.9 (C<sub>NAc</sub>), 102.3 (C-1<sub>D</sub>,  $^1J_{CH} = 157.0$  Hz), 101.7 (C-1<sub>B</sub>,  $^1J_{CH} = 173.2$  Hz), 98.6 (C-1<sub>A</sub>,  $^1J_{CH} = 170.6$  Hz), 95.0 (C-1<sub>E</sub>,  $^1J_{CH} = 167.0$  Hz), 79.5 (C-2<sub>B</sub>), 76.4, 74.6 (2C, C-4<sub>D</sub>, C-4<sub>B</sub>), 74.4 (C-2<sub>A</sub>), 74.1 (C-3<sub>A</sub>), 73.5 (C-3<sub>E</sub>), 72.6 (C-3<sub>D</sub>), 71.8 (C-5<sub>E</sub>), 71.7 (C-2<sub>E</sub>), 71.2 (C-4<sub>A</sub>), 70.4 (C-3<sub>B</sub>), 70.2 (C-5<sub>D</sub>), 70.0 (C<sub>Pr</sub>), 69.9 (2C, C-4<sub>E</sub>, C-5<sub>A</sub>), 69.1 (C-5<sub>B</sub>), 61.0 (C-6<sub>D</sub>), 60.8 (C-6<sub>E</sub>), 56.0 (C-2<sub>D</sub>), 23.0 (C<sub>NAc</sub>), 22.3 (CH<sub>2</sub>), 17.2, 17.0 (2C, C-6<sub>A</sub>, C-6<sub>B</sub>), 10.2 (CH<sub>3</sub>). HRMS (ESI<sup>+</sup>) for C<sub>29</sub>H<sub>51</sub>NO<sub>19</sub> ([M + H]<sup>+</sup>, 718.3134) found *m/z* 718.3198 ([M + Na]<sup>+</sup>, 740.2953) found *m/z* 740.3021.

**Allyl (3-O-Acetyl-2-deoxy-4,6-O-isopropylidene-2-trichloroacetamido-β-D-glucopyranosyl)-(1→2)-[2,3,4,6-tetra-O-benzyl-α-D-glucopyranosyl-(1→3)]-(4-O-benzyl-α-L-rhamnopyranosyl)-(1→2)-3,4-di-O-benzyl-α-L-rhamnopyranoside (19).** TMSOTf (69 μL, 380 μmol, 0.3 equiv) was added to a solution of acceptor **12** (1.5 g, 1.3 mmol) and trichloroacetimidate **18**<sup>27</sup> (1.2 g, 2.2 mmol 1.7 equiv) in toluene (30 mL) containing 4 Å MS (1.1 g), stirred at –40 °C. The reaction mixture was stirred for 1 h while slowly coming back to rt. TLC (Tol/EtOAc, 85:15) showed the complete disappearance of **12** and the presence of a major less polar product. Et<sub>3</sub>N (1 mL) was added and the mixture was filtered, and



concentrated to dryness. Chromatography of the residue (Tol/EtOAc, 95:5 → 1:1) gave **19** (1.9 g, 95%) as a white foam. Tetrasaccharide **19** had  $R_f = 0.45$  (Tol/EtOAc, 85:15);  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.54–7.09 (m, 36H, NH,  $\text{CH}_{\text{Ph}}$ ), 5.91 (m, 1H,  $\text{CH}=\text{}$ ), 5.30 (m, 1H,  $J_{\text{trans}} = 17.2$  Hz,  $=\text{CH}_2$ ), 5.23 (m, 1H,  $J_{\text{cis}} = 10.5$  Hz,  $=\text{CH}_2$ ), 5.23–5.19 (m, 3H, H-1<sub>E</sub>, H<sub>Bn</sub>), 5.15 (bs, 1H, H-1<sub>A</sub>), 5.11 (d, 1H,  $J = 10.9$  Hz, H<sub>Bn</sub>), 5.08 (d, 1H,  $J = 11.0$  Hz, H<sub>Bn</sub>), 4.96 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.88–4.84 (m, 2H,  $J_{1,2} = 8.5$  Hz,  $J_{2,3} = J_{3,4} = 9.8$  Hz, H-1<sub>D</sub>, H-3<sub>D</sub>), 4.83 (d, 1H,  $J = 11.0$  Hz, H<sub>Bn</sub>), 4.81 (bs, 1H, H-1<sub>B</sub>), 4.79 (d, 1H,  $J = 10.3$  Hz, H<sub>Bn</sub>), 4.74 (d, 1H,  $J = 11.8$  Hz, H<sub>Bn</sub>), 4.69 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.68 (d, 1H,  $J = 11.8$  Hz, H<sub>Bn</sub>), 4.52 (d, 1H,  $J = 11.0$  Hz, H<sub>Bn</sub>), 4.36 (d, 1H,  $J = 12.0$  Hz, H<sub>Bn</sub>), 4.25–4.12 (m, 6H, H-2<sub>D</sub>, H-3<sub>E</sub>, H-2<sub>A</sub>, H-3<sub>A</sub>, H<sub>All</sub>, H-5<sub>E</sub>), 4.03 (dd, 1H,  $J_{1,2} = 1.9$  Hz,  $J_{2,3} = 2.8$  Hz, H-2<sub>B</sub>), 4.01–3.92 (m, 3H, H<sub>All</sub>, H-2<sub>E</sub>, H-3<sub>B</sub>), 3.88–3.80 (m, 2H, H-4<sub>E</sub>, H-5<sub>A</sub>), 3.75 (dq, 1H,  $J_{4,5} = 9.4$  Hz, H-5<sub>B</sub>), 3.68 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.5$  Hz, H-4<sub>D</sub>), 3.57–3.46 (m, 5H, H-6<sub>AD</sub>, H-4<sub>A</sub>, H-4<sub>B</sub>, H-6<sub>AE</sub>, H-6<sub>BE</sub>), 3.43 (pt, 1H,  $J_{5,6a} = J_{6a,6b} = 10.4$  Hz, H-6<sub>BD</sub>), 2.83 (ddd, 1H,  $J_{5,6b} = 5.3$  Hz, H-5<sub>D</sub>), 2.11 (s, 3H, H<sub>Ac</sub>), 1.49 (s, 3H, H<sub>IPr</sub>), 1.44 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>A</sub>), 1.41 (s, 3H, H<sub>IPr</sub>), 1.36 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>B</sub>);  $^{13}\text{C NMR}$  ( $\text{CDCl}_3$ )  $\delta$  170.8 (C<sub>Ac</sub>), 162.2 (C<sub>NtCA</sub>), 138.6–137.6 (C<sub>Ph</sub>), 133.9 (CH=), 129.1–127.4 (CH<sub>Ph</sub>), 117.2 (=CH<sub>2</sub>), 101.4 (C-1<sub>D</sub>),  $J_{\text{CH}} = 164.1$  Hz), 101.0 (C-1<sub>A</sub>),  $J_{\text{CH}} = 172.5$  Hz), 99.7 (C<sub>IPr</sub>), 97.9 (C-1<sub>B</sub>),  $J_{\text{CH}} = 167.9$  Hz), 94.6 (C-1<sub>E</sub>),  $J_{\text{CH}} = 167.1$  Hz), 92.8 (CCl<sub>3</sub>), 83.5 (C-3<sub>E</sub>), 80.5 (C-4<sub>B</sub>), 79.8 (C-4<sub>A</sub>), 79.6 (C-3<sub>B</sub>), 78.6 (C-4<sub>E</sub>), 78.2 (C-2<sub>E</sub>), 76.2, 75.3, 75.2, 75.0 (4C, C<sub>Bn</sub>), 74.9 (C-2<sub>B</sub>), 74.4 (C-3<sub>A</sub>), 74.0 (C<sub>Bn</sub>), 73.9 (C-2<sub>A</sub>), 73.5 (C<sub>Bn</sub>), 72.6 (C-3<sub>D</sub>), 72.2 (C<sub>Bn</sub>), 71.3 (C-4<sub>D</sub>), 70.0 (C-5<sub>E</sub>), 68.7 (C-5<sub>A</sub>), 68.0 (C-5<sub>B</sub>), 67.9 (C-6<sub>E</sub>), 67.7 (C<sub>All</sub>), 67.4 (C-5<sub>D</sub>), 61.8 (C-6<sub>D</sub>), 56.6 (C-2<sub>D</sub>), 29.0 (C<sub>IPr</sub>), 20.9 (C<sub>Ac</sub>), 19.1 (C<sub>IPr</sub>), 18.0 (C-6<sub>B</sub>), 17.9 (C-6<sub>A</sub>); HRMS (ESI<sup>+</sup>) for C<sub>83</sub>H<sub>94</sub>Cl<sub>3</sub>NO<sub>20</sub> ([M + Na]<sup>+</sup>, 1552.5332) found  $m/z$  1552.5430, ([M + NH<sub>4</sub>]<sup>+</sup>, 1547.5779) found  $m/z$  1547.5880

**Allyl (2-Acetamido-2-deoxy-4,6-O-isopropylidene-β-D-glucopyranosyl)-(1→2)-[2,3,4,6-tetra-O-benzyl-α-D-glucopyranosyl-(1→3)]-(4-O-benzyl-α-L-rhamnopyranosyl)-(1→2)-3,4-di-O-benzyl-α-L-rhamnopyranoside (21).** Methanolic MeONa (0.5 mL, 5.7 mmol, 6.9 equiv) was added to a solution of fully protected tetrasaccharide **19** (630 mg, 412 μmol) in CH<sub>2</sub>Cl<sub>2</sub> (45 mL), and the mixture was stirred for 6.5 h. TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 98:2 and Tol/EtOAc, 85:15) showed the complete disappearance of the monoacetate and the presence of a single more polar product which reacted with ninhydrin, thus corresponding to intermediate **24** (MS (ESI<sup>+</sup>) of C<sub>79</sub>H<sub>93</sub>NO<sub>18</sub> ([M + H]<sup>+</sup>, 1344.6) found  $m/z$  1344.7). MeOH (91 mL) and acetic anhydride (400 μL) were added. After 2 h at rt, TLC (Tol/EtOAc, 85:15) showed the complete disappearance of **24**. Evaporation of the filtrate gave a syrup which was purified by chromatography (Tol/EtOAc, 6:4 → 4:6) to give compound **21** (470 mg, 83%) as a white foam. Tetrasaccharide **21** had  $R_f = 0.35$  (Tol/EtOAc, 6:4);  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.43–7.07 (m, 36H, NH,  $\text{CH}_{\text{Ph}}$ ), 5.89 (m, 1H,  $\text{CH}=\text{}$ ), 5.27 (m, 1H,  $J_{\text{trans}} = 17.2$  Hz,  $=\text{CH}_2$ ), 5.20 (m, 1H,  $J_{\text{cis}} = 11.7$  Hz,  $=\text{CH}_2$ ), 5.18 (d, 1H,  $J_{1,2} = 3.7$  Hz, H-1<sub>E</sub>), 5.12 (bs<sub>overlapped</sub>, 1H, H-1<sub>A</sub>), 5.11 (d<sub>po</sub>, 1H, H<sub>Bn</sub>), 5.01 (d, 1H,  $J = 11.8$  Hz, H<sub>Bn</sub>), 4.95 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.91 (d, 1H,  $J = 11.0$  Hz, H<sub>Bn</sub>), 4.89 (d, 1H,  $J = 11.7$  Hz, H<sub>Bn</sub>), 4.82 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.80 (d, 1H,  $J = 9.9$  Hz, H<sub>Bn</sub>), 4.77 (d, 1H,  $J_{1,2} = 1.6$  Hz, H-1<sub>B</sub>), 4.71 (d, 1H,  $J = 11.9$  Hz, H<sub>Bn</sub>), 4.66–4.63 (m, 3H, H<sub>Bn</sub>), 4.60 (d, 1H,  $J = 10.1$  Hz, H<sub>Bn</sub>), 4.55 (d, 1H,  $J = 10.9$  Hz, H<sub>Bn</sub>), 4.40–4.35 (m, 2H, H<sub>Bn</sub>, H-1<sub>D</sub>), 4.19–4.11 (m, 4H, H-3<sub>A</sub>, H-5<sub>E</sub>, H<sub>All</sub>, H-3<sub>E</sub>), 4.06 (dd, 1H,  $J_{1,2} = 2.1$  Hz,  $J_{2,3} = 2.4$  Hz, H-2<sub>A</sub>), 4.02 (dd, 1H,  $J_{2,3} = 2.9$  Hz, H-2<sub>B</sub>), 3.95–3.79 (m, 6H, H<sub>All</sub>, H-4<sub>E</sub>, H-3<sub>B</sub>, H-2<sub>E</sub>, H-2<sub>D</sub>, H-5<sub>A</sub>), 3.73 (dq, 1H,  $J_{4,5} = 9.4$  Hz, H-5<sub>B</sub>), 3.59–3.46 (m, 7H, H-6<sub>AD</sub>, H-6<sub>BD</sub>, H-6<sub>AE</sub>, H-6<sub>BE</sub>, H-4<sub>D</sub>, H-4<sub>A</sub>, H-4<sub>B</sub>), 2.83 (ddd, 1H,  $J_{4,5} = 9.9$  Hz,  $J_{5,6a} = 5.7$  Hz,  $J_{5,6b} = 9.8$  Hz, H-5<sub>D</sub>), 2.78 (pt, 1H,  $J_{2,3} = J_{3,4} = 9.2$  Hz, H-3<sub>D</sub>), 2.40 (s, 3H, H<sub>NAC</sub>), 1.50 (s, 3H, H<sub>IPr</sub>), 1.48 (s, 3H, H<sub>IPr</sub>), 1.40 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>A</sub>), 1.35 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>B</sub>);  $^{13}\text{C NMR}$  ( $\text{CDCl}_3$ )  $\delta$  172.9 (C<sub>NAC</sub>), 138.5–137.1 (C<sub>Ph</sub>), 133.8 (CH=), 129.2–127.4 (CH<sub>Ph</sub>), 117.2 (=CH<sub>2</sub>), 102.4 (C-1<sub>D</sub>),  $J_{\text{CH}} = 162.8$  Hz), 101.0 (C-1<sub>A</sub>),  $J_{\text{CH}} = 177.5$  Hz), 99.6 (C<sub>IPr</sub>), 97.8 (C-1<sub>B</sub>),  $J_{\text{CH}} = 169.0$  Hz), 94.1 (C-1<sub>E</sub>),  $J_{\text{CH}} = 167.0$  Hz), 83.3 (C-3<sub>E</sub>), 80.5 (C-4<sub>B</sub>), 80.0 (C-4<sub>A</sub>), 79.4 (C-2<sub>E</sub>), 79.2 (C-3<sub>B</sub>), 78.7

(C-4<sub>E</sub>), 76.3 (C<sub>Bn</sub>), 75.9 (C-2<sub>A</sub>), 75.7 (C<sub>Bn</sub>), 75.5 (C-2<sub>B</sub>), 75.4 75.3, 75.0 (3C, C<sub>Bn</sub>), 74.2 (C-3<sub>A</sub>), 74.1 (C-4<sub>D</sub>), 74.0 (C-3<sub>D</sub>), 73.5, 72.0 (2C, C<sub>Bn</sub>), 70.1 (C-5<sub>E</sub>), 68.8 (C-5<sub>A</sub>), 67.9 (2C, C-5<sub>B</sub>, C-6<sub>E</sub>), 67.7 (C<sub>All</sub>), 66.9 (C-5<sub>D</sub>), 61.7 (C-6<sub>D</sub>), 58.5 (C-2<sub>D</sub>), 29.1 (C<sub>IPr</sub>), 23.2 (C<sub>NAC</sub>), 19.0 (C<sub>IPr</sub>), 18.1 (C-6<sub>B</sub>), 17.7 (C-6<sub>A</sub>); HRMS (ESI<sup>+</sup>) of C<sub>81</sub>H<sub>95</sub>NO<sub>19</sub> ([M + H]<sup>+</sup>, 1386.6577) found  $m/z$  1386.6625, ([M + Na]<sup>+</sup>, 1408.6396) found  $m/z$  1408.6478.

**Allyl (2-O-Acetyl-3,4-di-O-benzyl-α-L-rhamnopyranosyl)-(1→3)-(2-acetamido-2-deoxy-4,6-O-isopropylidene-β-D-glucopyranosyl)-(1→2)-[2,3,4,6-tetra-O-benzyl-α-D-glucopyranosyl-(1→3)]-(4-O-benzyl-α-L-rhamnopyranosyl)-(1→2)-3,4-di-O-benzyl-α-L-rhamnopyranoside (25).** TFOH (2 μL, 23 μmol, 0.3 equiv) was added to a solution of acceptor **21** (128 mg, 94 μmol) and trichloroacetimidate **22**<sup>33</sup> (97 mg, 188 μmol, 2 equiv) in toluene (1.5 mL) containing 4 Å MS (200 mg), stirred at –40 °C. The reaction mixture was stirred for 1 h at this temperature, then for 2 h at rt. TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 97.5:2.5) showed the presence of a major less polar product together with traces of **21**. Et<sub>3</sub>N (200 μL) was added. The mixture was filtered, and concentrated to dryness. Chromatography of the residue (Chex/EtOAc, 100:0 → 50:50) gave the allyl glycoside **25** (116 mg, 71%) as a white solid. Pentasaccharide **25** had  $R_f = 0.58$  (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 98.5:2.5);  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  7.44–7.06 (m, 45H,  $\text{CH}_{\text{Ph}}$ ), 6.40 (d, 1H,  $J_{1,2} = 8.9$  Hz, NH), 5.85 (m, 1H,  $\text{CH}=\text{}$ ), 5.24 (d, 1H,  $J_{\text{trans}} = 17.2$  Hz,  $=\text{CH}_2$ ), 5.17 (m, 2H, H-1<sub>E</sub>,  $=\text{CH}_2$ ), 5.13 (dd, 1H,  $J_{1,2} = 2.2$  Hz,  $J_{2,3} = 3.0$  Hz, H-2<sub>C</sub>), 5.07–5.04 (m, 2H, H<sub>Bn</sub>, H-1<sub>A</sub>), 5.01–4.97 (m, 5H, H<sub>Bn</sub>), 4.81 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.76 (d, 1H,  $J = 10.1$  Hz, H<sub>Bn</sub>), 4.72 (bs, 1H, H-1<sub>B</sub>), 4.69–4.55 (m, 9H, H-1<sub>C</sub>, H<sub>Bn</sub>), 4.48 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.33 (d, 1H,  $J = 12.0$  Hz, H<sub>Bn</sub>), 4.27 (d, 1H,  $J_{1,2} = 8.6$  Hz, H-1<sub>D</sub>), 4.19–4.08 (m, 5H, H-3<sub>E</sub>, H-5<sub>E</sub>, H-2<sub>D</sub>, H-3<sub>A</sub>, H<sub>All</sub>), 3.98–3.87 (m, 8H, H-2<sub>A</sub>, H-3<sub>C</sub>, H-5<sub>C</sub>, H-2<sub>B</sub>, H<sub>All</sub>, H-3<sub>B</sub>, H-2<sub>E</sub>, H-4<sub>E</sub>), 3.75 (dq, 1H,  $J_{4,5} = 9.3$  Hz,  $J_{5,6} = 6.6$  Hz, H-5<sub>A</sub>), 3.69 (dq, 1H,  $J_{4,5} = 9.4$  Hz,  $J_{5,6} = 6.1$  Hz, H-5<sub>B</sub>), 3.52–3.41 (m, 8H, H-6<sub>AD</sub>, H-6<sub>BD</sub>, H-6<sub>AE</sub>, H-6<sub>BE</sub>, H-4<sub>A</sub>, H-4<sub>B</sub>, H-4<sub>D</sub>, H-4<sub>C</sub>), 2.74 (dt, 1H,  $J_{5,6a} = 5.3$  Hz,  $J_{4,5} = J_{5,6b} = 9.8$  Hz, H-5<sub>D</sub>), 2.62 (pt, 1H,  $J_{2,3} = J_{3,4} = 9.4$  Hz, H-3<sub>D</sub>), 2.28 (s, 3H, H<sub>Ac</sub>), 2.08 (s, 3H, H<sub>NAC</sub>), 1.39 (s, 6H, H<sub>IPr</sub>), 1.33–1.30 (m, 6H, H-6<sub>A</sub>, H-6<sub>B</sub>), 1.24 (d, 3H,  $J_{5,6} = 6.3$  Hz, H-6<sub>C</sub>);  $^{13}\text{C NMR}$  ( $\text{CDCl}_3$ )  $\delta$  170.9 (C<sub>Ac</sub>), 170.5 (C<sub>NAC</sub>), 139.2–18.7 (C<sub>Ph</sub>), 134.2 (CH=), 129.7–127.8 (CH<sub>Ph</sub>), 117.6 (=CH<sub>2</sub>), 103.8 (C-1<sub>D</sub>), 101.5 (C-1<sub>A</sub>),  $J_{\text{CH}} = 172.8$  Hz), 99.6 (C<sub>IPr</sub>), 98.8 (C-1<sub>B</sub>),  $J_{\text{CH}} = 172.9$  Hz), 98.2 (C-1<sub>C</sub>), 94.4 (C-1<sub>E</sub>), 83.7 (C-3<sub>E</sub>), 80.8–78.9 (7C, C-3<sub>D</sub>, C-4<sub>B</sub>, C-2<sub>E</sub>, C-4<sub>C</sub>, C-4<sub>A</sub>, C-3<sub>B</sub>, C-4<sub>E</sub>), 78.1 (C-3<sub>C</sub>), 76.6 (C<sub>Bn</sub>), 76.4 (C-2<sub>A</sub>), 75.6 (C-2<sub>B</sub>), 76.0–75.4 (5C, C<sub>Bn</sub>), 74.6 (C-3<sub>A</sub>), 73.9 (C<sub>Bn</sub>), 72.4 (C-4<sub>D</sub>), 72.3, 71.8 (2C, C<sub>Bn</sub>), 70.3 (2C, C-2<sub>C</sub>, C-5<sub>E</sub>), 69.2 (C-5<sub>A</sub>), 68.2 (C-5<sub>B</sub>), 68.0 (2C, C-6<sub>E</sub>, C<sub>All</sub>), 67.8 (C-5<sub>C</sub>), 67.5 (C-5<sub>D</sub>), 62.3 (C-6<sub>D</sub>), 55.7 (C-2<sub>D</sub>), 29.5 (C<sub>IPr</sub>), 24.4 (C<sub>NAC</sub>), 21.5 (C<sub>Ac</sub>), 19.5 (C<sub>IPr</sub>), 18.4, 18.3, 18.0 (3C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>C</sub>); HRMS (ESI<sup>+</sup>) for C<sub>103</sub>H<sub>119</sub>NO<sub>24</sub> ([M + H]<sup>+</sup>, 1754.8201) found  $m/z$  1755.8307.

**Propyl (2-O-Acetyl-α-L-rhamnopyranosyl)-(1→3)-(2-acetamido-2-deoxy-β-D-glucopyranosyl)-(1→2)-[α-D-glucopyranosyl-(1→3)]-α-L-rhamnopyranosyl-(1→2)-α-L-rhamnopyranoside (3).** TFA (2 mL 50% aq) was added, at 0 °C, to a solution of pentasaccharide **25** (105 mg, 84 μmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) at 0 °C, and the biphasic mixture was stirred vigorously at this temperature for 1 h. TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 95:5) showed the complete disappearance of **25** and the presence of a major more polar product. Repeated coevaporation with toluene provided crude diol **26** (103 mg). Checking the  $^1\text{H NMR}$  spectrum ensured the complete disappearance of the isopropylidene signals. The crude material was directly engaged in the next step. Crude diol **26** (102 mg, 60 μmol) was dissolved in EtOH (4 mL), treated with 10% Pd/C catalyst (100 mg), and the suspension was stirred at rt for 6 h, under a hydrogen atmosphere. TLC (*i*PrOH/H<sub>2</sub>O/NH<sub>3</sub>, 7:1:2) showed that the starting material had been transformed into a major polar product. The suspension was filtered on a pad of Celite, and the filtrate was concentrated. HPLC purification ( $t_R = 14.6$  min, C-18 Kromasil column, 0.01 M aq TFA/CH<sub>3</sub>CN, 100:0 → 70:30 over 20 min, 5.5 mL min<sup>-1</sup>, 215 nm) of the residue, followed by freeze-drying, gave the target pentasaccharide **3** (36 mg, 67%) as a white

foam. Pentasaccharide **3** was isolated as a mixture of regioisomers resulting from the migration of the acetyl group. Pentasaccharide **3** had  $R_f = 0.5$  (iPrOH/H<sub>2</sub>O/NH<sub>3</sub>, 7:1:2); HPLC (215 nm):  $t_R = 14.7$  min (52.3%), 15.4 min (29.4%), 15.9 min (16.9%) (Kromasil 5  $\mu$ m C-18 100Å 4.6  $\times$  250 mm analytical column, using a 0–35% linear gradient over 20 min of CH<sub>3</sub>CN in 0.01 M aq TFA at 1 mL min<sup>-1</sup> flow rate); <sup>1</sup>H NMR (D<sub>2</sub>O, selected signals)  $\delta$  5.18 (d, 1H,  $J_{1,2} = 3.5$  Hz, H-1<sub>E</sub>), 5.08 (bs, 1H, H-1<sub>A</sub>), 4.93 (dd, 0.3H,  $J_{2,3} = 3.3$  Hz,  $J_{3,4} = 10.0$  Hz, H-3<sub>C</sub>), 4.90–4.87 (m, 3H, H-1<sub>C</sub>, H-2<sub>C</sub>, H-1<sub>B</sub>), 4.84–4.79 (m, 1H,  $J_{1,2} = 8.7$  Hz, H-1<sub>D</sub>), 4.42 (bs, 1H, H-2<sub>A</sub>), 4.15–4.00 (m, 2H, H-5<sub>C</sub>, H-5<sub>E</sub>), 3.94–3.90 (m, 4H, H-3<sub>C</sub>, H-3<sub>A</sub>, H-2<sub>B</sub>, H-6<sub>aD</sub>), 3.87–3.58 (m, 11H, H-2<sub>D</sub>, H-3<sub>E</sub>, H-3<sub>B</sub>, H-6<sub>aE</sub>, H-6<sub>bE</sub>, H-5<sub>A</sub>, H-6<sub>bD</sub>, H-5<sub>B</sub>, H-2<sub>E</sub>, H-4<sub>C</sub>, H<sub>Pr</sub>), 3.53–3.38 (m, 7H, H-4<sub>D</sub>, H-4<sub>E</sub>, H<sub>Pr</sub>, H-4<sub>C</sub>, H-4<sub>B</sub>, H-3<sub>D</sub>, H-5<sub>D</sub>), 3.34 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.7$  Hz, H-4<sub>A</sub>), 2.14, 2.13 (s, 3H, H<sub>NAC</sub>), 2.10, 2.07 (s, 3H, H<sub>Ac</sub>), 1.64–1.55 (m, 2H, CH<sub>2</sub>), 1.27 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>B</sub>), 1.25–1.20 (m, 5H, H-6<sub>A</sub>, H-6<sub>C</sub>), 1.10 (d, 1H,  $J_{5,6} = 6.2$  Hz, H-6<sub>C</sub>), 0.89 (t, 3H,  $J = 7.9$  Hz, CH<sub>3</sub>). <sup>13</sup>C NMR (D<sub>2</sub>O, selected signals)  $\delta$  173.3 (C<sub>NAC</sub>), 101.5 (C-1<sub>D</sub>), <sup>1</sup>J<sub>CH</sub> = 164.0 Hz), 101.2 (C-1<sub>A</sub>), <sup>1</sup>J<sub>CH</sub> = 172.8 Hz), 98.5, 98.2 (2C, C-1<sub>B</sub>), <sup>1</sup>J<sub>CH</sub> = 172.9 Hz, C-1<sub>C</sub>), <sup>1</sup>J<sub>CH</sub> = 169.8 Hz), 94.3 (C-1<sub>E</sub>), 82.3 (C-3<sub>D</sub>), 79.1 (C-2<sub>B</sub>), 76.1 (C-5<sub>D</sub>), 74.3 (C-2<sub>A</sub>), 73.6–73.0 (2C, C-3<sub>A</sub>, C-3<sub>E</sub>), 72.2 (2C, C-2<sub>C</sub>, C-4<sub>B</sub>), 71.5, 71.4 (2C, C-2<sub>E</sub>, C-5<sub>E</sub>), 70.8 (C-4<sub>A</sub>), 69.3 (C-3<sub>B</sub>), 69.8 (C<sub>Pr</sub>), 69.6, 69.5 (3C, C-4<sub>C</sub>, C-4<sub>E</sub>, C-5<sub>A</sub>), 68.9–68.3 (4C, C-3<sub>C</sub>, C-5<sub>B</sub>, C-5<sub>C</sub>, C-4<sub>D</sub>), 60.7 (C-6<sub>D</sub>), 60.4 (C-6<sub>E</sub>), 55.5 (C-2<sub>D</sub>), 22.8, 22.5 (C<sub>NAC</sub>), 21.9 (CH<sub>2</sub>), 20.4, 20.9 (C<sub>Ac</sub>), 16.8, 16.6, 16.5, 16.4 (3C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>C</sub>), 9.8 (CH<sub>3</sub>); HRMS (ESI<sup>+</sup>) for C<sub>37</sub>H<sub>63</sub>NO<sub>24</sub> ([M + Na]<sup>+</sup>, 928.3638) found  $m/z$  928.3660.

**Propyl  $\alpha$ -L-Rhamnopyranosyl-(1 $\rightarrow$ 3)-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[ $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\alpha$ -L-rhamnopyranoside (**4**).** A mixture of regioisomers **3** (10 mg, 60  $\mu$ mol) was dissolved in water (1 mL) and methanolic sodium methoxide (0.5 M, 100  $\mu$ L) was added. The reaction mixture was stirred for 3 h at rt. Following RP-HPLC control (C18 Kromasil column, 4.6  $\times$  150, CH<sub>3</sub>CN/0.01 M aq TFA 0  $\rightarrow$  30% over 20 min, 215 nm), the reaction mixture was purified by preparative RP-HPLC (C18 Kromasil column, 10  $\times$  250) using the same elution system. Freeze-drying gave the target pentasaccharide **4** (6 mg, 63%) as a white foam. Pentasaccharide **4** had HPLC (215 nm):  $t_R = 14.6$  min (Kromasil 5  $\mu$ m C-18 100Å 4.6  $\times$  250 mm analytical column, using a 0–40% linear gradient over 20 min of CH<sub>3</sub>CN in 0.01 M aq TFA at 1 mL min<sup>-1</sup> flow rate); <sup>1</sup>H NMR (D<sub>2</sub>O)  $\delta$  5.18 (d, 1H,  $J_{1,2} = 3.7$  Hz, H-1<sub>E</sub>), 5.08 (d, 1H,  $J_{1,2} = 1.7$  Hz, H-1<sub>A</sub>), 4.88 (d, 1H,  $J_{1,2} = 1.2$  Hz, H-1<sub>B</sub>), 4.85 (d, 1H,  $J_{1,2} = 1.4$  Hz, H-1<sub>C</sub>), 4.82 (d, 1H,  $J_{1,2} = 8.6$  Hz, H-1<sub>D</sub>), 4.25 (dd, 1H,  $J_{2,3} = 2.2$  Hz, H-2<sub>A</sub>), 4.03 (ddd, 1H,  $J_{4,5} = 10.1$  Hz,  $J_{5,6a} = 2.5$  Hz,  $J_{5,6b} = 4.4$  Hz, H-5<sub>E</sub>), 3.96 (dq, 1H,  $J_{4,5} = 9.7$  Hz, H-5<sub>C</sub>), 3.94–3.90 (m, 2H, H-3<sub>A</sub>, H-2<sub>B</sub>), 3.89 (dd, 1H,  $J_{6a,6b} = 12.2$  Hz,  $J_{5,6a} = 1.8$  Hz, H-6<sub>aD</sub>), 3.85–3.76 (m, 5H, H-3<sub>B</sub>, H-2<sub>D</sub>, H-3<sub>E</sub>, H-6<sub>aE</sub>, H-2<sub>C</sub>), 3.75–3.69 (m, 6H, H-6<sub>bE</sub>, H-5<sub>A</sub>, H-6<sub>bD</sub>, H-3<sub>C</sub>, H-2<sub>E</sub>, H-5<sub>B</sub>), 3.62 (dt, 1H,  $J = 7.0$  Hz,  $J = 9.8$  Hz, H<sub>Pr</sub>), 3.51–3.41 (m, 6H, H<sub>Pr</sub>, H-4<sub>D</sub>, H-4<sub>E</sub>, H-3<sub>D</sub>, H-4<sub>B</sub>, H-5<sub>D</sub>), 3.40 (pt, 1H,  $J_{3,4} = 9.7$  Hz, H-4<sub>C</sub>), 3.35 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.7$  Hz, H-4<sub>A</sub>), 2.10 (s, 3H, H<sub>NAC</sub>), 1.62–1.54 (m, 2H, CH<sub>2</sub>), 1.27 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>B</sub>), 1.24 (d, 3H,  $J_{5,6} = 6.3$  Hz, H-6<sub>A</sub>), 1.21 (d, 3H,  $J_{5,6} = 6.3$  Hz, H-6<sub>C</sub>), 0.89 (t, 3H,  $J = 7.4$  Hz, CH<sub>3</sub>). <sup>13</sup>C NMR (D<sub>2</sub>O)  $\delta$  174.2 (C<sub>NAC</sub>), 101.4 (C-1<sub>D</sub>), 101.3 (C-1<sub>C</sub>), 101.2 (C-1<sub>A</sub>), 98.2 (C-1<sub>B</sub>), <sup>1</sup>J<sub>CH</sub> = 171.8 Hz), 94.4 (C-1<sub>E</sub>), <sup>1</sup>J<sub>CH</sub> = 171.8 Hz), 81.4 (C-3<sub>D</sub>), 79.1 (C-2<sub>B</sub>), 76.1 (C-5<sub>D</sub>), 74.3 (C-2<sub>A</sub>), 73.5 (C-3<sub>A</sub>), 73.2 (C-3<sub>E</sub>), 72.2 (C-4<sub>B</sub>), 71.9 (C-4<sub>C</sub>), 71.5, 71.4 (2C, C-2<sub>E</sub>, C-5<sub>E</sub>), 70.8 (C-2<sub>C</sub>), 70.7 (C-4<sub>A</sub>), 70.2 (C-3<sub>C</sub>), 70.0 (C-3<sub>B</sub>), 69.8 (C<sub>Pr</sub>), 69.5 (2C, C-5<sub>A</sub>, C-4<sub>E</sub>), 69.0 (C-5<sub>C</sub>), 68.7 (C-5<sub>B</sub>), 68.4 (C-4<sub>D</sub>), 60.7 (C-6<sub>D</sub>), 60.4 (C-6<sub>E</sub>), 55.6 (C-2<sub>D</sub>), 22.5 (C<sub>NAC</sub>), 21.9 (CH<sub>2</sub>), 16.8 (C-6<sub>A</sub>), 16.6 (C-6<sub>B</sub>), 16.5 (C-6<sub>C</sub>), 9.8 (CH<sub>3</sub>); HRMS (ESI<sup>+</sup>) for C<sub>35</sub>H<sub>61</sub>NO<sub>23</sub> ([M + H]<sup>+</sup>, 864.3713) found  $m/z$  864.3704, ([M + Na]<sup>+</sup>, 886.3532) found  $m/z$  886.3489.

**Allyl (2-Deoxy-4,6-O-isopropylidene-3-O-levulinoyl-2-trichloroacetamido- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-O-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-4-O-benzyl- $\alpha$ -L-rhamnopyranoside (**28**).** DCC (532 mg, 2.6 mmol, 1.5 equiv) and levulinic acid

(317  $\mu$ L, 3.0 mmol, 1.8 equiv) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (15 mL) was added to a solution of alcohol **27**<sup>15</sup> (2.0 g, 1.7 mmol) and DMAP (420 mg, 3.4 mmol, 2 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (15 mL). The mixture was stirred for 1 h at rt. TLC (Tol/EtOAc, 7:3) showed the complete disappearance of the starting material and the presence of one less polar product. DCU was filtered and the reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, then washed with water, saturated aq NaHCO<sub>3</sub> (3  $\times$  20 mL), brine (3  $\times$  20 mL), and water (3  $\times$  20 mL). The organic layer was dried on Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated under reduced pressure. Chromatography of the residue (Tol/EtOAc, 85:15  $\rightarrow$  80:20) gave levulinate **28** (1.9 g, 90%) as a colorless syrup. Compound **28** had  $R_f = 0.5$  (Tol/EtOAc, 7:3); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.47–7.05 (m, 26H, NH, CH<sub>Ph</sub>), 5.94 (m, 1H, CH=), 5.29 (m, 1H,  $J_{trans} = 17.2$  Hz, =CH<sub>2</sub>), 5.21 (m, 1H,  $J_{cis} = 10.4$  Hz, =CH<sub>2</sub>), 5.15–5.09 (m, 3H, H-1<sub>E</sub>, H<sub>Bn</sub>), 5.07 (s, 2H, H<sub>Bn</sub>), 4.80–4.69 (m, 5H, H-1<sub>A</sub>, H<sub>Bn</sub>, H-3<sub>D</sub>, H<sub>Bn</sub>, H-1<sub>D</sub>), 4.59–4.54 (m, 2H, H<sub>Bn</sub>), 4.90 (d, 1H,  $J = 11.0$  Hz, H<sub>Bn</sub>), 4.32 (d, 1H,  $J = 12.0$  Hz, H<sub>Bn</sub>), 4.18–4.07 (m, 5H, H<sub>All</sub>, H-2<sub>D</sub>, H-3<sub>E</sub>, H-3<sub>A</sub>), 4.08 (ddd, 1H,  $J_{4,5} = 10.2$  Hz, H-5<sub>E</sub>), 3.99 (dd<sub>po</sub>, 1H,  $J_{1,2} = 2.3$  Hz, H-2<sub>A</sub>), 3.94 (m<sub>po</sub>, 1H, H<sub>All</sub>), 3.90 (dd, 1H,  $J_{1,2} = 3.6$  Hz,  $J_{2,3} = 9.7$  Hz, H-2<sub>E</sub>), 3.85 (dd, 1H,  $J_{5,6a} = 5.4$  Hz,  $J_{6a,6b} = 10.7$  Hz, H-6<sub>aD</sub>), 3.79 (dd, 1H,  $J_{3,4} = 9.0$  Hz, H-4<sub>E</sub>), 3.74–3.69 (m, 2H, H-6<sub>bD</sub>, H-5<sub>A</sub>), 3.67 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.6$  Hz, H-4<sub>D</sub>), 3.45 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.6$  Hz, H-4<sub>A</sub>), 3.45–3.39 (m, 2H, H-6<sub>aE</sub>, H-6<sub>bE</sub>), 2.76–2.63 (m, 5H, H-5<sub>D</sub>, 4H<sub>Lev</sub>), 2.20 (s, 3H, CH<sub>3Lev</sub>), 1.47 (s, 3H, H<sub>IPr</sub>), 1.42 (s, 3H, H<sub>IPr</sub>), 1.39 (d, 3H,  $J_{5,6} = 6.3$  Hz, H-6<sub>A</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  206.4 (C<sub>Lev</sub>), 172.6 (C<sub>Lev</sub>), 162.5 (C<sub>NTCA</sub>), 139.0–138.5 (C<sub>Ph</sub>), 134.2 (CH=), 129.6–127.7 (CH<sub>Ph</sub>), 117.6 (=CH<sub>2</sub>), 101.8 (C-1<sub>D</sub>), <sup>1</sup>J<sub>CH</sub> = 163.0 Hz), 100.1 (C<sub>IPr</sub>), 98.7 (C-1<sub>A</sub>), <sup>1</sup>J<sub>CH</sub> = 171.6 Hz), 95.1 (C-1<sub>E</sub>), <sup>1</sup>J<sub>CH</sub> = 165.2 Hz), 93.2 (CCl<sub>3</sub>), 83.8 (C-3<sub>E</sub>), 80.2 (C-4<sub>A</sub>), 79.3 (C-2<sub>E</sub>), 79.1 (C-4<sub>E</sub>), 76.4, 75.5, 75.3 (3C, C<sub>Bn</sub>), 75.0 (C-3<sub>A</sub>), 74.7 (C<sub>Bn</sub>), 74.6 (C-2<sub>A</sub>), 73.8 (C<sub>Bn</sub>), 73.3 (C-3<sub>D</sub>), 71.6 (C-4<sub>D</sub>), 70.3 (C-5<sub>E</sub>), 68.7 (C-5<sub>A</sub>), 68.3, 68.2 (2C, C<sub>All</sub>, C-6<sub>E</sub>), 67.5 (C-5<sub>D</sub>), 62.4 (C-6<sub>D</sub>), 56.9 (C-2<sub>D</sub>), 38.4 (CH<sub>2Lev</sub>), 30.2 (CH<sub>3Lev</sub>), 29.4 (C<sub>IPr</sub>), 28.5 (CH<sub>2Lev</sub>), 19.3 (C<sub>IPr</sub>), 18.3 (C-6<sub>A</sub>); HRMS (ESI<sup>+</sup>) for C<sub>66</sub>H<sub>76</sub>Cl<sub>3</sub>NO<sub>17</sub> ([M + Na]<sup>+</sup>, 1282.4076) found  $m/z$  1282.4189, ([M + NH<sub>4</sub>]<sup>+</sup>, 1277.4523) found  $m/z$  1277.4634.

**(2-Deoxy-4,6-O-isopropylidene-3-O-levulinoyl-2-trichloroacetamido- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-O-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-4-O-benzyl- $\alpha$ -L-rhamnopyranoside (**29**).** 1,5-Cyclooctadiene-bis(methyldiphenylphosphine)-iridium hexafluorophosphate (30 mg) was dissolved in THF (11 mL) and the resulting red solution was degassed under an argon stream. Hydrogen was bubbled through the solution, causing the color to change to yellow. The solution was then degassed again under an argon stream. A solution of allyl glycoside **28** (1.8 g, 1.5 mmol) in THF (15 mL) was added. The mixture was stirred overnight at rt. TLC (Tol/EtOAc, 8:2) showed the complete disappearance of **28** and the presence of a single less polar product. The mixture was treated with a solution of iodine (740 mg, 2.9 mmol) in THF/water (10 mL, 4:1 v/v) for 1 h at rt. TLC (Tol/EtOAc, 8:2 and CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 98:2) showed the complete disappearance of the intermediate and the presence of a single more polar product. Excess iodine was destroyed by adding a solution of freshly prepared 5% aq sodium bisulphite (4 mL). CH<sub>2</sub>Cl<sub>2</sub> (50 mL) was added, and the organic phase was washed with brine (3  $\times$  30 mL), water (3  $\times$  30 mL), dried on Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to dryness. Chromatography of the residue (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 99:1  $\rightarrow$  90:10) gave **29** (1.4 g, 80%) as a yellow syrup. Hemiacetal **29** had  $R_f = 0.3$  (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 98:2); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.45–7.05 (m, 26H, NH, CH<sub>Ph</sub>), 5.16 (dd, 1H,  $J_{1,2} = 2.2$  Hz, H-1<sub>A</sub>), 5.15–5.07 (m, 5H, H-1<sub>E</sub>, H<sub>Bn</sub>), 4.78–4.68 (m, 4H, H-3<sub>D</sub>, H<sub>Bn</sub>, H-1<sub>D</sub>, H<sub>Bn</sub>), 4.58–4.44 (m, 3H, H<sub>Bn</sub>), 4.30 (d, 1H,  $J = 12.0$  Hz, H<sub>Bn</sub>), 4.21 (dd, 1H,  $J_{2,3} = 2.9$  Hz,  $J_{3,4} = 9.6$  Hz, H-3<sub>A</sub>), 4.16–4.07 (m, 3H, H-2<sub>D</sub>, H-3<sub>E</sub>, H-5<sub>E</sub>), 4.01 (dd, 1H, H-2<sub>A</sub>), 3.95 (dq, 1H,  $J_{4,5} = 9.4$  Hz, H-5<sub>A</sub>), 3.90 (dd, 1H,  $J_{1,2} = 2.3$  Hz,  $J_{2,3} = 9.6$  Hz, H-2<sub>E</sub>), 3.85 (dd, 1H,  $J_{5,6a} = 5.1$  Hz,  $J_{6a,6b} = 10.4$  Hz, H-6<sub>aD</sub>), 3.76 (dd, 1H,  $J_{3,4} = 9.2$  Hz,  $J_{4,5} = 9.9$  Hz, H-4<sub>E</sub>), 3.70 (dd, 1H,  $J_{5,6b} = 1.6$  Hz, H-6<sub>bD</sub>), 3.66 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.5$  Hz, H-4<sub>D</sub>), 3.45 (pt, 1H, H-4<sub>A</sub>), 3.39 (m, 2H, H-6<sub>aE</sub>,



H-6b<sub>E</sub>), 3.30 (d, 1H,  $J_{1,OH} = 3.4$  Hz, OH-1<sub>A</sub>), 2.77–2.61 (m, 5H, H-5<sub>D</sub>, 4H<sub>Lev</sub>), 2.19 (s, 3H, CH<sub>3Lev</sub>), 1.47 (s, 3H, H<sub>IPr</sub>), 1.42 (s, 3H, H<sub>IPr</sub>), 1.38 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>A</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 206.2 (C<sub>Lev</sub>), 172.2 (C<sub>Lev</sub>), 162.1 (C<sub>NTCA</sub>), 138.6–137.6 (C<sub>Ph</sub>), 129.3–127.2 (CH<sub>Ph</sub>), 101.4 (C-1<sub>D</sub>), <sup>1</sup>J<sub>CH</sub> = 167.7 Hz), 99.7 (C<sub>IPr</sub>), 94.6 (C-1<sub>E</sub>, <sup>1</sup>J<sub>CH</sub> = 167.2 Hz), 93.9 (C-1<sub>A</sub>, <sup>1</sup>J<sub>CH</sub> = 171.7 Hz), 92.8 (CCl<sub>3</sub>), 83.4 (C-3<sub>E</sub>), 79.7 (C-4<sub>A</sub>), 78.8 (C-2<sub>E</sub>), 78.7 (C-4<sub>E</sub>), 76.2, 75.9, 75.1 (3C, C<sub>Bn</sub>), 74.5 (C-2<sub>A</sub>), 74.4 (C<sub>Bn</sub>), 74.1 (C-3<sub>A</sub>), 73.3 (C<sub>Bn</sub>), 72.8 (C-3<sub>D</sub>), 71.2 (C-4<sub>D</sub>), 69.9 (C-5<sub>E</sub>), 68.3 (C-5<sub>A</sub>), 67.8 (C-6<sub>E</sub>), 67.1 (C-5<sub>D</sub>), 62.0 (C-6<sub>D</sub>), 56.4 (C-2<sub>D</sub>), 38.0 (CH<sub>2Lev</sub>), 29.8 (CH<sub>3Lev</sub>), 28.9 (C<sub>IPr</sub>), 28.0 (CH<sub>2Lev</sub>), 18.9 (C<sub>IPr</sub>), 17.9 (C-6<sub>A</sub>); HRMS (ESI<sup>+</sup>) for C<sub>63</sub>H<sub>72</sub>Cl<sub>3</sub>NO<sub>17</sub> ([M + Na]<sup>+</sup>, 1242.3763) found *m/z* 1242.3856, ([M + NH<sub>4</sub>]<sup>+</sup>, 1237.4209) found *m/z* 1237.4213.

**(2-Deoxy-4,6-O-isopropylidene-3-O-levulinoyl-2-trichloroacetamido-β-D-glucopyranosyl)-(1→2)-[2,3,4,6-tetra-O-benzyl-α-D-glucopyranosyl-(1→3)]-4-O-benzyl-α-L-rhamnopyranose Trichloroacetimidate (30).** Hemiactal **29** (1.3 g, 1.1 mmol) was dissolved in DCE (10 mL), placed under argon, and cooled to –5 °C. Trichloroacetonitrile (525 μL, 5.2 mmol, 5 equiv) and DBU (44 μL, 308 μmol, 0.28 equiv) were added. The mixture was stirred at –5 °C for 10 min. TLC (Chex/EtOAc + Et<sub>3</sub>N, 7:3) showed the complete disappearance of **29** and the presence of a less polar product. The mixture was directly chromatographed (Chex/EtOAc + 5% Et<sub>3</sub>N, 7:3 → 1:1) to give **30** (1.2 g, 85%) as a yellow syrup. Trichloroacetimidate **30** (α anomer) had *R<sub>f</sub>* = 0.45 (Chex/EtOAc, 6:4); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.65 (s, 1H, NH), 7.45–7.06 (m, 26H, NH, CH<sub>Ph</sub>), 6.22 (s, 1H, H-1<sub>A</sub>), 5.11–5.01 (m, 5H, H-1<sub>E</sub>, 4H<sub>Bn</sub>), 4.82–4.75 (m, 3H, H-3<sub>D</sub>, H<sub>Bn</sub>), 4.73 (d, 1H,  $J_{1,2} = 8.4$  Hz, H-1<sub>D</sub>), 4.57–4.49 (m, 3H, H<sub>Bn</sub>), 4.33 (d, 1H,  $J = 12.1$  Hz, H<sub>Bn</sub>), 4.22–4.07 (m, 5H, H-2<sub>D</sub>, H-3<sub>A</sub>, H-2<sub>A</sub>, H-3<sub>E</sub>, H-5<sub>E</sub>), 3.92 (dq, 1H,  $J_{4,5} = 9.5$  Hz, H-5<sub>A</sub>), 3.90–3.85 (m, 2H, H-2<sub>E</sub>, H-6<sub>aD</sub>), 3.82–3.74 (m, 2H, H-4<sub>E</sub>, H-6<sub>bD</sub>), 3.71 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.6$  Hz, H-4<sub>D</sub>), 3.56 (pt, 1H,  $J_{3,4} = 9.6$  Hz, H-4<sub>A</sub>), 3.47 (dd, 1H,  $J_{5,6a} = 2.5$  Hz,  $J_{6a,6b} = 11.0$  Hz, H-6<sub>aE</sub>), 3.37 (bd, 1H, H-6<sub>bE</sub>), 2.85 (ddd, 1H,  $J_{5,6a} = 5.2$  Hz, H-5<sub>D</sub>), 2.77–2.58 (m, 4H, 2CH<sub>2Lev</sub>), 2.20 (s, 3H, CH<sub>3Lev</sub>), 1.49 (s, 3H, H<sub>IPr</sub>), 1.43 (s, 3H, H<sub>IPr</sub>), 1.42 (d, 3H,  $J_{5,6} = 6.0$  Hz, H-6<sub>A</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 206.0 (C<sub>Lev</sub>), 172.3 (C<sub>Lev</sub>), 162.1 (C<sub>NTCA</sub>), 160.2 (C=NH), 138.5–137.4 (C<sub>Ph</sub>), 129.2–127.4 (CH<sub>Ph</sub>), 101.3 (C-1<sub>D</sub>, <sup>1</sup>J<sub>CH</sub> = 159.2 Hz), 99.8 (C<sub>IPr</sub>), 96.9 (C-1<sub>A</sub>, <sup>1</sup>J<sub>CH</sub> = 182.6 Hz), 95.0 (C-1<sub>E</sub>, <sup>1</sup>J<sub>CH</sub> = 165.6 Hz), 92.7 (CCl<sub>3</sub>), 91.1 (CCl<sub>3</sub>), 83.3 (C-3<sub>E</sub>), 79.2 (C-4<sub>A</sub>), 78.7 (C-2<sub>E</sub>), 78.6 (C-4<sub>E</sub>), 76.3, 75.3, 75.0, 74.4 (4C, C<sub>Bn</sub>), 74.3 (C-3<sub>A</sub>), 73.3 (C<sub>Bn</sub>), 72.7 (C-3<sub>D</sub>), 72.2 (C-2<sub>A</sub>), 71.2 (C-4<sub>D</sub>), 71.1 (C-5<sub>A</sub>), 70.1 (C-5<sub>E</sub>), 67.9 (C-6<sub>E</sub>), 67.3 (C-5<sub>D</sub>), 61.9 (C-6<sub>D</sub>), 56.5 (C-2<sub>D</sub>), 38.0 (CH<sub>2Lev</sub>), 29.8 (CH<sub>3Lev</sub>), 29.0 (C<sub>IPr</sub>), 28.1 (CH<sub>2Lev</sub>), 18.9 (C<sub>IPr</sub>), 17.9 (C-6<sub>A</sub>).

**Allyl (2-Deoxy-4,6-O-isopropylidene-3-O-levulinoyl-2-trichloroacetamido-β-D-glucopyranosyl)-(1→2)-[2,3,4,6-tetra-O-benzyl-α-D-glucopyranosyl-(1→3)]-4-O-benzyl-α-L-rhamnopyranosyl-(1→2)-3,4-di-O-benzyl-α-L-rhamnopyranoside (31) and Allyl (2-Deoxy-4,6-O-isopropylidene-3-O-levulinoyl-2-trichloroacetamido-β-D-glucopyranosyl)-(1→2)-[2,3,4,6-tetra-O-benzyl-α-D-glucopyranosyl-(1→3)]-4-O-benzyl-β-L-rhamnopyranosyl-(1→2)-3,4-di-O-benzyl-α-L-rhamnopyranoside (32).** TMSOTf (10 μL, 56 μmol, 0.3 equiv) was added to a solution of acceptor **7**<sup>23</sup> (71 mg, 185 μmol) and trichloroacetimidate **30** (380 mg, 280 μmol, 1.5 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) containing 4Å MS (160 mg), stirred at –78 °C. The reaction mixture was stirred for 15 min while slowly coming back to rt. TLC (Tol/EtOAc, 7:3) showed the complete disappearance of the acceptor and the presence of two new compounds. Et<sub>3</sub>N (1 mL) was added and the mixture was filtered, and concentrated to dryness. Chromatography of the residue (Tol/EtOAc, 9:1 → 1:1) gave, by order of elution, first **31** (130 mg, 44%), and then the β anomer **32** (130 mg, 44%), both as colorless syrups. Tetrasaccharide **31** had *R<sub>f</sub>* = 0.65 (Tol/EtOAc, 7:3); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.49–7.07 (m, 36H, NH, CH<sub>Ph</sub>), 5.89 (m, 1H, CH=), 5.27 (m, 1H,  $J_{trans} = 17.2$  Hz, =CH<sub>2</sub>), 5.21 (m, 1H,  $J_{cis} = 10.3$  Hz, =CH<sub>2</sub>), 5.16 (d, 1H,  $J_{1,2} = 3.6$  Hz, H-1<sub>E</sub>), 5.12 (bs, 3H, H-1<sub>A</sub>, 2H<sub>Bn</sub>), 5.09 (d, 1H,  $J = 11.5$  Hz, H<sub>Bn</sub>), 5.07 (d, 1H,  $J = 11.4$  Hz, H<sub>Bn</sub>), 4.93 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.83–4.76 (m, 5H, H-3<sub>D</sub>,

H<sub>Bn</sub>, H-1<sub>D</sub>, H-1<sub>B</sub>, H<sub>Bn</sub>), 4.71 (d, 1H,  $J = 11.8$  Hz, H<sub>Bn</sub>), 4.67 (d, 1H,  $J = 10.8$  Hz, H<sub>Bn</sub>), 4.66 (d, 1H,  $J = 11.8$  Hz, H<sub>Bn</sub>), 4.60–4.54 (m, 3H, H<sub>Bn</sub>), 4.33 (d, 1H,  $J = 12.0$  Hz, H<sub>Bn</sub>), 4.20–4.09 (m, 6H, H-3<sub>E</sub>, H-2<sub>D</sub>, H-2<sub>A</sub>, H-3<sub>A</sub>, H<sub>All</sub>, H-5<sub>E</sub>), 4.00 (dd, 1H,  $J_{1,2} = 2.1$  Hz,  $J_{2,3} = 2.9$  Hz, H-2<sub>B</sub>), 3.98 (m, 1H, H<sub>All</sub>), 3.93–3.89 (m, 2H, H-3<sub>B</sub>, H-2<sub>E</sub>), 3.85–3.77 (m, 2H, H-4<sub>E</sub>, H-5<sub>B</sub>), 3.73 (dq, 1H,  $J_{4,5} = 9.4$  Hz, H-5<sub>A</sub>), 3.64 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.5$  Hz, H-4<sub>D</sub>), 3.54–3.42 (m, 5H, H-6<sub>aD</sub>, H-4<sub>A</sub>, H-4<sub>B</sub>, H-6<sub>aE</sub>, H-6<sub>bE</sub>), 3.39 (pt, 1H,  $J_{5,6b} = J_{6a,6b} = 10.4$  Hz, H-6<sub>bD</sub>), 2.80–2.72 (m, 3H, H-5<sub>D</sub>, 2H<sub>Lev</sub>), 2.68–2.62 (m, 2H, H<sub>Lev</sub>), 2.20 (s, 3H, CH<sub>3Lev</sub>), 1.44 (s, 3H, H<sub>IPr</sub>), 1.40 (d, 3H,  $J_{5,6} = 6.3$  Hz, H-6<sub>B</sub>), 1.39 (s, 3H, H<sub>IPr</sub>), 1.32 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6<sub>A</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 206.2 (C<sub>Lev</sub>), 172.3 (C<sub>Lev</sub>), 162.2 (C<sub>NTCA</sub>), 138.6–137.5 (C<sub>Ph</sub>), 133.8 (CH=), 129.2–127.3 (CH<sub>Ph</sub>), 117.2 (=CH<sub>2</sub>), 101.3 (C-1<sub>D</sub>, <sup>1</sup>J<sub>CH</sub> = 162.0 Hz), 100.9 (C-1<sub>A</sub>, <sup>1</sup>J<sub>CH</sub> = 171.3 Hz), 99.7 (C<sub>IPr</sub>), 97.9 (C-1<sub>B</sub>, <sup>1</sup>J<sub>CH</sub> = 170.2 Hz), 94.6 (C-1<sub>E</sub>, <sup>1</sup>J<sub>CH</sub> = 165.1 Hz), 92.8 (CCl<sub>3</sub>), 83.4 (C-3<sub>E</sub>), 80.4 (C-4<sub>B</sub>), 79.8 (C-4<sub>A</sub>), 79.5 (C-2<sub>E</sub>), 78.6 (C-4<sub>E</sub>), 78.4 (C-3<sub>B</sub>), 76.1, 75.3, 75.2, 74.9 (4C, C<sub>Bn</sub>), 74.8 (C-2<sub>B</sub>), 74.4 (C-3<sub>A</sub>), 74.1 (C<sub>Bn</sub>), 73.8 (C-2<sub>A</sub>), 73.5 (C<sub>Bn</sub>), 72.9 (C-3<sub>D</sub>), 72.1 (C<sub>Bn</sub>), 71.3 (C-4<sub>D</sub>), 69.9 (C-5<sub>E</sub>), 68.6 (C-5<sub>B</sub>), 68.0 (C-5<sub>A</sub>), 67.9 (C-6<sub>E</sub>), 67.7 (C<sub>All</sub>), 67.2 (C-5<sub>D</sub>), 61.7 (C-6<sub>D</sub>), 56.6 (C-2<sub>D</sub>), 38.0 (CH<sub>2Lev</sub>), 29.7 (CH<sub>3Lev</sub>), 29.0 (C<sub>IPr</sub>), 28.1 (CH<sub>2Lev</sub>), 19.0 (C<sub>IPr</sub>), 18.0 (C-6<sub>A</sub>), 17.9 (C-6<sub>B</sub>); HRMS (ESI<sup>+</sup>) for C<sub>86</sub>H<sub>98</sub>Cl<sub>3</sub>NO<sub>21</sub> ([M + Na]<sup>+</sup>, 1608.5594) found *m/z* 1608.5730, ([M + NH<sub>4</sub>]<sup>+</sup>, 1603.6041) found *m/z* 1603.6112.

Tetrasaccharide **32** had *R<sub>f</sub>* = 0.6 (Tol/EtOAc, 7:3); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.67 (d, 1H,  $J_{NH,2} = 8.9$  Hz, NH), 7.53–7.05 (m, 35H, CH<sub>Ph</sub>), 5.89 (m, 1H, CH=), 5.31 (m, 1H,  $J_{trans} = 17.2$  Hz, =CH<sub>2</sub>), 5.22 (m, 1H,  $J_{cis} = 10.3$  Hz, =CH<sub>2</sub>), 5.20 (s, 2H, H<sub>Bn</sub>), 5.11 (d, 1H,  $J_{1,2} = 3.5$  Hz, H-1<sub>E</sub>), 5.07 (d, 1H,  $J = 11.0$  Hz, H<sub>Bn</sub>), 5.03 (d, 1H,  $J = 12.2$  Hz, H<sub>Bn</sub>), 4.95–4.86 (m, 5H, H-1<sub>D</sub>, H-3<sub>D</sub>, 2H<sub>Bn</sub>, H-1<sub>B</sub>), 4.84–4.79 (m, 2H, H<sub>Bn</sub>), 4.59–4.48 (m, 5H, 3H<sub>Bn</sub>, H-1<sub>A</sub>, 2H<sub>Bn</sub>), 4.34 (d, 1H,  $J = 12.2$  Hz, H<sub>Bn</sub>), 4.24–4.11 (m, 6H, H-2<sub>D</sub>, H<sub>All</sub>, H-2<sub>B</sub>, H-2<sub>A</sub>, H-3<sub>E</sub>, H-5<sub>E</sub>), 4.00 (m, 1H, H<sub>All</sub>), 3.98–3.93 (m, 2H, H-3<sub>B</sub>, H-2<sub>E</sub>), 3.88 (dd, 1H,  $J_{5,6a} = 5.2$  Hz,  $J_{6a,6b} = 10.6$  Hz, H-6<sub>aD</sub>), 3.80 (dd, 1H,  $J_{2,3} = 2.5$  Hz,  $J_{3,4} = 9.5$  Hz, H-3<sub>A</sub>), 3.78–3.67 (m, 3H, H-5<sub>B</sub>, H-4<sub>E</sub>, H-6<sub>bD</sub>), 3.60–3.40 (m, 5H, H-4<sub>D</sub>, H-4<sub>A</sub>, H-4<sub>B</sub>, H-6<sub>aE</sub>, H-6<sub>bE</sub>), 3.28 (dq, 1H,  $J_{4,5} = 9.2$  Hz,  $J_{5,6} = 6.2$  Hz, H-5<sub>A</sub>), 2.83 (ddd, 1H,  $J_{5,6b} = 9.8$  Hz, H-5<sub>D</sub>), 2.78–2.75 (m, 2H, H<sub>Lev</sub>), 2.68–2.64 (m, 2H, H<sub>Lev</sub>), 2.18 (s, 3H, CH<sub>3Lev</sub>), 1.37 (s, 3H, H<sub>IPr</sub>), 1.39–1.37 (m, 9H, H<sub>IPr</sub>, H-6<sub>A</sub>, H-6<sub>B</sub>), 1.32 (s, 3H, H<sub>IPr</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 206.1 (C<sub>Lev</sub>), 172.2 (C<sub>Lev</sub>), 162.0 (C<sub>NTCA</sub>), 139.1–137.5 (C<sub>Ph</sub>), 134.0 (CH=), 129.1–127.3 (CH<sub>Ph</sub>), 117.2 (=CH<sub>2</sub>), 100.7 (C-1<sub>D</sub>, <sup>1</sup>J<sub>CH</sub> = 163.8 Hz), 99.7 (C<sub>IPr</sub>), 97.5 (C-1<sub>A</sub>, <sup>1</sup>J<sub>CH</sub> = 155.1 Hz), 96.9 (C-1<sub>B</sub>, <sup>1</sup>J<sub>CH</sub> = 167.2 Hz), 94.1 (C-1<sub>E</sub>, <sup>1</sup>J<sub>CH</sub> = 167.7 Hz), 92.8 (CCl<sub>3</sub>), 83.7 (C-3<sub>E</sub>), 81.0 (C-4<sub>B</sub>), 79.7 (C-4<sub>A</sub>), 78.7 (C-4<sub>E</sub>), 77.8 (C-2<sub>E</sub>), 77.4 (C-3<sub>B</sub>), 76.4 (C-3<sub>A</sub>), 76.2, 75.3, 75.1, 74.6, 73.6 (5C, C<sub>Bn</sub>), 73.3 (3C, C-2<sub>A</sub>, C-2<sub>B</sub>, C-3<sub>D</sub>), 73.5 (C<sub>Bn</sub>), 72.4 (C-5<sub>A</sub>), 71.3 (C-4<sub>D</sub>), 71.0 (C<sub>Bn</sub>), 69.9 (C-5<sub>E</sub>), 68.0 (2C, C-5<sub>B</sub>, C-6<sub>E</sub>), 67.8 (C<sub>All</sub>), 67.0 (C-5<sub>D</sub>), 62.4 (C-6<sub>D</sub>), 56.7 (C-2<sub>D</sub>), 38.0 (CH<sub>2Lev</sub>), 29.8 (CH<sub>3Lev</sub>), 29.0 (C<sub>IPr</sub>), 28.1 (CH<sub>2Lev</sub>), 19.9 (C<sub>IPr</sub>), 18.4 (C-6<sub>A</sub>), 18.0 (C-6<sub>B</sub>); HRMS (ESI<sup>+</sup>) for C<sub>86</sub>H<sub>98</sub>Cl<sub>3</sub>NO<sub>21</sub> ([M + Na]<sup>+</sup>, 1608.5594) found *m/z* 1608.5914, ([M + NH<sub>4</sub>]<sup>+</sup>, 1603.6041) found *m/z* 1603.6248.

**Allyl (2-Acetamido-2-deoxy-β-D-glucopyranosyl)-(1→2)-[2,3,4,6-tetra-O-benzyl-α-D-glucopyranosyl-(1→3)]-4-O-benzyl-α-L-rhamnopyranoside (36) and Allyl (2-Deoxy-2-methylcarbamate-α-D-glucopyranosyl)-(1→2)-[2,3,4,6-tetra-O-benzyl-α-D-glucopyranosyl-(1→3)]-4-O-benzyl-α-L-rhamnopyranoside (37).** Methanolic MeONa (0.5 M, 4.8 mL, 2.4 mmol, 6 equiv) was added to a solution of triacetate **33**<sup>27</sup> (0.5 g, 0.4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (38 mL), and the mixture was stirred for 9 h. TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 92:8) showed the complete disappearance of **33** and the presence of a more polar product. MeOH (77 mL) and acetic anhydride (300 μL) were added. After 2 h, TLC (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 92:8) showed the complete disappearance of the aminotriol intermediate **35**. Evaporation of the filtrate gave a syrup which was purified by chromatography (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 98:2 → 95:5) to give, by order of elution, first carbamate **37** (20 mg, 5%), and then acetamide **36**<sup>15</sup> (371 mg, 90%) as white foams. Carbamate **37** had *R<sub>f</sub>* = 0.45 (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 92:

8);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.42–7.08 (m, 25H,  $\text{CH}_{\text{Ph}}$ ), 6.52 (d, 1H,  $J_{\text{NH},2} = 7.0$  Hz, NH), 5.90 (m, 1H, CH=), 5.29 (m, 1H,  $J_{\text{trans}} = 17.2$  Hz, = $\text{CH}_2$ ), 5.20 (m, 1H,  $J_{\text{cis}} = 10.4$  Hz, = $\text{CH}_2$ ), 5.11–4.96 (m, 4H,  $2\text{H}_{\text{Bn}}$ , H-1 $_E$ ,  $\text{H}_{\text{Bn}}$ ), 4.92 (bs, 1H, H-1 $_A$ ), 4.83 (d, 1H,  $J = 10.9$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.81 (d, 1H,  $J = 11.8$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.73 (d, 1H,  $J = 10.4$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.63–4.55 (m, 2H,  $\text{H}_{\text{Bn}}$ ), 4.45–4.42 (m, 2H, H-1 $_D$ ,  $\text{H}_{\text{Bn}}$ ), 4.29 (dd, 1H,  $J_{2,3} = 9.3$  Hz,  $J_{3,4} = 9.6$  Hz, H-3 $_E$ ), 4.27 (d, 1H,  $J = 12.0$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.15 (m, 1H,  $\text{H}_{\text{All}}$ ), 4.12–4.09 (m, 2H, H-3 $_A$ , H-5 $_E$ ), 3.98 (m, 1H,  $\text{H}_{\text{All}}$ ), 3.87 (bs, 1H, H-2 $_A$ ), 3.84–3.70 (m, 8H, H-4 $_E$ , OMe, H-6 $_{\text{AD}}$ , H-6 $_{\text{BD}}$ , H-2 $_E$ , H-5 $_A$ ), 3.63 (m, 1H,  $J_{1,2} = 8.1$  Hz, H-2 $_D$ ), 3.51 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.5$  Hz, H-4 $_A$ ), 3.49 (d, 1H,  $J_{3,4} = J_{4,5} = 9.8$  Hz, H-4 $_D$ ), 3.33 (bs, 2H, H-6 $_{\text{AE}}$ , H-6 $_{\text{BE}}$ ), 2.90 (m, 1H, H-5 $_D$ ), 2.61 (pt, 1H,  $J_{2,3} = 9.8$  Hz, H-3 $_D$ ), 1.40 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6 $_A$ );  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  159.4 ( $\text{C}_{\text{NOMe}}$ ), 139.7–137.8 ( $\text{C}_{\text{Ph}}$ ), 134.2 (CH=), 129.4–127.9 ( $\text{CH}_{\text{Ph}}$ ), 117.9 (=CH $_2$ ), 103.2 (C-1 $_D$ ,  $J_{\text{CH}} = 161.6$  Hz), 98.6 (C-1 $_A$ ,  $J_{\text{CH}} = 171.5$  Hz), 95.8 (C-1 $_E$ ,  $J_{\text{CH}} = 170.5$  Hz), 83.1 (C-3 $_E$ ), 80.1 (C-2 $_E$ ), 79.9 (C-4 $_A$ ), 78.7 (C-4 $_E$ ), 78.1 (C-2 $_A$ ), 77.1 (C-3 $_D$ ), 76.1 ( $\text{C}_{\text{Bn}}$ ), 76.0 (C-3 $_A$ ), 75.7, 75.6, 75.4 (3C,  $\text{C}_{\text{Bn}}$ ), 75.2 (C-5 $_D$ ), 73.8 ( $\text{C}_{\text{Bn}}$ ), 71.5 (C-4 $_D$ ), 70.6 (C-5 $_E$ ), 69.0 (C-5 $_A$ ), 68.3 ( $\text{C}_{\text{All}}$ ), 68.1 (C-6 $_E$ ), 62.7 (C-6 $_D$ ), 58.4 (C-2 $_D$ ), 52.9 (OMe), 18.1 (C-6 $_A$ ); HRMS (ESI $^+$ ) for  $\text{C}_{58}\text{H}_{69}\text{NO}_{16}$  ( $[\text{M} + \text{Na}]^+$ , 1058.4514) found  $m/z$  1058.4514, ( $[\text{M} + \text{NH}_4]^+$ , 1053.4960) found  $m/z$  1053.4946.

**Allyl (2-Acetamido-2-deoxy-4,6-O-isopropylidene- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-O-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-4-O-benzyl- $\alpha$ -L-rhamnopyranoside (40).**<sup>15</sup> **Route a.** Methanolic MeONa (0.5 M, 3.7 mL, 1.9 mmol, 6 equiv) was added to a solution of fully protected trisaccharide **39**<sup>15,27</sup> (375 mg, 310  $\mu\text{mol}$ ) in  $\text{CH}_2\text{Cl}_2$  (30 mL), and the mixture was stirred for 9 h. TLC (Chex/EtOAc, 1:1) showed the complete disappearance of **39**, and the presence of a single more polar product which reacted with ninhydrin. MeOH (60 mL) and acetic anhydride (300  $\mu\text{L}$ ) were added. After 2 h, TLC (Chex/EtOAc, 1:1) showed the complete disappearance of the aminotriol intermediate. Evaporation of the filtrate gave a syrup which was purified by chromatography (Chex/EtOAc, 1:1) to give compound **40** (258 mg, 90%) as a white foam.

**Route b.** Alternatively, starting from acceptor **41**<sup>24</sup> (2.1 g, 2.6 mmol) and donor **14** (1.8 g, 3.1 mmol, 1.2 equiv), the condensation, transesterification, and acetalation steps were run without any intermediate purification. Column chromatography (Chex/EtOAc, 9:1  $\rightarrow$  1:1) of the residue gave the expected **40** (2.4 g, 87%) over three steps.

**Route c.** Alternatively, starting from acceptor **41** (250 mg, 310  $\mu\text{mol}$ ) and donor **18**<sup>27</sup> (220 mg, 400  $\mu\text{mol}$ , 1.3 equiv), the condensation and transesterification steps were run without any intermediate purification. Column chromatography (Chex/EtOAc, 1:1) of the residue gave the expected **40** (258 mg, 80%) over two steps. Data for alcohol **40** were as described.<sup>15</sup>

**Allyl (2-Deoxy-4,6-O-isopropylidene-2-trichloroacetamido- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-O-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-(4-O-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-O-benzyl- $\alpha$ -L-rhamnopyranoside (20).** Anhydrous  $\text{K}_2\text{CO}_3$  (175 mg, 1.3 mmol, 1 equiv) was added to a stirred solution of **19** (1.9 g, 1.3 mmol) in dry MeOH (20 mL). The mixture was stirred at rt for 1 night, at which time TLC (Tol/EtOAc, 7:3) indicated total conversion of **19** into a single product. Volatiles were removed under reduced pressure to give crude alcohol **20** (1.7 g, 92%) as a white foam. Tetrasaccharide **20** had  $R_f = 0.45$  (Tol/EtOAc, 7:3);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.43–7.08 (m, 36H, NH,  $\text{CH}_{\text{Ph}}$ ), 5.88 (m, 1H, =CH), 5.29 (m, 1H,  $J_{\text{trans}} = 17.2$  Hz, = $\text{CH}_2$ ), 5.23–5.20 (m, 2H, H-1 $_E$ , = $\text{CH}_2$ ), 5.13 (d, 1H,  $J = 11.0$  Hz,  $\text{H}_{\text{Bn}}$ ), 5.12 (bs, 1H, H-1 $_A$ ), 5.08 (d, 1H,  $J = 11.9$  Hz,  $\text{H}_{\text{Bn}}$ ), 5.06 (d, 1H,  $J = 11.0$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.95 (d, 1H,  $J = 10.8$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.90 (d, 1H,  $J = 12.0$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.82 (d, 1H,  $J = 11.0$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.79–4.77 (m, 2H,  $\text{H}_{\text{Bn}}$ , H-1 $_B$ ), 4.72 (d, 1H,  $J = 11.8$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.70–4.61 (m, 4H,  $2\text{H}_{\text{Bn}}$ , H-1 $_D$ ,  $\text{H}_{\text{Bn}}$ ), 4.56 (d, 1H,  $J = 10.2$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.53 (d, 1H,  $J = 11.0$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.37 (d, 1H,  $J = 12.0$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.19–4.13 (m, 5H, H-3 $_A$ , H-2 $_A$ , H-3 $_E$ ,  $\text{H}_{\text{All}}$ , H-5 $_E$ ), 4.02 (dd, 1H,  $J_{1,2} = 2.0$  Hz,  $J_{2,3} = 2.8$  Hz, H-2 $_B$ ), 4.01–3.92 (m, 5H,  $\text{H}_{\text{All}}$ , H-3 $_B$ , H-2 $_D$ , H-4 $_E$ , H-2 $_E$ ), 3.80 (dq,

1H,  $J_{4,5} = 9.2$  Hz, H-5 $_A$ ), 3.73 (dq, 1H,  $J_{4,5} = 9.4$  Hz, H-5 $_B$ ), 3.56 (dd, 1H,  $J_{5,6} = 5.3$  Hz,  $J_{6\text{a},6\text{b}} = 10.8$  Hz, H-6 $_{\text{AD}}$ ), 3.52–3.47 (m, 4H, H-6 $_{\text{AE}}$ , H-6 $_{\text{BE}}$ , H-4 $_A$ , H-4 $_D$ ), 3.41 (pt, 1H,  $J_{3,4} = J_{4,5} = 9.2$  Hz, H-4 $_D$ ), 3.38 (pt, 1H,  $J_{5,6\text{b}} = J_{6\text{a},6\text{b}} = 10.0$  Hz, H-6 $_{\text{BD}}$ ), 2.85 (m, 1H, OH), 2.82 (ddd, 1H,  $J_{5,6\text{a}} = 5.4$  Hz, H-5 $_D$ ), 2.45 (ddd, 1H,  $J_{2,3} = 9.6$  Hz,  $J_{3,\text{OH}} = 2.1$  Hz, H-3 $_D$ ), 1.47 (s, 3H,  $\text{H}_{\text{IPr}}$ ), 1.46 (s, 3H,  $\text{H}_{\text{IPr}}$ ), 1.41 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6 $_A$ ), 1.35 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6 $_B$ );  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  163.7 ( $\text{C}_{\text{NTCA}}$ ), 138.6–137.5 ( $\text{C}_{\text{Ph}}$ ), 133.8 (CH=), 129.3–127.4 ( $\text{CH}_{\text{Ph}}$ ), 117.2 (=CH $_2$ ), 101.0 (C-1 $_D$ ,  $J_{\text{CH}} = 159.1$  Hz), 101.0 (C-1 $_A$ ,  $J_{\text{CH}} = 170.3$  Hz), 99.7 ( $\text{C}_{\text{IPr}}$ ), 97.9 (C-1 $_B$ ,  $J_{\text{CH}} = 168.6$  Hz), 94.1 (C-1 $_E$ ,  $J_{\text{CH}} = 166.2$  Hz), 92.7 ( $\text{CCl}_3$ ), 83.3 (C-3 $_E$ ), 80.5 (C-4 $_B$ ), 79.8 (C-4 $_A$ ), 79.7 (C-2 $_E$ ), 79.5 (C-3 $_B$ ), 78.7 (C-4 $_E$ ), 76.3, 75.4, 75.3, 74.9 (5C,  $\text{C}_{\text{Bn}}$ ), 74.7 (C-2 $_B$ ), 74.1 (C-3 $_A$ ), 74.0 (C-4 $_D$ ), 73.5 ( $\text{C}_{\text{Bn}}$ ), 73.3 (C-2 $_A$ ), 73.0 (C-3 $_D$ ), 72.1 ( $\text{C}_{\text{Bn}}$ ), 70.0 (C-5 $_E$ ), 68.7 (C-5 $_A$ ), 68.0 (C-5 $_B$ ), 67.9 (C-6 $_E$ ), 67.7 ( $\text{C}_{\text{All}}$ ), 67.1 (C-5 $_D$ ), 61.4 (C-6 $_D$ ), 58.8 (C-2 $_D$ ), 29.1 ( $\text{C}_{\text{IPr}}$ ), 19.0 ( $\text{C}_{\text{IPr}}$ ), 18.0 (C-6 $_B$ ), 17.8 (C-6 $_A$ ); HRMS (ESI $^+$ ) for  $\text{C}_{81}\text{H}_{92}\text{Cl}_3\text{NO}_{19}$  ( $[\text{M} + \text{Na}]^+$ , 1510.5227) found  $m/z$  1510.5243, ( $[\text{M} + \text{NH}_4]^+$ , 1505.5673) found  $m/z$  1505.5669.

**(3,4-Di-O-benzyl-2-O-levulinoyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-2-O-acetyl-4-O-benzyl- $\alpha$ / $\beta$ -L-rhamnopyranose (43).** 1,5-Cyclooctadiene-bis(methyldiphenylphosphine)-iridium hexafluorophosphate (175 mg) was dissolved in THF (90 mL) and the resulting red solution was degassed under an argon stream. Hydrogen was bubbled through the solution, causing the color to change to yellow. The solution was then degassed again under an argon stream. A solution of **42**<sup>16</sup> (8.0 g, 10.6 mmol) in THF (15 mL) was added. The mixture was stirred overnight at rt. TLC (Tol/EtOAc, 8:2) showed the complete disappearance of **42** and the presence of a less polar product. The mixture was treated with a solution of iodine (5.4 g, 21.2 mmol) in THF/water (30 mL, 4:1 v/v), and stirred for 1 h at rt. TLC (Tol/EtOAc, 8:2 and  $\text{CH}_2\text{Cl}_2/\text{MeOH}$ , 98:2) showed the conversion of the intermediate into a more polar product. Excess iodine was destroyed by adding a solution of freshly prepared 5% aq sodium bisulphite (25 mL).  $\text{CH}_2\text{Cl}_2$  (200 mL) was added, and the organic phase was washed with brine (3  $\times$  50 mL), water (3  $\times$  50 mL), dried on  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated to dryness. Chromatography of the residue ( $\text{CH}_2\text{Cl}_2/\text{MeOH}$ , 99:1  $\rightarrow$  90:10) gave **43** (7.1 g, 93%) as a yellow syrup. Hemiacetal **43** had  $R_f = 0.45$  ( $\text{CH}_2\text{Cl}_2/\text{MeOH}$ , 98:2);  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.39–7.28 (m, 15H,  $\text{CH}_{\text{Ph}}$ ), 5.44 (dd, 1H,  $J_{1,2} = 1.9$  Hz, H-2 $_B$ ), 5.16 (dd, 1H,  $J_{1,2} = 1.8$  Hz, H-2 $_C$ ), 5.12 (dd, 1H,  $J_{1,\text{OH}} = 3.9$  Hz, H-1 $_C$ ), 5.05 (d, 1H, H-1 $_B$ ), 4.92 (d, 1H,  $J = 11.0$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.83 (d, 1H,  $J = 10.9$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.66 (d, 1H,  $J = 11.1$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.63–4.60 (m, 2H,  $\text{H}_{\text{Bn}}$ ), 4.45 (d, 1H,  $J = 11.3$  Hz,  $\text{H}_{\text{Bn}}$ ), 4.20 (dd, 1H,  $J_{2,3} = 3.3$  Hz,  $J_{3,4} = 9.5$  Hz, H-3 $_C$ ), 3.98 (dq, 1H,  $J_{4,5} = 9.5$  Hz, H-5 $_C$ ), 3.89 (dd, 1H,  $J_{2,3} = 3.3$  Hz,  $J_{3,4} = 9.3$  Hz, H-3 $_B$ ), 3.83 (dq, 1H,  $J_{4,5} = 9.3$  Hz, H-5 $_B$ ), 3.77 (d, 1H, OH-1 $_C$ ), 3.47 (pt, 1H, H-4 $_C$ ), 3.43 (pt, 1H, H-4 $_B$ ), 2.70 (m, 4H,  $4\text{H}_{\text{Lev}}$ ), 2.18 (s, 3H,  $\text{CH}_3\text{Lev}$ ), 2.14 (s, 3H,  $\text{H}_{\text{Ac}}$ ), 1.29 (d, 3H,  $J_{5,6} = 6.3$  Hz, H-6 $_B$ ), 1.30 (d, 3H,  $J_{5,6} = 6.2$  Hz, H-6 $_C$ );  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  206.8 ( $\text{C}_{\text{Lev}}$ ), 172.6 ( $\text{C}_{\text{Lev}}$ ), 171.6 ( $\text{C}_{\text{Ac}}$ ), 138.8–138.4 ( $\text{C}_{\text{Ph}}$ ), 128.9–128.1 ( $\text{CH}_{\text{Ph}}$ ), 99.9 (C-1 $_B$ ,  $J_{\text{CH}} = 173.3$  Hz), 92.0 (C-1 $_C$ ,  $J_{\text{CH}} = 170.0$  Hz), 80.7 (C-4 $_C$ ), 80.2 (C-4 $_B$ ), 78.0 (C-3 $_B$ ), 77.7 (C-3 $_C$ ), 75.8, 75.6 (2C,  $\text{C}_{\text{Bn}}$ ), 73.2 (C-2 $_C$ ), 72.0 ( $\text{C}_{\text{Bn}}$ ), 69.7 (C-2 $_B$ ), 69.0 (C-5 $_B$ ), 68.0 (C-5 $_C$ ), 38.4 ( $\text{CH}_2\text{Lev}$ ), 30.2 ( $\text{CH}_3\text{Lev}$ ), 28.5 ( $\text{CH}_2\text{Lev}$ ), 21.5 ( $\text{C}_{\text{Ac}}$ ), 18.5, 18.4 (2C, C-6 $_B$ , C-6 $_C$ ); HRMS (ESI $^+$ ) for  $\text{C}_{40}\text{H}_{48}\text{O}_{12}$  ( $[\text{M} + \text{Na}]^+$ , 743.3043) found  $m/z$  743.3173, ( $[\text{M} + \text{NH}_4]^+$ , 738.3489) found  $m/z$  738.3627.

**(3,4-Di-O-benzyl-2-O-levulinoyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-2-O-acetyl-4-O-benzyl- $\alpha$ / $\beta$ -L-rhamnopyranose Trichloroacetimidate (23).** Hemiacetal **43** (8.7 g, 12.1 mmol) was dissolved in DCE (50 mL), placed under argon, and cooled to  $-5^\circ\text{C}$ . Trichloroacetimidate (6.1 mL, 60.7 mmol, 5 equiv) and DBU (508  $\mu\text{L}$ , 3.3 mmol, 0.28 equiv) were added. The mixture was stirred at  $-5^\circ\text{C}$  for 10 min. TLC (Chex/EtOAc +  $\text{Et}_3\text{N}$ , 7:3) showed the complete disappearance of **43** and the presence of a single less polar product. The mixture was directly chromatographed (Chex/EtOAc + 5%  $\text{Et}_3\text{N}$ , 7:3  $\rightarrow$  1:1) to give **23** (8.7 g, 84%) as a yellow syrup. Trichloroacetimidate **23** ( $\alpha$  anomer) had  $R_f = 0.4$  (Chex/EtOAc,



7:3); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.72 (s, 1H, NH), 7.42–7.29 (m, 15H, CH<sub>Ph</sub>), 6.22 (d, 1H, *J*<sub>1,2</sub> = 1.9 Hz, H-1<sub>C</sub>), 5.48 (dd, 1H, *J*<sub>1,2</sub> = 1.8 Hz, H-2<sub>B</sub>), 5.31 (dd, 1H, H-2<sub>C</sub>), 5.10 (d, 1H, H-1<sub>B</sub>), 4.92 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.85 (d, 1H, *J* = 10.8 Hz, H<sub>Bn</sub>), 4.68–4.62 (m, 3H, H<sub>Bn</sub>), 4.51 (d, 1H, *J* = 11.3 Hz, H<sub>Bn</sub>), 4.26 (dd, 1H, *J*<sub>2,3</sub> = 3.3 Hz, *J*<sub>3,4</sub> = 9.5 Hz, H-3<sub>C</sub>), 3.96 (dq, 1H, *J*<sub>4,5</sub> = 9.5 Hz, H-5<sub>C</sub>), 3.89 (dd, 1H, *J*<sub>2,3</sub> = 3.3 Hz, *J*<sub>3,4</sub> = 9.3 Hz, H-3<sub>B</sub>), 3.82 (dq, 1H, *J*<sub>4,5</sub> = 9.4 Hz, H-5<sub>B</sub>), 3.58 (pt, 1H, H-4<sub>C</sub>), 3.44 (pt, 1H, H-4<sub>B</sub>), 2.72 (m, 4H, 4H<sub>Lev</sub>), 2.18 (s, 3H, CH<sub>3Lev</sub>), 2.17 (s, 3H, H<sub>Ac</sub>), 1.36 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>C</sub>), 1.29 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>B</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 206.5 (C<sub>Lev</sub>), 172.3 (C<sub>Lev</sub>), 170.3 (C<sub>Ac</sub>), 160.5 (C=NH), 138.8–138.0 (C<sub>Ph</sub>), 129.2–127.9 (CH<sub>Ph</sub>), 99.9 (C-1<sub>B</sub>), <sup>1</sup>J<sub>CH</sub> = 168.1 Hz), 94.5 (C-1<sub>C</sub>, <sup>1</sup>J<sub>CH</sub> = 178.7 Hz), 91.2 (CCl<sub>3</sub>), 80.2 (C-4<sub>C</sub>), 80.1 (C-4<sub>B</sub>), 77.9 (C-3<sub>B</sub>), 76.5 (C-3<sub>C</sub>), 76.1, 75.7, 72.0 (3C, C<sub>Bn</sub>), 71.2 (C-2<sub>C</sub>), 71.1 (C-5<sub>C</sub>), 69.6 (C-2<sub>B</sub>), 69.1 (C-5<sub>B</sub>), 38.4 (CH<sub>2Lev</sub>), 30.2 (CH<sub>3Lev</sub>), 28.5 (CH<sub>2Lev</sub>), 21.3 (C<sub>Ac</sub>), 18.4, 18.3 (2C, C-6<sub>B</sub>, C-6<sub>C</sub>).

**Allyl (3,4-Di-*O*-benzyl-2-*O*-levulinoyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-*O*-acetyl-4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-acetamido-2-deoxy-4,6-*O*-isopropylidene- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-4-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (44).** TfOH (23  $\mu$ L, 263  $\mu$ mol, 0.9 equiv) was added to a solution of trisaccharide acceptor **40** (310 mg, 290  $\mu$ mol) and trichloroacetimidate **23** (379 mg, 440  $\mu$ mol, 1.5 equiv) in toluene (10 mL) containing 4Å MS (1.7 g), stirred at 0 °C. After 15 min, TLC (Chex/EtOAc, 1:1) showed the presence of a single more polar product. Et<sub>3</sub>N (0.2 mL) was added. The mixture was filtered, and concentrated to dryness. Chromatography of the residue (Tol/EtOAc, 8:2  $\rightarrow$  6:4) gave **44** (383 mg, 74%) as a white foam slightly contaminated. Pentasaccharide **44** had *R*<sub>f</sub> = 0.35 (Chex/EtOAc, 1:1); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.45–7.07 (m, 40H, CH<sub>Ph</sub>), 6.47 (d, 1H, *J*<sub>NH,2</sub> = 9.0 Hz, NH), 5.90 (m, 1H, CH=), 5.46 (dd, 1H, *J*<sub>2,3</sub> = 3.0 Hz, H-2<sub>B</sub>), 5.29 (m, 1H, *J*<sub>trans</sub> = 17.2 Hz, =CH<sub>2</sub>), 5.20 (m, 1H, *J*<sub>cis</sub> = 10.4 Hz, =CH<sub>2</sub>), 5.15 (d, 1H, *J*<sub>1,2</sub> = 3.5 Hz, H-1<sub>E</sub>), 5.09–4.98 (m, 5H, H-1<sub>B</sub>, H<sub>Bn</sub>), 4.94 (dd, 1H, *J*<sub>1,2</sub> = 2.1 Hz, *J*<sub>2,3</sub> = 2.7 Hz, H-2<sub>C</sub>), 4.93–4.88 (m, 3H, H<sub>Bn</sub>), 4.84 (bs, 1H, H-1<sub>A</sub>), 4.79–4.77 (bs, 2H, H-1<sub>C</sub>, H<sub>Bn</sub>), 4.67–4.60 (m, 5H, H<sub>Bn</sub>), 4.52 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.42 (d, 1H, *J* = 11.4 Hz, H<sub>Bn</sub>), 4.35 (d, 1H, *J* = 11.9 Hz, H<sub>Bn</sub>), 4.28 (d, 1H, *J*<sub>1,2</sub> = 8.6 Hz, H-1<sub>D</sub>), 4.21–4.13 (m, 6H, H-3<sub>E</sub>, H-2<sub>D</sub>, H-5<sub>E</sub>, H-3<sub>C</sub>, H<sub>All</sub>, H-3<sub>A</sub>), 4.00 (dq, 1H, *J*<sub>4,5</sub> = 9.6 Hz, H-5<sub>C</sub>), 3.97–3.83 (m, 6H, H<sub>All</sub>, H-3<sub>B</sub>, H-2<sub>E</sub>, H-2<sub>A</sub>, H-4<sub>E</sub>, H-6a<sub>D</sub>), 3.79 (dq, 1H, *J*<sub>4,5</sub> = 9.4 Hz, H-5<sub>B</sub>), 3.73 (bd, 1H, *J*<sub>5,6b</sub> = 10.3 Hz, H-6b<sub>D</sub>), 3.70 (dq, 1H, *J*<sub>4,5</sub> = 9.5 Hz, H-5<sub>A</sub>), 3.52 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.3 Hz, H-4<sub>D</sub>), 3.58–3.40 (m, 5H, H-4<sub>A</sub>, H-6a<sub>E</sub>, H-6b<sub>E</sub>, H-4<sub>C</sub>, H-4<sub>B</sub>), 2.7–2.68 (m, 6H, H-5<sub>D</sub>, 4H<sub>Lev</sub>, H-3<sub>D</sub>), 2.35 (s, 3H, H<sub>NAC</sub>), 2.19 (s, 3H, CH<sub>3Lev</sub>), 2.06 (s, 3H, H<sub>Ac</sub>), 1.48 (s, 3H, H<sub>IPr</sub>), 1.45 (s, 3H, H<sub>IPr</sub>), 1.38 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>A</sub>), 1.34 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>B</sub>), 1.27 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>C</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 206.1 (C<sub>Lev</sub>), 171.8 (C<sub>Lev</sub>), 170.5 (C<sub>Ac</sub>), 170.1 (C<sub>NAC</sub>), 138.7–137.5 (C<sub>Ph</sub>), 134.0 (CH=), 129.4–127.8 (CH<sub>Ph</sub>), 117.1 (=CH<sub>2</sub>), 103.7 (C-1<sub>D</sub>, <sup>1</sup>J<sub>CH</sub> = 159.9 Hz), 99.5 (C-1<sub>B</sub>, <sup>1</sup>J<sub>CH</sub> = 169.7 Hz), 99.3 (C<sub>IPr</sub>), 98.4 (C-1<sub>A</sub>, <sup>1</sup>J<sub>CH</sub> = 174.4 Hz), 97.5 (C-1<sub>C</sub>, <sup>1</sup>J<sub>CH</sub> = 170.6 Hz), 94.4 (C-1<sub>E</sub>, <sup>1</sup>J<sub>CH</sub> = 170.1 Hz), 83.5 (C-3<sub>E</sub>), 80.7 (C-3<sub>D</sub>), 80.4 (C-2<sub>E</sub>), 80.1, 80.0 (2C, C-4<sub>B</sub>, C-4<sub>C</sub>), 79.8 (C-4<sub>A</sub>), 78.5 (C-4<sub>E</sub>), 78.2 (C-3<sub>C</sub>), 77.9 (C-3<sub>B</sub>), 76.7 (C-2<sub>A</sub>), 76.1, 75.9, 75.5, 75.4, 75.1, 75.0 (6C, C<sub>Bn</sub>), 74.8 (C-3<sub>A</sub>), 73.4 (C<sub>Bn</sub>), 73.3 (C-2<sub>C</sub>), 72.4 (C-4<sub>D</sub>), 71.3 (C<sub>Bn</sub>), 70.0 (C-5<sub>E</sub>), 69.3 (C-2<sub>B</sub>), 68.6 (C-5<sub>A</sub>), 68.5 (C-5<sub>B</sub>), 67.9 (C-6<sub>E</sub>), 67.8 (C<sub>All</sub>), 67.4 (C-5<sub>D</sub>), 67.3 (C-5<sub>C</sub>), 62.2 (C-6<sub>D</sub>), 55.0 (C-2<sub>D</sub>), 38.1 (CH<sub>2Lev</sub>), 29.8 (CH<sub>3Lev</sub>), 29.2 (C<sub>IPr</sub>), 28.3 (CH<sub>2Lev</sub>), 24.1 (C<sub>NAC</sub>), 21.1 (C<sub>Ac</sub>), 19.1 (C<sub>IPr</sub>), 17.9, 17.8, 17.7 (3C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>C</sub>); HRMS (ESI<sup>+</sup>) for C<sub>101</sub>H<sub>119</sub>NO<sub>26</sub> ([M + H]<sup>+</sup>, 1762.8098) found *m/z* 1762.8198, ([M + Na]<sup>+</sup>, 1784.7917) found *m/z* 1784.7988.

**Allyl (3,4-Di-*O*-benzyl-2-*O*-levulinoyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-*O*-acetyl-4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-acetamido-2-deoxy-4,6-*O*-isopropylidene- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-(4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (45).** TMSOTf (18  $\mu$ L, 100  $\mu$ mol, 0.9 equiv) was added to a solution of acceptor **21** (150 mg, 110  $\mu$ mol) and trichloroacetimidate **23** (140 mg, 160  $\mu$ mol, 1.5 equiv) in toluene

(5 mL) containing 4Å MS (620 mg), stirred at 0 °C. After 45 min, TLC (Tol/EtOAc, 75:25; Chex/EtOAc, 1:1) showed the presence of a single less polar product. Et<sub>3</sub>N (0.2 mL) was added and the mixture was filtered, and concentrated to dryness. Chromatography of the residue (Tol/EtOAc, 75:25  $\rightarrow$  1:1) gave **45** (149 mg, 65%) as a white foam slightly contaminated. Hexasaccharide **45** had *R*<sub>f</sub> = 0.55 (Chex/EtOAc, 1:1); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.47–7.06 (m, 50H, CH<sub>Ph</sub>), 6.38 (d, 1H, *J*<sub>NH,2</sub> = 9.1 Hz, NH), 5.86 (m, 1H, CH=), 5.44 (dd, 1H, *J*<sub>2,3</sub> = 3.2 Hz, H-2<sub>B</sub>), 5.26 (m, 1H, *J*<sub>trans</sub> = 17.2 Hz, =CH<sub>2</sub>), 5.21–5.17 (m, 2H, H=CH<sub>2</sub>, H-1<sub>E</sub>), 5.07 (d, 1H, *J*<sub>1,2</sub> = 1.6 Hz, H-1<sub>A</sub>), 5.05 (d, 1H, *J*<sub>1,2</sub> = 1.8 Hz, H-1<sub>B</sub>), 5.04–4.87 (m, 9H, 4H<sub>Bn</sub>, H-2<sub>C</sub>, 2H<sub>Bn</sub>), 4.84 (d, 2H, *J* = 11.2 Hz, H<sub>Bn</sub>), 4.78 (d, 1H, *J* = 10.2 Hz, H<sub>Bn</sub>), 4.75 (d, 1H, *J*<sub>1,2</sub> = 1.4 Hz, H-1<sub>B</sub>), 4.70 (bs, 1H, H-1<sub>C</sub>), 4.68–4.57 (m, 9H, H<sub>Bn</sub>), 4.50 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.41 (d, 1H, *J* = 11.3 Hz, H<sub>Bn</sub>), 4.35 (d, 1H, *J* = 12.0 Hz, H<sub>Bn</sub>), 4.28 (d, 1H, *J*<sub>1,2</sub> = 8.6 Hz, H-1<sub>D</sub>), 4.21–4.08 (m, 6H, H-3<sub>E</sub>, H-5<sub>E</sub>, H-2<sub>D</sub>, H-3<sub>A</sub>, H<sub>All</sub>, H-3<sub>C</sub>), 3.99 (dd, 1H, *J*<sub>2,3</sub> = 2.6 Hz, H-2<sub>A</sub>), 3.96–3.84 (m, 7H, H-5<sub>C</sub>, H-2<sub>B</sub>, H<sub>All</sub>, H-3<sub>B</sub>, H-3<sub>B</sub>, H-2<sub>E</sub>, H-4<sub>E</sub>), 3.81–3.73 (m, 2H, H-5<sub>A</sub>, H-5<sub>B</sub>), 3.71 (dq, 1H, *J*<sub>4,5</sub> = 9.4 Hz, *J*<sub>5,6</sub> = 6.2 Hz, H-5<sub>B</sub>), 3.54–3.38 (m, 8H, H-6a<sub>D</sub>, H-6b<sub>D</sub>, H-6a<sub>E</sub>, H-6b<sub>E</sub>, H-4<sub>A</sub>, H-4<sub>B</sub>, H-4<sub>D</sub>, H-4<sub>C</sub>, H-4<sub>B</sub>), 2.75–2.64 (m, 6H, H-5<sub>D</sub>, 4H<sub>Lev</sub>, H-3<sub>D</sub>), 2.31 (s, 3H, H<sub>NAC</sub>), 2.18 (s, 3H, CH<sub>3Lev</sub>), 2.05 (s, 3H, H<sub>Ac</sub>), 1.42 (s, 3H, H<sub>IPr</sub>), 1.39 (s, 3H, H<sub>IPr</sub>), 1.35–1.31 (m, 9H, H-6<sub>A</sub>, H-6<sub>B</sub>, H-6<sub>C</sub>), 1.24 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>C</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 206.1 (C<sub>Lev</sub>), 171.7 (C<sub>Lev</sub>), 170.4 (C<sub>Ac</sub>), 170.0 (C<sub>NAC</sub>), 138.6–137.5 (C<sub>Ph</sub>), 133.8 (CH=), 129.3–127.7 (CH<sub>Ph</sub>), 117.1 (=CH<sub>2</sub>), 103.6 (C-1<sub>D</sub>, <sup>1</sup>J<sub>CH</sub> = 159.0 Hz), 101.1 (C-1<sub>A</sub>, <sup>1</sup>J<sub>CH</sub> = 172.8 Hz), 99.5 (C-1<sub>B</sub>, <sup>1</sup>J<sub>CH</sub> = 167.8 Hz), 99.2 (C<sub>IPr</sub>), 97.9 (C-1<sub>B</sub>, <sup>1</sup>J<sub>CH</sub> = 172.8 Hz), 97.8 (C-1<sub>C</sub>, <sup>1</sup>J<sub>CH</sub> = 172.8 Hz), 94.2 (C-1<sub>E</sub>, <sup>1</sup>J<sub>CH</sub> = 165.3 Hz), 83.4 (C-3<sub>E</sub>), 80.5 (C-3<sub>D</sub>), 80.4 (C-4<sub>B</sub>), 80.2 (C-2<sub>E</sub>), 80.1 (C-4<sub>B</sub>), 80.0 (C-4<sub>C</sub>), 79.8 (C-4<sub>A</sub>), 79.2 (C-3<sub>B</sub>), 78.5 (C-4<sub>E</sub>), 78.2 (C-3<sub>C</sub>), 77.9 (C-3<sub>B</sub>), 76.2 (C-2<sub>A</sub>), 76.1, 75.7, 75.4 (3C, C<sub>Bn</sub>), 75.3 (C-2<sub>B</sub>), 75.2, 75.1, 75.0, 74.5 (4C, C<sub>Bn</sub>), 74.5 (C-3<sub>A</sub>), 73.5 (C<sub>Bn</sub>), 73.2 (C-2<sub>C</sub>), 72.1 (C-4<sub>D</sub>), 71.9, 71.4 (2C, C<sub>Bn</sub>), 70.0 (C-5<sub>E</sub>), 69.3 (C-2<sub>B</sub>), 68.9 (C-5<sub>A</sub>), 68.5 (C-5<sub>B</sub>), 67.9 (C-6<sub>B</sub>), 67.8 (C-5<sub>B</sub>), 67.7 (C<sub>All</sub>), 67.3 (C-5<sub>C</sub>), 67.2 (C-5<sub>D</sub>), 62.0 (C-6<sub>D</sub>), 55.0 (C-2<sub>D</sub>), 38.1 (CH<sub>2Lev</sub>), 29.8 (CH<sub>3Lev</sub>), 29.2 (C<sub>IPr</sub>), 28.2 (CH<sub>2Lev</sub>), 24.1 (C<sub>NAC</sub>), 21.0 (C<sub>Ac</sub>), 19.1 (C<sub>IPr</sub>), 18.0, 17.9, 17.8, 17.7 (4C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>C</sub>, C-6<sub>C</sub>); HRMS (ESI<sup>+</sup>) for C<sub>121</sub>H<sub>141</sub>NO<sub>30</sub> ([M + H]<sup>+</sup>, 2088.9617) found *m/z* 2088.9619, ([M + Na]<sup>+</sup>, 2110.9436) found *m/z* 2110.9497.

**Allyl (2-Deoxy-4,6-*O*-isopropylidene-2-trichloroacetamido-3-*O*-trimethylsilyl- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-4-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (47).** TMSOTf (61.0  $\mu$ L, 340  $\mu$ mol, 0.3 equiv) was added to a solution of acceptor **27** (1.3 g, 1.1 mmol) and trichloroacetimidate **23** (1.5 g, 1.7 mmol, 1.5 equiv) in toluene (30 mL) containing 4Å MS (960 mg), stirred at –78 °C. After 15 min, TLC (Tol/EtOAc, 8:2) showed the absence of **27**. Et<sub>3</sub>N (0.2 mL) was added and the mixture was filtered, and concentrated to dryness. Chromatography of the residue (Tol/EtOAc, 9:1  $\rightarrow$  7:3) gave first **47** (233 mg, 11%), and then pentasaccharide **48** (1.1 g, 52%), both as white foams. Trisaccharide **47** had *R*<sub>f</sub> = 0.5 (Tol/EtOAc, 8:2); <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.47–7.08 (m, 26H, NH, CH<sub>Ph</sub>), 5.89 (m, 1H, CH=), 5.32 (m, 1H, *J*<sub>trans</sub> = 17.2 Hz, =CH<sub>2</sub>), 5.23 (m, 1H, *J*<sub>cis</sub> = 10.6 Hz, =CH<sub>2</sub>), 5.12 (d, 1H, *J*<sub>1,2</sub> = 3.6 Hz, H-1<sub>E</sub>), 5.11–5.05 (m, 3H, H<sub>Bn</sub>), 4.93 (d, 1H, *J* = 12.8 Hz, H<sub>Bn</sub>), 4.86 (d, 1H, *J*<sub>1,2</sub> = 1.6 Hz, H-1<sub>A</sub>), 4.83–4.76 (m, 3H, H<sub>Bn</sub>, H-1<sub>D</sub>, H<sub>Bn</sub>), 4.61–4.56 (m, 2H, H<sub>Bn</sub>), 4.50 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.32 (d, 1H, *J* = 12.0 Hz, H<sub>Bn</sub>), 4.22–4.10 (m, 4H, H-3<sub>E</sub>, H<sub>All</sub>, H-3<sub>A</sub>, H-5<sub>E</sub>), 4.08–3.97 (m, 3H, H-2<sub>A</sub>, H-2<sub>D</sub>, H<sub>All</sub>), 3.90 (dd, 1H, *J*<sub>5,6a</sub> = 5.3 Hz, *J*<sub>6a,6b</sub> = 10.7 Hz, H-6a<sub>D</sub>), 3.83–3.72 (m, 4H, H-2<sub>E</sub>, H-4<sub>E</sub>, H-6b<sub>D</sub>, H-5<sub>A</sub>), 3.58–3.52 (m, 2H, H-4<sub>D</sub>, H-4<sub>A</sub>), 3.45 (dd, 1H, *J*<sub>5,6a</sub> = 1.5 Hz, *J*<sub>6a,6b</sub> = 10.8 Hz, H-6a<sub>E</sub>), 3.40 (dd, 1H, *J*<sub>5,6b</sub> = 1.5 Hz, H-6b<sub>E</sub>), 3.36 (pt, 1H, *J*<sub>2,3</sub> = *J*<sub>3,4</sub> = 9.2 Hz, H-3<sub>D</sub>), 2.96 (ddd, 1H, *J*<sub>5,6b</sub> = *J*<sub>4,5</sub> = 9.8 Hz, H-5<sub>D</sub>), 1.51 (s, 3H, H<sub>IPr</sub>), 1.45 (s, 3H, H<sub>IPr</sub>), 1.44 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>A</sub>), 0.13 (s, 9H, H<sub>Si</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 161.7 (C<sub>NITCA</sub>), 138.7–137.7 (C<sub>Ph</sub>), 134.0 (CH=), 129.1–127.3 (CH<sub>Ph</sub>), 117.1 (=CH<sub>2</sub>), 101.5 (C-1<sub>D</sub>, <sup>1</sup>J<sub>CH</sub> = 163.6 Hz), 99.4 (C<sub>IPr</sub>), 98.5 (C-1<sub>A</sub>, <sup>1</sup>J<sub>CH</sub> = 174.0 Hz), 95.2 (C-1<sub>E</sub>, <sup>1</sup>J<sub>CH</sub> = 167.5 Hz), 93.3 (CCl<sub>3</sub>), 83.3 (C-3<sub>E</sub>), 79.8 (C-4<sub>A</sub>), 78.7, 78.6 (2C, C-2<sub>E</sub>, C-4<sub>E</sub>), 75.9 (C<sub>Bn</sub>), 75.4 (C-3<sub>A</sub>), 75.3,

74.9 (2C, C<sub>Bn</sub>), 74.3 (C-3<sub>D</sub>), 74.3 (C<sub>Bn</sub>), 74.2 (C-4<sub>D</sub>), 74.1 (C-2<sub>A</sub>), 73.4 (C<sub>Bn</sub>), 70.0 (C-5<sub>E</sub>), 68.5 (C-5<sub>A</sub>), 67.9 (C-6<sub>E</sub>), 67.8 (C<sub>All</sub>), 67.2 (C-5<sub>D</sub>), 62.1 (C-6<sub>D</sub>), 58.6 (C-2<sub>D</sub>), 29.1 (C<sub>IPr</sub>), 19.0 (C<sub>IPr</sub>), 18.0 (C-6<sub>A</sub>), 0.7 (C<sub>Si</sub>); HRMS (ESI<sup>+</sup>) for C<sub>64</sub>H<sub>76</sub>Cl<sub>3</sub>NO<sub>15</sub>Si ([M + Na]<sup>+</sup>, 1256.4104) found *m/z* 1256.4188, ([M + NH<sub>4</sub>]<sup>+</sup>, 1251.4550) found *m/z* 1251.4548.

**Allyl (3,4-Di-*O*-benzyl-2-*O*-levulinoyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-*O*-acetyl-4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-deoxy-4,6-*O*-isopropylidene-2-trichloroacetamido- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-(4-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (48).** TMSOTf (49  $\mu$ L, 270  $\mu$ mol, 0.3 equiv) was added to a solution of acceptor **27** (1.1 g, 0.9 mmol) and trichloroacetimidate **23** (1.2 g, 1.4 mmol, 1.5 equiv) in toluene (30 mL) containing 4Å MS (1.1 g), stirred at -40 °C. After 15 min, TLC (Tol/EtOAc, 8:2) showed the absence of **27**. Et<sub>3</sub>N (0.2 mL) was added. The mixture was filtered, and concentrated to dryness. Chromatography of the residue (Tol/EtOAc, 9:1  $\rightarrow$  7:3) gave **48** (1.4 g, 75%) as a white foam. Pentasaccharide **48** had *R<sub>f</sub>* = 0.35 (Tol/EtOAc, 8:2); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.51–7.13 (m, 40H, CH<sub>Ph</sub>), 6.94 (d, 1H, *J*<sub>NH,2</sub> = 8.9 Hz, NH), 5.94 (m, 1H, CH=), 5.48 (dd, 1H, H-2<sub>B</sub>), 5.33 (m, 1H, *J*<sub>trans</sub> = 17.2 Hz, =CH<sub>2</sub>), 5.24 (m, 1H, *J*<sub>cis</sub> = 10.4 Hz, =CH<sub>2</sub>), 5.24 (d, 1H, *J*<sub>1,2</sub> = 3.5 Hz, H-1<sub>E</sub>), 5.16 (m, 2H, H<sub>Bn</sub>), 5.12 (dd, 1H, *J*<sub>2,3</sub> = 2.8 Hz, H-2<sub>C</sub>), 5.11 (d, 1H, *J*<sub>1,2</sub> = 1.3 Hz, H-1<sub>B</sub>), 5.07 (d, 1H, *J* = 12.2 Hz, H<sub>Bn</sub>), 4.98–4.94 (m, 2H, H<sub>Bn</sub>), 4.89 (d, 2H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.84 (d, 1H, *J*<sub>1,2</sub> = 1.7 Hz, H-1<sub>A</sub>), 4.79 (d, 1H, *J*<sub>1,2</sub> = 1.6 Hz, H-1<sub>C</sub>), 4.76 (d, 1H, *J* = 10.2 Hz, H<sub>Bn</sub>), 4.71 (dd, 1H, *J* = 11.2 Hz, H<sub>Bn</sub>), 4.67–4.54 (d, 6H, H<sub>Bn</sub>, H-1<sub>D</sub>), 4.46 (d, 1H, *J* = 11.3 Hz, H<sub>Bn</sub>), 4.36 (d, 1H, *J* = 12.0 Hz, H<sub>Bn</sub>), 4.24–4.15 (m, 5H, H-3<sub>E</sub>, H-3<sub>C</sub>, H<sub>All</sub>, H-3<sub>A</sub>, H-5<sub>E</sub>), 4.09–3.96 (m, 5H, H-2<sub>D</sub>, H-2<sub>A</sub>, H-5<sub>C</sub>, H<sub>All</sub>, H-5<sub>B</sub>), 3.95 (dd, 1H, *J*<sub>2,3</sub> = 3.2 Hz, *J*<sub>3,4</sub> = 9.2 Hz, H-3<sub>B</sub>), 3.91–3.85 (m, 3H, H-6<sub>AD</sub>, H-2<sub>E</sub>, H-4<sub>E</sub>), 3.78–3.73 (m, 2H, H-5<sub>A</sub>, H-6<sub>BD</sub>), 3.57 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.2 Hz, H-4<sub>D</sub>), 3.52 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.6 Hz, H-4<sub>A</sub>), 3.50–3.44 (m, 4H, H-6<sub>AE</sub>, H-6<sub>BE</sub>, H-4<sub>C</sub>, H-4<sub>B</sub>), 2.87 (ddd, 1H, *J*<sub>5,6a</sub> = 5.2 Hz, *J*<sub>5,6b</sub> = 9.8 Hz, H-5<sub>D</sub>), 2.77 (m, 5H, H-3<sub>D</sub>, 4H<sub>Lev</sub>), 2.21 (s, 3H, CH<sub>3Lev</sub>), 2.16 (s, 3H, H<sub>Ac</sub>), 1.52 (s, 3H, H<sub>IPr</sub>), 1.50 (s, 3H, H<sub>IPr</sub>), 1.46 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>A</sub>), 1.40 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>B</sub>), 1.31 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>C</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  206.5 (C<sub>Lev</sub>), 172.1 (C<sub>Lev</sub>), 169.9 (C<sub>Ac</sub>), 162.4 (C<sub>NNTCA</sub>), 139.1–138.8 (C<sub>Ph</sub>), 134.4 (CH=), 129.5–127.8 (CH<sub>Ph</sub>), 117.5 (=CH<sub>2</sub>), 101.7 (C-1<sub>D</sub>, <sup>1</sup>*J*<sub>CH</sub> = 163.9 Hz), 99.9 (C<sub>IPr</sub>), 99.4 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 174.7 Hz), 98.8 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 171.5 Hz), 98.1 (C-1<sub>C</sub>, <sup>1</sup>*J*<sub>CH</sub> = 167.6 Hz), 94.9 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 165.3 Hz), 93.5 (CCl<sub>3</sub>), 83.5 (C-3<sub>E</sub>), 80.9 (C-2<sub>E</sub>), 80.8 (C-4<sub>B</sub>), 80.3 (C-4<sub>C</sub>), 80.1 (C-4<sub>A</sub>), 79.2 (C-3<sub>D</sub>), 79.1 (C-4<sub>E</sub>), 78.0 (C-3<sub>B</sub>), 77.1 (C-3<sub>C</sub>), 76.4, 75.8, 75.7, 75.4, 75.3, 75.2 (6C, C<sub>Bn</sub>), 75.1 (C-3<sub>A</sub>), 74.2 (C-2<sub>A</sub>), 73.8 (C<sub>Bn</sub>), 73.1 (C-4<sub>D</sub>), 72.8 (C-2<sub>C</sub>), 71.9 (C<sub>Bn</sub>), 70.3 (C-5<sub>E</sub>), 69.9 (C-2<sub>B</sub>), 69.0 (C-5<sub>B</sub>), 68.8 (C-5<sub>A</sub>), 68.4 (C-6<sub>E</sub>), 68.3 (C-5<sub>C</sub>), 68.2 (C<sub>All</sub>), 67.4 (C-5<sub>D</sub>), 62.5 (C-6<sub>D</sub>), 57.8 (C-2<sub>D</sub>), 38.6 (CH<sub>2Lev</sub>), 30.3 (CH<sub>3Lev</sub>), 29.6 (C<sub>IPr</sub>), 28.7 (CH<sub>2Lev</sub>), 21.5 (C<sub>Ac</sub>), 19.4 (C<sub>IPr</sub>), 18.5 (C-6<sub>C</sub>), 18.4 (C-6<sub>A</sub>), 18.3 (C-6<sub>B</sub>); HRMS (ESI<sup>+</sup>) for C<sub>101</sub>H<sub>116</sub>Cl<sub>3</sub>NO<sub>26</sub> ([M + Na]<sup>+</sup>, 1886.6749) found *m/z* 1886.6445, ([M + NH<sub>4</sub>]<sup>+</sup>, 1881.7195) found *m/z* 1881.6897.

**Allyl (3,4-Di-*O*-benzyl-2-*O*-levulinoyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-*O*-acetyl-4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-deoxy-4,6-*O*-isopropylidene-2-trichloroacetamido- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-(4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (50).** TMSOTf (61  $\mu$ L, 340  $\mu$ mol, 0.3 equiv) was added to a solution of acceptor **20** (1.7 g, 1.1 mmol) and trichloroacetimidate **23** (1.5 g, 1.7 mmol, 1.5 equiv) in toluene (35 mL) containing 4Å MS (975 mg), stirred at -40 °C. After 1 h, TLC (Tol/EtOAc, 8:2) showed the absence of **20**. Et<sub>3</sub>N (0.2 mL) was added. The mixture was filtered, and concentrated to dryness. Chromatography of the residue (Tol/EtOAc, 95:5  $\rightarrow$  70:30) gave, by order of elution, first **50** (1.45 g, 59%), and then diol **51** (505 mg, 21%), both as white foams. Hexasaccharide **50** had *R<sub>f</sub>* = 0.45 (Tol/EtOAc, 8:2); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.47–7.10 (m, 50H, CH<sub>Ph</sub>), 6.94 (d, 1H, *J*<sub>NH,2</sub> = 8.8 Hz, NH), 5.92 (m, 1H, CH=), 5.46 (dd, 1H, *J*<sub>1,2</sub> = 1.9 Hz, *J*<sub>2,3</sub> = 3.0 Hz, H-2<sub>B</sub>), 5.30 (m, 1H, *J*<sub>trans</sub> = 17.2 Hz, =CH<sub>2</sub>), 5.25–5.22 (m, 2H, H-1<sub>E</sub>, =CH<sub>2</sub>), 5.15–5.12 (m, 3H,

H-1<sub>A</sub>, 2H<sub>Bn</sub>), 5.11–5.09 (m, 2H, H-1<sub>B</sub>, H-2<sub>C</sub>), 5.06 (d, 1H, *J* = 12.2 Hz, H<sub>Bn</sub>), 4.98–4.93 (m, 3H, H<sub>Bn</sub>), 4.89–4.83 (m, 3H, H<sub>Bn</sub>), 4.81 (d, 1H, *J*<sub>1,2</sub> = 1.4 Hz, H-1<sub>B</sub>), 4.72–4.58 (m, 10H, H-1<sub>C</sub>, H<sub>Bn</sub>, H-1<sub>D</sub>), 4.54 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.45 (d, 1H, *J* = 11.3 Hz, H<sub>Bn</sub>), 4.37 (d, 1H, *J* = 12.0 Hz, H<sub>Bn</sub>), 4.24–4.14 (m, 5H, H-3<sub>E</sub>, H-3<sub>C</sub>, H<sub>All</sub>, H-3<sub>A</sub>, H-2<sub>A</sub>, H-5<sub>E</sub>), 4.07–3.86 (m, 9H, H-2<sub>B</sub>, H-2<sub>D</sub>, H<sub>All</sub>, H-5<sub>C</sub>, H-5<sub>B</sub>, H-3<sub>B</sub>, H-3<sub>B</sub>, H-2<sub>E</sub>, H-4<sub>E</sub>), 3.80 (dq, 1H, *J*<sub>4,5</sub> = 9.5 Hz, *J*<sub>5,6</sub> = 6.2 Hz, H-5<sub>A</sub>), 3.74 (dq, 1H, *J*<sub>4,5</sub> = 9.4 Hz, H-5<sub>B</sub>), 3.58 (dd, 1H, *J*<sub>5,6a</sub> = 5.2 Hz, H-6<sub>AD</sub>), 3.56–3.43 (m, 8H, H-4<sub>A</sub>, H-4<sub>B</sub>, H-6<sub>AE</sub>, H-6<sub>BE</sub>, H-4<sub>D</sub>, H-4<sub>B</sub>, H-4<sub>C</sub>), 3.35 (m, 1H, *J*<sub>6a,6b</sub> = 10.4 Hz, H-6<sub>BD</sub>), 2.82 (ddd, 1H, *J*<sub>5,6b</sub> = 1.5 Hz, *J*<sub>4,5</sub> = 9.7 Hz, H-5<sub>D</sub>), 2.75–2.69 (m, 5H, 4H<sub>Lev</sub>, H-3<sub>D</sub>), 2.21 (s, 3H, CH<sub>3Lev</sub>), 2.14 (s, 3H, H<sub>Ac</sub>), 1.44–1.43 (m, 9H, H<sub>IPr</sub>, H-6<sub>A</sub>), 1.39 (d<sub>po</sub>, 3H, H-6<sub>B</sub>), 1.37 (d<sub>po</sub>, 3H, H-6<sub>B</sub>), 1.37 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>C</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  206.3 (C<sub>Lev</sub>), 171.8 (C<sub>Lev</sub>), 169.6 (C<sub>Ac</sub>), 162.0 (C<sub>NNTCA</sub>), 138.7–137.6 (C<sub>Ph</sub>), 133.9 (CH=), 129.4–127.5 (CH<sub>Ph</sub>), 117.2 (=CH<sub>2</sub>), 101.2 (C-1<sub>D</sub>, <sup>1</sup>*J*<sub>CH</sub> = 163.9 Hz), 100.9 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 172.0 Hz), 99.4 (C<sub>IPr</sub>), 99.0 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 168.3 Hz), 98.0 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 168.7 Hz), 97.6 (C-1<sub>C</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.4 Hz), 94.3 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 167.5 Hz), 93.1 (CCl<sub>3</sub>), 83.1 (C-3<sub>E</sub>), 80.4, 80.1, 79.9, 79.7, 79.6, 79.0 (7C, C-3<sub>B</sub>, C-3<sub>B</sub>, C-4<sub>A</sub>, C-4<sub>B</sub>, C-4<sub>B</sub>, C-4<sub>C</sub>, C-4<sub>E</sub>), 78.7 (C-3<sub>D</sub>), 77.6 (C-2<sub>E</sub>), 76.8 (C-3<sub>C</sub>), 76.2, 75.4, 75.3, 75.2, 75.0, 74.9, 74.8 (7C, C<sub>Bn</sub>), 74.3 (C-2<sub>B</sub>), 74.2 (C-3<sub>A</sub>), 73.5 (C<sub>Bn</sub>), 73.4 (C-2<sub>A</sub>), 72.6 (C-4<sub>D</sub>), 72.1 (C-2<sub>C</sub>), 72.0, 71.5 (2C, C<sub>Bn</sub>), 69.9 (C-5<sub>E</sub>), 69.4 (C-2<sub>B</sub>), 68.7 (C-5<sub>A</sub>), 68.5 (C-5<sub>B</sub>), 68.0 (C-5<sub>B</sub>), 67.9 (C-6<sub>E</sub>), 67.8 (C-5<sub>C</sub>), 67.7 (C<sub>All</sub>), 67.0 (C-5<sub>D</sub>), 61.8 (C-6<sub>D</sub>), 57.4 (C-2<sub>D</sub>), 38.2 (CH<sub>2Lev</sub>), 29.9 (CH<sub>3Lev</sub>), 29.2 (C<sub>IPr</sub>), 28.2 (CH<sub>2Lev</sub>), 21.2 (C<sub>Ac</sub>), 19.1 (C<sub>IPr</sub>), 18.3, 18.1, 18.0, 17.9 (4C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>B</sub>, C-6<sub>C</sub>); HRMS (ESI<sup>+</sup>) for C<sub>121</sub>H<sub>138</sub>Cl<sub>3</sub>NO<sub>30</sub> ([M + H]<sup>+</sup>, 2190.8447) found *m/z* 2190.8403, ([M + Na]<sup>+</sup>, 2212.8267) found *m/z* 2212.8215.

**Allyl (3,4-Di-*O*-benzyl-2-*O*-levulinoyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-*O*-acetyl-4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-deoxy-2-trichloroacetamido- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-*O*-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-(4-*O*-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-*O*-benzyl- $\alpha$ -L-rhamnopyranoside (51).** TFA (50% aq, 8 mL) was added, at 0 °C, to a solution of hexasaccharide **50** (1.9 g, 880  $\mu$ mol) in CH<sub>2</sub>Cl<sub>2</sub> (30 mL), and the biphasic mixture was stirred vigorously at rt for 1 h. TLC (Tol/EtOAc, 7:3) showed the complete disappearance of **50** and the presence of a major more polar product. Repeated coevaporation with toluene, followed by chromatography of the residue (Tol/EtOAc, 8:2  $\rightarrow$  6:4) provided diol **51** (1.7 g, 89%) as a white foam. The latter had *R<sub>f</sub>* = 0.4 (Tol/EtOAc, 7:3); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.62–7.13 (m, 50H, CH<sub>Ph</sub>), 7.04 (d, 1H, *J*<sub>NH,2</sub> = 8.6 Hz, NH), 5.94 (m, 1H, CH=), 5.53 (dd, 1H, *J*<sub>1,2</sub> = 2.0 Hz, *J*<sub>2,3</sub> = 3.0 Hz, H-2<sub>B</sub>), 5.44 (bs, 1H, H-1<sub>A</sub>), 5.34 (m, 1H, *J*<sub>trans</sub> = 17.2 Hz, =CH<sub>2</sub>), 5.34 (d, 1H, *J*<sub>1,2</sub> = 3.6 Hz, H-1<sub>E</sub>), 5.26 (m, 1H, *J*<sub>cis</sub> = 10.4 Hz, =CH<sub>2</sub>), 5.21 (d, 1H, *J* = 11.4 Hz, H<sub>Bn</sub>), 5.19–5.11 (m, 3H, H<sub>Bn</sub>, H-1<sub>B</sub>, H-2<sub>C</sub>), 5.13 (d, 1H, *J* = 12.7 Hz, H<sub>Bn</sub>), 5.07–4.88 (m, 7H, H<sub>Bn</sub>), 4.82–4.78 (m, 3H, 2H<sub>Bn</sub>, H-1<sub>B</sub>), 4.75–4.58 (m, 9H, H<sub>Bn</sub>, H-1<sub>C</sub>), 4.56 (d, 1H, *J*<sub>1,2</sub> = 8.8 Hz, H-1<sub>D</sub>), 4.45 (d, 1H, (d, 1H, *J* = 11.4 Hz, H<sub>Bn</sub>), 4.43 (d, 1H, *J* = 11.9 Hz, H<sub>Bn</sub>), 4.30–4.18 (m, 7H, H-3<sub>C</sub>, H-2<sub>A</sub>, H-3<sub>A</sub>, H-3<sub>E</sub>, H-5<sub>E</sub>, H-2<sub>B</sub>, H<sub>All</sub>), 4.05–3.70 (m, 7H, H<sub>All</sub>, H-2<sub>D</sub>, H-3<sub>B</sub>, H-3<sub>B</sub>, H-5<sub>C</sub>, H-4<sub>E</sub>, H-2<sub>E</sub>), 3.88 (dq, 1H, *J*<sub>4,5</sub> = 9.4 Hz, H-5<sub>B</sub>), 3.81 (dq, 1H, *J*<sub>4,5</sub> = 9.6 Hz, H-5<sub>B</sub>), 3.78 (dq, 1H, *J*<sub>4,5</sub> = 10.0 Hz, H-5<sub>A</sub>), 3.72 (bd, 1H, *J*<sub>6a,6b</sub> = 11.3 Hz, H-6<sub>AD</sub>), 3.61–3.52 (m, 5H, H-4<sub>B</sub>, H-6<sub>AE</sub>, H-6<sub>BE</sub>, H-4<sub>A</sub>, H-4<sub>C</sub>), 3.48 (pt, 1H, *J*<sub>3,4</sub> = 9.4 Hz, H-4<sub>B</sub>), 3.12–3.08 (m, 2H, H-4<sub>D</sub>, H-5<sub>D</sub>), 3.05 (m, 1H, H-6<sub>BD</sub>), 2.82–2.73 (m, 4H, H<sub>Lev</sub>), 2.24 (s, 3H, CH<sub>3Lev</sub>), 2.23 (s, 3H, H<sub>Ac</sub>), 2.19 (m, 1H, H-3<sub>D</sub>), 1.50 (d, 3H, *J*<sub>5,6</sub> = 6.3 Hz, H-6<sub>A</sub>), 1.46 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>B</sub>), 1.38 (d, 3H, *J*<sub>5,6</sub> = 6.1 Hz, H-6<sub>B</sub>), 1.37 (d, 3H, *J*<sub>5,6</sub> = 6.1 Hz, H-6<sub>C</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  206.1 (C<sub>Lev</sub>), 171.7 (C<sub>Lev</sub>), 170.0 (C<sub>Ac</sub>), 162.2 (C<sub>NNTCA</sub>), 138.8–137.6 (C<sub>Ph</sub>), 133.9 (CH=), 129.4–127.5 (CH<sub>Ph</sub>), 117.1 (=CH<sub>2</sub>), 101.1 (C-1<sub>D</sub>, <sup>1</sup>*J*<sub>CH</sub> = 165.4 Hz), 100.2 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 175.4 Hz), 99.1 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 171.8 Hz), 98.5 (C-1<sub>C</sub>, <sup>1</sup>*J*<sub>CH</sub> = 173.3 Hz), 98.3 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 168.3 Hz), 94.1 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 166.9 Hz), 93.3 (CCl<sub>3</sub>), 87.2 (C-3<sub>D</sub>), 83.1 (C-3<sub>E</sub>), 80.6 (C-3<sub>B</sub>), 80.5 (2C, C-2<sub>E</sub>, C-4<sub>B</sub>), 80.1 (C-4<sub>B</sub>), 79.7 (C-4<sub>A</sub>), 79.5 (C-4<sub>C</sub>), 78.8 (C-4<sub>E</sub>), 77.7 (C-3<sub>B</sub>), 77.3 (C-3<sub>C</sub>), 76.3 (C<sub>Bn</sub>), 75.6 (C-5<sub>D</sub>), 75.5, 75.4, 75.3, 75.0 (6C, C<sub>Bn</sub>), 74.1, 74.0 (C-2<sub>B</sub>, C-3<sub>A</sub>), 73.5, 72.9 (2C, C<sub>Bn</sub>), 72.1 (C-2<sub>C</sub>), 71.4 (C<sub>Bn</sub>),



71.3 (2C, C-2<sub>A</sub>, C-4<sub>D</sub>), 70.1 (C-5<sub>E</sub>), 69.4 (2C, C-2<sub>B'</sub>, C-5<sub>C</sub>), 68.9 (C-5<sub>A</sub>), 68.7 (C-5<sub>B'</sub>), 68.4 (C-5<sub>B</sub>), 68.0 (C-6<sub>E</sub>), 67.6 (C<sub>All</sub>), 62.8 (C-6<sub>D</sub>), 55.5 (C-2<sub>D</sub>), 38.2 (CH<sub>2Lev</sub>), 29.9 (CH<sub>3Lev</sub>), 28.3 (CH<sub>2Lev</sub>), 21.1 (C<sub>Ac</sub>), 18.1, 18.0, 17.8, 17.9 (4C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>B'</sub>, C-6<sub>C</sub>); HRMS (ESI<sup>+</sup>) for C<sub>118</sub>H<sub>134</sub>Cl<sub>3</sub>NO<sub>30</sub> ([M + Na]<sup>+</sup>, 2172.7954) found *m/z* 2172.8081.

**Allyl (3,4-Di-O-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-O-acetyl-4-O-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-deoxy-2-trichloroacetamido- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-O-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-(4-O-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-O-benzyl- $\alpha$ -L-rhamnopyranoside (52).** A solution of hydrazine hydrate (158  $\mu$ L, 3.2 mmol, 10 equiv) in pyridine/acetic acid (3:2, v/v, 5 mL) was added to diol **51** (700 mg, 320  $\mu$ mol) in pyridine (3 mL). After 25 min at 0 °C, TLC (Tol/EtOAc, 7:3) showed the complete disappearance of the diol and the presence of a major more polar product. Water (10 mL) and CH<sub>2</sub>Cl<sub>2</sub> (50 mL) were added, and the organic phase was washed with brine (3  $\times$  20 mL), water (3  $\times$  20 mL), dried on Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated to dryness. Chromatography of the residue (Tol/EtOAc, 9:1  $\rightarrow$  7:3) gave triol **52** (560 mg, 84%) as a white foam. Hexasaccharide **52** had *R<sub>f</sub>* = 0.35 (Tol/EtOAc, 7:3); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.48–7.07 (m, 50H, CH<sub>Ph</sub>), 7.02 (d, 1H, *J*<sub>NH,2</sub> = 8.6 Hz, NH), 5.90 (m, 1H, CH=), 5.40 (bs, 1H, H-1<sub>A</sub>), 5.30 (d, 1H, *J*<sub>1,2</sub> = 3.6 Hz, H-1<sub>E</sub>), 5.29 (m, 1H, *J*<sub>trans</sub> = 17.2 Hz, =CH<sub>2</sub>), 5.22 (m, 1H, *J*<sub>cis</sub> = 10.4 Hz, =CH<sub>2</sub>), 5.17 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 5.16–5.12 (m, 3H, H-1<sub>B'</sub>, H-2<sub>C</sub>, H<sub>Bn</sub>), 5.10 (d, 1H, *J* = 13.0 Hz, H<sub>Bn</sub>), 5.04 (d, 1H, *J* = 12.8 Hz, H<sub>Bn</sub>), 5.01 (d, 1H, *J* = 10.4 Hz, H<sub>Bn</sub>), 4.93 (d, 1H, *J* = 10.9 Hz, H<sub>Bn</sub>), 4.85–4.81 (m, 2H, H<sub>Bn</sub>), 4.77–4.66 (m, 7H, 3H<sub>Bn</sub>, H-1<sub>B</sub>, 3H<sub>Bn</sub>), 4.62–4.54 (m, 6H, 5H<sub>Bn</sub>, H-1<sub>C</sub>), 4.51 (d, 1H, *J*<sub>1,2</sub> = 8.4 Hz, H-1<sub>D</sub>), 4.38 (d, 1H, *J* = 11.9 Hz, H<sub>Bn</sub>), 4.23–4.13 (m, 7H, H-2<sub>B</sub>, H-3<sub>C</sub>, H-3<sub>E</sub>, H-3<sub>A</sub>, H<sub>All</sub>, H-5<sub>E</sub>, H-2<sub>A</sub>), 4.07 (dd, 1H, H-2<sub>B'</sub>), 4.00–3.87 (m, 6H, H<sub>All</sub>, H-2<sub>D</sub>, H-3<sub>B</sub>, H-5<sub>C</sub>, H-4<sub>E</sub>, H-2<sub>E</sub>), 3.84–3.81 (m, 1H, H-3<sub>B'</sub>, *J*<sub>2,3</sub> = 2.4 Hz, *J*<sub>3,4</sub> = 9.1 Hz, H-5<sub>B'</sub>), 3.78–3.69 (m, 2H, H-5<sub>B</sub>, H-5<sub>A</sub>), 3.65 (bd, 1H, *J*<sub>6a,6b</sub> = 11.2 Hz, H-6<sub>aD</sub>), 3.55–3.49 (m, 5H, H-6<sub>aE</sub>, H-6<sub>bE</sub>, H-4<sub>B</sub>, H-4<sub>A</sub>, H-4<sub>B'</sub>), 3.47 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.5 Hz, H-4<sub>C</sub>), 3.06–3.04 (m, 2H, H-4<sub>D</sub>, H-5<sub>D</sub>), 2.94 (dd, 1H, *J*<sub>5,6b</sub> = 5.7 Hz, H-6<sub>bD</sub>), 2.20 (s, 3H, H<sub>Ac</sub>), 2.15 (m, 1H, H-3<sub>D</sub>), 1.46 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>A</sub>), 1.42 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>B</sub>), 1.33–1.30 (m, 6H, H-6<sub>C</sub>, H-6<sub>B'</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  170.0 (C<sub>Ac</sub>), 162.3 (C<sub>NtCA</sub>), 138.7–137.4 (C<sub>Ph</sub>), 133.8 (CH=), 129.4–127.4 (CH<sub>Ph</sub>), 117.1 (=CH<sub>2</sub>), 101.1 (C-1<sub>D</sub>, <sup>1</sup>*J*<sub>CH</sub> = 161.0 Hz), 100.6 (C-1<sub>B'</sub>, <sup>1</sup>*J*<sub>CH</sub> = 171.0 Hz), 100.1 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 174.7 Hz), 98.3 (C-1<sub>C</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.4 Hz), 98.2 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.4 Hz), 94.0 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 163.1 Hz), 93.1 (CCl<sub>3</sub>), 87.1 (C-3<sub>D</sub>), 83.1 (C-3<sub>E</sub>), 80.5 (C-3<sub>B</sub>), 80.4 (C-4<sub>B</sub>), 80.3 (C-2<sub>E</sub>), 80.0 (C-4<sub>A</sub>), 79.7 (2C, C-3<sub>B'</sub>, C-4<sub>B'</sub>), 79.4 (C-4<sub>C</sub>), 78.7 (C-4<sub>E</sub>), 77.7 (C-3<sub>C</sub>), 76.3 (C<sub>Bn</sub>), 75.5 (C-5<sub>D</sub>), 75.4, 75.4, 75.3, 75.0, 74.9 (6C, C<sub>Bn</sub>), 74.1 (C-2<sub>A</sub>), 73.9 (C-3<sub>A</sub>), 73.5, 72.8 (2C, C<sub>Bn</sub>), 72.0 (C-2<sub>C</sub>), 71.7 (C-2<sub>B</sub>), 71.3 (C-4<sub>D</sub>), 71.0 (C-2<sub>B</sub>), 70.0 (C-5<sub>E</sub>), 69.1 (C-5<sub>C</sub>), 68.8 (2C, C-6<sub>B</sub>, C-5<sub>A</sub>), 68.4, 68.3 (2C, C-5<sub>B</sub>, C-5<sub>B'</sub>), 67.8 (C-6<sub>E</sub>), 67.6 (C<sub>All</sub>), 62.7 (C-6<sub>D</sub>), 55.3 (C-2<sub>D</sub>), 21.2 (C<sub>Ac</sub>), 18.0, 17.8 (4C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>B'</sub>, C-6<sub>C</sub>); HRMS (ESI<sup>+</sup>) for C<sub>113</sub>H<sub>128</sub>Cl<sub>3</sub>NO<sub>28</sub> ([M + Na]<sup>+</sup>, 2074.7585) found *m/z* 2074.7581, ([M + NH<sub>4</sub>]<sup>+</sup>, 2069.8032) found *m/z* 2069.7925.

**Propyl  $\alpha$ -L-Rhamnopyranosyl-(1 $\rightarrow$ 3)-(2-O-acetyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[ $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\alpha$ -L-rhamnopyranoside (5).** Hexasaccharide **52** (400 mg, 195  $\mu$ mol) was dissolved in EtOH (20 mL), treated with 10% Pd/C catalyst (400 mg), and the suspension was stirred at rt for 10 days, under a hydrogen atmosphere (45 bar). TLC (*i*PrOH/H<sub>2</sub>O/NH<sub>3</sub>, 4:1:0.5) and Tol/EtOAc, 7:3) showed that **52** had been transformed into a more polar product. The suspension was filtered on Acrodisc LC 25 mm, and the filtrate was concentrated. Reverse phase chromatography (H<sub>2</sub>O/CH<sub>3</sub>CN, 100:0  $\rightarrow$  70:30) of the residue, followed by freeze-drying, gave acetate **5** (131 mg, 64%) as a white foam. Hexasaccharide **5** had *R<sub>f</sub>* = 0.65 (*i*PrOH/H<sub>2</sub>O/NH<sub>3</sub>, 4:1:0.5); HPLC (215 nm): *t<sub>R</sub>* = 13.3 min (Kromasil 5  $\mu$ m C-18 100 $\text{\AA}$  4.6  $\times$  250 mm analytical column, using a 0–40% linear gradient over 20 min of CH<sub>3</sub>CN in 0.01 M aq TFA at 1 mL min<sup>-1</sup> flow rate); <sup>1</sup>H NMR

(D<sub>2</sub>O)  $\delta$  5.10 (d, 1H, *J*<sub>1,2</sub> = 3.7 Hz, H-1<sub>E</sub>), 5.00 (d, 1H, *J*<sub>1,2</sub> = 1.6 Hz, H-1<sub>A</sub>), 4.91 (dd, 1H, *J*<sub>1,2</sub> = 1.9 Hz, *J*<sub>2,3</sub> = 3.1 Hz, H-2<sub>C</sub>), 4.89 (d, 1H, *J*<sub>1,2</sub> = 1.4 Hz, H-1<sub>B'</sub>), 4.80 (bs, 2H, H-1<sub>B</sub>, H-1<sub>C</sub>), 4.71 (d, 1H, H-1<sub>D</sub>), 4.35 (dd, 1H, *J*<sub>2,3</sub> = 2.6 Hz, H-2<sub>A</sub>), 4.00 (dq, 1H, *J*<sub>4,5</sub> = 9.7 Hz, *J*<sub>5,6</sub> = 6.2 Hz, H-5<sub>C</sub>), 3.95 (m, 1H, H-5<sub>E</sub>), 3.93 (dd, 1H, *J*<sub>2,3</sub> = 3.4 Hz, H-2<sub>B'</sub>), 3.89–3.51 (m, 14H, H-3<sub>C</sub>, H-2<sub>B</sub>, H-3<sub>A</sub>, H-6<sub>aD</sub>, H-2<sub>D</sub>, H-3<sub>B</sub>, H-3<sub>E</sub>, H-6<sub>aE</sub>, H-6<sub>bE</sub>, H-5<sub>A</sub>, H-6<sub>bD</sub>, H-2<sub>E</sub>, H-5<sub>B</sub>, H-3<sub>B'</sub>, H<sub>Pr</sub>), 3.49 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.8 Hz, H-4<sub>C</sub>), 3.46–3.30 (m, 8H, H-4<sub>D</sub>, H<sub>Pr</sub>, H-5<sub>B'</sub>, H-4<sub>E</sub>, H-4<sub>B</sub>, H-4<sub>B'</sub>, H-5<sub>D</sub>, H-3<sub>D</sub>), 3.26 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.6 Hz, H-4<sub>A</sub>), 2.08 (s, 3H, H<sub>Ac</sub>), 2.03 (s, 3H, H<sub>NAC</sub>), 1.54–1.47 (m, 2H, CH<sub>2</sub>), 1.22–1.13 (m, 12H, H-6<sub>B</sub>, H-6<sub>A</sub>, H-6<sub>B'</sub>, H-6<sub>C</sub>), 0.82 (t, 3H, *J* = 7.4 Hz, CH<sub>3</sub>); <sup>13</sup>C NMR (D<sub>2</sub>O)  $\delta$  174.4 (C<sub>NAC</sub>), 173.2 (C<sub>Ac</sub>), 102.7 (C-1<sub>B'</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.6 Hz), 101.4 (C-1<sub>D</sub>, <sup>1</sup>*J*<sub>CH</sub> = 166.0 Hz), 101.2 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 173.5 Hz), 98.4 (C-1<sub>C</sub>, <sup>1</sup>*J*<sub>CH</sub> = 167.8 Hz), 98.1 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 171.5 Hz), 94.2 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 172.8 Hz), 82.6 (C-3<sub>D</sub>), 79.1 (C-2<sub>B</sub>), 77.0 (C-3<sub>C</sub>), 76.0 (C-5<sub>D</sub>), 74.1 (C-2<sub>A</sub>), 73.3 (C-3<sub>A</sub>), 73.0 (C-3<sub>E</sub>), 72.4 (C-2<sub>C</sub>), 71.9 (C-4<sub>B</sub>), 71.7 (C-4<sub>B'</sub>), 71.3, 71.2, 71.1 (C-5<sub>E</sub>, C-2<sub>E</sub>, C-4<sub>C</sub>), 70.7 (C-4<sub>A</sub>), 70.1 (C-3<sub>B'</sub>), 69.9 (2C, C-2<sub>B'</sub>, C-3<sub>B</sub>), 69.7 (C<sub>Pr</sub>), 69.4, 69.3 (3C, C-4<sub>E</sub>, C-5<sub>A</sub>, C-5<sub>B'</sub>), 68.8 (C-5<sub>C</sub>), 68.6 (C-5<sub>B</sub>), 68.1 (C-4<sub>D</sub>), 60.6 (C-6<sub>D</sub>), 60.3 (C-6<sub>E</sub>), 55.3 (C-2<sub>D</sub>), 22.7 (C<sub>NAC</sub>), 21.9 (CH<sub>2</sub>), 20.2 (C<sub>Ac</sub>), 16.8, 16.6, 16.5, 16.3 (4C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>B'</sub>, C-6<sub>C</sub>), 9.8 (CH<sub>3</sub>); HRMS (ESI<sup>+</sup>) for C<sub>43</sub>H<sub>73</sub>NO<sub>28</sub> ([M + H]<sup>+</sup>, 1052.4397) found *m/z* 1052.4360, ([M + Na]<sup>+</sup>, 1074.4216) found *m/z* 1074.4202.

**Allyl (3,4-Di-O-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(4-O-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 3)-(2-deoxy-2-trichloroacetamido- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[2,3,4,6-tetra-O-benzyl- $\alpha$ -D-glucopyranosyl-(1 $\rightarrow$ 3)]-(4-O-benzyl- $\alpha$ -L-rhamnopyranosyl)-(1 $\rightarrow$ 2)-3,4-di-O-benzyl- $\alpha$ -L-rhamnopyranoside (53).** Methanolic MeONa (0.5 M, 745  $\mu$ L, 370  $\mu$ mol, 1 equiv) was added to a solution of diol **51** (800 mg, 370  $\mu$ mol) in MeOH (20 mL), and the mixture was refluxed for 1 h. TLC (Tol/EtOAc, 6:4) showed the complete disappearance of the diol and the presence of a single more polar product. The mixture was neutralized by addition of Dowex X8-200 ion-exchange resin (H<sup>+</sup>), and filtered. Evaporation of the filtrate gave a syrup which was chromatographed (Tol/EtOAc, 8:2  $\rightarrow$  7:3) to give **53** (688 mg, 92%) as a white foam. Hexasaccharide **53** had *R<sub>f</sub>* = 0.45 (Tol/EtOAc, 6:4); <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.44–7.05 (m, 50H, CH<sub>Ph</sub>), 6.86 (d, 1H, *J*<sub>NH,2</sub> = 8.6 Hz, NH), 5.89 (m, 1H, CH=), 5.36 (bs, 1H, H-1<sub>A</sub>), 5.31–5.25 (m, 2H, =CH<sub>2</sub>, H-1<sub>E</sub>), 5.21 (m, 1H, *J*<sub>cis</sub> = 10.4 Hz, =CH<sub>2</sub>), 5.17 (d, 1H, *J*<sub>1,2</sub> = 1.6 Hz, H-1<sub>B'</sub>), 5.14 (d, 1H, *J* = 11.1 Hz, H<sub>Bn</sub>), 5.08 (m, 1H, *J* = 11.1 Hz, H<sub>Bn</sub>), 5.03–4.90 (m, 4H, H<sub>Bn</sub>), 4.82 (d, 1H, *J* = 11.0 Hz, H<sub>Bn</sub>), 4.74–4.52 (m, 13H, H-1<sub>B</sub>, H<sub>Bn</sub>), 4.50 (d, 1H, *J*<sub>1,2</sub> = 8.4 Hz, H-1<sub>D</sub>), 4.47 (d, 1H, *J*<sub>1,2</sub> = 1.6 Hz, H-1<sub>C</sub>), 4.38 (d, 1H, *J* = 11.9 Hz, H<sub>Bn</sub>), 4.19 (dd, 1H, *J*<sub>1,2</sub> = 2.1 Hz, *J*<sub>2,3</sub> = 2.4 Hz, H-2<sub>B</sub>), 4.17–4.12 (m, 5H, H<sub>All</sub>, H-5<sub>E</sub>, H-3<sub>E</sub>, H-3<sub>A</sub>, H-2<sub>A</sub>), 4.06–4.03 (m, 2H, H-3<sub>C</sub>, H-2<sub>B</sub>), 3.99–3.82 (m, 9H, H<sub>All</sub>, H-2<sub>C</sub>, H-2<sub>D</sub>, H-3<sub>B</sub>, H-5<sub>C</sub>, H-5<sub>B'</sub>, H-4<sub>E</sub>, H-2<sub>E</sub>, H-3<sub>B'</sub>), 3.78–3.70 (m, 2H, H-5<sub>B</sub>, H-5<sub>A</sub>), 3.65 (bd, 1H, *J*<sub>6a,6b</sub> = 11.4 Hz, H-6<sub>aD</sub>), 3.59–3.45 (m, 6H, H-4<sub>B</sub>, H-6<sub>aE</sub>, H-6<sub>bE</sub>, H-4<sub>B</sub>, H-4<sub>A</sub>, H-4<sub>C</sub>), 3.08–3.03 (m, 2H, H-5<sub>D</sub>, H-4<sub>D</sub>), 2.94 (m, 1H, H-6<sub>bD</sub>), 2.23 (dd, 1H, *J*<sub>2,3</sub> = 9.7 Hz, *J*<sub>3,4</sub> = 7.7 Hz, H-3<sub>D</sub>), 1.44 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>B</sub>), 1.43 (d, 3H, *J*<sub>5,6</sub> = 6.1 Hz, H-6<sub>A</sub>), 1.40 (d, 3H, *J*<sub>5,6</sub> = 6.2 Hz, H-6<sub>B</sub>), 1.32 (d, 3H, *J*<sub>5,6</sub> = 6.1 Hz, H-6<sub>C</sub>); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  161.9 (C<sub>NtCA</sub>), 138.7–137.5 (C<sub>Ph</sub>), 133.9 (CH=), 129.1–127.3 (CH<sub>Ph</sub>), 117.1 (=CH<sub>2</sub>), 101.0 (C-1<sub>D</sub>, <sup>1</sup>*J*<sub>CH</sub> = 162.7 Hz), 100.8 (C-1<sub>C</sub>, <sup>1</sup>*J*<sub>CH</sub> = 169.2 Hz), 100.1 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 180.3 Hz), 100.0 (C-1<sub>B'</sub>, <sup>1</sup>*J*<sub>CH</sub> = 180.3 Hz), 98.3 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.3 Hz), 94.0 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 168.6 Hz), 93.2 (CCl<sub>3</sub>), 86.5 (C-3<sub>D</sub>), 83.0 (C-3<sub>E</sub>), 80.5 (C-3<sub>B</sub>), 80.4 (2C, C-2<sub>E</sub>, C-4<sub>B</sub>), 79.8, 79.7, 79.6 (4C, C-4<sub>A</sub>, C-3<sub>B'</sub>, C-4<sub>B'</sub>, C-4<sub>C</sub>), 78.8 (C-4<sub>E</sub>), 78.3 (C-3<sub>C</sub>), 76.3 (C<sub>Bn</sub>), 75.5 (C-5<sub>D</sub>), 75.4, 75.3, 75.2, 74.9 (6C, C<sub>Bn</sub>), 74.0 (C-3<sub>A</sub>), 73.8 (C-2<sub>A</sub>), 73.5, 72.8, 72.0 (3C, C<sub>Bn</sub>), 71.3 (C-2<sub>B</sub>), 71.1 (C-4<sub>D</sub>), 70.4 (C-2<sub>C</sub>), 70.0 (C-5<sub>E</sub>), 68.9 (C-5<sub>C</sub>), 68.8 (2C, C-5<sub>A</sub>, C-2<sub>B'</sub>), 68.7 (C-5<sub>B'</sub>), 68.3 (C-5<sub>B</sub>), 67.9 (C-6<sub>E</sub>), 67.6 (C<sub>All</sub>), 62.7 (C-6<sub>D</sub>), 55.8 (C-2<sub>D</sub>), 18.1, 18.0, 17.8 (4C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>B'</sub>, C-6<sub>C</sub>); HRMS (ESI<sup>+</sup>) for C<sub>111</sub>H<sub>126</sub>Cl<sub>3</sub>NO<sub>27</sub> ([M + Na]<sup>+</sup>, 2032.7480) found *m/z* 2032.7448.

**Propyl  $\alpha$ -L-Rhamnopyranosyl-(1 $\rightarrow$ 3)- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 3)-(2-acetamido-2-deoxy- $\beta$ -D-glucopyranosyl)-(1 $\rightarrow$ 2)-[ $\alpha$ -D-**

**glucopyranosyl-(1→3)]- $\alpha$ -L-rhamnopyranosyl-(1→2)- $\alpha$ -L-rhamnopyranoside (6).** Hexasaccharide **53** (580 mg, 289  $\mu$ mol) was dissolved in EtOH (15 mL), treated with 10% Pd/C catalyst (340 mg), and the suspension was stirred at rt for 10 days, under a hydrogen atmosphere (45 bar). TLC (*i*PrOH/H<sub>2</sub>O/NH<sub>3</sub>, 4:1:0.5 and Tol/EtOAc, 6:4) showed that **53** had been transformed into a more polar product. The suspension was filtered on Acrodisc LC 25 mm, and the filtrate was concentrated. Reverse phase chromatography (H<sub>2</sub>O/CH<sub>3</sub>CN, 100:0  $\rightarrow$  70:30) of the residue, followed by freeze-drying, gave the target **6** (240 mg, 82%) as a white foam. Hexasaccharide **6** had *R<sub>f</sub>* = 0.2 (*i*PrOH/H<sub>2</sub>O/NH<sub>3</sub>, 4:1:0.5); HPLC (215 nm): *t<sub>R</sub>* = 13.4 min (Kromasil 5  $\mu$ m C-18 100 $\text{\AA}$  4.6  $\times$  250 mm analytical column, using a 0–40% linear gradient over 20 min of CH<sub>3</sub>CN in 0.01 M aq TFA at 1 mL min<sup>-1</sup> flow rate); <sup>1</sup>H NMR (D<sub>2</sub>O)  $\delta$  5.20 (d, 1H, *J*<sub>1,2</sub> = 3.7 Hz, H-1<sub>E</sub>), 5.10 (d, 1H, *J*<sub>1,2</sub> = 1.6 Hz, H-1<sub>A</sub>), 5.01 (d, 1H, *J*<sub>1,2</sub> = 1.2 Hz, H-1<sub>B</sub>), 4.90 (bs, 1H, H-1<sub>B</sub>), 4.84 (d, 1H, *J*<sub>1,2</sub> = 1.9 Hz, H-1<sub>C</sub>), 4.83 (d, 1H, *J*<sub>1,2</sub> = 9.5 Hz, H-1<sub>D</sub>), 4.44 (dd, 1H, *J*<sub>2,3</sub> = 2.1 Hz, H-2<sub>A</sub>), 4.07–3.99 (m, 3H, H-2<sub>B</sub>, H-5<sub>E</sub>, H-5<sub>C</sub>), 3.96–3.93 (m, 2H, H-2<sub>B</sub>, H-3<sub>A</sub>), 3.92–3.68 (m, 14H, H-6<sub>aD</sub>, H-2<sub>C</sub>, H-2<sub>D</sub>, H-3<sub>B</sub>, H-3<sub>E</sub>, H-3<sub>B</sub>, H-6<sub>aE</sub>, H-6<sub>bE</sub>, H-3<sub>C</sub>, H-5<sub>B</sub>, H-5<sub>A</sub>, H-6<sub>bD</sub>, H-2<sub>E</sub>, H-5<sub>B</sub>), 3.64 (dt, *J* = 9.8 Hz, *J* = 6.9 Hz, H<sub>Pr</sub>), 3.55–3.38 (m, 8H, H<sub>Pr</sub>, H-4<sub>C</sub>, H-4<sub>D</sub>, H-4<sub>E</sub>, H-3<sub>D</sub>, H-4<sub>B</sub>, H-4<sub>B</sub>, H-5<sub>D</sub>), 3.35 (pt, 1H, *J*<sub>3,4</sub> = *J*<sub>4,5</sub> = 9.7 Hz, H-4<sub>A</sub>), 2.11 (s, 3H, H<sub>NAc</sub>), 1.65–1.56 (m, 2H, CH<sub>2</sub>), 1.31 (d, 3H, *J*<sub>5,6</sub> = 6.3 Hz, H-6<sub>B</sub>), 1.28 (d, 3H, *J*<sub>5,6</sub> = 6.3 Hz, H-6<sub>B</sub>), 1.26 (d, 3H, *J*<sub>5,6</sub> = 6.3 Hz, H-6<sub>A</sub>), 1.23 (d, 3H, *J*<sub>5,6</sub> = 6.3 Hz, H-6<sub>C</sub>), 0.91 (t, 3H, *J* = 7.4 Hz, CH<sub>3</sub>); <sup>13</sup>C NMR (D<sub>2</sub>O)  $\delta$  174.0 (C<sub>NAc</sub>), 102.4 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 171.2 Hz), 101.5 (C-1<sub>D</sub>, <sup>1</sup>*J*<sub>CH</sub> = 163.6 Hz), 101.3 (C-1<sub>C</sub>, <sup>1</sup>*J*<sub>CH</sub> = 169.8 Hz),

101.2 (C-1<sub>A</sub>, <sup>1</sup>*J*<sub>CH</sub> = 169.8 Hz), 98.2 (C-1<sub>B</sub>, <sup>1</sup>*J*<sub>CH</sub> = 170.5 Hz), 94.4 (C-1<sub>E</sub>, <sup>1</sup>*J*<sub>CH</sub> = 171.2 Hz), 81.7 (C-3<sub>D</sub>), 79.1 (C-2<sub>B</sub>), 78.1 (C-3<sub>C</sub>), 76.0 (C-5<sub>D</sub>), 74.2 (C-2<sub>A</sub>), 73.5 (C-3<sub>A</sub>), 73.2 (C-3<sub>E</sub>), 72.2 (C-4<sub>B</sub>), 72.0 (C-4<sub>B</sub>), 71.4 (C-5<sub>E</sub>), 71.3 (C-2<sub>E</sub>), 71.3 (C-4<sub>C</sub>), 70.8 (C-4<sub>A</sub>), 70.5 (C-2<sub>C</sub>), 70.2 (C-3<sub>B</sub>), 70.2 (C-2<sub>B</sub>), 70.0 (C-3<sub>B</sub>), 69.8 (C<sub>Pr</sub>), 69.5 (C-4<sub>E</sub>), 69.4 (C-5<sub>A</sub>), 69.0 (2C, C-5<sub>C</sub>, C-5<sub>B</sub>), 68.7 (C-5<sub>B</sub>), 68.3 (C-4<sub>D</sub>), 60.7 (C-6<sub>D</sub>), 60.4 (C-6<sub>E</sub>), 55.5 (C-2<sub>D</sub>), 22.6 (C<sub>NAc</sub>), 21.9 (CH<sub>2</sub>), 16.8, 16.7, 16.5 (4C, C-6<sub>A</sub>, C-6<sub>B</sub>, C-6<sub>B</sub>, C-6<sub>C</sub>), 9.9 (CH<sub>3</sub>); HRMS (ESI<sup>+</sup>) for C<sub>41</sub>H<sub>71</sub>NO<sub>27</sub> ([M + H]<sup>+</sup>, 1010.4292) found *m/z* 1010.4295, ([M + Na]<sup>+</sup>, 1032.4111) found *m/z* 1032.4116.

**Acknowledgment.** The authors thank the Agence Nationale pour la Recherche (ANR, Grant NT05-1\_42479), the Institut Pasteur, and the Ministère de l'Éducation Nationale, de la Recherche et de la Technologie (MENRT, fellowship to J.B.) for financial support to this work. The authors also thank F. Bonhomme (CNRS URA 2128) for running the NMR and mass spectra. Dr. Yves Janin (CNRS URA 2128) is acknowledged for proofreading this manuscript.

**Supporting Information Available:** General experimental procedures, <sup>1</sup>H, <sup>13</sup>C, and HMBC NMR spectra for compounds **1–6**, **10–12**, **15**, **16**, **19–21**, **23**, **25**, **28–32**, **37**, **44**, **45**, **47**, **48**, **50–53**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO802127Z