


# Near-Infrared Absorbing Azo Dyes: Synthesis and X-ray Crystallographic and Spectral Characterization of Monoazopyrroles, Bisazopyrroles, and a Boron–Azopyrrole Complex

Yan Li, Brian O. Patrick, and David Dolphin\*

Department of Chemistry, The University of British Columbia, 2036 Main Mall, Vancouver, B.C., V6T 1Z1, Canada

david.dolphin@ubc.ca

Received February 11, 2009



Symmetric 2,5-bisazopyrroles **2(a–d)** were synthesized by a one-step reaction of substituted phenyl diazonium salts  $[\text{R}'(\text{Ph})\text{N}_2^+\text{Cl}^-]$  [**a**,  $\text{R}' = 4\text{-N}(\text{CH}_3)_2$ ; **b**,  $\text{R}' = 2\text{-OH}$ ; **c**,  $\text{R}' = 2\text{-CO}_2\text{H}$ ; **d**,  $\text{R}' = 4\text{-NO}_2$ ] with pyrrole under basic conditions. Asymmetric 2,5-bisazopyrroles **3(a–d)** were synthesized by reacting substituted phenyl diazonium salts  $[\text{R}''(\text{Ph})\text{N}_2^+\text{Cl}^-]$  (**a**,  $\text{R}'' = 4\text{-OCH}_3$ ; **b**,  $\text{R}'' = \text{H}$ ; **c**,  $\text{R}'' = 4\text{-Br}$ ; **d**,  $\text{R}'' = 4\text{-NO}_2$ ) with 2-(4-dimethylaminophenylazo)-1*H*-pyrrole (**1a**) under the same conditions. The reactions of **2a** with boron trifluoride and iodomethane provided a  $\text{BF}_2$ -azopyrrole complex of 1*H*-pyrrolo[2,1-*c*]-1,2,4,5-boratriazole (**4**) and 2,5-bisazo-1-methylpyrrole **5**. X-ray crystallographic and spectral analysis of **1a**, **2a**, **2b**, and **4** showed that **1a** has three crystal forms: **1a(I)**, **1a(II)**, and **1a(III)**, the latter two bearing a bicyclic ring system formed via intermolecular hydrogen bonding. Complex **4** was found to be the most planar due to a rigid trans-azo configuration and has the longest N=N bond distances (1.322 and 1.300 Å) and wavelength of maximum absorption (754 nm). The N=N bond distances increase in the sequence of monoazopyrrole [**1a(I)**: 1.253 Å], bisazopyrrole (**2a**: 1.283 Å), bisazopyrrole with intramolecular hydrogen bonding (**2b**: 1.293 and 1.293 Å), and the  $\text{BF}_2$ -azopyrrole complex. Their maximum absorptions shift bathochromically in the sequence of monoazopyrrole (**1a**: 443 nm), bisazopyrroles [**2(a–d)**, **3(a–d)**, **5**: 486–615 nm], and the  $\text{BF}_2$ -azopyrrole complex. These results are important for the design of near-infrared absorbing azo dyes and suggest an efficient path for the preparation of near-infrared absorbing azo dyes by effectively enhancing  $\pi$ -electron delocalization.

## Introduction

Azo dyes are a class of compounds containing a N=N double bond and, due to their ability to absorb visible light, and ease of synthesis, have been extensively used in the textile, fiber, leather, paint and printing industries for more than a century.<sup>1</sup> Among the known azo dyes, five-membered heterocyclic azo dyes, such as azothiazole, azothiophene, azopyrrole, and azofuran, are important since they have pronounced bathochromic absorptions compared to azo-

benzene dyes<sup>2</sup> and are environmentally friendly.<sup>3</sup> Near-infrared absorbing azo dyes, based on five-membered heterocyclic rings, have been made effectively by using three methods: (1) introducing electron-donating or -accepting groups to the azo-linked aromatic ring to increase molecular

(1) Herbst, W.; Hunger, K. *Industrial Organic Pigments: Production, Properties, Applications* 2nd; VCH: Weinheim, Germany, 1997.

(2) (a) Weaver, M. A.; Shuttleworth, L. *Dyes Pigm.* **1982**, *3*, 81. (b) Sekar, N. *Colourage* **1994**, *41*, 38. (c) Towns, A. D. *Dyes Pigm.* **1999**, *42*, 3. (d) Ledoux, I.; Zyss, J.; Barni, E.; Barolo, C.; Diulgheroff, N.; Quagliotto, P.; Viscardi, G. *Synth. Met.* **2000**, *115*, 213. (e) Li, Z.; Zhao, Y.; Zhou, J.; Shen, Y. *Eur. Polym. J.* **2000**, *36*, 2417.

(3) (a) Freeman, H. S.; Hinks, D.; Esancy, J. *Physico-Chemical Principles of Color Chemistry*; Peters, A. T., Freeman, H. S., Eds.; Blackie Academic and Professional: London, UK, 1996; 254 (b) Sekar, N. *Colourage* **2004**, *51*, 58.

donor–acceptor polarization;<sup>4</sup> (2) expanding the azo-linked aromatic ring from benzene to naphthalene and from thiazole to benzothiazole;<sup>5</sup> and (3) extending the conjugated N=N bond from monoazo to bisazo dyes to enhance molecular  $\pi$ -resonance effects.<sup>6</sup> The near-infrared absorbing azo heterocycles have been found to be useful as optical data carriers,<sup>4c,4f,6b</sup> thermal transfer recording media,<sup>4c,5a,5b,6c</sup> electrophotographic toners,<sup>4b,4d,6f</sup> and semiconductor lasers.<sup>5d,6e</sup>

While numerous syntheses and applications of five-membered heterocyclic azo dyes have been reported, studies on the relationship between molecular structure and color by theoretical and experimental methods have also received much interest. Most of the theoretical work addresses molecular donor–acceptor polarization,<sup>7–9</sup> with relatively less interest in the donor–acceptor polarization enhanced by  $\pi$ -resonance effects of expanded aromatic rings and extended N=N bonds.<sup>9</sup> Experimental studies with X-ray crystallography have been limited since it has been difficult to compare differences in crystallographic structures and the resulting change in color. In 2004, McNelis and co-workers<sup>10</sup> successfully investigated the relationship between the donor–acceptor polarization of azothiophenes and their colors using X-ray crystallography. 2-(4-Diethylaminophenylazo) thiophene has a N=N bond distance of 1.226 Å; when the strong electron-accepting tricyanovinyl group was introduced, the resulting products, 2-(4-diethylamino)phenylazo-5-tricyanovinylthiophene and 2-(4-tricyanovinyl)-phenylazo-5-diethylaminothiophene, exhibited N=N bond distances of 1.297 and 1.314 Å, with maximum absorptions at 719 and 708 nm, respectively. Regrettably, the relationship between  $\pi$ -resonance, planarity, and colors for five-membered heterocyclic azo dyes has not been explored theoretically nor via X-ray crystallography. In 2004, Centore and co-workers<sup>11</sup> investigated [4-di(hydroxyethyl)aminophenyl]azobenzenes containing a benzimidazole group. When molecular planarity was

significantly improved by the insertion of a vinylene group between [4-di(hydroxyethyl)aminophenyl]azobenzene and benzimidazole, the  $\lambda_{\max}$  was bathochromically shifted from 466 to 487 nm. This result indicated, not surprisingly, the importance of  $\pi$ -resonance effects on the bathochromic absorptions.

Most azo dyes are prepared via coupling of aromatic diazonium salts with a nucleophilic component. While this is true for 2,5-bisazathiophenes which are prepared from thiophen-2-yl diazonium salts,<sup>12</sup> the corresponding analogues of 2-aminothiophene<sup>13</sup> and 2,5-diaminothiophene<sup>14</sup> require multicomponent condensations; pyrrole, on the other hand, is more reactive than thiophene or even aniline (which is  $10^{20}$  times more reactive than benzene). Few 2,5-bisazopyrroles have been reported<sup>15–17</sup> and we report here the coupling to pyrrole to produce 2,5-bis(aryloxy)pyrroles.

A series of symmetric 2,5-bis(aryloxy)-1*H*-pyrroles [**2(a–d)**] and asymmetric 2,5-bis(aryloxy)-1*H*-pyrroles [**3(a–d)**] have been prepared and their maximum absorptions appear between 486 and 615 nm in dichloromethane, which are longer than that of monoazopyrrole **1a** (447 nm). When **2a** was reacted with iodomethane, symmetric 2,5-bis(aryloxy)-1-methylpyrrole (**5**) was obtained, which has a similar absorption at 563 nm. However, when **2a** was reacted with boron trifluoride, the resulting boron–azopyrrole complex **4** displayed a near-infrared absorption at 754 nm. Comparisons of the X-ray crystallographic structures of **1a**, **2a**, **2b**, and **4** show that **4** is the most planar and has the longest N=N bond distance (1.322 and 1.300 Å), whereas **1a**, **2a**, and **2b** have N=N bond distances at 1.253 [**1a(I)**], 1.286 [**1a(II)**], 1.289 [**1a(III)**], 1.283 (**2a**), and 1.293 and 1.293 Å (**2b**).

## Results and Discussion

**Synthesis of 2,5-Bisazopyrrole and Derivatives.** Following the general procedure to prepare azo dyes, substituted phenyl diazonium salts were formed by reacting aromatic amines [ $R'PhNH_2$ ;  $R' = 4-N(CH_3)_2$ , 4-OCH<sub>3</sub>, 2-OH, H, 4-Br, 2-CO<sub>2</sub>H, 4-NO<sub>2</sub>] with sodium nitrite/aqueous HCl at temperatures lower than 0 °C. Neutralization with pyridine and treatment with a half equivalent of pyrrole in methanol gave symmetric bisazopyrroles **2(a–d)** [ $R' = 4-N(CH_3)_2$ , 2-OH, 2-CO<sub>2</sub>H, 4-NO<sub>2</sub>] in 61–82% yields (Scheme 1, path 1). Monoazopyrrole **1a** was the major product when the 4-dimethylaminophenyl diazonium salt reacted with excess pyrrole under the same conditions (Scheme 1, path 2). Further reactions of **1a** with substituted phenyl diazonium salts provided asymmetric bisazopyrroles **3(a–d)** ( $R'' = 4-OCH_3$ , H, 4-Br, 4-NO<sub>2</sub>)

(4) (a) Bello, K. A.; Griffiths, J. *J. Chem. Soc., Chem. Commun.* **1986**, 1639. (b) Takiguchi, T. JP 63029761, 1988, CAN 109:83367. (c) JP 07214928, 1995, CAN 124:18488. (d) Maeda, S.; Mito, K. JP 08029993, 1996, CAN 124:356154. (e) Nishimoto, T.; Misawa, T.; Sugimoto, K.; Tsuda, T.; Umehara, H.; Takuma, H. JP 09040659, 1997, CAN 126:218690. (f) Berneth, H.; Bruder, F.-K.; Haese, W.; Hagen, R.; Hassenrueck, K.; Kostromine, S.; Landenberger, P.; Oser, R.; Sommermann, T.; Stawitz, J.-W.; Bieringer, T. WO 2002080152, 2002, CAN 137:286559. (g) Griffiths, J.; Lee, W. J. *Adv. Colloid Sci. Technol.* **2002**, 5, 99.

(5) (a) Nakayama, N.; Komamura, T.; Tanaka, T. JP 05005969, 1993, CAN 119:37592. (b) Kawakami, S.; Nakajima, A.; Maejima, K.; Komamura, T. JP 07172059, 1995, CAN 123:343738. (c) Hall, N.; Patel, P. GB 2237804, 1991, CAN 117:9806. (d) Emandi, A.; Serban, I.; Bandula, R. *Dyes Pigm.* **1999**, 41, 63. (e) Hattori, R. JP 11119415, 1999, CAN 130:345070.

(6) (a) Gregory, P. EP 280434, 1988, CAN 110:116654. (b) Umehara, H.; Yanagimachi, M.; Taniguchi, Y.; Hirose, S.; Misawa, T.; Takuma, H. JP 08108625, 1996, CAN 125:127897. (c) Slark, A. T.; Fox, J. E. *Polymer* **1997**, 38, 2989. (d) Matsumoto, S.; Yagisawa, T.; Matsumoto, S.; Shimabukuro, K. JP 11269136, 1999, CAN 131:272843. (e) Hattori, R. JP 11119415, 1999, CAN 130:345070. (f) Yoshida, S.; Nakamura, Y.; Takesawa, S. JP 2005099419, 2005, CAN 142:400497.

(7) (a) Albert, I. D. L.; Morley, J. O.; Pugh, D. *J. Phys. Chem.* **1995**, 99, 8024. (b) Astrand, P.-O.; Sommer-Larsen, P.; Hvilsted, S.; Ramanujam, P. S.; Bak, K. L.; Sauer, S. P. A. *Chem. Phys. Lett.* **2000**, 325, 115. (c) Astrand, P.-O.; Ramanujam, P. S.; Hvilsted, S.; Bak, K. L.; Sauer, S. P. A. *J. Am. Chem. Soc.* **2000**, 122, 3482. (d) Astrand, P.-O.; Bak, K. L.; Sauer, S. P. A. *Chem. Phys. Lett.* **2001**, 343, 171.

(8) Morley, J. O. *J. Phys. Chem.* **1994**, 98, 13177.

(9) (a) Morley, J. O.; Pugh, D. *J. Chem. Soc., Faraday Trans.* **1991**, 87, 3021. (b) Morley, J. O.; Naji, M.; Hutchings, M. G.; Hall, N. *J. Phys. Chem. A* **1998**, 102, 5802.

(10) Moylan, C. R.; McNelis, B. J.; Nathan, L. C.; Marques, M. A.; Hermstad, E. L.; Brichler, B. A. *J. Org. Chem.* **2004**, 69, 8239.

(11) Carella, A.; Centore, R.; Fort, A.; Peluso, A.; Sirigu, A.; Tuzi, A. *Eur. J. Org. Chem.* **2004**, 12, 2620.

(12) (a) Mitsubishi Chem. Ind. JP 57111356, 1982, CAN 98:18401. (b) Niwa, T.; Himeno, K.; Maeda, S.; Shimizu, Y. DE 3301822, 1983, CAN 99:177495. (c) Ito, N.; Aiga, H.; Nishihara, M. JP 61028556, 1986, CAN 105:135409. (d) Etzbach, K. H.; Hansen, G.; Schefczik, E.; Reichelt, H. DE 3622297, 1988, CAN 108:169160. (e) Haehnke, M.; Mueller, J.; Boos, M.; Hett, H. EP 305887, 1989, CAN 111:155818.


(13) Sabnis, R. W.; Rangnekar, D. W.; Sonawane, N. D. *J. Heterocycl. Chem.* **1999**, 36, 333.

(14) (a) Middleton, W. J.; Engelhardt, V. A.; Fisher, B. S. *J. Am. Chem. Soc.* **1958**, 80, 2822. (b) Skene, W. G.; Trefz, T. *Polym. Prepr.* **2004**, 45, 563.

(15) Rozovskii, Y. T.; Savvin, S. B.; Khakimkhodzhaev, N.; Pachadzhanov, D. N.; Yusupov, M. *Dokl. Akad. Nauk Tadzh. SSR* **1968**, 11, 40.

(16) Wagner-Wysiecka, E.; Luboch, E.; Kowalczyk, M.; Biernat, J. F. *Tetrahedron* **2003**, 59, 4415.

(17) (a) Savvin, S. B.; Rozovskii, Y. G.; Propistsova, R. F.; Likhonina, E. A. *Izv. Akad. Nauk SSSR, Ser. Khim* **1969**, 6, 1364. (b) Depraetere, S.; Dehaen, W. *Tetrahedron Lett.* **2003**, 44, 345. (c) Takahashi, M.; Suzuki, T.; Ikemizu, H.; Ikesu, S. JP 2004307633, 2004, CAN 141:396980.

**SCHEME 1. Synthesis of Symmetric 2,5-Bisazopyrroles 2(a–d) and Asymmetric 2,5-Bisazopyrroles 3(a–d)**



in 62–79% yields (Scheme 1, path 3). **1a** can be converted into 2-(4-dimethylaminophenylazo)-5-nitro-1*H*-pyrrole (**1b**) by tritiation at  $-50^{\circ}\text{C}$ .<sup>18</sup>

When **2a** was mixed with boron trifluoride and diisopropylethylamine, under nitrogen, and refluxed in dichloromethane for 1 h, the corresponding  $\text{BF}_2$ -azopyrrole complex **4** was formed in 85% yield (Scheme 2, path 1). Complex **4** was purified by chromatography and was found to be stable in nonprotic solvents such as acetone, tetrahydrofuran, acetonitrile, dichloromethane, benzene, etc., but unstable in protic solvents such as methanol. When **2a** was refluxed with iodomethane and sodium carbonate in acetone for 3 h, 2,5-bisazo-1-methylpyrrole **5** was obtained in 91% yield (Scheme 2, path 2).

While monoazopyrrole **1a** is yellow, these new bisazopyrroles [**2(a–d)**, **3(a–d)**, and **5**] are red to blue, and the boron–azopyrrole complex **4** is green in dichloromethane.

**X-ray Crystal Structures.** The structures of **1a**, **2a**, **2b**, and **4** were determined by X-ray crystallography and are shown in Figure 1. Their selected bond lengths and angles are listed in Table 1. Crystal data and structure refinements are summarized in the Supporting Information. While 2-phenylazo-1-vinylpyrrole (**6**)<sup>19</sup> and a macrocyclic 2,3-bisazopyrrole<sup>16</sup> have been reported with X-ray crystal structures, and some boron–azo complexes have been identified by NMR, IR, and MS spectra,<sup>20,21</sup> **1a**, **2a**, **2b**, and **4** are the first examples of monoazo-1*H*-pyrroles, 2,5-bisazo heterocycles (thiophene, pyrrole, and furan), and boron–azo complexes reported with X-ray crystal structures.

Monoazopyrrole **1a** has three forms within a unit cell: **1a(I)** shows no hydrogen bonding while **1a(II)** and **1a(III)** are both intermolecularly hydrogen bonded to each other between the nitrogen-1 of pyrrole and the nitrogen of an azo bond with  $\text{N}-\text{H}\cdots\text{N}$  bond distances of 3.020 ( $\text{N1}\cdots\text{N7}$ ) and 3.087 Å ( $\text{N3}\cdots\text{N5}$ ). Compound **2a** has crystallographic mirror symmetry with identical  $\text{H4n}-\text{N4}-\text{C9}$  and  $\text{H4n}-\text{N4}-\text{C9}^*$  angles of  $125.8^{\circ}$ . Compound **2b** is slightly asymmetric, it has two intramolecular hydrogen bonds between

**SCHEME 2. Reactions of 2,5-Bisazopyrrole (2a) with Boron Trifluoride and Iodomethane**


the oxygen of the *o*-hydroxy groups and the nitrogen of the azo bond ( $=\text{N}$ -pyrrole) with  $\text{O}-\text{H}\cdots\text{N}$  bond distances of 2.629 ( $\text{O1}\cdots\text{N2}$ ) and 2.601 Å ( $\text{O2}\cdots\text{N4}$ ). Intramolecular hydrogen bonding has been reported for *o*-hydroxy monoazo dyes, where the azophenol tautomers have  $\text{H}\cdots\text{N}(\text{azo})$  bond distances of 1.66–1.92 Å<sup>22</sup> and the hydrazoketone tautomers have  $\text{H}-\text{N}(\text{hydrazone})$  bond distances of 0.84–1.24 Å.<sup>22,23</sup> The  $\text{H}\cdots\text{N}$  bond distances of 1.839 ( $\text{H1o}\cdots\text{N2}$ ) and 1.819 Å ( $\text{H2o}\cdots\text{N4}$ ) indicate that **2b** principally exists as an azophenol tautomer. Intramolecular hydrogen bonding between N-1 of pyrrole and a nitrogen of the azo bond for **1a(I)**, **2a**, and **2b** should be very weak if it exists at all as the distances between the N-1 proton of pyrrole and the nitrogen of the azo bond [**1a(I)**: 2.592 Å; **2a**: 2.518 and 2.518 Å; **2b**: 2.458 and 2.568 Å] are slightly smaller than the sum of the van der Waals radii of hydrogen (1.20 Å) and nitrogen (1.50 Å) atoms.<sup>24</sup> Surprisingly, the N-1 proton of **1a(I)** is rotated away from the nitrogen of the azo bond, since the  $\text{H9n}-\text{N9}-\text{C28}$  angle ( $127.7^{\circ}$ ) is larger than the  $\text{H9n}-\text{N9}-\text{C25}$  angle ( $122.8^{\circ}$ ). For the  $\text{BF}_2$ -azopyrrole complex **4**, the nitrogen of one azo bond ( $\text{Ar}-\text{N}=\text{N}$ ) is coordinated to the N-1- $\text{BF}_2$  substituent at the pyrrole to form the novel structure 1*H*-pyrrolo[2,1-*c*]-1,2,4,5-boratriazole. The boron atom is  $\text{sp}^3$  hybridized with an  $\text{F1}-\text{B1}-\text{F2}$  angle ( $112.0^{\circ}$ ) divided by the  $\text{B1}/\text{N2}/\text{N3}/\text{C9}/\text{N4}$  plane. The B–N bond distances are 1.537 ( $\text{B1}-\text{N4}$ ) and 1.646 Å ( $\text{B1}-\text{N2}$ ).

The structure of **1a(I)** is nearly planar, where the dihedral angle between the pyrrole and phenyl ( $\text{C29}/\text{C30}/\text{C31}/\text{C32}/\text{C33}/\text{C34}$ ) planes is only  $5.36^{\circ}$ . As shown by side-view drawings in Figure 2, bisazopyrrole **2a** is less coplanar and butterfly like, with identical dihedral angles between the pyrrole and phenyl planes ( $\text{C3}/\text{C4}/\text{C5}/\text{C6}/\text{C7}/\text{C8}$  and  $\text{C3}^*/\text{C4}^*/\text{C5}^*/\text{C6}^*/\text{C7}^*/\text{C8}^*$ ) being  $16.59^{\circ}$ , which are larger than the dihedral angles of the phenyl planes in bisazo aromatic dyes ( $2.3$ – $14.1^{\circ}$ ).<sup>25</sup> Intramolecular hydrogen-bonded **2b** is slightly ruffled, with the angles of the corresponding two

(22) (a) Olivieri, A. C.; Wilson, R. B.; Paul, I. C.; Curtin, D. Y. *J. Am. Chem. Soc.* **1989**, *111*, 5525. (b) Conner, J. A.; Kennedy, R. J.; Dawes, H. M.; Hursthouse, M. B.; Walker, N. P. C. *J. Chem. Soc., Perkin Trans. 2* **1990**, *1*, 203.

(23) (a) Alder, M. J.; Flower, K. R.; Pritchard, R. G. *Tetrahedron Lett.* **1998**, *39*, 3571. (b) You, X.-L.; Lu, C.-R.; Huang, Z.-L.; Zhang, D.-C. *Dyes Pigm.* **2004**, *63*, 217. (c) Luboch, E.; Kravtsov, V. C. *J. Mol. Struct.* **2004**, *699*, 9. (d) Gilli, P.; Bertolasi, V.; Pretto, L.; Lycka, A.; Gilli, G. *J. Am. Chem. Soc.* **2002**, *124*, 13554.

(24) Ladd, M. F. C. *Structure and Bonding in Solid State Chemistry*; Ellis Horwood Ltd: New York, 1979; p 253.

(25) (a) Gilardi, R. D.; Karle, I. L. *Acta Crystallogr.* **1972**, *B28*, 1635. (b) Freeman, H. S.; McIntosh, S. A.; Singh, P. *Dyes Pigm.* **1997**, *35*, 11. (c) Freeman, H. S.; McIntosh, S. A.; Singh, P. *Dyes Pigm.* **1997**, *35*, 149.

(18) Morgan, K. J.; Morrey, D. P. *Tetrahedron* **1966**, *22*, 57.

(19) Trofimov, B. A.; Schmidt, E. Y.; Mikhaleva, A. I.; Vasil'tsov, A. M.; Zaitsev, A. B.; Smolyanina, N. S.; Senotrusova, E. Y.; Afonin, A. V.; Ushakov, I. A.; Petrusenko, K. B.; Kazheva, O. N.; Dyachenko, A. A.; Smirnov, V. V.; Schmidt, A. F.; Markova, M. V.; Morozova, L. V. *Eur. J. Org. Chem.* **2006**, 4021.

(20) Hohaues, E.; Wessendorf, K. Z. *Naturforsch. B: Anorg. Chem., Org. Chem.* **1980**, *35*, 319.

(21) (a) Das, M. K.; Bandyopadhyay, S. N. *J. Chem. Res., Synop.* **1991**, *4*, 83. (b) Dedkov, Y. M.; Lozinskaya, E. F. *Izv. Vyssh. Uchebn. Zaved., Khim. Khim. Tekhnol.* **2002**, *45*, 118.




FIGURE 1. ORTEP views of the structures **1a**, **2a**·CH<sub>2</sub>Cl<sub>2</sub>, **2b**, and **4** (hydrogen atoms of **1a** are not shown).

TABLE 1. Selected Bond Distances (Å) and Angles (deg) of the Structures **1a**, **2a**·CH<sub>2</sub>Cl<sub>2</sub>, **2b**, and **4**

	<b>1a(I)</b>	<b>1a(II)</b>	<b>1a(III)</b>	<b>2a</b> ·CH <sub>2</sub> Cl <sub>2</sub>	<b>2b</b>	<b>4</b>
H–N(pyrrole)	0.84(2)	0.90(2),	0.88(2)	0.84(3)	0.87(2)	
B–N(pyrrole,azo)						1.537(2), 1.646(2)
H...N(azo)		2.142	2.226		1.839, 1.819	
N–N	1.253(2)	1.286(2)	1.289(2)	1.283(2)	1.293(1), 1.293(1)	1.300(2), 1.322(2)
N(azo)–C(pyrrole)	1.4269(2)	1.386(2)	1.390(2)	1.380(2)	1.379(2), 1.380(2)	1.361(2), 1.377(2),
N(azo)–C(phenyl)	1.453(2)	1.423(2)	1.420(2)	1.404(2)	1.408(2), 1.412(2)	1.392(2), 1.398(2)
N–N–C(pyrrole)	110.9(2)	114.2(1)	113.8(1)	113.4(1)	113.1(1), 114.2(1)	112.5(1), 106.7(1),
N–N–C(phenyl)	112.1(2)	114.8(1)	115.0(1)	113.6(1)	116.1(1), 115.0(1)	114.4(1), 119.9(1)
X(B,C)–N4–C						143.2(1), 109.8(1), 106.9(1)
X(B,N)–N2–Y(C,N)						113.8(1), 126.3(1), 119.9(1)

planes being 5.66° (pyrrole and C1/C2/C3/C4/C5/C6 plane) and 6.43° (pyrrole and C11/C12/C13/C14/C15/C16 plane). Significantly, complexation improves the planarity of bisazopyrrole molecules; complex **4** is the most planar where the dihedral angles between the pyrrole and phenyl planes are 4.73° (pyrrole and C3/C4/C5/C6/C7/C8 plane) and 3.53° (pyrrole and C13/C14/C15/C16/C17/C18 plane). The mean deviations of carbon and nitrogen atoms from the  $\pi$ -delocalized skeletons containing azo, phenyl, and pyrrole groups are 0.047 Å in **1a**, 0.203 Å in **2a**, 0.065 Å in **2b**, and 0.045 Å in **4**. The small proton chemical shift differences of **2a** between 25 and –80 °C further indicate that the torsion angles

between the aromatic planes do not significantly change [0.048, 0.096, 0.048, 0.087, and 0.044 ppm for HN (pyrrole), HC (benzene), HC (pyrrole), HC (benzene), and H<sub>3</sub>C, respectively; see the Supporting Information].

All configurations for N=N–C(pyrrole) are in a trans-cis form. **1a(I)** exhibits a N=N bond distance of 1.253 Å, which is longer than those of simple monoazo dyes with thiophene (1.226 Å)<sup>10</sup> and shorter than those of **6** with enhanced  $\pi$ -resonance effects (1.271 Å)<sup>19</sup> and monoazothiophenes with electron-withdrawing groups (1.297–1.314 Å).<sup>10</sup> Inter- and intramolecular hydrogen bonds and the second azo bond extend the N=N bond distances to 1.286 Å for




FIGURE 2. Side views of the structures of **1a(I)**, **2a**, **2b**, and **4**.

TABLE 2. Absorption Data of Azopyrroles in Dichloromethane and Methanol

comps	$\lambda_{\max}$ (log $\epsilon$ ) in CH <sub>2</sub> Cl <sub>2</sub>	$\lambda_{\max}$ (log $\epsilon$ ) in MeOH
<b>1a</b>	447 (4.79)	443
<b>1b</b>	530	498
<b>2a</b>	575 (4.77)	574 (4.79)
<b>2b</b>	519 (4.62)	507 (4.66)
<b>2c</b>	486	466 (4.64)
<b>2d</b>	490 (4.60)	487 (4.57)
<b>3a</b>	548 (4.69)	543 (4.74)
<b>3b</b>	558 (4.69)	543 (4.71)
<b>3c</b>	572 (4.68)	554 (4.71)
<b>3d</b>	615 (4.69)	589 (4.70)
<b>4</b>	754 (4.78)	
<b>5</b>	563 (4.81)	580

**1a(II)**, 1.289 Å for **1a(III)**, 1.283 Å for **2a**, and 1.293 and 1.293 Å for **2b**. The boron–azopyrrole complex **4** has distinguishingly long N=N bond lengths. The free N=N bond distance is 1.300 Å and the rigid N=N bond distance is 1.322 Å, longer than the calculated N–N bond distance for 2-arylhydrazone–pyrrole structures (1.316 Å).<sup>8</sup> In contrast, **1a(I)** clearly has long N–C(pyrrole) and N–C(Ar) bond distances. The N–C(pyrrole) bond distances are 1.426 Å for **1a(I)**, 1.386 Å for **1a(II)**, 1.390 Å for **1a(III)**, 1.380 Å for **2a**, 1.391 and 1.388 Å for **2b**, and 1.361 and 1.377 Å for **4**, and the N–C(Ar) bond distances are 1.453 Å for **1a(I)**, 1.423 Å for **1a(II)**, 1.420 Å for **1a(III)**, 1.404 Å for **2a**, 1.408 and 1.412 Å for **2b**, and 1.398 and 1.392 Å for **4**, decreasing in the sequence of monoazo, bisazo, and boron–azo complex, suggesting that complex **4** has the best  $\pi$ -electron delocalization.

**UV–Vis Absorption Spectra of 2,5-Bisazopyrrole and Derivatives.** The maximum absorptions of azopyrroles **1(a,b)**, **2(a–d)**, **3(a–d)**, **4**, and **5** are listed in Table 2. For monoazo dyes, the visible absorption spectra are dominated by molecular donor–acceptor polarization. 2-(4-Dimethylaminophenyl) azo-1-*H*-pyrrole (**1a**) has a maximum absorption at 447 nm in dichloromethane or 443 nm in methanol, longer than those for (4-dimethylaminophenyl)azobenzene (408 nm)<sup>26</sup> and


2-(4-nitrophenyl)azo-1-*H*-pyrrole (420 nm),<sup>27</sup> which indicate that pyrrole acts as an effective acceptor in 2-arylazopyrrole chromophores. When the donor–acceptor polarization of **1a** is increased by introducing an electron-accepting nitro group into the 5-position of the pyrrole, the maximum absorption of **1b** is bathochromically shifted to 530 nm in dichloromethane and to 498 nm in methanol. Furthermore, when the  $\pi$ -resonance is enhanced by extending the monoazopyrrole to bisazopyrroles, asymmetric 2,5-bisazopyrroles **3a**, **3b**, **3c**, and **3d** have their maximum absorptions at 548, 558, 572, and 615 nm in dichloromethane and at 543, 543, 554, and 589 nm in methanol, respectively. The maximum absorption shifts bathochromically along with the increase in the electron-accepting ability of substituents in the sequence of *p*-OCH<sub>3</sub> (**3a**) < *p*-H (**3b**) < *p*-Br (**3c**) < *p*-NO<sub>2</sub> (**3d**), consistent with an increase in molecular donor–acceptor polarization. Note that the difference of the maximum absorptions of bisazopyrroles from **3b** to **3d** is smaller than that of monoazopyrroles from **1a** to **1b**. Symmetric 2,5-bisazopyrroles **2a**, **2b**, **2c**, **2d**, and **5** also exhibit maximum absorptions at 575, 519, 486, 490, and 563 nm in dichloromethane and at 574, 507, 466, 487, and 580 nm in methanol, respectively, larger than those of **1a**. However, the maximum absorption shifts bathochromically along with the increase of the electron-donating ability of substituents on both sides in the sequence *p*-N(CH<sub>3</sub>)<sub>2</sub> (**2a**) > *o*-OH (**2b**) > *p*-NO<sub>2</sub> (**2d**) > *o*-CO<sub>2</sub>H (**2c**).

Typically, as shown in Figure 3 for **3a** and **3c**, the maximum absorption peaks for some bisazopyrroles overlap with a minor absorption peak. Similar spectral characteristics have been found for *o*-hydroxy azo dyes and reflect the azohydrazone tautomeric equilibria in solution,<sup>28</sup> where the absorption peaks to the red belong to the hydrazone tautomer and the other to the azo tautomer. Similar tautomeric equilibria, through intramolecular N(azo)···H···N (pyrrole) bonding, should exist for monoazo- and bisazo-1-*H*-pyrroles as depicted in Scheme 3. The positions of tautomeric equilibria are affected by solvents and substituents; in addition, protic solvents can weaken the intramolecular

(26) Sawicki, E. *J. Org. Chem.* **1957**, *22*, 915.

(27) Ahmad, A. K.; Hassan, Y. I.; Bashir, W. A. *Analyst* **1987**, *112*, 97.

(28) Isak, S. J.; Eyring, E. M.; Spikes, J. D.; Meekins, P. A. *J. Photochem. Photobiol., A* **2000**, *134*, 77.




**FIGURE 3.** Absorption spectra of **1a** (orange line), bisazopyrrole **3a** (red line) and **3c** (blue line), and boron–azopyrrole complex **4** (green line) in dichloromethane.

hydrogen bonding and shift the equilibrium toward the azo tautomer.

For dichloromethane solutions of **3c** and **3d**, minor absorption peaks appear as shoulders on the blue side of the maximum absorption peaks, and tend to strengthen with a decrease in the electron-accepting ability of substituents in the sequence of **3b** (H) > **3c** (Br) > **3d** (NO<sub>2</sub>). In contrast, for **2a** and **3a**, the minor absorption peaks appear as a shoulder on the red side of the maximum absorption peaks, and tend to weaken with an increase in the electron-donating ability of substituents in the sequence **3a** (OCH<sub>3</sub>) < **3b** (H). Compared to those in dichloromethane, the maximum absorptions in methanol solutions show large hypsochromic shifts of 15 nm for **3b**, 18 nm for **3c**, 26 nm for **3d**, and 32 nm for **1b**, versus small shifts of 1 nm for **2a**, 3 nm for **2d**, 4 nm for **1a**, and 5 nm for **3a**. The hypsochromic shifts of the maximum absorptions by 12 nm for **2b** and 20 nm for **2c** are due to the disruption of the intramolecular N(azo)···H···O bonding. These spectral characteristics indicate that the hydrazone tautomers predominate in dichloromethane solution for monoazo- and unsymmetric 2,5-bisazo-1*H*-pyrroles with strong donor–acceptor polarization, such as **1b**, **3b**, **3c**, and **3d**, and the azo tautomers are predominant for monoazo- and 2,5-bisazo-1*H*-pyrroles with non- and weak donor–acceptor polarization, such as **1a**, **2a**, **2b**, **2d**, and **3a**, which are consistent with the X-ray crystal structures. An attempt to capture the hydrazone tautomer by reacting **2a** with MeI failed; only 2,5-bisazo-1-methylpyrrole **5** was isolated in high yield. The spectrum of **5** in dichloromethane also has a minor absorption peak appearing as a shoulder on the red side of the maximum absorption peak, similar to that of 2-aryloxy-1-vinylpyrroles.<sup>19</sup>

Significantly, the boron–azopyrrole complex **4** has its maximum absorption at 754 nm in dichloromethane (Figure 3), bathochromically shifted by 179 nm relative to that of **2a**. This presumably results from significant  $\pi$ -electron delocalization in the planar structure dominated

### SCHEME 3. Tautomerization Proposed for Monoazo- and Bisazo-1*H*-pyrroles



by complexation of the electron-withdrawing BF<sub>2</sub> group. Since symmetric bisazobenzidines<sup>28</sup> and BF<sub>2</sub> complexes of dipyrromethene<sup>29</sup> and tetraarylazadipyrromethene<sup>30</sup> have been investigated as photosensitizers for photodynamic therapy, the BF<sub>2</sub>–azo complex described here is a potential photosensitizer in this field.

### Conclusion

Symmetric 2,5-bisazopyrroles **2(a–d)** and **5**, asymmetric 2,5-bisazopyrroles **3(a–d)**, and the novel BF<sub>2</sub>–azopyrrole complex of 1*H*-pyrrolo[2,1-*c*]-1,2,4,5-borotriazole (**4**) have been synthesized and fully characterized. The near-infrared absorption of BF<sub>2</sub>–azo complex **4** is dominated by  $\pi$ -resonance effects which was achieved by extending the conjugation around the N=N bond and forming a rigid azo configuration. These results are important for the design of near-infrared absorbing azo dyes and provide an effective and simple method to synthesize such dyes by enhancing  $\pi$ -electron delocalization. The synthesis of proton-inert complexes of 1*H*-pyrrolo[2,1-*c*]-1,2,4,5-borotriazoles and their application as photosensitizers for photodynamic therapy are under progress.

### Experimental Section

**Preparation of 2,5-Bis(4-dimethylaminophenylazo)-1*H*-pyrrole (**2a**).** To a suspension of the aromatic amine (8.0 mmol) in water (15 mL) was added concentrated hydrochloric acid (24 mmol, 2 mL) until the mixture was homogeneous. The solution was cooled and kept at 0–5 °C in an ice bath and diazotized by addition of a solution of sodium nitrite (8.2 mmol, 0.566 g) in water (5 mL), followed by stirring for 30 min at 0–5 °C. To a solution of pyrrole (4 mmol, 0.28 mL) and pyridine (50 mmol, 4 mL) in methanol (150 mL) was slowly added a solution of the diazonium salt at 0–5 °C. The resulting mixture was stirred for 5 h and then evaporated under vacuum to dryness. The residues were purified by column chromatography, eluting with a mixture of dichloromethane and hexane (1:1). Yield 82%. Mp 228–229 °C. Anal. Calcd for C<sub>20</sub>H<sub>23</sub>N<sub>7</sub>·<sup>1</sup>/<sub>3</sub> CH<sub>2</sub>Cl<sub>2</sub> (solvent of crystallization determined by NMR): C, 62.66; H, 6.12; N, 25.16. Found: C, 62.41; H, 6.52; N, 25.52. <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>)  $\delta$  9.77 (s, 1H), 7.80 (s, 4H), 6.88 (s, 2H), 6.77 (d, *J* = 8.9 Hz, 4H), 3.10 (s, 12H). <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  9.77 (s, 1H),

(29) (a) Yogo, T.; Urano, Y.; Ishitsuka, Y.; Maniwa, F.; Nagano, T. *J. Am. Chem. Soc.* **2005**, *127*, 12162. (b) Atilgan, S.; Ekmekci, Z.; Dogan, A. L.; Guç, D.; Akkaya, E. U. *Chem. Commun.* **2006**, *42*, 4398.

(30) Gorman, A.; Killoran, J.; O'Shea, C.; Kenna, T.; Gallagher, W. M.; O'Shea, D. F. *J. Am. Chem. Soc.* **2004**, *126*, 10619.

7.78 (d,  $J=8.3$  Hz, 4H), 6.89 (s, 2H), 6.73 (d,  $J=8.3$  Hz, 4H), 3.09 (s, 12H).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  7.74 (d,  $J=9.1$  Hz, 4H), 6.81 (s, 2H), 6.80 (d,  $J=9.2$  Hz, 4H), 3.07 (s, 12H).  $^{13}\text{C}$  NMR ( $\text{CD}_2\text{Cl}_2$ )  $\delta$  152.7, 146.5, 144.5, 124.9, 114.3, 112.3, 40.3. IR  $\nu$  ( $\text{N}=\text{N}$ )  $1377\text{ cm}^{-1}$ . UV-vis ( $1.03 \times 10^{-5}$  M,  $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 411 (4.34), 575 (4.77). UV-vis ( $1.41 \times 10^{-5}$  M,  $\text{CH}_3\text{OH}$ )  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 407 (4.22), 574 (4.79). EI MS  $m/z$  361 ( $\text{M}^+$ ).

**Preparation of 2-(4-Dimethylaminophenylazo)-1H-pyrrole (1a).**<sup>31</sup> **1a** was prepared by using the same procedure as for **2a** but with 2 equiv of pyrrole and it was purified by column chromatography eluting with a mixture of dichloromethane and hexane (1:2). Mp 147–148 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  9.33 (s, 1H), 7.75 (d,  $J=9.2$  Hz, 2H), 6.84 (d,  $J=3.6$  Hz, 1H), 6.79 (s, 1H), 6.73 (d,  $J=9.2$  Hz, 2H), 6.32 (t,  $J=3.2$  Hz, 1H), 3.03 (s, 6H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  151.8, 146.3, 143.6, 124.1, 119.9, 112.1, 112.0, 111.0, 40.6. IR  $\nu$  ( $\text{N}=\text{N}$ )  $1377\text{ cm}^{-1}$ . UV-vis ( $1.64 \times 10^{-5}$  M,  $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$  447 (4.52). UV-vis ( $\text{CH}_3\text{OH}$ )  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 443. EI MS  $m/z$  214 ( $\text{M}^+$ ).

**Preparation of 2-(4-Dimethylaminophenylazo)-5-nitro-1H-pyrrole (1b).** This compound was prepared according to the literature<sup>18</sup> by nitration of **1a** with a mixture of acetic anhydride and nitric acid at  $-50$  °C. Dark crystals. Yield 32%. Mp 208–209 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  9.93 (s, 1H), 7.80 (d,  $J=9.2$  Hz, 2H), 7.17 (d,  $J=4.4$  Hz, 1H), 6.74 (d,  $J=4.0$  Hz, 1H), 6.20 (d,  $J=8.97$  Hz, 2H), 3.11 (s, 6H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ ) 153.5, 146.7, 143.4, 126.1, 114.9, 112.8, 111.9, 110.5, 40.5. IR:  $\nu$  ( $\text{N}=\text{N}$ )  $1376\text{ cm}^{-1}$ . UV-vis ( $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$  530. UV-vis ( $\text{CH}_3\text{OH}$ )  $\lambda_{\text{max}}$  498. HR-EI MS ( $\text{M}^+$ )  $m/z$  calcd for  $\text{C}_{12}\text{H}_{13}\text{N}_5\text{O}_2$  259.1069, found 259.1071.

**Preparation of 2-(4-Methoxyphenylazo)-5-(4-dimethylamino-phenylazo)-1H-pyrrole (3a).** **3a** was prepared by using the same procedure as for **2a** but with 1 equiv of **1a** instead of pyrrole and purified by column chromatography eluting with a mixture of dichloromethane and hexane (2:3). Yield 79%. Mp 166–167 °C. Anal. Calcd for  $\text{C}_{19}\text{H}_{20}\text{N}_6\text{O}$ : C, 65.50; H, 5.79; N, 24.12. Found: C, 65.15; H, 5.85; N, 24.39.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  9.74 (s, 1H), 7.83 (d,  $J=8.8$  Hz, 2H), 7.80 (d,  $J=8.2$  Hz, 2H), 7.01–6.95 (m, 3H), 6.92 (d,  $J=4.1$  Hz, 1H), 6.76–6.71 (m, 2H), 3.87 (s, 3H), 3.08 (s, 6H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  161.6, 152.4, 147.6, 146.7, 145.5, 144.2, 125.0, 124.3, 116.2, 114.6, 114.0, 112.0, 55.8, 40.5. IR  $\nu$  ( $\text{N}=\text{N}$ )  $1380\text{ cm}^{-1}$ . UV-vis ( $1.31 \times 10^{-5}$  M,  $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$

(log  $\epsilon$ ) 384 (4.20), 548 (4.69). UV-vis ( $1.24 \times 10^{-5}$  M,  $\text{CH}_3\text{OH}$ )  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 382 (4.18), 543 (4.74). EI MS  $m/z$  348 ( $\text{M}^+$ ).

**Preparation of 2,5-Bis(4-dimethylaminophenylazo)pyrrole-1-boron Difluoride (4).** To a solution of **2a** (0.28 mmol, 100 mg) and diisopropylethylamine (4.2 mmol, 0.73 mL) in dry dichloromethane (50 mL) was slowly added boron trifluoride diethyl etherate (1.4 mmol, 0.18 mL) under nitrogen. The resulting solution was refluxed for 1 h and then evaporated under vacuum to dryness. The residue was purified by column chromatography eluting with dichloromethane and hexane (1:1) to give a green product. Yield 85%. Mp 205–206 °C. Anal. Calcd for  $\text{C}_{20}\text{H}_{22}\text{BF}_2\text{N}_7$ : C, 58.70; H, 5.42; N, 23.96. Found: C, 58.99; H, 5.59; N, 24.14.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.89 (d,  $J=9.20$  Hz, 4H), 6.94 (s, 2H), 6.72 (d,  $J=9.30$  Hz, 4H), 3.10 (s, 12H).  $^{11}\text{B}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.62. IR  $\nu$  ( $\text{N}=\text{N}$ )  $1422\text{ cm}^{-1}$ . UV-vis ( $1.12 \times 10^{-5}$  M,  $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 466 (4.26), 754 (4.78). ESI MS  $m/z$  409 ( $\text{M}^+$ ).

**Preparation of 2,5-Bis(4-dimethylaminophenylazo)-1-methylpyrrole (5).** To a mixture of **2a** (0.28 mmol, 100 mg) and potassium carbonate (1.68 mmol, 232 mg) in dry acetone (50 mL) was added iodomethane (0.84 mmol, 52  $\mu\text{L}$ ) under nitrogen. The resulting solution was refluxed for 3 h and then evaporated under vacuum to dryness. The residue was purified by column chromatography eluting with a mixture of dichloromethane and hexane (1:1). Yield 91%. Mp 242–243 °C. Anal. Calcd for  $\text{C}_{21}\text{H}_{25}\text{N}_7$ : C, 67.18; H, 6.71; N, 26.11. Found: C, 66.87; H, 6.70; N, 26.28.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.81 (d,  $J=8.98$  Hz, 4H), 6.73 (d,  $J=9.0$  Hz, 6H), 4.18 (s, 3H), 3.07 (s, 12H).  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta$  151.8, 148.4, 145.2, 124.4, 112.0, 101.0, 40.6, 29.5. IR  $\nu$  ( $\text{N}=\text{N}$ )  $1378\text{ cm}^{-1}$ . UV-vis ( $1.24 \times 10^{-5}$  M,  $\text{CH}_2\text{Cl}_2$ )  $\lambda_{\text{max}}$  (log  $\epsilon$ ) 563 (4.81), 767 (4.13). UV-vis ( $\text{CH}_3\text{OH}$ )  $\lambda_{\text{max}}$  580. EI MS  $m/z$  375 ( $\text{M}^+$ ).

**Acknowledgment.** This work is supported by QLT Inc., Vancouver, BC, and the Natural Sciences and Engineering Council (NSERC) of Canada. We thank the NMR, Mass Spectroscopy and Elemental Analysis Laboratories of the Chemistry Department at the University of British Columbia.

**Supporting Information Available:** Experimental procedures, NMR and UV-vis spectra, and detailed X-ray crystallographic characterizations. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(31) Kalle, A.-G. NL 6407379, 1965, CAN 62:79569.