


# Perylenequinone Natural Products: Enantioselective Synthesis of the Oxidized Pentacyclic Core<sup>†</sup>

Carol A. Mulrooney, Barbara J. Morgan, Xiaolin Li, and Marisa C. Kozlowski\*

Department of Chemistry, Roy and Diana Vagelos Laboratories, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323

marisa@sas.upenn.edu

Received June 30, 2009



An enantioselective approach to the perylenequinone core found in the mold perylenequinone natural products is outlined. Specifically, the first asymmetric syntheses of helical chiral perylenequinones absent any additional stereogenic centers are described. Key elements of the synthetic venture include a catalytic enantioselective biaryl coupling, a PIFA-induced naphthalene hydroxylation, and a palladium-mediated aromatic decarboxylation. Transfer of the binaphthalene axial stereochemistry to the perylenequinone helical stereochemistry proceeded with good fidelity. Furthermore, the resultant perylenequinones were shown to possess sufficient atropisomeric stability to be viable intermediates in the biogenesis of the perylenequinone natural products. This stability supports the use of the helical axis as a stereochemical relay in synthesis of the natural products containing additional stereochemical centers.

## Introduction

The parent perylenequinone **1** was first prepared by Zinke in 1929,<sup>1</sup> but it was not until 1954 that the correct structure was elucidated (Figure 1).<sup>2</sup> This compound was subsequently isolated from the fungus *Daldinia concentrica* as blue-black crystals and is the simplest of the naturally occurring perylenequinones.<sup>3</sup> Most members of this family of natural products fall into three classes (examples of each in Figure 1): (a) C<sub>20</sub> compounds without carbon substituents,<sup>4</sup> (b) mold perylenequinones containing carbon substituents, and (c) perylenequinones from aphids. Of the three, class B is the most prevalent. Due to their diversity, interesting stereochemistries, and anticancer activities we have undertaken the syntheses of these compounds via dimerization of achiral naphthols.




FIGURE 1. Representative structures of the perylenequinone classes.

The archetypical cercosporin **6** (Figure 2) was among the first of the mold perylenequinones to be isolated in 1957.<sup>5</sup> The

<sup>†</sup> Dedicated to the memory of Ralph Hirschmann.

(1) Zinke, A.; Hirsch, W.; Brozek, E. *Monatsch. Chem.* **1929**, *51*, 205–220.  
(2) Calderbank, A.; Johnson, A. W.; Todd, A. R. *J. Chem. Soc.* **1954**, 1285–1289.

(3) Anderson, J. M.; Murray, J. *Chem. Ind.* **1956**, 376–376.

(4) This class also includes some partially reduced compounds from the molds of *Alternaria* and *Stemphylium*.

(5) Kuyama, S.; Tamura, T. *J. Am. Chem. Soc.* **1957**, *79*, 5725–5726.




FIGURE 2. Mold perylenequinones 2–17.

unusual features of this molecule hindered a complete structural determination until the 1970s.<sup>6</sup> The majority of the structural details were determined by classical redox quinone chemistry, but the most significant finding was the transformation involving the thermal atropisomerization of (*M*)-cercosporin into (*P*)-*epi*-cercosporin. The opposite helical configuration of the perylenequinone epimers was confirmed by circular dichroism (CD) analysis,<sup>6b</sup> and the helical nature along with the absolute and relative configuration of **6** was established crystallographically.<sup>7</sup> Since the discovery of **6**, more than 20 perylenequinones have been isolated as dark red pigments from a variety of fungi within the *Ascomycota* phylum (Figure 2).<sup>8</sup> Correlation with the CD and NMR spectra of **6** has allowed determination of the helical configurations and relative stereochemistries of most of the remaining natural products.

Notably, all of the mold perylenequinones (Figure 2) are derivatives of **1** (Figure 1). With the exception of hypomycin B (**15**), the C4,C4'- or C5,C5'-phenols are not alkylated. These acidic phenols ( $pK_a \sim 7.3\text{--}8.3$ )<sup>9</sup> are characteristic of the structures and form strong internal hydrogen bonds with the quinone carbonyls.<sup>10</sup> Many of the mold perylenequinones are  $C_2$ -symmetric, having C7,C7'-substitution with centrochiral stereocenters of the same absolute configuration. The remainder (hypocrellin, hypocrellin A–B, shiraiachrome A, elsinochrome B<sub>1</sub>–B<sub>2</sub>, and hypomycin B<sup>11</sup>) likely originated from  $C_2$ -symmetric intermediates.

Regardless of the varied C7,C7'-substitution, all class B perylenequinones contain a conjugated pentacyclic core in a highly oxidized state. Furthermore, the distinct oxygenation pattern at the C2,C2',C4,C4',C5,C5',C6,C6'-centers (Figure 2) is conserved throughout the series. In the past two decades, the construction of this highly oxygenated core en route to

the perylenequinone natural products has been examined by several groups.<sup>12–19</sup> Prior to these achievements, there had been relatively few synthetic efforts devoted to this family of natural products. As mentioned above, **1** (class A, Figure 1) was first unknowingly prepared by Zinke and co-workers (ca. 1929) from tetranitroperylene **18** upon exposure to sulfuric acid at 140 °C under air.<sup>1</sup> Due to the atypical stability of the proposed product **19**, Calderbank and co-workers repeated the procedure (Scheme 1),<sup>2</sup> and the structure of 4,4',5,5'-tetraquinone **19** reported by Zinke was correctly reassigned as the parent perylenequinone **1**.

Surprisingly, it was not until 1972 that there was further progress when Weisgraber and Weiss attempted to synthesize the oxygenation pattern present in the mold perylenequinones (Figure 2). The initial assignments of both the coupling product **21** and the subsequent perylenequinone **22** were discovered to be incorrect,<sup>20,21</sup> and the final cyclized product was reassigned as dinaphthofuranedione **24** (Scheme 2).

In work by Dallacker and Leidig, a different approach was used in the synthesis of 2,2',4,4'-tetramethoxyperylenequinone, commencing with formation of bisphenyl substrate **25**

(12) Chao, C.; Zhang, P. *Tetrahedron Lett.* **1988**, *29*, 225–226.

(13) Diwu, Z.; Lown, J. W. *Tetrahedron* **1992**, *48*, 45–54. (b) Liu, J.; Diwu, Z.; Lown, J. W. *Tetrahedron* **1993**, *49*, 10785–10792.

(14) Broka, C. A. *Tetrahedron Lett.* **1991**, *32*, 859–862.

(15) Hauser, F. M.; Sengupta, D.; Corlett, S. A. *J. Org. Chem.* **1994**, *59*, 1967–1969.

(16) (a) Coleman, R. S.; Grant, E. B. *J. Am. Chem. Soc.* **1994**, *116*, 8795–8796. (b) Coleman, R. S.; Grant, E. B. *J. Am. Chem. Soc.* **1995**, *117*, 10889–10904.

(17) (a) Merlic, C. A.; Aldrich, C. C.; Albaneze-Walker, J.; Saghatelian, A. *J. Am. Chem. Soc.* **2000**, *122*, 3224–3225. (b) Merlic, C. A.; Aldrich, C. C.; Albaneze-Walker, J.; Saghatelian, A.; Mammen, J. *J. Org. Chem.* **2001**, *66*, 1297–1309.

(18) O'Brien, E. M.; Morgan, B. J.; Kozlowski, M. C. *Angew. Chem., Int. Ed.* **2008**, *47*, 6877–6880.

(19) Morgan, B. J.; Dey, S.; Johnson, S. W.; Kozlowski, M. C. *J. Am. Chem. Soc.* **2009**, *131*, 9413–9425.

(20) (a) Weisgraber, K. H.; Weiss, U. *J. Chem. Soc., Perkin Trans. 1* **1972**, 83–88. (b) Weiss, U.; Fales, H. M.; Weisgraber, K. H. *Liebigs Ann. Chem.* **1979**, 914–919. (c) Fales, H. M.; Weiss, U.; Jaouni, T. *Liebigs Ann. Chem.* **1983**, 367–371.

(21) The proposed structure of **22** is the keto–enol tautomer of the mold perylenequinones seen in Figure 2, with the exception of hypomycin B, **15**. The perylenequinones actually exist as a mixture of the two tautomers, though the predominance of each tautomeric form is different for each natural product: Arnone, A.; Merlini, L.; Mondelli, R.; Nasini, G.; Ragg, E.; Scaglioni, L.; Weiss, U. *J. Chem. Soc., Perkin Trans. 2* **1993**, 1447–1454.

(6) (a) Lousberg, R. J. J.; Weiss, U.; Salemink, C. A.; Arnone, A.; Merlini, L.; Nasini, G. *Chem. Commun.* **1971**, 1463–1464. (b) Yamazaki, S.; Ogawa, T. *Agric. Biol. Chem.* **1972**, *36*, 1707–1718.

(7) Nasini, G.; Merlini, L.; Andreatti, G. D.; Bocelli, G.; Sgarabotto, P. *Tetrahedron* **1982**, *38*, 2787–2796.


(8) For a review, see: Daub, M. E.; Herrero, S.; Chung, K.-R. *FEMS Microbiol. Lett.* **2005**, *252*, 197–206. and references cited therein.

(9) Hongyu, Z.; Zhiyi, Z. *Chin. Sci. Bull.* **1997**, *42*, 2005–2009.


(10) The chelation allows for keto–enol tautomerization; see ref 21.

(11) (a) Liu, W. Z. Fermentation and Structure Determination of Hypomycin B. M.D. thesis, Yunnan University, **2000**. (b) Liu, W. Z.; Chen, Y. T.; Xie, J. L. *J. Yunnan Univ., Nat. Sci. Ed.* **2000**, *22*, 389–391. (c) Liu, W. Z.; Ma, L. Y.; Li, C.; Chen, Y. T.; Xie, J. L. *Acta Pharmacol. Sin.* **2001**, *36*, 313–314.


## SCHEME 1. Previous Synthesis of Parent Perylenequinone 1



## SCHEME 2. Weiss Attempted Synthesis of Perylenequinone 22 (PPA = Polyphosphoric Acid)



## SCHEME 3. Dallacker Synthesis of Perylenequinone 27




(Scheme 3). Unfortunately, the synthesis was lengthy and lacked the necessary C6,C6'-oxygenation.<sup>22</sup>

Prior to the report by Broka on the calphostins,<sup>14</sup> syntheses of the highly oxidized perylenequinone core were either racemic or yielded optically inactive products. Thus, these early efforts did not address a key architectural element of the perylenequinones; the chiral helical pentacyclic core. Due to steric interactions of the C7,C7'- and C2,C2'-substitution, the pentacyclic core is forced to adopt a chiral helical structure (dihedral angle of 20°). With larger C7,C7'- and C2,C2'-groups, the rate of interconversion of the two possible enantiomeric conformational isomers is slow leading to stable atropisomers. As discussed above, the helicity of most perylenequinones was determined by correlation to cercosporin. In the cases of hypocrellin B (4) and scutiaquinone B (17), there is an absence of helicity in the CD spectra. However, calculations indicate that the lowest energy ground-state structures are in fact helical (Scheme 4). Presumably, the double bond of hypocrellin B (4) and the lack of substituents at C2,C7'-positions of scutiaquinone B (17)

lower the atropisomerization barrier such that the two helical forms rapidly interconvert.<sup>23,24</sup>

The helicity of the perylenequinone natural products is not uniform. Rather, compounds exhibiting both (*M*)- and (*P*)-helicity have been isolated, leaving the origin of the helical stereochemistry an interesting and unsolved issue. Since different helical stereochemistries arise from different organisms (hypocrellin from *Hypocrella bambusae*<sup>25</sup> and hypocrellin A from *Shiraia bambusicola*<sup>23</sup>), it seems likely that an enzyme-mediated enantioselective dimerization occurs in the biosynthetic routes. The fact that horseradish peroxidase, a metalloporphyrin enzyme, has been found to elicit asymmetric naphthol couplings with good yields and modest enantioselectivities indicates that enzymes are capable of catalyzing this transformation (eq 1).<sup>26</sup> Presumably, the peroxidase catalyzes the production of a stabilized radical to enable the homocoupling.<sup>27</sup> It is entirely probable, therefore, that helical chiral perylenequinones, such as 30, absent any centrochiral elements (i.e., R<sup>2</sup> not containing further stereocenters) can be produced enzymatically. Such production likely occurs via an axial chiral binaphthalene with transfer of axial stereochemistry to helical stereochemistry in the perylenequinone-forming step (Scheme 5).



In an examination of prior approaches, the installation of the helical chirality has been the most difficult aspect and thus the focal point of the syntheses.<sup>14–17</sup> Poor diastereocontrol in the couplings of chiral C7,C7'-substituted naphthalenes also points away from such a pathway in the biosynthesis (i.e., 38 to 8a–d, Scheme 6).<sup>14–17</sup> Thus, we formulated a plan revolving around diketone perylenequinone 33. Notably, perylenequinone 33 is a retrosynthetic intermediate to four different natural product classes: the calphostins (8a–d) and cercosporins (6) via reduction, the hypocrellins (2) via aldol cyclization, and the elsinochromes (10) via oxidative enol coupling. In contrast, there is no direct route from monomer 38 to the hypocrellins (2) and elsinochromes (10); oxidation and loss of the two alcohol stereocenters would be required first. We proposed that helical perylenequinone 33 could be produced from cyclization of axial chiral binaphthalene 34 which would in turn be produced by facile enantioselective oxidative coupling of the corresponding monomer 35. The illustrated monomer 35 is generated in a few simple steps from phenylacetic acid 37, in contrast to the laborious preparations of enantiopure monomers 38.

A central element of this synthetic plan is the atropisomeric stability of the key intermediate 33. The calphostins (8a–d) exhibit good atropisomeric stability, but the simple change of introducing a bridging methyldene as in cercosporin

(24) Ayers, S.; Zink, D. L.; Mohn, K.; Powell, J. S.; Brown, C. M.; Murphy, T.; Brand, R.; Pretorius, S.; Stevenson, D.; Thompson, D.; Singh, S. B. *J. Nat. Prod.* **2007**, *70*, 425–427.



(25) Chen, W. S.; Chen, Y. T.; Wang, X. Y.; Friedrichs, E.; Puff, H.; Breitmaier, E. *Liebigs Ann. Chem.* **1981**, 1880–1885.

(26) Sridhar, M.; Vadivel, S. K.; Bhalerao, U. T. *Tetrahedron Lett.* **1997**, *38*, 5695–5696.

(27) Feringa, B.; Wynberg, H. *Bioorg. Chem.* **1978**, *7*, 397–408.

(22) Dallacker, F.; Leidig, H. *Chem. Ber.* **1979**, *112*, 2672–2679.

(23) Kishi, T.; Tahara, S.; Tsuda, M.; Tanaka, C.; Takahashi, S. *Planta Med.* **1991**, *57*, 376–379.

**SCHEME 4. Conformational Isomers of Hypocrellin B and Scutiaquinone B**

**SCHEME 5. Proposed Enzymatic Perylenequinone Formation**


(6) abrogates much of this stability with atropisomerization occurring at 37 °C.<sup>7</sup> Due to the sterically larger C7,C7'-hydroxypropyl groups in the calphostins (**8a–d**) compared to the 2-oxopropyl groups of **33**, we anticipated that **33** would exhibit lower atropisomeric stability. Since no enantiopure perylenequinones with only the helical stereochemistry as a chiral element had been reported previously, we set out to establish the degree of configurational stability of such species. In this paper, we detail the synthesis of the enantiopure perylenequinone core, addressing installation of the necessary oxygenation pattern as well as the formation and stability of the helical stereochemistry in the absence of other stereochemical substitution.<sup>28</sup>

**Results and Discussion**

**Retrosynthetic Analysis.** With an initial goal of producing an enantiopure perylenequinone possessing only the helical stereochemical component, we elected to undertake the synthesis of analogue **39** possessing simple *n*-propyl groups at the C7, C7'-positions. Based on prior reports, we identified two intermediates (**40** and **42**) that could lead to the desired perylenequinone **39**, both devolving to common chiral biaryl

intermediate **41** (Scheme 7). Merlic has reported diastereoselective oxidative couplings to either provide the calphostin perylenequinone directly or via a bis-*o*-quinone intermediate,<sup>17,29</sup> such as **40** (path *a*). Similarly, Broka<sup>14</sup> and Coleman<sup>16</sup> utilized a diastereoselective biaryl coupling to yield an axial chiral biaryl as an intermediate in their calphostin syntheses. The perylenequinone arose from the cyclization of the corresponding 5,5'-bisanthol, similar to debenzylated **42** (path *b*). For both paths *a* and *b*, effective stereochemistry transfer of the axial stereochemistry to the ultimate helical stereochemistry was observed.<sup>14–17</sup> However, the role of the bulky protected C7,C7'-hydroxypropyl groups in stabilizing this stereochemistry transfer was unknown.

Due to both the reported success of Merlic<sup>29</sup> and also the precedent for the oxidation of naphthalenes to bis-*o*-quinones,<sup>30</sup> we initially chose to pursue path *a*. We commenced this effort by exploring the ability of our catalyst system in the enantioselective coupling of a series of highly substituted naphthols to provide the required axial chiral precursor **41**.<sup>28</sup>

**Monomer Synthesis and Asymmetric Oxidative Coupling.** Based on prior methods for the formation of highly substituted naphthols,<sup>20a</sup> we developed a protocol to generate naphthols **45a–d** with high efficiency (Schemes 8 and 9).<sup>28</sup> Commencing with simple commercially available phenylacetic acids (**43a–c**), the requisite naphthols were generated rapidly via acid chloride formation, malonate acylation, and cationic cyclization. After diacetylation, the less hindered C2-acetate was hydrolyzed with NaOMe to provide the desired coupling substrates **45a–c** (Scheme 8). The flexibility of this route to quickly generate substrates with a variety of substitution patterns has proven invaluable in identifying optimal substrates for the enantioselective oxidative biaryl coupling and for the perylenequinone syntheses.

In the formation of biaryl intermediate **41** (Scheme 7), the first naphthol proposed was C7-allyl **48** (Scheme 9). The allyl group would provide a platform both to intermediate **41** (Scheme 7) and also to introduce functional groups present in the natural products. The synthesis began with an allylation of the hydroxyl and carboxyl groups of commercially available **46** followed by a Claisen rearrangement and methylation of the revealed phenol (Scheme 9). Unfortunately, all attempts to form the desired **48** from **47** were unsuccessful, most likely due to incompatibility of the allyl group with the strongly acidic cyclization conditions. In light of this result, the hydrogenation of **47** was performed to yield the inert propyl derivative. After saponification to supply **49**, the naphthol **45d** was successfully prepared using the optimized conditions for **45a–c**.

In previous efforts, we have assessed the versatility of our diaza-*cis*-decalin catalyst **51** in catalytic asymmetric oxidative naphthol couplings.<sup>18,19,28,31</sup> Table 1 outlines the


(28) For a preliminary communication of this work, see: Mulrooney, C. A.; Li, X.; DiVirgilio, E. S.; Kozlowski, M. C. *J. Am. Chem. Soc.* **2003**, *125*, 6856–6857.

(29) In prior work, Zhang (ref 12) and Lown (ref 13) have shown similar transformations in the racemic syntheses of perylenequinones. Hauser (ref 15) also utilized dimerization of an *o*-quinone but found the pathway from the bis-*o*-quinone to the perylenequinone to be unproductive.


(30) Broadhurst, M. J.; Hassal, C. H.; Thomas, G. J. *J. Chem. Soc., Perkin Trans. 1* **1982**, 2239–2248.

(31) (a) Li, X.; Yang, J.; Kozlowski, M. C. *Org. Lett.* **2001**, *3*, 1137–1140. (b) Kozlowski, M. C.; Li, X.; Carroll, P. J.; Xu, Z. *Organometallics* **2002**, *21*, 4513–4522. (c) Xie, X.; Phuan, P.-W.; Kozlowski, M. C. *Angew. Chem., Int. Ed.* **2003**, *42*, 2168–2170. (d) Li, X.; Hewgley, J. B.; Mulrooney, C.; Yang, J.; Kozlowski, M. C. *J. Org. Chem.* **2003**, *68*, 5500–5511. (e) Morgan, B. J.; Xie, X.; Phuan, P.-W.; Kozlowski, M. C. *J. Org. Chem.* **2007**, *72*, 6171–6182. (f) DiVirgilio, E. S.; Dugan, E. C.; Mulrooney, C. A.; Kozlowski, M. C. *Org. Lett.* **2007**, *9*, 385–388. (g) Kozlowski, M. C.; Dugan, E. C.; DiVirgilio, E. S.; Maksimenka, K.; Bringmann, G. *Adv. Synth. Catal.* **2007**, *349*, 583–594. (h) Hewgley, J. B.; Stahl, S. S.; Kozlowski, M. C. *J. Am. Chem. Soc.* **2008**, *130*, 12232–12233.


## SCHEME 6. Retrosynthesis of the Perylenequinone Natural Products



## SCHEME 7. Perylenequinone Retrosynthetic Analysis




## SCHEME 8. Synthesis of Naphthol Substrates



significant results pertinent to this study.<sup>32</sup> Notably, an electron-withdrawing, coordinating group at C3 enhances selectivity,<sup>31</sup> presumably by via coordination to the copper complex together with the C2-phenol. The effects of the substrate oxidation potential are illustrated in entries 1–3 (Table 1),<sup>28</sup> where the electron-rich nature of the trimethoxy substrate (entry 2) led to lower optical purities due to rapid atropisomerization of the product **52e** via further oxidation. Replacement of the donating C4-methoxy group with an electron-withdrawing acetoxy group (entry 3) reduced the oxidation potential and furnished the pentasubstituted naphthalene **52b** in high enantioselectivity. The introduction of an additional electron-donating group (C5-methoxy)


## SCHEME 9. Synthesis of C7-Propyl Substrate



counteracted the effect of the C4-acetoxy group and facilitated enantiomerization of **52c** (entry 4).<sup>28</sup> Additionally, the steric gearing effects prevented the substrate from adopting the conformation needed for catalyst coordination, explaining the slow conversion as well as the low enantioselectivity. With these results in hand, it was straightforward to design substrates with a suitable oxidation potential (sufficiently electron rich for the substrate to oxidize but not too electron rich to allow product oxidation) to obtain high reactivity and selectivity. For example, **52a,b,d** (entries 1, 3, 5) furnish functionalized binaphthols in high yield and enantioselectivity. The *M*-helicity of these products was assigned on the basis of trends observed with related substrates; in the over 30 cases where the stereochemistry has been assigned via correlation, crystallization, or calculation, the *S,S*-catalyst provides the *M*-helicity and the *R,R*-catalyst the *P*-helicity.<sup>31</sup> Significantly, the enantioenriched naphthol **52d** for this investigation is generated with the necessary functionalization at the C2,C3,C4,C7-centers.

**Perylenequinone via a Bis-*o*-quinone.** Following the success of the enantioselective biaryl couplings, investigations began into transforming biaryl **52d**, arising from **45d**, to perylenequinone **39** via bis-*o*-quinone **40** (Scheme 7). As previously mentioned, there is precedent for oxidation of naphthalenes with a phenol at the position *peri* to the reaction center using various oxidants.<sup>30</sup> Our efforts commenced with the synthesis of model biaryl **53**, containing the necessary C4,C4'-phenols, but possessing C7,C7'-methoxy groups instead of propyl groups. Direct benzylation of the C2,C2'-phenols to

(32) Further investigations of the biaryl coupling reaction are examined in the subsequent paper in this series. Morgan, B. J.; Mulrooney, C.A.; O'Brien, E. M.; Kozlowski, M. C. *J. Org. Chem.* **2010**, *75*, (DOI 10.1021/jo901384h).


**TABLE 1. Enantioselective Biaryl Coupling with Diaza-*cis*-decalin (*S,S*)-**51****


entry	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	product	yield (%), ee (%)
1	H	H	Ac	<b>a</b>	71, 86
2	OMe	H	Me	<b>e</b>	81, 35 ( <i>P</i> ) <sup>a</sup>
3	OMe	H	Ac	<b>b</b>	72, 90 ( <i>P</i> ) <sup>b</sup>
4	OMe	OMe	Ac	<b>c</b>	41, 27
5	<i>n</i> -propyl	H	Ac	<b>d</b>	85, 87


<sup>a</sup>(*M*)-Enantiomer formed initially. <sup>b</sup>(*R,R*)-**51** catalyst complex was used.

generate **53** was unsuccessful, a trend that was observed throughout the series. Apparently, the C2,C2'-positions are relatively hindered, which slows alkylation significantly. On the other hand, Mitsunobu reaction generally proceeded well on these centers and was used here to install the C2,C2'-benzyl ethers. Subsequent hydrolysis of the C4,C4'-acetates afforded **53** (Scheme 10). After screening a number of oxidants, the hypervalent iodine reagent bis(trifluoroacetoxy)iodobenzene [PIFA = PhI(O<sub>2</sub>CCF<sub>3</sub>)<sub>2</sub>] following McKillop's protocol was found to generate the desired bis-*o*-quinone **54** as confirmed by the crystal structure.

With the model bis-*o*-quinone **54** in hand, we next examined formation of the perylenequinone core. Merlic has reported the transformation of a similar bis-*o*-quinone **55** to perylenequinone **56** using trifluoroacetic acid (TFA) under O<sub>2</sub> (eq 2).<sup>17</sup> When applied to methylated **57**, this procedure yielded only decomposition products (eq 3). We attributed this result to the differences between the two *o*-quinones (**55** vs **57**), specifically the presence of C2, C2'-benzyl ether, C3,C3'-ester, and C7,C7'-methyl ether groups in **57**.



Reasoning that the C7,C7'-methoxy groups of **57** (eq 3) changed the electronic characteristics of the *o*-quinone relative to **55** (eq 2), we moved to biaryl **59** (Scheme 11) with C7,C7'-*n*-propyl groups as well as C2,C2'-methoxy groups to


**SCHEME 10. Synthesis of Bis-*o*-quinone **54** (ORTEP Drawing with 30% Probability Thermal Ellipsoids)**


more closely approximate Merlic's system (eq 2) in formation of the pentacyclic core. Coupling of **45d** with CuCl(OH)-TMEDA under O<sub>2</sub> provided large quantities of the racemic biaryl **52d**, which we employed to develop a viable route to **39**. Biaryl **52d** was treated with NaH and MeI to methylate the C2,C2'-phenols. As has been seen throughout the series, alkylation of the hindered C2,C2'-phenols is difficult, and this transformation required the use of anhydrous DMF<sup>33</sup> to succeed. Subsequent hydrolysis of C4,C4'-acetates efficiently afforded **59** (Scheme 11). Oxidation utilizing PIFA yielded the desired bis-*o*-quinone **60** in quantitative yield, but this compound was unstable to chromatography or prolonged exposure to air. Immediate methylation of the C4, C4'-phenols with *n*-Bu<sub>4</sub>NF and MeI provided the stable **61**, which could be purified. Surprisingly, **61** could not be isomerized to the perylenequinone **62** and only decomposition occurred following the procedure developed by Merlic. Presumably, the isomerization requires one of the *o*-quinone moieties to reduce so that a nucleophilic phenol can undergo a transannular addition to the remaining electrophilic *o*-quinone. Subsequent tautomerization and air oxidation would then furnish the perylenequinone. In an attempt to reproduce this manifold, bis-*o*-quinone **61** was treated under conditions to induce reduction (NaBH<sub>4</sub> and O<sub>2</sub> or DBU and O<sub>2</sub>); however, only decomposition was observed.

The presence of the C3,C3'-methyl ester groups accounts for the last major difference between the previously isomerized bis-*o*-quinone **55** (eq 2) and the structures **57** (eq 3) and **61** (Scheme 11). The electron-withdrawing nature of the methyl esters may destabilize the perylenequinone **62** and/or interfere with the isomerization. To test the hypothesis, decarboxylation of the biaryl was examined. To generate a biaryl that would be stable to saponification and decarboxylation, the previously formed bis-*o*-quinone **61** (Scheme 11) was reduced with sodium dithionite (Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub>) and methylated (Scheme 12). Unfortunately, mixtures of the desired octamethyl ether **63**, partially reduced hexamethyl ether **63a**

(33) In a later synthesis involving substrate **52d**, *wet* DMF is used as the solvent. These conditions cleave the acetates (presumably due to NaOH formed from trace H<sub>2</sub>O combined with the NaH) revealing a tetraphenol, which is exhaustively alkylated with the MeI to form a tetramethyl ether (**82**, Scheme 18).

SCHEME 11. Attempted Synthesis of C7,C7'-Bispropyl Perylenequinone




and starting material **61** were obtained with the highest yield of 46% for **63**. Saponification of **63** proceeded smoothly with aqueous LiOH in refluxing dioxane, to afford the bisacid. Subsequent decarboxylation with Cu-quinoline at 180 °C furnished **64** in a modest 30% yield over two steps (Scheme 12). Thereafter, treatment of **64** with BCl<sub>3</sub> in an attempt to selectively cleave the C5,C5'-methyl ethers resulted in a mixture of demethylated compounds, which upon treatment with K<sub>3</sub>Fe(CN)<sub>6</sub> yielded only a complex mixture from which none of the desired **65** was isolated. Fortunately, removal of the C5,C5'-methyl ethers was not necessary for oxidation and direct exposure of **64** to ceric ammonium nitrate (CAN) afforded the oxidized bis-*o*-quinone **40** (Scheme 12). Longer reaction times in the CAN oxidation did not afford the desired perylenequinone **65**, and all attempts to isomerize **40** to **65** led only to decomposition. This result was surprising due to the close structural similarity of **40** (Scheme 12) to **55** (eq 2).


Since an examination of the transformation proposed by Hauser and Merlic<sup>15,17</sup> indicated that a direct isomerization of **40** to **65** (Scheme 12) is not viable without a reducing agent, protocols involving a reduction were examined. Strong precedent for such an approach is seen in the work of Lown<sup>13</sup> and Zhang<sup>12</sup> on the syntheses of a perylenequinones from bis-*o*-quinones via a two-step reduction–oxidation sequence (FeCl<sub>2</sub>, then FeCl<sub>3</sub>; SO<sub>2</sub>–H<sub>2</sub>O, then FeCl<sub>3</sub>). Due to the length of the synthetic route needed to obtain large quantities of bis-*o*-quinone **40**, **61** was used to examine these new isomerization procedures. Treatment of **61** with the reducing agent Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> followed by FeCl<sub>3</sub> afforded a complex mixture that was determined to contain the undesired isomer **61** as the major product as well as the desired product **62** (Scheme 13).<sup>34</sup>

(34) Compound **62** proved unstable to SiO<sub>2</sub> chromatography, but a small portion was purified by semipreparative reversed-phase HPLC to confirm the structure via <sup>1</sup>H NMR spectroscopy.

SCHEME 12. Decarboxylation and Attempted Synthesis of a Perylenequinone



With confidence that the synthesis of perylenequinone **62** via this route could be optimized the reactions were examined with enantioenriched **52d** to verify the optical purity of the synthetic intermediates. Formation of enantioenriched **59** (70% ee) following the racemic synthesis seen in Scheme 11, permitted the PIFA-mediated oxidation of the optically active intermediate to bis-*o*-quinone **61** in 70% yield (eq 4). Disappointingly, when evaluated by both optical rotation and HPLC, compound **61** was found to be racemic.



The mechanism of the oxidation reaction was studied to provide an explanation of the atropisomerization. The use of PIFA and phenyliodonium diacetate [PIDA = PhI(OAc)<sub>2</sub>] for phenolic oxidation has been well reviewed,<sup>35</sup> and notably, biphenol **67**, upon oxidation leads to the extended quinone **68** (eq 5).<sup>36</sup> When only 2 equiv of PIFA (7.5 equiv used in eq 4) were used in the oxidation of **59**, a binaphthone (**69**, Figure 3) was isolated as a dark green material from

(35) (a) Stang, P. J.; Zhdankin, V. V. *Chem. Rev.* **1996**, *96*, 1123–1178.

(b) Moriarty, R. M.; Prakash, O. *Acc. Chem. Res.* **1986**, *19*, 244–250.

(c) Pohnert, G. *J. Prakt. Chem.* **2000**, *342*, 731–734.

(36) Pelter, A.; Ward, R. S. *Tetrahedron* **2001**, *57*, 273–282.






FIGURE 3. Strained extended quinone structures.

SCHEME 13. Two-Step Synthesis of Perylenequinone from Bis-*o*-quinone



the reaction mixture. In our assignment of **69**, the propyl groups are placed *anti* to each other because the  $^1\text{H}$  NMR spectrum does not indicate any restricted rotation about the propyl groups (i.e., diastereotopic protons), which is seen in the  $^1\text{H}$  NMR spectra of perylenequinones where the propyl groups are in close proximity. Similar to **69**, the highly strained bianthrone **70** has been reported in the literature.<sup>37,38</sup> Interestingly, bianthrone **70** was found to equilibrate between twisted and stack conformations (Figure 3). Analogously, it is clear that **69** is a chiral compound, but the facile interconversion of the twisted and stacked conformations provides a ready racemization pathway (Figure 4).



The use of the less reactive PIDA allowed a full evaluation of the oxidation pathway of **59** (Scheme 14). While all efforts to optimize formation of binaphthone **69** with PIFA as the oxidant proved to be unsuccessful, PIDA allows **69** to be isolated in a 91% yield (Scheme 14) with minimal amounts of decomposition during the reaction. The treatment of **69** with 2 equiv of PIFA yielded racemic **60**, strongly supporting the

(37) The dinaphthofuranedione **24** in Scheme 2 (ref 20) is also similar to the strained extended quinone structure of **69**.

(38) (a) Kral, A.; Laatsch, H. *Z. Naturforsch.* **1993**, *48b*, 1401–1407. (b) Tapuhi, Y.; Kalisky, O.; Agranat, I. *J. Org. Chem.* **1979**, *44*, 1949–1952. (c) Evans, D. H.; Fitch, A. *J. Am. Chem. Soc.* **1984**, *106*, 3039–3041.


FIGURE 4. Proposed racemization pathway of **69**.SCHEME 14. Isolated Intermediate in the Oxidation to Bis-*o*-Quinone **60**

intermediacy of **69** during the PIFA-oxidation of **59** to bis-*o*-quinone **61** (eq 4).

Based upon the above evidence, an oxidative mechanism for conversion of binaphthol **59** to *o*-quinone **60** is proposed in Scheme 15. One equivalent of PIFA oxidizes **59** to the binaphthone **69** via adduct **71**. The trifluoroacetic acid liberated from the PIFA protonates the C4-carbonyl, promoting conjugate addition of  $\text{H}_2\text{O}$  to the C5-center. After tautomerization, another oxidation forms the binaphthone **74**. A second conjugate addition of  $\text{H}_2\text{O}$  at the C5'-position is followed by tautomerization to intermediate **75** that can then be oxidized twice further. Final hydrolysis of the alkylated carbonyls of **76** with the excess  $\text{H}_2\text{O}$  present furnishes bis-*o*-quinone **60**. This mechanism requires 4 equiv of an oxidant to form **60** from **59**, which is consistent with the amount employed in eq 4.


**Perylenequinone via an Oxidized Binaphthalene.** Due to the racemization of biaryl **59** during the formation of bis-*o*-quinone **60**, path *a* (Scheme 7) was not a viable route to enantiopure **39**. Shifting to path *b*, we examined the formation of **39** via an advanced biaryl intermediate **42**, containing C5,C5'-benzyl ethers (Scheme 7). The first undertaking was the hydroxylation of the C5,C5'-centers to provide **42** from available precursor **41**. Based upon the above results (Scheme 15), the C4,C4'-phenols needed to be masked as ethers during this oxidation to prevent formation of binaphthone **69**. Thus, an oxidation was required that did not rely on hydroxyl activation in oxidizing the C5,C5'-centers. One such method is the oxygenation of aryl lithiums with electrophilic oxygen sources. Since the requisite binaphthalene (see **82** below) is very electron rich and the C5,C5'-centers are hindered, a naphthalene model system **78** was utilized to



SCHEME 15. Proposed Mechanism for Bis-*o*-quinone Formation


explore the stability and reactivity of the corresponding C5-lithium species. Model **77** was prepared by C5-bromination of **77** followed by hydrolysis of the C2,C4-acetates and methylation to provide trimethyl ether **78** (Scheme 16). With *tert*-butyllithium, lithium-halogen exchange provided intermediate **79** as confirmed by quenching with D<sub>2</sub>O which generated the corresponding C5-deuterated arene and minimal amounts of C3-*tert*-butyl ketone. Unfortunately, reaction with a series of oxidants and electrophiles (oxygen gas, Davis oxaziridine, lithium salt of *tert*-butyl hydroperoxide, and B(OMe)<sub>3</sub>)<sup>39</sup> was unsuccessful, which was attributed to the hindered nature of the C5 center. Given this challenge, an alternate aryl hydroxylation reaction was sought.

Notably, Kita has utilized PIFA to induce the nucleophilic substitution of phenyl ethers to afford functionalized arenes.<sup>40,41</sup> His studies indicate that the reaction proceeds by a single-electron transfer (SET) resulting in the formation of a radical cation [ArH<sup>•+</sup>] as the reactive intermediate (**81**, eq 6).<sup>40</sup> This mechanism contrasts with PIFA-oxidations




described above (Scheme 10, 11, 14, and 15; eq 4), which proceeded via phenolic C4,C4'-hydroxyl activation (Scheme 15). Based on the reported success of the acetoxy group as a nucleophile in the PIFA-induced SET pathway, we set out to synthesize an intermediate containing the necessary perylenequinone oxygenation in this manner. The greatest concern in this transformation was the regioselection in the addition to the radical cation. In considering the radical cation, many resonance structures are possible, but the two that are pertinent (cations at

## SCHEME 16. Attempted C5-Oxidation of a Naphthalene Model System



## SCHEME 17. Regioselection in the PIFA-Mediated Hydroxylation



(39) (a) Parker, K. A.; Koziski, K. A. *J. Org. Chem.* **1987**, *52*, 674–676. (b) Davis, F. A.; Wei, J.; Sheppard, A. C.; Gubernick, S. *Tetrahedron Lett.* **1987**, *28*, 5115–5118. (c) Green, K. *J. Org. Chem.* **1991**, *56*, 4325–4326.

(40) Kita, Y.; Tohma, H.; Hatanaka, K.; Takada, T.; Fujita, S.; Mitoh, S.; Sakurai, H.; Oka, S. *J. Am. Chem. Soc.* **1994**, *116*, 3684–3691.

(41) Arisawa, M.; Ramesh, N. G.; Nakajima, M.; Tohma, H.; Kita, Y. *J. Org. Chem.* **2001**, *66*, 59–65.

unsubstituted positions where reaction will allow regeneration of aromaticity via deprotonation) are illustrated in Scheme 17. The stabilization of the cations is expected to

be similar at C5 and C8. However, the radical would need to reside at C6 or C7 to maintain the greatest amount of conjugation in the system. The C6-methoxy would be expected to stabilize radical character at the C6-position to a greater degree. Thus, resonance structure **82b** is expected to predominate leading to reaction at the C5-center. In addition, the C5-position is less hindered by the neighboring C4,C6-methoxys than the C8-position with larger flanking groups (C1-aryl group and C7-propyl group). The requisite substrate to test this hypothesis was prepared from **52d** using reagent DMF during methylation to conveniently prepare hexamethyl ether **82** in one step.<sup>33</sup> After initial C2,C2'-methylation, the small amount of water in the DMF was converted to NaOH, cleaving the C4,C4'-acetates; subsequent methylation afforded **82** (Scheme 18). Treatment of **82** with PIFA and TMSOAc in (CF<sub>3</sub>)<sub>2</sub>CHOH afforded a mixture of compounds, from which bisphenol **87** and the C5,C5'-bistrifluoroacetate **86** were isolated. The reaction also proceeded well without TMSOAc; presumably, trifluoroacetate from the PIFA is sufficiently nucleophilic to trap the radical cation. When the reaction mixture (without TMSOAc) was treated with aqueous NaOH, the bistrifluoroacetate intermediate **86** was efficiently cleaved furnishing **87** in 89% yield (Scheme 18). However, there was erosion in the optical purity, from 86% ee in **82** to 68–70% ee in bisnaphthol **87**. We propose that this attrition arises from racemization of radical cation or cation intermediates where the atropisomerization barrier is lower.<sup>28</sup> To mitigate this effect, hexafluoro-2-propanol was replaced with the less polar trifluoroethanol, which should be sufficiently polar to permit the initial SET to the radical cation but will reduce its lifetime. Pleasingly, bisnaphthol **87** was obtained with improved enantiopurity (82% ee) but in a diminished 51% yield.<sup>42</sup>


The oxidative cyclization of **87** with MnO<sub>2</sub> proceeded smoothly to yield perylenequinone **88**,<sup>16</sup> representing the first enantioselective synthesis of a perylenequinone with the helical chirality as the only stereochemical element. Since the optical purity of **88** could not be measured directly, the C4,C4'-methyl ethers were selectively removed using MgI<sub>2</sub> to provide **89**.<sup>15,17</sup> An HPLC assay verified the stereochemical integrity of the axial to helical chirality transfer (Scheme 19).

**Decarboxylation and Atropisomeric Stability.** This initial result on the atropisomeric stability of a perylenequinone with no stereogenic elements aside from the helical axis was encouraging. However, it was unclear that the corresponding compounds lacking the C3,C3'-ester groups, which were needed as outlined in the retrosynthesis (see **33** in Scheme 6), would possess the same degree of atropisomeric stability. Removal of the C3,C3'-ester groups alleviates steric gearing interactions, thereby decreasing the effective size of the C2, C2'-groups so that they might more easily slide by one another and thereby lower the atropisomerization barrier.


Thus, the synthesis of **39**, which contains the necessary substitution pattern of the natural products perylenequinone core, was our next goal (Scheme 20). To accomplish this goal,

(42) Improvements to the hydroxylation reaction were later developed, mitigating the loss of yield and enantioenrichment. As seen in the subsequent papers in this series on more complex intermediates, no erosion of enantioenrichment was observed when NaOAc was used to quench the reagents after oxidation. The NaOAc was added prior to solvent evaporation and subsequent NaOH treatment removes excess oxidant and TFA that can cause degradation.

### SCHEME 18. PIFA-Mediated Hydroxylation



### SCHEME 19. Synthesis of Optically Active Perylenequinone **89**




**87** was treated with benzyl bromide and NaH to mask the reactive C5,C5'-phenols. Hydrolysis was affected by LiOH to supply the diacid **90**. Disappointingly, the high temperatures of conventional aromatic decarboxylation protocols, such as Cu-quinoline, caused not only atropisomerization of the biaryl but also significant decomposition resulting in modest yields of **42** (31% after optimization). As a consequence, we developed a palladium decarboxylation protocol involving stoichiometric palladium(II) trifluoroacetate and silver carbonate followed by the reduction of the resultant aryl palladium species. The described procedure worked well on these systems and supplied the key intermediate **42** in 87% yield without racemization.<sup>19,43</sup> Further examination of this strategy led to the development of a single-step decarboxylation using catalytic palladium and excess of trifluoroacetic acid as the proton source to provide a variety of arenes in good yield.<sup>43</sup> After hydrogenolysis of bisbenzyl ether **42**, the axial chiral bisphenol was oxidized by MnO<sub>2</sub> to afford the helical chiral perylenequinone as a bright red resin (Scheme 20). The directed, selective cleavage of the C4,C4'-methyl ethers was accomplished by MgI<sub>2</sub> to complete the first synthesis of optically active perylenequinone **39**.<sup>44,45</sup>

(43) Dickstein, J. S.; Mulrooney, C. A.; O'Brien, E. M.; Morgan, B. J.; Kozlowski, M. C. *Org. Lett.* **2007**, *9*, 2441–2444.

(44) Merlic, in ref 17, synthesized perylenequinone **39** as the racemate via the dimerization of an *o*-quinone, but no logical entry to the enantioenriched variant is available with such a route.

(45) Following this work, the synthesis of **39** evolved further, culminating in a 16-step protocol with 4.2% overall yield of 98% ee material. The evolution is described briefly in ref 19 and in detail in the third paper in this series. Morgan, B. J.; Mulrooney, C.; Kozlowski, M. C. *J. Org. Chem.* **2010**, *75* (DOI 10.1021/jo9013854).

**SCHEME 20. Palladium-Mediated Decarboxylation and Synthesis of Optically Active 39**


Pleasingly, perylenequinone **39** was also found to exhibit considerable atropisomeric stability with an atropisomerization half-life of 4 d at 60 °C in benzene.<sup>19</sup>

**Summary.** As summarized in Scheme 21, the final route to **39** proceeded in 18 steps and a 5.4% overall yield (92% average yield per step). The key steps in this synthesis were an enantioselective biaryl coupling, a PIFA-induced phenol formation, and a palladium-mediated aromatic decarboxylation. The development and optimization of these transformations was crucial not only to this investigation but also for the syntheses of the perylenequinone natural products detailed in the subsequent papers in this series.

**Conclusions.** The synthesis of **39** provided the first helical chiral perylenequinone containing the natural product substitution pattern but absent the conventional stereogenic C7, C7'-substitution. The central issues addressed here including generation of the necessary oxygenation pattern in a complex binaphthol and formation of the helical stereochemistry. Importantly, such perylenequinones were shown to possess sufficient atropisomeric stability to be viable intermediates in the biogenesis of the perylenequinone natural products. Furthermore, this stability supports their use as key intermediates in biomimetic total synthesis ventures (see **33** in the retrosynthesis in Scheme 6). Thus, these discoveries laid the groundwork for the total syntheses of (+)-calphostin D (**8d**) and (+)-phleichrome (**9**)<sup>19</sup> and the first total syntheses of cercosporin (**6**)<sup>19</sup> and hypocrellin A (*ent-2*),<sup>18</sup> which are described in the following papers in this series (DOIs 10.1021/jo901384h, 10.1021/jo9013854, and 10.1021/jo901386d). Importantly, the versatility of this strategy enabled selective and convergent synthesis of all stereoisomeric combinations permitting comparison of their biological effects for the first time.<sup>19</sup>

**Experimental Section**

**2-(4-Methoxy-3-propylphenyl)acetic Acid (49).** To a solution of **47** (1.0 g, 4.0 mmol) in MeOH (50 mL) was added 10% Pd/C (106 mg), and the mixture was stirred under H<sub>2</sub> (1 atm). After 2 d, the reaction was filtered through Celite, rinsing with CH<sub>2</sub>Cl<sub>2</sub>. This solution was filtered through SiO<sub>2</sub>, and the solvent was evaporated to yield the dipropyl product as an oil (1.0 g, 100%): <sup>1</sup>H NMR (360 MHz, CDCl<sub>3</sub>) δ 0.90–0.97 (m, 6H),


1.57–1.67 (m, 4H), 2.57 (t, *J* = 7.5 Hz, 2H), 3.54 (s, 2H), 3.80 (s, 3H), 4.05 (t, *J* = 6.5 Hz, 2H), 6.78 (d, *J* = 8.1 Hz, 1H), 7.06 (m, 2H).

A mixture of propyl ester (329 mg, 1.31 mmol) and 1 N NaOH (5 mL, 5 mmol) in THF (5 mL) was stirred for 16 h. The reaction was diluted with 1 N NaOH and hexanes, and the aqueous phase was extracted with hexanes. The aqueous phase was then acidified with 1 N HCl and extracted with EtOAc. The organic phase was washed with H<sub>2</sub>O and brine and dried (Na<sub>2</sub>SO<sub>4</sub>), and the solvent was evaporated to yield **49** as a white solid (250 mg, 92%): mp 58–60 °C; IR (thin film) 3076 (br), 3003, 2961, 1710, 1251 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.94 (t, *J* = 7.4 Hz, 3H), 1.56–1.62 (m, 2H), 2.56 (t, *J* = 7.6 Hz, 2H), 3.57 (s, 2H), 3.80 (s, 3H), 6.79 (d, *J* = 8.2 Hz, 1H), 7.03 (s, 1H), 7.07 (d, *J* = 8.3 Hz, 1H), 11.0 (br s, 1H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 178.5, 157.0, 131.8, 131.3, 128.0, 125.3, 110.8, 55.8, 40.7, 32.6, 23.3, 14.5; HRMS (CI) [*M*<sup>+</sup>] calcd for C<sub>12</sub>H<sub>16</sub>O<sub>3</sub> 208.1099, found 208.1104.

**Methyl 1,3-Dihydroxy-7-methoxy-6-propyl-2-naphthoate (50).** To a solution of acid **49** (6.4 g, 30.7 mmol) in PhH (30 mL) was added SOCl<sub>2</sub> (9.0 mL, 123 mmol). The reaction was heated to reflux under Ar for 2 h. The solvent was evaporated in vacuo to yield a brown oil. To a suspension of NaH (60% in mineral oil, 3.7 g, 92.1 mmol) in THF (200 mL) under Ar was added dimethyl malonate (10.5 mL, 92.1 mmol). After being stirred for 30 min, the mixture was chilled in an ice–H<sub>2</sub>O bath, and a solution of the above acid chloride in THF (50 mL) was added. The reaction was stirred for 3 h with warming to room temperature. The mixture was poured into a mixture of ice and 1 N HCl and extracted into EtOAc. The organic phase was washed with H<sub>2</sub>O and brine and dried (Na<sub>2</sub>SO<sub>4</sub>), and the solvent was evaporated. The oil was filtered through SiO<sub>2</sub>, rinsing with 0–20% EtOAc/hexanes. The solvent was evaporated, the oil was dissolved in MeSO<sub>3</sub>H (55 mL), and P<sub>2</sub>O<sub>5</sub> (5.0 g) was added. After being stirred under Ar for 3 h, the mixture was poured over ice–H<sub>2</sub>O and filtered. The yellow solid was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, was washed with H<sub>2</sub>O and brine, and dried (Na<sub>2</sub>SO<sub>4</sub>), and the solvent was evaporated to yield **50** (6.9 g, 77%), which was carried on to the next step without further purification: IR (thin film) 3443, 2957, 1670, 1644, 1252, 1217, 1158 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.99 (t, *J* = 7.3 Hz, 3H), 1.64–1.71 (m, 2H), 2.69 (t, *J* = 7.4 Hz, 2H), 3.92 (s, 3H), 4.10 (s, 3H), 6.71 (s, 1H), 7.29 (s, 1H), 7.43 (s, 1H), 8.63 (br s, 1H), 11.32 (br s, 1H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 171.0, 160.4, 155.2, 152.4, 138.5, 133.7, 126.7, 118.9, 102.4, 101.2, 97.2, 55.7, 53.3, 33.3, 23.0, 14.5; HRMS (CI) [*M*<sup>+</sup>] calcd for C<sub>16</sub>H<sub>18</sub>O<sub>5</sub> 290.1154, found 290.1156.

**Methyl 1-Acetoxy-3-hydroxy-7-methoxy-6-propyl-2-naphthoate (45d).** A solution of naphthol **50** (6.0 g, 20.7 mmol), Ac<sub>2</sub>O (60 mL), and pyridine (60 mL) was stirred for 3 h. The reaction mixture was poured over ice–1 N HCl and stirred for 30 min. The solid was filtered, rinsed with H<sub>2</sub>O, and dried by pulling air through the filter cake to yield the diacetate (6.9 g, 89%): IR (thin film) 2955, 1776, 1724, 1316, 1195 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.97 (t, *J* = 7.3 Hz, 3H), 1.64–1.69 (m, 2H), 2.32 (s, 3H), 2.44 (s, 3H), 2.72 (t, *J* = 7.4 Hz, 2H), 3.90 (s, 3H), 3.91 (s, 3H), 7.01 (s, 1H), 7.37 (s, 1H), 7.52 (s, 1H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 170.0, 169.2, 164.6, 158.0, 146.4, 143.8, 137.1, 130.6, 128.2, 125.6, 118.4, 116.5, 99.5, 55.7, 52.8, 33.1, 22.9, 21.3, 21.2, 14.4; HRMS (ESI) calcd for C<sub>20</sub>H<sub>22</sub>O<sub>7</sub> (*M*<sup>+</sup>) 374.1366, found 374.1364.

To a solution of this acetate (2.5 g, 6.7 mmol) in MeOH (150 mL) and CH<sub>2</sub>Cl<sub>2</sub> (15 mL) under Ar was added K<sub>2</sub>CO<sub>3</sub> (913 mg, 6.6 mmol). After being stirred for 15 min, the green reaction mixture was poured over ice–1 N HCl. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> and the organic extracts were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and the solvent was evaporated. Purification was accomplished with chromatography (10–20% EtOAc/hexanes) to yield product **45d** as a yellow

SCHEME 21. Enantioselective Total Synthesis of (*M*)-39

solid (1.7 g, 76%): mp 121.5–125.5 °C; IR (thin film) 3155, 2957, 1773, 1675, 1223, 1197 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.98 (t, *J* = 7.4 Hz, 3H), 1.65–1.69 (m, 2H), 2.48 (s, 3H), 2.69 (t, *J* = 7.5 Hz, 2H), 3.89 (s, 3H), 4.00 (s, 3H), 6.91 (s, 1H), 7.17 (s, 1H), 7.39 (s, 1H), 10.48 (s, 1H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 169.9, 169.6, 156.3, 155.2, 147.6, 138.5, 133.8, 127.0, 121.6, 110.1, 107.2, 99.1, 55.5, 53.4, 33.2, 22.9, 21.3, 14.4; HRMS (CI) calcd for C<sub>18</sub>H<sub>20</sub>O<sub>6</sub> (M<sup>+</sup>) 332.1260, found 332.1250.

**General Procedure for the Oxidative Biaryl Coupling.** To a solution of the 2-naphthol substrate dissolved in the appropriate solvent was added the copper catalyst (10 mol %). The mixture was sonicated to yield a clear green, blue, purple, or red solution, which was stirred under O<sub>2</sub> (1 atm) at the indicated temperature and time. After cooling, the reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with 1 N HCl. The aqueous phase was back extracted with CH<sub>2</sub>Cl<sub>2</sub> and the combined organic solutions were dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration and concentration afforded the crude product. Purification was accomplished by SiO<sub>2</sub> chromatography.

**(*M*)-Dimethyl 4,4'-Diacetoxy-2,2'-dihydroxy-6,6'-dimethoxy-7,7'-dipropyl-1,1'-binaphthyl-3,3'-dicarboxylate (52d).** A mixture of naphthol **45d** (103 mg, 0.309 mmol), 4 Å molecular sieves (86 mg), and CuI-(*S,S*)-**51** (11.4 mg, 0.0329 mmol) in MeCN (1.5 mL) was stirred at room temperature under an O<sub>2</sub> atmosphere for 24 h. The reaction was quenched with 1 N HCl, and the aqueous phase was extracted with EtOAc. The organic extracts were washed with H<sub>2</sub>O and brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and filtered through SiO<sub>2</sub> with 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub>. Purification was accomplished via chromatography to yield the product as a resin (86.7 mg, 85%) in 87% ee: [α]<sub>D</sub><sup>20</sup> +20.9 (*c* 0.23, MeOH, 87% ee); IR (thin film) 3130, 2958, 1770, 1672, 1234, 1195 cm<sup>-1</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 0.79 (t, *J* = 7.4 Hz, 6H), 1.39–1.44 (m, 4H), 2.44–2.55 (m, 10H), 3.90 (s, 6H), 4.01 (s, 6H), 6.91 (s, 2H), 7.05 (s, 2H), 10.64 (s, 2H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 170.1, 169.7, 156.3, 152.9, 147.8, 138.7, 133.2, 125.8, 121.8, 115.4, 107.3, 99.7, 55.5, 53.4, 33.5, 23.2, 21.9, 14.3; HRMS (ESI) calcd for C<sub>36</sub>H<sub>38</sub>O<sub>12</sub>Na (MNa<sup>+</sup>) 685.2261, found 685.2276; CSP HPLC (Chiralpak AD, 1.0 mL/min, 70:30 hexanes/*i*-PrOH) t<sub>R</sub>(*S*) = 10.5 min, t<sub>R</sub>(*R*) = 13.3 min.

**Dimethyl 2,2'-Bis(benzyloxy)-4,4'-dihydroxy-7,7'-dimethoxy-5,5',6,6'-tetraoxo-5,5',6,6'-tetrahydro-1,1'-binaphthyl-3,3'-dicarboxylate (54).** Bisphenol **53** (51 mg, 0.07 mmol, 74% ee) was dissolved in MeCN/CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O (6:1:1, 8 mL), and PhI(OCOCF<sub>3</sub>)<sub>2</sub> (300 mg, 0.7 mmol) was added. After 2 h at room temperature, the deep red solution was diluted with CH<sub>2</sub>Cl<sub>2</sub> and

washed with H<sub>2</sub>O. Concentration afforded an oil which was purified by SiO<sub>2</sub> chromatography (2–3.5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to provide **54** as a red solid (45 mg, 88%); IR (thin film) 3683, 2954, 1733, 1693, 1636, 1608, 1274, 1261, 1227, 1115 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CD<sub>2</sub>Cl<sub>2</sub>/DMSO-*d*<sub>6</sub>) δ 3.60 (s, 6H), 3.99 (s, 6H), 4.84 (d, *J* = 9.0 Hz, 2H), 5.08 (d, *J* = 9.0 Hz, 2H), 5.84 (s, 2OH), 7.09 (m, 4H), 7.24 (m, 6H), 12.9 (s, 2OH); <sup>13</sup>C NMR (125 MHz, CD<sub>2</sub>Cl<sub>2</sub>/DMSO-*d*<sub>6</sub>) δ 179.8, 175.3, 165.4, 163.6, 152.8, 137.1, 135.6, 128.5, 128.4, 128.1, 127.7, 126.7, 119.7, 115.3, 109.2, 109.1, 76.2, 56.0, 53.0; HRMS (ESI) calcd for C<sub>40</sub>H<sub>30</sub>O<sub>14</sub>Na (MNa<sup>+</sup>) 757.1558, found 757.1533.

**Dimethyl 2,2',4,4'-Tetramethoxy-5,5',6,6'-tetraoxo-7,7'-dipropyl-5,5',6,6'-tetrahydro-1,1'-binaphthyl-3,3'-dicarboxylate (61).** To a solution of binaphthol **59** (230 mg, 0.379 mmol) in MeCN (50 mL), CH<sub>2</sub>Cl<sub>2</sub> (20 mL), and H<sub>2</sub>O (10 mL) was added PhI(OCOCF<sub>3</sub>)<sub>2</sub> (1.2 g, 2.8 mmol). The dark blue reaction mixture was stirred for 45 min and then diluted with H<sub>2</sub>O and EtOAc. The phases were separated, and the aqueous phase was washed with EtOAc. The organic phase was extracted with H<sub>2</sub>O and brine and dried (Na<sub>2</sub>SO<sub>4</sub>), and the solvent was evaporated to yield **60** as an orange solid that was carried on to the next step without further purification: <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.87 (t, *J* = 7.3 Hz, 6H), 1.39–1.44 (m, 4H), 2.32 (t, *J* = 6.7 Hz, 4H), 3.80 (s, 6H), 3.99 (s, 6H), 6.60 (s, 2H), 12.8 (br s, 2H); CSP HPLC (Chiralpak AD, 1.0 mL/min, 70:30 hexanes/*i*-PrOH) t<sub>R</sub>(*S*) = 12.3 min, t<sub>R</sub>(*R*) = 22.2 min.

To a solution of *o*-quinone (0.379 mmol) and MeI (0.20 mL, 3.2 mmol) in THF (30 mL) was added TBAF (1 M in THF, 1.5 mL, 1.5 mmol). The black reaction mixture was stirred for 16 h and then diluted with EtOAc and H<sub>2</sub>O. The phases were separated, and the aqueous phase was washed with EtOAc. The organic phase was washed with H<sub>2</sub>O and brine and dried (Na<sub>2</sub>SO<sub>4</sub>), and the solvent was evaporated to yield an orange solid. Purification by chromatography (30% EtOAc/hexanes) yielded product **61** as an orange solid (168 mg, 70%): mp 95–97 °C; IR (thin film) 2955, 1737, 1667, 1224 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.87 (t, *J* = 7.3 Hz, 6H), 1.41 (m, 4H), 2.31 (m, 4H), 3.71 (s, 6H), 3.94 (s, 6H), 3.97 (s, 6H), 6.71 (s, 2H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 180.8, 177.3, 165.7, 163.6, 161.1, 143.0, 138.0, 137.0, 122.9, 122.7, 118.9, 63.8, 60.8, 53.5, 32.0, 21.9, 14.0; HRMS (ESI) calcd for C<sub>34</sub>H<sub>34</sub>O<sub>12</sub>Na (MNa<sup>+</sup>) 657.1948, found 657.1936; CSP HPLC (Chiralpak AD, 1.0 mL/min, 70:30 hexanes/*i*-PrOH) t<sub>R</sub>(*S*) = 14.7 min, t<sub>R</sub>(*R*) = 36.3 min.

**2,2',4,4'-Tetramethoxy-7,7'-dipropyl-1,1'-binaphthyl-5,5',6,6'-tetraone (40).** To a solution of **64** (3.0 mg, 0.0052 mmol) in MeCN (0.5 mL) were added H<sub>2</sub>O (1 drop) and CAN (31 mg, 0.057 mmol). The red reaction mixture was stirred for 10 min, diluted with H<sub>2</sub>O, and extracted with EtOAc. The organic phase was washed with H<sub>2</sub>O and brine and dried (Na<sub>2</sub>SO<sub>4</sub>), and the solvent was

(46) The high wavenumber of the C=O in the acetate groups has been observed in simple acetate protected phenols: *The Aldrich Library of 13C and 1H FTNMR Spectra*, 1st ed.; Pouchert, C. J., Behnke, J., Eds.; Aldrich Chemical Co.: Milwaukee, WI, 1993.

evaporated to yield **40** as an orange oil that was not stable to further purification:  $^1\text{H NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.83 (t,  $J = 7.4$  Hz, 6H), 1.35–1.40 (m, 4H), 2.26 (t,  $J = 6.4$  Hz, 4H), 3.83 (s, 6H), 4.08 (s, 6H), 6.52 (s, 2H), 6.69 (s, 2H).

**2,2',6,6'-Tetramethoxy-4,4'-dioxo-7,7'-dipropyl-4H,4'H-[1,1']-binaphthalenylidene-3,3'-dicarboxylic Acid Dimethyl Ester (69).** A mixture of MeCN (7 mL),  $\text{CH}_2\text{Cl}_2$  (3 mL), and  $\text{H}_2\text{O}$  (1 mL) was degassed with Ar for 15 min. Naphthol **59** (32.3 mg, 0.0532 mmol) was added followed by PIDA (44.2 mg, 0.118 mmol). The reaction mixture turned a dark blue color and was stirred for 1 h. The reaction mixture was diluted with EtOAc and  $\text{H}_2\text{O}$ , the phases were partitioned, and the organic layer was washed with  $\text{H}_2\text{O}$  and brine. After drying ( $\text{Na}_2\text{SO}_4$ ), the solvent was evaporated. Purification was accomplished with chromatography to yield 29 mg (91%) of a dark green resin that was not stable in solution over extended periods of time: IR (thin film) 2957, 1737, 1586, 1227  $\text{cm}^{-1}$ ;  $^1\text{H NMR}$  (500 MHz,  $\text{CD}_2\text{Cl}_2$ )  $\delta$  0.86 (t,  $J = 7.4$  Hz, 6H), 1.47–1.50 (m, 2H), 2.52 (m, 4H), 3.58 (s, 6H), 3.90 (s, 6H), 3.97 (s, 6H), 7.34 (s, 2H), 7.50 (s, 2H); HRMS (ES) calcd for  $\text{C}_{34}\text{H}_{37}\text{O}_{10}$  ( $\text{MH}^+$ ) 605.2387, found 605.2412. Bianthrone **69** was unstable in solution over time precluding  $^{13}\text{C NMR}$  characterization.

**(M)-Dimethyl 2,2',4,4',6,6'-Hexamethoxy-7,7'-dipropyl-1,1'-binaphthyl-3,3'-dicarboxylate (82).** To a solution of binaphthol (*M*)-**52d** (70 mg, 0.11 mmol) in DMF (2 mL) under an Ar atmosphere was added NaH (60%, 82 mg, 2.1 mmol). The mixture was stirred for 10 min, and then  $\text{CH}_3\text{I}$  (0.2 mL, 3.2 mmol) was added. The reaction was stirred at room temperature for 18 h. The yellow mixture was quenched with  $\text{H}_2\text{O}$ , and the aqueous phase was extracted with EtOAc. The organics were washed with brine and dried ( $\text{MgSO}_4$ ), and the solvent was evaporated in vacuo to yield a yellow oil. Purification was accomplished via chromatography (25% EtOAc/hexanes) to yield **82** as a resin (52 mg, 75%):  $[\alpha]_{\text{D}}^{20} -26.4$  ( $c$  0.22, MeOH, 86% ee); IR (thin film) 2956, 1731, 1592, 1493, 1454, 1224, 1085, 1015, 733  $\text{cm}^{-1}$ ;  $^1\text{H NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.74 (t,  $J = 7.3$  Hz, 6H), 1.37–1.41 (m, 4H), 2.44–2.49 (m, 4H), 3.31 (s, 6H), 3.95 (s, 6H), 3.97 (s, 6H), 4.13 (s, 6H), 6.90 (s, 2H), 7.40 (s, 2H);  $^{13}\text{C NMR}$  (125 MHz,  $\text{CDCl}_3$ )  $\delta$  167.9, 156.7, 153.6, 151.8, 135.5, 130.8, 127.1, 124.9, 120.4, 120.2, 100.2, 63.1, 62.3, 55.8, 52.9, 33.2, 23.2, 14.2; HRMS (ESI) calcd for  $\text{C}_{36}\text{H}_{42}\text{O}_{10}\text{Na}$  ( $\text{MNa}^+$ ) 657.2676, found 657.2666.

**(M)-Dimethyl 5,5'-Dihydroxy-2,2',4,4',6,6'-hexamethoxy-7,7'-dipropyl-1,1'-binaphthyl-3,3'-dicarboxylate (87).** To a solution of binaphthalene **82** (131 mg, 0.269 mmol) in  $\text{CF}_3\text{CH}_2\text{OH}$  (10 mL) under an Ar atmosphere was added  $\text{PhI}(\text{OC}(\text{O})\text{CF}_3)_2$  (199 mg, 0.463 mmol). After the purple mixture was stirred for 30 min, the solvent was evaporated in vacuo. The resultant mixture was treated with a mixture of 1 N NaOH (0.1 mL, 0.1 mmol) in  $\text{H}_2\text{O}/\text{THF}/\text{EtOH}$  (1:1:1, v/v/v) for 30 min, followed by acidification with 1 N HCl. The aqueous phase was extracted with EtOAc, the organics were washed with brine and dried ( $\text{Na}_2\text{SO}_4$ ), and the solvent was evaporated. Purification was accomplished by chromatography (25% EtOAc/hexanes) to yield product **87** as a yellow resin (70 mg, 51%, 82% ee):  $[\alpha]_{\text{D}}^{20} -12.1$  ( $c$  0.22, MeOH, 76% ee); IR (film) 3369, 2957, 1732, 1311  $\text{cm}^{-1}$ ;  $^1\text{H NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.74 (t,  $J = 7.2$  Hz, 6H), 1.35–1.42 (m, 4H), 2.42–2.51 (m, 4H), 3.32 (s, 6H), 3.93 (s, 6H), 3.98 (s, 6H), 4.16 (s, 6H), 6.42 (s, 2H), 9.24 (s, 2H);  $^{13}\text{C NMR}$  (125 MHz,  $\text{CDCl}_3$ )  $\delta$  167.1, 154.4, 152.6, 146.0, 142.6, 139.8, 132.9, 120.8, 119.2, 117.6, 114.2, 64.6, 62.3, 60.9, 53.2, 33.2, 23.9, 14.2; HRMS (ESI) calcd for  $\text{C}_{36}\text{H}_{42}\text{O}_{12}\text{Na}$  ( $\text{MNa}^+$ ) 689.2574, found 689.2599; CSP HPLC (Chiralpak AD, 1.0 mL/min, 98:2 hexanes/*i*-PrOH)  $t_{\text{R}}(\text{S}) = 12.3$  min,  $t_{\text{R}}(\text{R}) = 15.6$  min.

**(M)-5,5'-Bis(benzyloxy)-2,2',4,4',6,6'-hexamethoxy-7,7'-dipropyl-1,1'-binaphthyl-3,3'-dicarboxylic Acid (90).** To a solution of 68% ee binaphthol **87** (37 mg, 0.055 mmol) and benzyl bromide (46  $\mu\text{L}$ , 0.38 mmol) in DMF (1.0 mL) under an Ar atmosphere was

added 60% NaH (12.5 mg, 0.312 mmol). The dark green mixture was stirred for 15 h, quenched with  $\text{H}_2\text{O}$ , and washed with EtOAc. The organic phase was washed with  $\text{H}_2\text{O}$  and brine and dried ( $\text{Na}_2\text{SO}_4$ ), and the solvent was evaporated to yield an orange oil. Purification was accomplished by chromatography (15% EtOAc/hexanes) to yield the bisbenzyl ether (36 mg, 78%) as a yellow resin:  $[\alpha]_{\text{D}}^{20} -24.7$  ( $c$  1.0,  $\text{CHCl}_3$ , 68% ee); IR (thin film) 2939, 1735, 1330, 1094  $\text{cm}^{-1}$ ;  $^1\text{H NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.75 (t,  $J = 7.4$  Hz, 6H), 1.39–1.43 (m, 4H), 2.49–2.54 (m, 4H), 3.39 (s, 6H), 3.95 (s, 6H), 3.98 (s, 6H), 4.00 (s, 6H), 5.09 (m, 4H), 6.76 (s, 2H), 7.38 (t,  $J = 7.3$  Hz, 2H), 7.44 (t,  $J = 7.3$  Hz, 4H), 7.63 (d,  $J = 7.0$  Hz, 4H);  $^{13}\text{C NMR}$  (125 MHz,  $\text{CDCl}_3$ )  $\delta$  167.8, 154.4, 152.6, 146.8, 138.9, 138.4, 133.8, 129.2, 128.8, 128.3, 127.4, 123.1, 120.7, 120.3, 114.2, 77.1, 64.6, 62.3, 60.9, 53.2, 33.2, 23.9, 14.2; HRMS (ESI) calcd for  $\text{C}_{50}\text{H}_{54}\text{O}_{12}\text{Na}$  ( $\text{MNa}^+$ ) 869.3513, found 869.3541.

To a suspension of the bisbenzyl ether (163 mg, 0.192 mmol) in dioxane (8 mL) and  $\text{H}_2\text{O}$  (9 mL) was added LiOH· $\text{H}_2\text{O}$  (402 mg, 9.57 mmol), and the mixture was heated to reflux for 24 h. The brown mixture was cooled to room temperature and diluted with  $\text{H}_2\text{O}$ . The aqueous phase was washed with EtOAc, the organic phase was washed with  $\text{H}_2\text{O}$  and brine and dried ( $\text{Na}_2\text{SO}_4$ ), and the solvent was evaporated to recover 73 mg (0.086 mmol, 45%) of starting material. The aqueous phase was acidified with 1 N HCl and extracted with EtOAc, the organic phase was washed with  $\text{H}_2\text{O}$  and brine and dried ( $\text{Na}_2\text{SO}_4$ ), and the solvent was evaporated to yield the diacid **90** (81 mg, 93% based on recovered starting material) as an orange resin that was used for the next step without further purification:  $[\alpha]_{\text{D}}^{20} -29.1$  ( $c$  1.0,  $\text{CHCl}_3$ , 82% ee); IR (thin film) 2937, 1736, 1707, 1327, 1107  $\text{cm}^{-1}$ ;  $^1\text{H NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.76 (t,  $J = 7.3$  Hz, 6H), 1.40–1.44 (m, 4H), 2.50–2.54 (m, 4H), 3.44 (s, 6H), 3.72 (s, 6H), 3.98 (s, 6H), 3.99 (s, 6H), 5.11 (dd,  $J = 9.8, 14.9$  Hz, 4H), 6.80 (s, 2H), 7.36 (t,  $J = 7.3$  Hz, 2H), 7.43 (t,  $J = 7.3$  Hz, 4H), 7.64 (d,  $J = 7.0$  Hz, 4H);  $^{13}\text{C NMR}$  (125 MHz,  $\text{CDCl}_3$ )  $\delta$  172.5, 155.2, 152.4, 150.7, 146.9, 139.3, 138.2, 134.5, 129.3, 128.8, 128.3, 122.9, 122.1, 120.6, 120.1, 67.5, 65.2, 62.4, 61.7, 33.2, 23.9, 14.2; HRMS (ESI) calcd for  $\text{C}_{48}\text{H}_{50}\text{O}_{12}\text{Na}$  ( $\text{MNa}^+$ ) 841.3200, found 841.3240.

**(M)-5,5'-Bis(benzyloxy)-2,2',4,4',6,6'-hexamethoxy-7,7'-dipropyl-1,1'-binaphthyl (42).** To a solution of diacid **90** (26.7 mg, 0.0326 mmol) in DMSO–DMF (0.3 mL) was added Pd( $\text{O}(\text{C}(\text{O})\text{CF}_3)_2$ ) (22.8 mg, 0.0739 mmol). The mixture was heated in a 90 °C oil bath for 1 h. After the brown reaction mixture was cooled to room temperature, 1 N HCl was added, and the mixture was extracted with EtOAc. The organic phase was washed with  $\text{H}_2\text{O}$  and brine and dried ( $\text{Na}_2\text{SO}_4$ ), and the solvent was evaporated to yield a brown oil. This material was dissolved in THF (1.0 mL) and stirred under a  $\text{H}_2$  atmosphere for 5 min. After filtration through Celite, purification was accomplished via chromatography (20% EtOAc/hexanes) to yield **42** (20.7 mg, 87%) as a resin:  $[\alpha]_{\text{D}}^{20} 0$  ( $c$  1.0,  $\text{CHCl}_3$ , 70% ee) (this compound was resynthesized but the optical rotation was still 0; the compound was carried on to the next step and enantioselectivity confirmed by CSP HPLC (69% ee)); IR (thin film) 2933, 1338  $\text{cm}^{-1}$ ;  $^1\text{H NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.75 (t,  $J = 7.2$  Hz, 6H), 1.35–1.42 (m, 4H), 2.46–2.53 (m, 4H), 3.74 (s, 6H), 3.92 (s, 6H), 3.99 (s, 6H), 5.10 (m, 4H), 6.64 (s, 2H), 6.76 (s, 2H), 7.37 (t,  $J = 7.3$  Hz, 2H), 7.43 (t,  $J = 7.3$  Hz, 4H), 7.62 (d,  $J = 7.0$  Hz, 4H);  $^{13}\text{C NMR}$  (125 MHz,  $\text{CDCl}_3$ )  $\delta$  157.1, 154.9, 148.8, 147.2, 138.9, 137.0, 133.7, 129.0, 128.5, 127.8, 122.1, 116.5, 112.5, 95.9, 76.2, 61.4, 57.5, 56.1, 32.8, 23.6, 14.0; HRMS (ESI) calcd for  $\text{C}_{46}\text{H}_{50}\text{O}_8\text{Na}$  ( $\text{MNa}^+$ ) 753.3403, found 753.3427.

**4,9-Dihydroxy-2,6,7,11-tetramethoxy-1,12-dipropylperylene-3,10-dione (39).** To a solution of binaphthalene **42** (6.4 mg, 0.0088 mmol) in MeOH (0.3 mL) and THF (0.3 mL) was added 10% Pd/C (2 mg). The mixture was stirred under an atmosphere of  $\text{H}_2$  for 3 h. The reaction was filtered through Celite, rinsing with MeOH and  $\text{CH}_2\text{Cl}_2$ . The solvents were evaporated to yield an unstable yellow oil (69% ee):  $^1\text{H NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.76

(t,  $J = 7.3$  Hz, 6H), 1.38–1.43 (m, 4H), 2.43–2.47 (m, 4H), 3.69 (s, 6H), 3.89 (s, 6H), 4.16 (s, 6H), 6.37 (s, 2H), 6.70 (s, 2H), 9.28 (s, 2H); CSP HPLC (Chiralpak AD, 1.0 mL/min, 98:2 hexanes/*i*-PrOH)  $t_{R(S)} = 37.9$  min,  $t_{R(R)} = 45.2$  min.

To a solution of bisnaphthol (11 mg, 0.020 mmol) in Et<sub>2</sub>O (1 mL) was added MnO<sub>2</sub> (60 mg, 0.69 mmol). The mixture was stirred for 30 min and filtered through Celite, and the solvent was evaporated to yield a red resin, which was chromatographed (2.5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to yield the perylenequinone as a red resin:  $[\alpha]_D^{20} -410$  (*c* 0.021, MeOH, 69% ee); <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.54 (t,  $J = 7.3$  Hz, 6H), 0.96–0.99 (m, 2H), 1.18–1.22 (m, 2H), 2.48–2.51 (m, 2H), 3.08–3.11 (m, 2H), 4.05 (s, 6H), 4.11 (s, 6H), 4.17 (s, 6H), 6.78 (s, 2H).

To a solution of the above perylenequinone product (12 mg, 0.0221 mmol) in THF (2 mL) under an argon atmosphere was added a solution of MgI<sub>2</sub> in Et<sub>2</sub>O (0.07 M, 665  $\mu$ L, 0.0464 mmol). The dark purple mixture was stirred for 10 min (until the mixture turns from purple to black), diluted with EtOAc, washed with saturated aq NH<sub>4</sub>Cl, and dried (Na<sub>2</sub>SO<sub>4</sub>). Concentration yielded a red residue, which was chromatographed (2.5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to yield product **39** as a red resin (10 mg, 87% (69% for the three steps)): <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.52 (t,  $J = 7.2$  Hz, 6H), 0.88 (m, 2H), 1.25 (m, 2H), 2.76 (m, 2H), 3.33 (m, 2H), 4.09 (s, 6H), 4.21 (s, 6H), 6.59 (s, 2H), 15.82 (s, 2H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  14.1, 24.3, 35.0, 56.3, 61.1,

101.1, 105.7, 116.5, 126.2, 127.5, 140.5, 151.0, 166.5, 174.1, 176.8; IR (film) 3383, 2927, 2858, 1607, 1460, 1274, 1213 cm<sup>-1</sup>; HRMS (ES) calcd for C<sub>30</sub>H<sub>31</sub>O<sub>8</sub> (MH<sup>+</sup>) 519.2019, found 519.2036; see the Supporting Information for the CD spectrum; CSP HPLC (Chiralpak AD-H, 0.5 mL/min, 95:5 hexanes/*i*-PrOH)  $t_{R(P)} = 15.9$  min,  $t_{R(M)} = 17.1$  min.

**Acknowledgment.** We are grateful to the NIH (CA-109164) and NSF (CHE-0911713) for financial support. Partial instrumentation support was provided by the NIH for MS (1S10RR023444) and NMR (1S10RR022442). We thank 3D Pharmaceuticals (C.A.M.), Novartis (B.J.M.), and the Division of Organic Chemistry of the American Chemical Society (C.A.M., B.J.M.) for graduate fellowships. We thank Feng Gai for assistance with CD measurements. Thank you to Genette McGrew for the cover art preparation. The background to the cover art is being used with permission of Wiley-Blackwell: Daub, M. E.; Herrero, S.; Chung, K.-R. *FEMS Microbiol. Lett.* 2005, 252, 197-206.

**Supporting Information Available:** Additional experimental descriptions, NMR spectra, and crystallographic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.