

**Correction to Synthesis of Tetrahydropyrans from Propargyl Alcohols and 1,1-Cyclopropanediester: A One-Pot Ring-Opening/Conia-ene Protocol** [*J. Org. Chem.* **2009**, *74*, 8414. DOI: 10.1021/jo9019122]. Andrew B. Leduc, Terry P. Lebold, and Michael A. Kerr\*

Page 8414. The following references regarding the use of propargyl alcohols in a Michael addition/Conia-ene process were regrettably omitted:

Marat, X.; Monteiro, N.; Balme, G. *Synlett* **1997**, 845. Cavicchioli, M.; Marat, X.; Monteiro, N.; Hartmann, B.; Balme, G. *Tetrahedron Lett.* **2002**, *43*, 2609. Nakamura, M.; Liang, C.; Nakamura, E. *Org. Lett.* **2004**, *6*, 2015. Morikawa, S.; Yamazaki, S.; Furusaki, Y.; Amano, N.; Zenke, K.; Kakiuchi, K. *J. Org. Chem.* **2006**, *71*, 3540. Morikawa, S.; Yamazaki, S.; Tsukada, M.; Izuhara, S.; Morimoto, T.; Kakiuchi, K. *J. Org. Chem.* **2007**, *72*, 6459.

DOI: 10.1021/jo101123d

Published on Web 06/21/2010