


Faulknerynes A–C from a Bahamian Sponge *Diplastrella* sp.:
Stereoassignment by Critical Application of Two Exciton
Coupled CD Methods

Jaeyoung Ko,[†] Brandon I. Morinaka,[†] and Tadeusz F. Molinski^{*,†,§}

[†]Department of Chemistry and Biochemistry and [§]Skaggs School of Pharmacy and Pharmaceutical Sciences,
University of California, San Diego, La Jolla, California 92093-0358, United States

tmolinski@ucsd.edu

Received November 2, 2010



Long-chain polyacetylene alcohols, faulknerynes A–C, along with known compounds diplynes A, C and E, were isolated from two specimens of the encrusting sponge, *Diplastrella* sp., collected from the surface of coral in the Bahamas. Two CD methods were critically evaluated for their suitability to terminal propargylic glycols and applied to assignment of configurations of faulkneryne A and diplyne C.

Introduction

Long chain polyacetylenic alcohols (PAAs) are characteristic natural products of Haplosclerid and Spirastrellid sponges, including the genera *Diplastrella*,¹ *Petrosia*, *Siphonochalina*, *Haliciona*,² and *Cribochalina*.³ Among the first to be reported were the highly cytotoxic duryne³ and the HIV reverse-transcriptase inhibitor petrosynol⁴ from *Cribochalina* and *Petrosia*, respectively. Many PAAs are exceedingly cytotoxic toward tumor cell lines (IC₅₀'s ≤ 30 ng/mL), a biological property that has been correlated with the presence of a terminal propargylic alcohol.² Faulkner and co-workers reported C₁₆ brominated diynes A, B, C (**1**), and E (**2**) and three related sulfate half-esters from the Philippines sponge *Diplastrella* sp.¹ Fusetani and co-workers described the C₂₃ polynols, callyspongiols A–E, along with

related hydrocarbons, from *Callyspongia truncata*⁵ and siphonodiol (**3**) from *Siphonochalina truncata*.^{6,7} In a screen for compounds with antiproliferative activity, extracts of two specimens of the Bahamian sponge *Diplastrella* sp. were found to elicit moderate in vitro cytotoxicity against the cultured tumor cell line HCT-116. We describe new brominated long-chain PAAs, faulknerynes A–C (**4a–c**), from these extracts and assignment of the absolute configuration of **1** and **4a** using two microscale methods based on exciton coupling circular dichroism (ECCD). We also establish that **1** obtained from *Diplastrella* sp. from the Bahamas and the Philippines are both (*R*).¹

Results and Discussion

A sample of the thinly encrusting *Diplastrella* sp., recovered from small patches laboriously scraped from coral substrate with a knife, was exhaustively extracted with

(1) Lerch, M. L.; Harper, M. K.; Faulkner, D. J. *J. Nat. Prod.* **2003**, *66*, 667–670.

(2) Zhou, G.-X.; Molinski, T. F. *Mar. Drugs* **2003**, *1*, 46–53.

(3) Wright, A. E.; McConnell, O. J.; Kohmoto, S.; Lui, M. S.; Thompson, W.; Snader, K. M. *Tetrahedron Lett.* **1987**, *28*, 1377–1379.

(4) Isaacs, S.; Kashman, Y.; Loya, S.; Hizi, A.; Loya, Y. *Tetrahedron* **1993**, *49*, 10435–10438.

(5) Tsukamoto, S.; Kato, H.; Hirota, H.; Fusetani, N. *J. Nat. Prod.* **1997**, *60*, 126–130.

(6) Tada, H.; Yasuda, F. *Chem. Lett.* **1984**, 779–780.

(7) Fusetani, N.; Sugano, M.; Matsunaga, S.; Hashimoto, K. *Tetrahedron Lett.* **1987**, *28*, 4311–4312.

MeOH. The MeOH extract was progressively adjusted in water content and partitioned against solvents of increasing polarity (hexane, CHCl_3 , and *n*-BuOH). The CHCl_3 -soluble fraction was further purified to give the known diplyne C (**1**, identified from NMR data¹) and new natural products, falknerynes A–C (**4a–c**).⁸ The formula $\text{C}_{16}\text{H}_{21}\text{BrO}_2$ for **4a** was secured from HRMS measurements ($[\text{M} + \text{Na}]^+$, 347.0633, $\Delta\text{mmu} = 1.6$). Examination of the ^1H NMR spectrum of **4a** (Table 1) readily revealed chemical shifts of the two chain termini: a 1,2-disubstituted vinyl bromide (δ 6.01, d, $J = 13.5$ Hz; 6.17, dt, J 13.5, 7.3 Hz) and a mutually coupled ABX system corresponding to a 1,2-glycol (δ 3.72, dd, $J = 11.4$, 6.2 Hz, H1a; δ 3.78, dd, $J = 11.4$, 3.8 Hz, H1b; 4.57, dd, $J = 6.2$, 3.8, H2). The formula required an additional five degrees of unsaturation, which were assigned to a conjugated (*Z*)-1,3,5-ene-diyne system, based on a red-shifted UV spectrum [**4a**; λ_{max} 271 nm] compared to **1** and **3** [**3**; λ_{max} 228 (ϵ 15,100)⁶], the presence of four acetylenic ^{13}C NMR sp signals (δ 71.1 s, 76.5 s, 76.8 s, 80.1 s) and ^1H NMR signals ascribed to a (*Z*)-1,2-disubstituted double bond, C7–C8 (δ 5.49, d, $J = 10.9$ Hz; δ 6.11, dt, $J = 10.9$, 7.6 Hz) conjugated to the diyne. HMBC crosspeaks from H7 to the acetylenic carbon signals of C3–C5 corroborated UV evidence that the (*Z*)-double bond was conjugated to the diyne. The remaining signals were assigned to allylic CH_2 groups and an unresolved methylene chain. Therefore, **4a** is (*Z*)-7,8-dehydro-**1**.




TABLE 1. NMR Data (CDCl_3) for Falkneryne A (**4a**)

no.	δ_{C} , mult ^a	δ_{H} , mult (J in Hz) ^b	COSY ^c	HMBC (H \rightarrow C)
1	66.4, CH	3.72, dd (11.4, 6.2)	H2	C2, C3
	66.4, CH	3.78, dd (11.4, 3.8)	H2	C3
2	64.0, CH	4.57, dd (6.2, 3.8)	H1, H7	C1, C3, C4, C5
3	80.1, C			
4	71.1, C			
5	76.8, C ^d			
6	76.5, C ^d			
7	107.8, CH	5.49, d (10.9)	H2, H8, H9	C3, C4, C5, C8, C9
8	149.5, CH	6.11, dt (10.9, 7.6)	H7, H9	C6, C7, C9
9	30.9, CH_2	2.33, (7.6)	H7, H8, H10	C6, C7, C8, C11
10	28.7, CH_2^e	1.41, m	H9, H11	C8, C9, C11
11	28.8, CH_2^e	1.30, m	H10, H12	C12
12	28.9, CH_2	1.31, m	H11, H13	C11
13	28.6, CH_2	1.39, m	H12, H14	C14, C15
14	33.0, CH_2	2.03, m (7.3)	H13, H15, H16	C12, C15, C16
15	138.3, CH	6.17, dt (13.5, 7.3)	H14, H16	C13, C14, C16
16	104.3, CH	6.01, d (13.5)	H14, H15	C14, C15

^a125 MHz. ^b600 MHz. ^cMultiplicity assigned from HSQC. ^dPeaks may be interchanged. ^ePeaks may be interchanged.

Falknerynes **B** (**4b**) and **C** (**4c**) ($\text{C}_{16}\text{H}_{17}^{79}\text{BrO}_2$, m/z 343.0304 [$[\text{M} + \text{Na}]^+$ $\Delta\text{mmu} = 0.5$]), obtained in amounts of 4 and 14 μg , respectively, are dehydro-derivatives of **2** and isomeric with each other. Analysis of COSY spectrum and vicinal J values of the vinyl proton signals reveals that the two compounds differ only in the location of the (*Z*)-carbon–carbon double bond; in **4b**, it is positioned at C-7–C-8 while compound **4c** has a 11-*Z* double bond in conjugation with the terminal ene-yne. Therefore, **4b** is C-13,14-didehydro falkneryne A and **4c** is the 11-*Z* geometrical isomer of 11-*E* diplyne E (**2**).

With diplyne C (**1**) from *Diplastrella* sp. collected in two different oceans, one from the Pacific and the other from the Atlantic, an opportunity presented itself for configuration analysis by chiroptical comparison, although not without some challenges. The optical rotation of natural **1** was not reported, although a total synthesis of (*S*)-(+)-**1** ($[\alpha]_{\text{D}} +13.3$)⁹ and unnatural (*S*)-(+)-diplyne A ($[\alpha]_{\text{D}} +9.6$)¹⁰ and chiroptical comparison with levorotatory diplyne A ($[\alpha]_{\text{D}} -8.7$)¹ would suggest an (*R*) configuration for both diplynes A and C (**1**). Insufficient amounts of **4a–c** were available to record rotations for reliable comparison of $[\alpha]_{\text{D}}$ with **1** or **3**, so attention was turned to alternative methods. Configurational assignment of secondary alcohols is frequently carried out by NMR using the modified Mosher's method;¹¹ however, the limited amount of available sample necessitated deployment of sensitive CD spectroscopy, a method well suited to “nanomole-scale” natural products investigation.¹² The stereochemistry of siphonodiol (*R*)-(-)-(**3**) was assigned by Fusetani and co-workers using the dibenzoate exciton coupling method; specifically, the observation of a negative

(8) The compounds are named in honor of the late D. John Faulkner (1940–2002) a pioneer in marine natural products who described diplynes A–E (ref 1), related polyacetylenic alcohols, and hundreds of other compounds. Andersen, R. J.; Ireland, C. M.; Molinski, T. F.; Bewley, C. B. *J. Nat. Prod.* **2004**, *67*, 1239–1251.

(9) Gung, B. W.; Gibeau, C.; Jones, A. *Tetrahedron: Asymmetry* **2005**, *16*, 3107–3114.

(10) Gung, B. W.; Gibeau, C.; Jones, A. *Tetrahedron: Asymmetry* **2004**, *15*, 3973–3977.

(11) Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. *J. Am. Chem. Soc.* **1991**, *113*, 4092–4096.

(12) (a) Molinski, T. F. *Curr. Opin. Drug Discovery Dev.* **2009**, *12*, 197–206. (b) Molinski, T. F. *Curr. Opin. Biotechnol.* **2010**, *21*, 819–826.

bisignate Cotton effect, CE [λ 324 nm ($\Delta\epsilon$ -25.3); 303 (+24.2)] in the exciton coupled CD spectrum of **3a**, the derived 4-dimethylaminobenzoate diester (DMB diester).⁷ From exciton coupling theory,¹³ the sign of the split CE is predicted from the helicity (twist) of the electronic dipole transition vectors associated with the prominent 1L_a charge-transfer transition in degenerate benzoate chromophores. These vectors are approximately aligned with the corresponding C–O bond directions; a negative sign for the long-wavelength band of the split CE for **3a** indicates a negative helicity of the two C–O vectors, and a positive split CE indicates a positive helicity. The strength of the ECCD signal depends upon several other factors and is conveniently reported as the difference between maximum and minimum of the bisignate split CE [e.g., for **3a**, $A = 24.2 - (-25.3) = 49.5$].¹³

Glycol dibenzoate ECCD is dependent upon the dihedral angle of O–C–C–O around C1–C2. Because *acyclic* 1,2-glycols have a freely rotating C1–C2 bond, the *absolute configurational* assignment requires knowledge of the limiting staggered conformers. We were concerned about two implicit assumptions used in the assignment of **3** and, consequently, our approach to **1** and **4a**: a low barrier to rotation of the C1–C2 bond and similar energies of staggered conformers due to the presence of the relatively sterically unencumbered acetylenic group and additional exciton coupling from the diyne chromophore with the propargylic DMB chromophore. Dibenzoate esters of **1–4** would be expected to exhibit both types of dichromophoric interactions: yne-benzoate and benzoate-benzoate.¹⁴ Consequently, room temperature CD measurements of propargylic benzoates would sample significant proportions of populated staggered conformers **a–c**, with **a** and **c** giving rise to opposing signs of ECCDs, and small structural differences may even contribute nonstaggered conformers that further influence the magnitude or even the sign of the ECCD. In order to quantify some of these expected differences, we calculated the energies of the conformers of propargylic 1,2-glycol dibenzoates (Spartan, PM3) and ordered the lowest lying conformers (Figure 1).

Perhaps not unexpectedly, the potential energy surface of the model propargylic 1,2-benzoate was found to be fairly shallow. Nevertheless, for the dibenzoates of (*R*)-**1**, the three lowest energy conformers depicted in Figure 1d,e comprising 47% of the Boltzmann distribution, subtended the same negative helicity corresponding to the idealized geometry (a). The torsional angle, θ , ranged from -55° to -84.1° about the C–O vectors of the (C=O)O groups and differed only in minor changes in orientation about the benzoate ester bonds. Surprisingly, the nearest conformers with antiperiplanar dispositions ($\theta \sim 180^\circ$) of the two benzoate groups and expected to give zero contribution to the ECCD (cf. idealized geometry of Figure 1a) were higher in energy and ranked fifth and sixth; together, they contribute only 13.2% to the population [$E = -0.54$ kcal mol $^{-1}$ (7.0%); $E = -0.27$




FIGURE 1. Conformers of propargylic 1,2-glycol *O*-dibenzoate analogues of **5**. (a–c) Idealized geometries, helicities, and signs of contributions to ECCD ($\Delta\epsilon$). (d–f) Calculated conformers of propargylic 1,2-glycol *O*-dibenzoate analogues of **5** ranked in relative energy, E (kcal mol $^{-1}$); θ , the dihedral angle of the C–O bond vectors; and % Boltzmann populations (Spartan 08, semi-empirical PM3, gas phase): (d) 0, -55.6° , (22.3%); (e) 1.29, -84.1° , (13.3%); (f) 1.67, -55.0° (11.4%). Only the three best ranked conformations are shown (see text). For ease of calculation, the chain is truncated to a 3,5-diyne.

kcal mol $^{-1}$ (6.2%)]. The *gauche–gauche* conformers (not shown, but cf. Figure 1b), which are expected to give ECCD of opposing sign to Figure 1c, were found to be of even lower ranking and less stable by more than 2.8 kcal mol $^{-1}$ than the favored conformers; they represented less than 3% in the conformer population.

ECCD of arenecarboxylate esters (acylates) of allylic and propargylic alcohols arise from electronic coupling of arene and C–C multiple bond π – π^* transitions. Reliable assignments of secondary allylic alcohols have been made from ECCD of the corresponding allylic *O*-benzoates¹⁵ *O*-naphthoates,¹⁶ and *O*-naphthoates of conjugated dienes.¹⁷ The model used for interpretation of ECCD in allylic benzoates/naphthoates¹⁵ follows from well-understood considerations of allylic strain¹⁸ leading to a dominant conformation that is supported by NMR. A large number of examples of chiral allylic benzoates were studied, which consistently revealed large vicinal coupling constants ($J \approx 9$ Hz) for vicinal protons on the adjacent sp^2 – sp^3 carbons that imply the vinyl H is eclipsed by the allylic methine H.^{15,17} In turn, this sets the torsional angle between the C=C bond and the benzoate C–O vector and the relative helicity of the respective chromophoric transition dipoles. The allylic ECCD method should be extendable to propargylic acylates; however, conformations of the latter are expected to exhibit subtle differences (Figure 2a,b). In both double and triple carbon–carbon bonds the π – π^* transition dipoles are polarized along the C–C direction. Unlike the planar C=C bonds in allylic acylates, the radially symmetrical triple bond in a propargylic acylate is essentially a free rotor. A consequence of this, as we

(13) Harada, N.; Nakanishi, K. *Circular Dichroic Spectroscopy: Exciton Coupling in Organic Stereochemistry*; University Science Books: Mill Valley, CA, 1983; p 162.

(14) Presumably, for this reason, DMB diesters were chosen for the dibenzoate assignment of **2** to move the bisignate split CE to longer wavelengths ($\lambda \sim 310$ nm) and diminish potential interference from diyne–benzoate interactions (ref 7).

(15) Gonnella, N. C.; Nakanishi, K.; Martin, V. S.; Sharpless, K. B. *J. Am. Chem. Soc.* **1982**, *104*, 3775–3776.

(16) Molinski, T. F.; Brzezinski, L. J.; Leahy, J. W. *Tetrahedron: Asymmetry* **2002**, *13*, 1013–1016.

(17) Schneider, C.; Schreier, P.; Humpf, H. U. *Chirality* **1997**, *9*, 563–567.

(18) Johnson, F. *Chem. Rev.* **1968**, *68*, 375–413.




FIGURE 2. Conformers of propargylic (*R*)-2-*O*-(2'-naphthoate) esters of terminal propargylic 1,2-glycols. (a, b) Idealized geometries (two views) (cf. 6) and the predicted sign of $\Delta\epsilon$ in the ECCD spectrum. ($R = \text{CH}_2\text{OAc}$, $R' = \text{alkenyl}$, alkynyl; Nph = 2'-naphthoyl; shaded bar is 2-naphthyl (C_{10}H_7)). (c, d) Two views of the energy minimized model (MMFF, Spartan 08) (cf. 6; however, the chain is truncated to a 3,5-diyne, $R' = \text{ethynyl}$), showing a *syn*-clinal arrangement of the conjugated diyne- and naphthoate plane with a positive helicity (dihedral angle of $\text{C3}-\text{C2}-\text{C2}'-\text{C6}'$, $\theta = +33.4^\circ$).

showed from molecular mechanics calculations (Spartan, MMFF94, Figure 2c,d), is that the dominant conformer places $\text{H}\alpha$ almost *syn*-coplanar with the conjugated ester $\text{Ar}(\text{C}=\text{O})\text{O}$ group and an angle of $\pm 20^\circ$ is subtended between the synclinal arenecarboxylate plane and the acetylenic vector. For propargylic esters of (*R*)-**1** (Figure 2c), the respective transition dipoles have a calculated positive helicity ($\theta = +33.4^\circ$), where the dihedral θ is approximated by the dihedral angle $\text{C3}-\text{C2}-\text{C2}'-\text{C6}'$ ($\text{C2}'$, $\text{C6}'$ are the *ipso* and *para* carbons of the first benzenoid ring of the naphthoate). In turn, this predicts that (*R*)-**1** will exhibit a positive $\Delta\epsilon$ in the CD spectrum. Harada has described similar arguments and assignments of alk-1-yn-3-ols by ECCD after a two step-derivatization: Sonogashira coupling with a terminal acetylene with an aryl iodide followed by esterification of the secondary alcohol to the corresponding arene-yne benzoate and measurement of CD spectra.¹⁹ The sign of the resulting split CEs conform to the rule; however, the major limitation of the method is its applicability to terminal acetylenes only.

An advantage of allylic *O*-naphthoates over *O*-benzoates is the very intense Cotton effects (A values of up to ~ 200 ¹⁷) associated with the 1B_b transition along the skew long axis of the naphthalene ring system, closely parallel with the $\text{C}-\text{O}$ ester bond, which is better suited for subnanomole CD measurements. Unlike vicinal propargylic dibenzoates, interpretation of CD in propargylic naphthoates benefits from the simplicity of only two participant chromophores in exciton coupling.

In order to assess the relative merits of the dibenzoate method and the propargylic *O*-naphthoate method, we chose to determine the configuration of **1** and **4a** using both approaches. Dipyne **1** was acylated (Scheme 1) (*N*-(4-dimethylaminobenzoyl)imidazole, DBU, CH_3CN , 25°C) to provide the DMB diester **5** ($\sim 34\ \mu\text{g}$),²⁰ which was purified by

(19) Naito, J.; Yamamoto, Y.; Akagi, M.; Sekiguchi, S.; Watanabe, M.; Harada, N. *Monatsh. Chem.* **2005**, *136*, 411–445.

(20) The yields of **5** and **6** were estimated by UV-vis assuming λ_{max} 313 nm ($\epsilon = 56,000$) and λ_{max} 239 nm ($\epsilon = 65,000$), respectively.






FIGURE 3. CD spectra (EtOH, 23°C) of (a) **5** and **16** and (b) **6**.

SCHEME 1. Derivatization of Polyacetylenic Alcohols to Chromophoric Derivatives **5**, **6**, and **16**



HPLC. The CD spectrum of **5** (Figure 3a) exhibited a pronounced ECCD [λ 301 nm ($\Delta\epsilon +22.4$), 326 (-21.5)] of the same sign and similar magnitude observed for the DMB diesters of **3**,⁷ indicating the same C2 configuration. Independently, **1** was converted to a C2-O -(2'-naphthoate) monoester **6** as follows (Scheme 1): the primary hydroxyl group was selectively acetylated in the presence of a lipase (Novozyme

SCHEME 2. Preparation of En-yne and Di-yne Propargylic 2-Naphthoate Esters (*S*)-7 and (*S*)-8^a



^aNote: CIP priorities of 7–15 differ from those of 1–4.

435, vinyl acetate, CH₃CN), and the free secondary OH group was subsequently acylated, as above, to give diester **6** (~24 μg) after HPLC purification. The CD (EtOH) spectrum of **6** (Figure 3b) revealed a positive long wavelength band CE [λ 238 nm (Δε +25.3)]²¹ that was interpreted as arising from a positive helicity between the diyne and naphthoate chromophores.²⁴ The negative component of the ECCD was observed clearly when the CD spectrum of **6** was recorded in CH₃CN [λ 238 nm (Δε +29.7), 188 nm (−50.5)]. Both methods lead to the same answer, and we assign the (*R*) configuration to **1**, **5**, and **6**.

To validate the outcome of assignment of **1** by the propargylic acylate ECCD method, CD spectra were obtained of 2-naphthoate esters (*S*)-7 and (*S*)-8 prepared from the known (*S*)-9 propargylic alcohol²² (note the change in CIP priorities) as shown in Scheme 2. Desilylation of (*S*)-9 to alcohol (*S*)-10 was followed by protection as the acetate ester and Sonogashira coupling with (*E*)-iododecene in the presence of bis(triphenylphosphine)palladium(II) dichloride (**11**) and CuI to give en-yne acetate ester **13**. Exchange of the *O*-acyl groups (acetyl for 2-naphthoate) through the intermediary alcohol **14** was achieved in two steps (NH₃, MeOH; 2-naphthoic acid, EDC, DMAP, 76% over two steps) to provide en-yne model ester (*S*)-7.²³ In parallel, (*S*)-10 was acylated to the naphthoate ester **15**, which was desilylated (TBAF, 86%) to give the propargylic *O*-naphthoate **15a** and transformed into the diyne naphthoate (*S*)-8 (CuCl, NH₂OH·HCl, *n*-BuNH₂, 1-bromoheptyne).

The CD spectra of (*S*)-7 and (*S*)-8 (CH₃CN) showed strong bisignate CEs [(*S*)-7 λ 188 nm (Δε −37.8); (*S*)-8 225 (−24.6), 242 (+35.3)] (Figures 4 and 5). Note that the

secondary alcohol (*S*)-14, corresponding to (*S*)-7 and lacking a dichromophoric exciton coupling, shows essentially only baseline CD (Figure 3). Because the π–π* transition of the conjugated diyne occurs at lower wavelenths [λ 214 nm (ε 37,300), Figure 5],¹ the short-wavelength component of the exciton couplet of (*S*)-8 [λ 188 nm (Δε −37.8)] is at the edge of instrument detection limits; however, in (*S*)-7 this high energy component reveals itself readily [λ 225 nm (Δε −24.6)] due to the presence of a red-shifted extended en-yne chromophore.²⁴ Nevertheless, as noted by Nakanishi




FIGURE 4. CD spectra at 25 °C of (a) (*S*)-7 in CH₃CN, (b) (*S*)-7 in MeOH, and (c) (*S*)-14 in hexane.

(21) Note that the remote C10–C14 diyne chromophore is “CD silent”.

(22) (a) Skepper, C. K. Marine Derived Heterocycles. Ph.D. Dissertation, University of California, San Diego, La Jolla, CA, November, 2009. (b) Supporting information of Skepper, C. K.; MacMillan, J. B.; Zhou, G. X.; Masuno, M. N.; Molinski, T. F. *J. Am. Chem. Soc.* **2007**, *129*, 4150–4151.

(23) The configuration of the secondary alcohol **14** was verified using the modified Mosher’s method.¹¹

(24) Generally, the short-wavelength component of the ECCD is obscured (λ < 200 nm) in most allylic benzoates with mono- and disubstituted double bonds. However, this component is red-shifted and may be observable at λ ~180–200 nm in allylic 2-naphthoates (see ref 16) with tri- and tetra-alkyl substituted double bonds and, of course, in allylic acylates of conjugated dienes (λ > 220 nm).




FIGURE 5. CD spectra for (*S*)-**8** in (a) CH₃CN, (b) MeOH ($\lambda < 200$ nm, obscured by solvent end-absorption), and (c) hexane at 25 °C.

and co-workers,¹⁵ only the sign of the long-wavelength component is necessary and sufficient to assign allylic benzoates which also applies for propargylic naphthoates. Due to the stronger oscillator strength of the ¹B_b transition, even acyclic nonconjugated propargylic *O*-naphthoates should be amenable to chiroptical analysis, unlike the corresponding *O*-benzoates that show only weak or nondetectable ECCD from the shorter wavelength acetylene chromophore ($\lambda < 190$ nm).¹⁹ This is demonstrated in the CD spectrum of **15a** (Figure 6), which displays a weaker yet still prominent positive component [λ 240 nm ($\Delta\epsilon +4.6$)] of the biphasic Cotton effect.

Since both the dibenzoate and the propargylic naphthoate ECCD methods lead to the same configurational assignment in **1**, it can be concluded that both can be reliably used for stereoassignments of other long-chain propargylic alcohols under appropriate conditions, although the one-step dibenzoate method is attractive for its operational simplicity.

A simple extension of the glycol benzoate method was applied to faulkneryne A (**4a**). Conversion of **4a**, by the aforementioned procedure, gave the di-DMB ester **16** which showed an CD spectrum (EtOH, Figure 3a) with a negative split CE [λ 302 nm ($\Delta\epsilon +17.4$), 327 (-9.0)], similar to that of **5**. As with **5**, it is fortuitous that both the short-wavelength component of the dibenzoate ECCD of **16** and the super-

(25) In fact, this is evident from the *relative* asymmetry of the split CEs in **16** and **5** due to differences of superposed paired propargylic *O*-DMB-alkyne ECCDs arising from the diyne $\pi-\pi^*$ transition in **5** versus the red-shifted 7-en-3,5-diyne $\pi-\pi^*$ transition in **16**.




FIGURE 6. CD spectrum of (*S*)-**15a** (CH₃CN, 25 °C).

posed ene-diyne benzoate ECCD interaction (cf. **6**, Figure 3b) are of the same sign (positive)²⁵ and reinforce at $\sim\lambda$ 302 nm.²⁶ Therefore, the configurations of **4a** and **16**, are revealed to be (*R*). Assuming congeneric **4b,c** are of the same configuration as **4a**, it would appear the biosynthesis of propargylic diols **1–4** from sponges across two different oceans conserves the C2 stereochemistry.²⁷

Diplyne C (**1**) exhibited cytotoxicity against cultured human colon tumor cells (HCT-116; LD50 3.6 $\mu\text{g}/\text{mL}$); however, insufficient amounts of **4a–c** were available for biological evaluation.

Conclusions

The complete structures of new long-chain polyacetylenic diols, faulknerynes A–C (**4a–c**), were elucidated with the aid of microcryoprobe NMR spectroscopy. The configurations of **4a** and the known compound diplyne C (**1**) were assigned the (*R*) configuration by two different microscale ECCD methods that converged upon the same answer. The ECCD spectrum of propargylic en-yne naphthoates are nicely suited for observation of both halves of the exciton couplets for microscale configurational analysis of polyacetylenic alcohols without the necessity for additional synthetic elaboration of the acetylenic chromophore.

Experimental Section

General Experimental Procedures. See Supporting Information.

Animal Material. The sponge *Diplastrella* sp., Family Spirastrellidae (Ridley & Dendy 1886) (08-05-026) was collected on 31 May 2008 at Sweetings Cay, Bahamas (26° 34.080' N, 77° 53.407' W) at a depth of -19 m using scuba and kept in EtOH at -20 °C until extraction. The tissue was pale pink to red in life, thinly encrusting on coral in small patches. Microscopic spicule analysis (SEM) showed tylostyles with rounded heads and microscleres with a preponderance of diplasters over spirasters

(26) Insufficient sample of **4a–c** remained to apply the ene-diyne naphthoate ECCD method.

(27) The *unnatural* (*S*) enantiomer of siphonodiol **2** is depicted in ref 1.

as is typical for *Diplastrella*.²⁸ A second sponge, *Diplastrella* sp. (08-07-038), was collected in the same vicinity at similar depths. Voucher samples for all animal material are archived at UCSD.

Extraction and Isolation. The EtOH extract of the sponge *Diplastrella* sp. (08-05-026, wet sponge vol ~2–5 mL) was concentrated and redissolved in MeOH. The MeOH extract was sequentially extracted with solvents of increasing polarity with adjustment of the H₂O content at each stage and removal of solvent from the organic layer: hexane, 0% H₂O (5.2 mg), CHCl₃, 40% H₂O (12.4 mg) The aqueous phase was concentrated under reduced pressure to remove MeOH and then extracted with *n*-BuOH (15.6 mg); the latter was shown to contain diplyne E.¹ Finally, the aqueous phase was concentrated under reduced pressure (106.9 mg). The CHCl₃-soluble phase was separated by flash chromatography (SiO₂, 1–100% MeOH–CHCl₃ step gradient) to give a diyne-containing fraction (2.1 mg), which was further separated by RP HPLC (C₁₈, 5 μm, 10 mm × 250 mm, CH₃CN–H₂O, 2 mL/min) to give diplyne C¹ (**1**, 500 μg) and falkneryne A (**4a**, 400 μg).

The EtOH extract of a second specimen of *Diplastrella* sp. (08-07-038; wet sponge vol ~2–5 mL) was concentrated and redissolved in MeOH. The MeOH extract was sequentially extracted with solvents of increasing polarity with adjustment of the H₂O content at each stage and removal of solvent from the organic layer: hexane, 0% H₂O (9.8 mg), CHCl₃, 40% H₂O (19.5 mg) The aqueous phase was concentrated under reduced pressure to remove MeOH and then partitioned against with *n*-BuOH, which was concentrated to give the *n*-BuOH-soluble fraction (13.3 mg). Finally, the aqueous phase was concentrated under reduced pressure (73.9 mg). The CHCl₃-soluble phase was separated by RP HPLC (C₁₈, 5 μm, 10 mm × 250 mm, CH₃CN–H₂O) to give falknerynes A–C (**4a**, 1 mg), (**4b**, 4.7 μg²⁹), (**4c**, 14.1 μg²⁹), diplyne A¹ (3 mg), and diplyne E (**2**, 100 μg).¹

Falkneryne A (4a). Colorless glass; UV (MeOH) λ_{max} 271 nm. ¹H NMR, ¹³C NMR, see Table 1. HRMS *m/z* 347.0633 [M + Na]⁺ calcd for C₁₆H₂₁⁷⁹BrNaO₂ 347.0623.

Falkneryne B (4b). Colorless glass; UV (MeOH) λ_{max} 273 nm ¹H NMR, see Table S1, Supporting Information. HREIMS: the compound failed to give pseudomolecular ions.

Falkneryne C (4c). Colorless glass; UV (MeOH) λ_{max} 272 nm. ¹H NMR, see Table S1, Supporting Information. HRMS *m/z* 343.0304 [M + Na]⁺ calcd for C₁₆H₁₇⁷⁹BrNaO₂ 343.0309

Preparation of 1,2-*O*-Bis(4'-dimethylaminobenzoyl)diplyne C (5). A solution of **1** (100 μg, 0.305 μmol) in CH₃CN (100 μL) was treated with *N*-(4-dimethylaminobenzoyl)imidazole (197 μg, 0.917 μmol) and DBU (140 μg, 0.917 μmol) and the mixture allowed to stir for 6 h, then concentrated to give a crude product, which was purified by normal phase HPLC (17:83 EtOAc/hexane) to give pure **5**. UV–vis (EtOH), λ_{max} 313 nm. CD (EtOH) λ 301 nm (Δε +22.4), 326 (–21.5); ¹H NMR (600 MHz, CDCl₃) 7.91 (d, 2H, *J* = 8.8 Hz), 7.88 (d, 2H, *J* = 8.8 Hz), 6.62 (d, 2H, *J* = 8.8 Hz), 6.60 (d, 2H, *J* = 8.8 Hz), 6.15 (dt, 1H, *J* = 13.9, 7.3 Hz), 5.99 (d, 1H, *J* = 13.7 Hz), 5.96 (m, 1H), 4.56 (m, 2H), 3.03 (s, 6H), 3.02 (s, 6H), 2.25 (t, 2H, *J* = 7.1 Hz), 2.01 (q, 2H, *J* = 7.4 Hz); HRMS *m/z* 621.2330 [M + H]⁺ calcd for C₃₄H₄₂⁷⁹BrN₂O₄ 621.2328.

Preparation of (R)-1-*O*-Acetyl-(2'-*O*-naphthoyl)diplyne C (6). A solution of **1** (100 μg, 0.305 μmol) in CH₃CN (100 μL) was treated with Novozyme 435 (7 mg) and vinyl acetate (263 μg, 3.05 μmol, 10 equiv), and the mixture stirred at 40–45 °C. After 2 h the reaction was quenched, the enzyme was filtered off, and the filtrate was concentrated under reduced pressure to give the

crude product. The latter material was redissolved in anhydrous CH₃CN (100 μL) and treated with *N*-(2'-naphthoyl)imidazole³⁰ (137 μg, 0.61 μmol) and DBU (93 μg, 0.61 μmol) and allowed to stir at rt for 6 h. The reaction mixture was concentrated under reduced pressure to give the crude product, which was purified by normal phase HPLC (9:91 EtOAc in hexane) to give pure (*R*)-**6**. CD (EtOH) λ 238 nm (Δε +25.3, see Figure 2b. CD (CH₃CN) λ 238 nm (Δε +29.7), 188 nm (–50.5). ¹H NMR (600 MHz) 8.63 (s, 1H), 8.05 (d, 1H, *J* = 8.5 Hz), 7.97 (d, 1H, *J* = 8.1 Hz), 7.90 (d, 1H, *J* = 8.6 Hz), 7.89 (d, 1H, *J* = 8.1 Hz), 7.62 (m, 1H), 7.56 (m, 1H), 6.15 (dt, 1H, *J* = 13.8, 7.3 Hz), 5.99 (d, 1H, *J* = 13.5 Hz), 5.95 (dd, 1H, *J* = 7.2, 4.1 Hz), 4.47 (dd, 1H, *J* = 11.8, 4.0 Hz), 4.44 (dd, 1H, *J* = 11.8, 7.4 Hz), 2.27 (t, 2H, *J* = 7.0 Hz), 2.08 (s, 3H), 2.02 (q, 2H, *J* = 7.2 Hz). HRMS *m/z* 545.1298 [M + Na]⁺ calcd for C₂₉H₃₁⁷⁹BrNaO₄ 545.1303.

(S)-1-Cyclohexylprop-2-ynyl Acetate (12). (*S*)-**9** (52.4 mg, 0.21 mmol) was dissolved in THF (2 mL), and TBAF (0.25 mL, 1 M in THF) was added dropwise at 0 °C. The mixture was stirred at 0 °C for 30 min, and the solvent was removed under reduced pressure to give a crude product that was immediately subject to acetylation (Ac₂O, 30 μL) in anhydrous pyridine (500 μL) at 25 °C for 48 h. The volatiles were removed and the mixture separated (silica cartridge, 3:7 hexane–CH₂Cl₂) to provide (–)-**12** (14.1 mg, 38%). [α]_D²⁴ –66.6 (*c* 0.011, CHCl₃) [lit. [α]_D²⁴ –65.5 (*c* 1.5, CHCl₃)].²² MS and NMR data were identical with literature values.

En-yne (13). A mixture of 1-iododecene (43 mg, 0.12 mmol), (Ph₃P)₂PdCl₂ (5.5 mg, 7.8 μmol), CuI (6.0 mg, 0.032 mmol) and Et₃N (2.5 mL) were stirred at rt for 30 min. This solution was then added dropwise to a solution of alkyne **12** in Et₃N (1 mL) and the mixture reaction stirred at rt for 1.5 h. The volatiles were removed and the residue separated by flash chromatography (silica cartridge, 3:97 EtOAc–hexanes) to yield acetoxy en-yne **13** as an oil (21.5 mg, 79%). [α]_D²⁴ –88.0 (*c* 0.004, CHCl₃); IR (KBr) ν_{max} 2925, 2854, 1743, 1451, 1369, 1229, 1017, 977, 955 cm^{–1}; ¹H NMR (400 MHz, CDCl₃) 0.86 (t, *J* = 7.2 Hz, 3H), 1.04–1.36 (m, 21H), 1.61–1.84 (m, 6H), 2.06 (s, 3H), 2.06 (dq, *J* = 1.6, 7.2 Hz, 5H), 5.30 (dd, *J* = 2.0, 6.4 Hz, 1H), 5.45 (dq, *J* = 2.0, 15.6 Hz, 1H), 6.15 (dt, *J* = 7.2, 15.6 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 14.1 (CH₃), 21.1 (CH₃), 22.7 (CH₂), 25.7 (CH₂), 25.8 (CH₂), 26.2 (CH₂), 28.1 (CH₂), 28.6 (CH₂), 29.1 (CH₂), 29.3 (CH₂), 29.4 (CH₂), 29.5 (CH₂), 29.6 (CH₂), 31.9 (CH₂), 33.1 (CH₂), 42.0 (CH), 68.9 (CH), 74.0 (C), 83.9 (C), 84.7 (C), 108.6 (CH), 146.0 (CH), 170.2 (C). HREIMS *m/z* 346.2869 [M]⁺, calcd for C₂₃H₃₈O₂ 346.2866.

En-yn-ol (14). A solution of **13** in MeOH (100 μL) was treated with anhydrous NH₃ (2 M in MeOH, 300 μL), stirred overnight, and then concentrated under a stream of N₂. The crude mixture was separated by chromatography (silica cartridge, 95:5 hexanes–EtOAc) to yield alcohol **14** as an oil (4.8 mg, 91%). [α]_D²⁴ +5.3 (*c* 0.02, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 0.86 (t, *J* = 6.8 Hz, 3H), 1.02–1.38 (m, 21H), 1.49–1.86 (m, 7H), 2.07 (dq, *J* = 1.2, 7.2 Hz, 5H), 4.22 (m, 1H), 5.47 (dq, *J* = 1.6, 15.6 Hz, 1H), 6.13 (dt, *J* = 7.2, 15.6 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 14.1 (CH₃), 22.7 (CH₃), 25.9 (CH₂), 26.3 (CH₂), 28.2 (CH₂), 28.5 (CH₂), 28.6, 29.1 (CH₂), 29.3 (CH₂), 29.4 (CH₂), 29.5 (CH₂), 29.6 (CH₂), 29.7 (CH₂), 31.9 (CH₂), 33.1 (CH₂), 44.3 (CH), 67.7 (CH), 84.5 (C), 87.5 (C), 108.8 (CH), 145.4 (CH). The configuration of the carbinol center in **14** was confirmed by the modified Mosher's method.¹¹ (*R*)-MTPA ester of compound **14**: compound **14** (0.8 mg, 2.6 μmol) was dissolved DCM–pyridine (1:1 200 μL). (*S*)-MTPA-Cl (5 μL, 0.27 mmol) was added and the mixture stirred for 1.5 h. The volatiles were removed and the crude product separated by chromatography (silica, 95:5 hexanes–EtOAc) to yield the (*R*)-MTPA ester of compound **14** (1.3 mg, 95%). ¹H NMR (400 MHz,

(28) Rützler, K., Family Spirastrellidae Ridley & Dendy, 1886. In *Systema Porifera*; Hooper, J., Van Soest, R. W. M., Eds.; Kluwer/Plenum: New York, 1994; Vol. 1.

(29) Submilligram yields were determined using quantitation by solvent ¹³C satellite (QSCS). Dalisay, D. S.; Molinski, T. F. *J. Nat. Prod.* **2009**, *72*, 739–744.

(30) Ikemoto, N.; Lo, L.-C.; Nakanishi, K. *Angew. Chem., Int. Ed.* **1992**, *31*, 890–891.

CDCl_3) δ 0.86 (t, $J = 6.8$ Hz, 3H), 1.05–1.36 (m, 21H), 1.63–1.74 (m, 5H), 1.81 (brd, $J = 12.4$ Hz, 1H), 2.08 (dq, $J = 1.2, 6.8$ Hz, 2H), 3.54 (s, 3H), 5.42 (d, $J = 5.6$ Hz, 1H), 5.43 (dq, $J = 1.6, 15.6$ Hz, 1H), 6.12 (dt, $J = 6.8, 15.6$ Hz, 1H). **(S)-MTPA ester of compound 14:** The same procedure was applied to compound **14** (1.0 mg, 3.3 μmol) with (*S*)-MTPA-Cl (5 μL , 0.27 mmol) to yield the corresponding (*S*)-MTPA ester (1.4 mg, 99%) yield. $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 0.86 (t, $J = 6.8$ Hz, 3H), 0.99–1.38 (m, 21H), 1.60–1.71 (m, 5H), 1.71 (m, 1H), 2.09 (dq, $J = 1.6, 7.2$ Hz, 2H), 3.57 (d, $J = 1.2$ Hz, 3H), 5.42 (d, $J = 5.6$ Hz, 1H), 5.46 (dq, $J = 1.6, 16.0$ Hz, 1H), 6.15 (dt, $J = 6.8, 16.0$ Hz, 1H).

Naphthoate Ester (S)-7. A solution of enyn-ol **14** (2 mg, 6.57 μmol) in CH_2Cl_2 (200 μL) was treated with 2-naphthoic acid (2.8 mg, 16.4 μmol), EDC (3.0 mg, 19.7 μmol), and a small crystal of DMAP. The resulting mixture was stirred for 48 h, concentrated, and separated by flash chromatography (SiO_2) to afford naphthoate ester (*S*)-**7** (2.4 mg, 82%). $[\alpha]_{\text{D}}^{24} +20.2$ (c 0.94, CHCl_3); $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.61 (bs, 1H), 8.07 (dd, 8.4, 1.2 Hz, 1H), 7.95 (d, 8.0 Hz, 1H), 7.86 (d, 8.4 Hz, 2H), 7.55 (m, 2H), 6.17 (dt, 16.0, 7.2 Hz, 1H), 5.62 (dd, 5.6, 1.2 Hz, 1H), 5.49 (dq, 16.0, 0.8 Hz, 1H), 2.07 (dq, 6.8, 1.6 Hz, 2H), 1.66–1.98 (m, 4H), 1.22 (m, 20H); HREIMS m/z 458.3176 $[\text{M}^+]$ calcd for $\text{C}_{32}\text{H}_{42}\text{O}_2$, 458.3179.

Naphthoate Ester (S)-15. Alkynol (*S*)-**9** (10.4 mg, 0.041 mmol) was converted into naphthoate ester **15** (13.5 mg, 82%) using the procedure described above. $[\alpha]_{\text{D}}^{24} +12.6$ (c 1.93, CHCl_3); $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.61 (bs, 1H), 8.07 (dd, $J = 8.4, 1.2$ Hz, 1H), 7.95 (d, $J = 8.0$ Hz, 1H), 7.86 (d, $J = 8.4$ Hz, 2H), 7.55 (m, 2H), 5.56 (d, $J = 6.0$ Hz, 1H), 1.67–1.99 (m, 6H), 1.13–1.32 (m, 4H), 0.98 (t, $J = 8.0$ Hz, 9H), 0.59 (q, $J = 8.0$ Hz, 6H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 165.7 (C), 135.5 (C), 132.4 (C), 131.2 (CH), 129.4 (CH), 128.3 (CH), 128.1 (CH), 127.7 (CH), 127.4 (C), 126.6 (CH), 125.4 (CH), 103.0 (C), 88.6 (C), 69.3 (CH), 42.1 (CH), 28.7 (CH₂), 28.1 (CH₂), 26.3 (CH₂), 25.8 (CH₂), 25.7 (CH₂), 7.8 (CH₃), 4.2 (CH₂); HREIMS m/z 406.2330 $[\text{M}^+]$ calcd for $\text{C}_{26}\text{H}_{34}\text{O}_2\text{Si}$, 406.2323.

Diyne Naphthoate Ester (S)-8. Naphthoate ester **15** (13.5 mg, 0.033 mmol) was dissolved in THF (1 mL), and the solution was cooled to 0 °C. TBAF (1 M in THF, 33.2 μL) was added dropwise, and the mixture was stirred for 10 min before removal of the volatiles. The residue was purified by flash chromatography (SiO_2 , 9:1 hexanes–ether) to afford essentially pure propargyl *O*-naphthoate **15a** (8.1 mg, 86%). $[\alpha]_{\text{D}}^{23} -12.8$ (c 1.19, CHCl_3); FTIR (KBr) ν_{max} 3298, 2929, 2853, 1719, 1281, 1225, 1195, 1129, 1088, 973, 777, 761 cm^{-1} ; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.61 (bs, 1H), 8.07 (dd, $J = 8.4, 1.2$ Hz, 1H), 7.95 (d, $J = 8.0$ Hz, 1H), 7.86 (d, $J = 8.4$ Hz, 2H), 7.55 (m, 2H), 5.51 (dd, $J = 6.0, 2.4$ Hz, 1H), 2.48 (d, $J = 2.4$ Hz, 1H), 1.68–2.00 (m, 6H), 1.18–1.32 (m, 4H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 165. (C), 135.6 (C), 132.4 (C), 131.3 (CH), 129.4 (CH), 128.4 (CH), 128.2 (CH), 127.8 (CH), 127.1 (C), 126.7 (CH), 125.3 (CH), 80.3 (C), 74.3 (CH), 68.6 (CH), 41.8 (CH), 28.5 (CH₂), 28.2 (CH₂), 26.2 (CH₂), 25.8 (CH₂), 25.7 (CH₂).

A mixture of **15a** (7.8 mg, 27 μmol), CuCl (0.5 mg, 5.3 μmol), and $\text{NH}_2\text{OH}\cdot\text{HCl}$ (2.8 mg, 40 μmol) was suspended in MeOH (150 μL) at 0 °C under N_2 and treated dropwise with neat *n*-butylamine (250 μL). The mixture was stirred for 10 min and treated dropwise with a solution of 1-bromoheptyne (4.6 mg, 26.7 μmol) in MeOH (100 μL). The mixture was stirred at 0 °C for 1 h and then poured into ice–water (2 mL) before acidification with 5% H_2SO_4 and extraction with ether (3 \times 4 mL). The combined ether extracts were washed with brine, dried with anhydrous MgSO_4 , and separated by flash chromatography

(silica cartridge, 8:1, hexanes–diethyl ether) to give diyne (*S*)-**8** as an oil (8.2 mg, 80%). $[\alpha]_{\text{D}}^{23} +58.8$ (c 1.24, CHCl_3); $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.60 (bs, 1H), 8.05 (dd, $J = 8.4, 1.2$ Hz, 1H), 7.95 (d, $J = 8.0$ Hz, 1H), 7.86 (d, $J = 8.4$ Hz, 2H), 7.55 (m, 2H), 5.55 (d, $J = 6$ Hz, 1H), 2.25 (t, $J = 6.8$ Hz, 2H), 1.67–1.99 (m, 6H), 1.50 (p, $J = 7.6$ Hz, 2H), 1.17–1.37 (m, 4H), 0.87 (t, $J = 7.2$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 165.6 (C), 135.6 (C), 132.4 (C), 131.3 (CH), 129.4 (CH), 128.4 (CH), 128.2 (CH), 127.8 (CH), 127.1 (C), 126.7 (CH), 125.3 (CH), 81.8 (C), 72.0 (C), 71.2 (C), 69.2 (CH), 64.5 (C), 42.2 (CH), 31.0 (CH₂), 28.6 (CH₂), 28.3 (CH₂), 27.8 (CH₂), 26.1 (CH₂), 25.8 (CH₂), 25.7 (CH₂), 22.1 (CH₂), 19.2 (CH₂), 13.9 (CH₃). HREIMS m/z 386.2238 $[\text{M}^+]$ calcd for $\text{C}_{37}\text{H}_{30}\text{O}_2$, 386.2240.

Preparation of (R)-1,2-O-Bis(4'-dimethylaminobenzoyl)faulkneryne (16). Using the procedure described above for **1**, faulkneryne A (**3**, 100 μg , 0.307 μmol) was converted into the corresponding DMB diester (*R*)-**16** (21 μg). UV–vis (EtOH), λ_{max} 312 nm. CD (EtOH) λ 302 nm ($\Delta\epsilon +17.4$), 327 (–9.0); $^1\text{H NMR}$ (600 MHz, CDCl_3) 7.92 (d, 2H, $J = 8.8$ Hz), 7.89 (d, 2H, $J = 9.0$ Hz), 6.63 (d, 2H, $J = 9.6$ Hz), 6.60 (d, 2H, $J = 9.4$ Hz), 6.15 (dt, 1H, $J = 13.3, 7.5$ Hz), 6.09 (dt, 1H, $J = 10.8, 7.7$ Hz), 6.02 (m, 1H), 5.98 (m, 2H), 5.48 (m, 1H), 4.60 (m, 1H), 2.35 (m, 2H), 2.03 (m, 2H). HRMS m/z 619.2141 $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{34}\text{H}_{39}^{79}\text{BrN}_2\text{O}_4$, 619.2172.

Molecular Modeling. An analogue of compound **5** (truncated to the 3,5-diyne) was minimized using molecular mechanics (MMFF94, Spartan 08, gas phase). Energies were calculated (semiempirical, PM3) and Monte Carlo conformational searching applied to obtain the lowest 25 models ranging in energy from –1.06 to 25 kcal mol^{-1} . For the three most stable conformers, see Figure 1d–f. An analogue of dibenzoate ester **6** (truncated to the 3,5-diyne) was minimized using semiempirical methods (PM3, Spartan 08) and the 25 lowest energy conformers determined by Monte Carlo methods. See Figure 2.

Cytotoxicity Assay. Cytotoxicity of **1** against cultured human colon tumor cells (HCT-116) incubated under 5% CO_2 in the presence and absence of compound, followed by colorimetric measurement of growth inhibition by the MTS method using a microplate reader as described elsewhere.² Dipylyne C (**1**) exhibited an LC_{50} of 3.6 $\mu\text{g}/\text{mL}$ (etoposide was used as a positive control).

Acknowledgment. We thank C. K. Skepper for assistance sponge collection and preparation of *S*-(+)-**9**. We're grateful to S. Zea and M-K. Harper (University of Utah) for helpful discussions on sponge taxonomy, J. Siegel (University of Zürich) for stereochemical advice, J. R. Pawlik (University of North Carolina, Wilmington), and the captain and crew of the *R/V* Seward Johnson for logistical support during collecting expeditions. The NSF Biological Oceanography Program (OCE-0095724, 0550468 to J.R.P.) is acknowledged for ship time. The 500 MHz NMR spectrometers were purchased with a grant from the NSF (CRIF, CHE0741968). This work was supported by grants from NIH (CA122256 and AI039987 to T.F.M.) and a Ruth L. Kirschstein National Research Service Award NIH/NCI (T32 CA009523 to B.I.M.).

Supporting Information Available: ^1H , ^{13}C NMR and 2D NMR spectra of **3**, and ^1H , ^{13}C NMR of **4** and all synthetic compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.