

# Correction to Synthesis of 3-Substituted Isocoumarins via Cascade Intramolecular Ullmann-Type Coupling–Rearrangement Process

Zhi-Yuan Ge, Xiang-Dong Fei, Ting Tang, Yong-Ming Zhu,\* and Jing-Kang Shen\*

*J. Org. Chem.* **2012**, *77*, 5736–5743. DOI: 10.1021/jo300916s.

Page 5738. Entries 12–19 of Table 2 were inadvertently omitted.

The corrected version of Table 2 follows:

**Table 2. Synthesis of 3-Substituted Isocoumarins<sup>a</sup>**

| Entry | Substrate | Product | Time | Yield [%] <sup>b</sup> | Entry | Substrate | Product | Time | Yield [%] <sup>b</sup> |
|-------|-----------|---------|------|------------------------|-------|-----------|---------|------|------------------------|
| 1     |           |         | 6h   | 81                     | 11    |           |         | 6h   | 81                     |
| 2     |           |         | 6h   | 87                     | 12    |           |         | 8h   | 69                     |
| 3     |           |         | 6h   | 84                     | 13    |           |         | 4h   | 59                     |
| 4     |           |         | 6h   | 79                     | 14    |           |         | 4h   | 85                     |
| 5     |           |         | 12h  | 62                     | 15    |           |         | 4h   | 53                     |
| 6     |           |         | 6h   | 86                     | 16    |           |         | 4h   | 93                     |
| 7     |           |         | 6h   | 80                     | 17    |           |         | 7h   | 84                     |
| 8     |           |         | 8h   | 74                     | 18    |           |         | 7h   | 79                     |
| 9     |           |         | 10h  | 61                     | 19    |           |         | 6h   | 68                     |
| 10    |           |         | 6h   | 81                     |       |           |         |      |                        |

<sup>a</sup>The reactions were performed in a sealed tube with **1** (0.5 mmol), CuI (0.05 mmol), 2-picolinic acid (0.1 mmol), and K<sub>2</sub>CO<sub>3</sub> (1.0 mmol) in toluene (2 mL) at 110 °C under nitrogen. <sup>b</sup>Isolated yield.