


# Urushiol Detection using a Profluorescent Nitroxide

Rebecca Braslau,\* Frank Rivera, III, Erin Lilie, and MariEllen Cottman

Department of Chemistry and Biochemistry, University of California, Santa Cruz, California, 95064

**S** Supporting Information


**ABSTRACT:** A method to visually detect minute amounts of urushiol, the toxic catechol from poison oak, poison ivy, and poison sumac, has been developed utilizing the reaction of a profluorescent nitroxide with the *B*-*n*-butylcatecholboronate ester formed in situ from urushiol and *B*-*n*-butylboronic acid. The resulting *N*-alkoxyamine is strongly fluorescent upon illumination with a fluorescent lamp, allowing the location of the toxic urushiol contamination to be visualized. This methodology constitutes the groundwork for the future development of a spray to detect urushiol to avoid contact dermatitis, as well as to detect catecholamines for biomedical applications.



## INTRODUCTION

Urushiol is the causative agent of allergic contact dermatitis from exposure to *Toxicodendron spp.* (poison oak, poison ivy, and poison sumac). For susceptible individuals, exposure to tiny amounts of this invisible oil elicits contact dermatitis, which is contracted by either direct exposure to the plant source, or indirectly by exposure to contaminated tools, clothing, and pets. The deleterious effects of *Toxicodendron* dermatitis are a large personal and economic burden to the United States, causing significant workers' compensation claims and lost time for those engaged in a myriad of outdoor jobs,<sup>1,2</sup> accounting for greater than 10% of the total U.S. Forest Services lost-time injuries.<sup>3</sup> In 1988, the National Institute for Occupational Safety and Health (NIOSH) estimated that 1.07–1.65 million occupational skin injuries occurred yearly, with an estimated annual rate of 1.4–2.2 cases per 100 workers.<sup>4</sup> Approximately 50–70% of the adult American population is clinically allergic to urushiol.<sup>1</sup> Outside of North America, significant occupational hazards arise from exposure to the sap (kiurushi) of Asian lacquer trees (ex. *Toxicodendron verniciflua* in Japan,<sup>5</sup> *Melanorrhœa usitata* in Thailand<sup>6</sup>) used as a varnish in lacquerware. Symptoms of allergic contact dermatitis from urushiol exposure vary from mild annoyance to weeks of irritation and pain. Occasionally, exposure can lead to nephropathy<sup>7</sup> and even to fatal systemic anaphylaxis.<sup>8</sup>

Chemically, urushiol (Figure 1) is the name given to a collection of related compounds consisting of 3-substituted catechols, in which the long hydrophobic chain is a linear C<sub>15</sub> or C<sub>17</sub> alkyl group containing 0–4 degrees of predominately *cis* unsaturation.<sup>9,10</sup> Each of the different members of the *Toxicodendron species* contain mixtures of urushiol bearing C<sub>15</sub> or C<sub>17</sub> alkyl chains, with various degrees of unsaturation. They all share the catechol functionality, and a long, greasy alkyl chain, which facilitates migration into the skin. The majority of allergic contact dermatitis cases from urushiol result from unexpected exposure. Since as little as 2 mg or less of urushiol can lead to contact dermatitis,<sup>1</sup> a very sensitive method for



**Figure 1.** Structure of the most abundant components of urushiol from *Toxicodendron species*.

detecting urushiol is warranted. The ability to visualize the distribution of this oil would allow sensitive individuals to avoid or wash away urushiol<sup>11</sup> to prevent or minimize contact dermatitis. Early work includes the development of a “black spot test” to help identify these plants in the field,<sup>12</sup> but this is not practical for detecting the distribution of contamination. Thus, a fluorescent detection method is attractive, given the strong signal generated from very small amounts of fluorophore. A dilute solution of nontoxic spray, which reacts selectively to form a highly fluorescent species in the presence of urushiol, would enable the location of urushiol contamination to be determined quickly, without the need for specialized instrumentation beyond a hand-held fluorescent lamp. Reactions based on oxidation of the catechol to the orthoquinone, followed by condensation with ethylenediamine<sup>13</sup> or benzylamine<sup>14</sup> to give fluorescent heteroaromatic derivatives, have been used traditionally to quantitate catechols and catecholamines. However, both ethylenediamine and benzylamine are considered very dangerous upon exposure to the skin.


Received: June 7, 2012

Published: August 17, 2012

## RESULTS AND DISCUSSION


To develop a system to selectively detect catechols in the presence of other alcohols and diols (such as sugars), a reaction that takes place with catechols but not with other alcohols is required. *B*-Alkylcatecholboronate esters have been used to selectively generate alkyl radicals upon reaction with oxygen radicals.<sup>15,16</sup> Early work by Brown and Negishi<sup>17,18</sup> entailed enol oxygen radical 2 addition to *B*-alkylcatecholboronate ester 1 to give an alkyl radical, which then adds to an unsaturated ketone in a chain reaction (Scheme 1). The efficacy of oxygen

**Scheme 1. Brown and Negishi's 1969 Sequence for Generating Carbon Radicals from *B*-Alkylcatecholboronate Ester 1**



radical addition specifically to *B*-alkylcatecholboronate esters 1 is due to stabilization of the intermediate perboryl radical 3, by delocalization of the unpaired electron into the aromatic ring (Scheme 2). Direct ESR evidence for a delocalized perboryl


**Scheme 2. Roberts Demonstrated that the Addition of an Oxygen Radical to an *B*-Alkylcatecholboronate Ester 1 Formed a Perboryl Radical 3, Visible by EPR**




radical 3 was observed below 270 K by Roberts.<sup>19</sup> A number of very useful synthetic methodologies have been developed from this chemistry.<sup>20</sup>

Key to this urushiol detection is the seminal work by Renaud,<sup>21–23</sup> in which addition of two equivalents of the oxygen radical TEMPO 4, a commercially available persistent nitroxide, results in formation of the carbon radical trapping product, *N*-alkoxyamine 5 (Scheme 3). We have found that trialkylboranes do not form a resonance-stabilized perboryl radical, and thus do not react with TEMPO. For example, trialkylborane 6 gave no reaction upon exposure to TEMPO, but did react with the “hotter” *t*-butoxy radical generated from stoichiometric di-*t*-butyl hyponitrite (Scheme 4).<sup>24</sup> Thus nitroxides react selectively with *B*-alkylcatecholboronate esters 1, to generate *N*-alkoxyamines 5.

**Scheme 3. Renaud's Method of Radical Generation by Addition of Nitroxide to a *B*-Alkylcatecholboronate Ester 1**



**Scheme 4. TEMPO Did Not Add to Trialkylborane 6, however *t*BuO• Generated from di-*t*-Butyl Hyponitrite Did Add to the Trialkylborane 6**



Profluorescent nitroxides<sup>25–28</sup> 7 (sometimes referred to as “pre-fluorescent nitroxides”<sup>29</sup>) are nitroxides bearing a short covalent tether to a fluorophore. The free nitroxide quenches the fluorescence (Figure 2). Upon reaction of the nitroxide




**Figure 2. Profluorescent nitroxides 7: the free nitroxide quenches fluorescence of a closely tethered fluorophore; fluorescence is restored upon reaction to form the *N*-alkoxyamine 8.**


with a carbon radical or reducing agent, to form an *N*-alkoxyamine 8 or an *N*-hydroxylamine, the fluorescence is restored. Reaction of a catechol with a *B*-alkyl boronic acid derivative to form a *B*-alkyl boronate ester 1 in situ, in the presence of a profluorescent nitroxide 7, is expected to form a highly fluorescent *N*-alkoxyamine 8 (Scheme 5). Generation of a fluorescent signal would be a positive indicator for the presence of the catechol functionality.

To explore the viability of this approach, a preformed *B*-alkylcatecholboronate ester 1 was prepared from *B*-*n*-butylboronic acid using Dean–Stark conditions in toluene. Although this was successful, a much easier procedure developed by Brown<sup>30</sup> provides the same *B*-*n*-butylcatecholboronate ester in minutes at room temperature: the reaction is run in pentane. Water generated under the condensation conditions phase separates, conveniently forming the catechol boronate 1 in good yield. Addition of two equivalents of orange-colored TEMPO 4 to an NMR tube containing one equivalent of the preformed *B*-*n*-butylcatecholboronate ester 1 resulted in the dissipation of the orange color, and formation of the *N*-*n*-butoxyamine of TEMPO 5 (Scheme 6), identified by the distinct *O*-methylene triplet at  $\delta$  3.73 in CDCl<sub>3</sub>. Authentic *N*-*n*-butoxyamine 5 was prepared from TEMPO 4 by the method

**Scheme 5. Fluorescence Generation by Condensation of a Catechol with a *B*-Alkyl Boronic Acid, Followed by Reaction with a Profluorescent Nitroxide 7, to Form a Stable Fluorescent *N*-Alkoxyamine 8**




**Scheme 6. Generation of the *N*-*n*-Butoxyamine 5 from TEMPO 4 using Preformed *B*-*n*-Butylcatecholboronate Ester 1**



of Whitesides<sup>31</sup> employing *n*-butyllithium, allowing confidence in the spectral assignment.


Mixing two equivalents of profluorescent nitroxide dansyl-TEMPO<sup>32</sup> 7 with one equivalent of the preformed *B*-*n*-butylcatecholboronate ester 1 formed the highly fluorescent *N*-*n*-butoxyamine 8; fluorescence was detected almost immediately. Even more gratifying was the generation of fluorescence from free catechol, via the reversible formation of the *B*-*n*-butylcatecholboronate ester 1 formed in situ. In this key experiment, two equivalents of dansyl-TEMPO 7 were mixed with one equivalent of *B*-*n*-butylboronic acid in acetone; addition of one equivalent of catechol under nitrogen immediately resulted in strong fluorescence upon illumination with a UV lamp (Scheme 7). The fluorescent *N*-alkoxyamine product 8 was isolated in 64% yield. It should be noted that

**Scheme 7. Fluorescent *N*-Alkoxyamine 8 Formed by Reaction with Catechol using Profluorescent Nitroxide Dansyl-TEMPO 7**



alkyl radicals react with molecular oxygen at a rate of  $5 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$  in hydrocarbon solvent,<sup>33</sup> whereas the rate of trapping of primary alkyl radicals by TEMPO is  $1.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ,<sup>34</sup> approximately four times slower. Thus for the detection of urushiol in air, a significant portion of *n*-butyl radicals will trap  $\text{O}_2$ , forming peroxides, which will most likely undergo autoxidation. Thus the nitroxide trapping method will underestimate the amount of urushiol present, but enough fluorescent alkoxyamine will be formed to indicate the spatial distribution of urushiol contamination.

Spot tests on a paper towel give a visual display of the efficacy of this procedure in detecting the presence of catechol. In Figure 3a, the profluorescent dansyl-TEMPO 7 was spotted




**Figure 3. (a)** Spot tests (circled in pencil) on a paper towel of acetone solutions of dansyl-TEMPO 7, the corresponding *N*-alkoxyamine 8, and a control mixture of dansyl-TEMPO 7 and boronic acid; **(b)** crystalline catechol on a paper towel upon addition of a dilute solution of  $n\text{BuB}(\text{OH})_2 + \text{FI-NitO}\bullet$  7 in acetone. Both are visualized by illumination under a TLC lamp at 365 nm.

against the fluorescent *N*-alkoxyamine 8. As a control, a mixture of *B*-*n*-butylboronic acid and dansyl-TEMPO 7 was also spotted; illumination with a hand-held TLC fluorescent lamp (365 nm) only showed the *N*-alkoxyamine 8 to be fluorescent as observed by the human eye. A small sample of crystalline catechol likewise generated a spectacular fluorescent display upon addition of a drop of dilute *B*-*n*-butylboronic acid and dansyl-TEMPO 7 in acetone (Figure 3b).


Quantification of the fluorescent response was determined by fluorimetry. The fluorescence excitation and emission spectra of the profluorescent nitroxide dansyl-TEMPO 7, and the corresponding *N*-alkoxyamine 8 are shown in Figure 4. The profluorescent nitroxide 7 does display a residual amount of fluorescent activity, but it is significantly suppressed compared to the *N*-alkoxyamine 8.

To detect urushiol contamination, the generation of a fluorescent species should be rapid. Time studies (Figure 5) were conducted utilizing a 2 mL solution of dansyl-TEMPO 7 ( $1.5 \times 10^{-3} \text{ M}$ ) and *B*-*n*-butylboronic acid ( $7.5 \times 10^{-4} \text{ M}$ ) in acetone (fluorescence of this solution is shown at time zero). To this was added two mL of catechol ( $1.5 \times 10^{-3} \text{ M}$ ) in acetone to give a  $3.75 \times 10^{-4} \text{ M}$  solution containing a 2:1:2 molar ratio of dansyl-TEMPO/ $n\text{Bu}(\text{OH})_2$ /catechol. The ratio of fluorescence intensity,  $F/F_0$  ( $F$  = fluorescence and  $F_0$  = initial fluorescence), was plotted as a function of time. The fluorescent signal increased over the first hour; by the second hour the fluorescence had reached a stable maximum that remained unchanged after 24 h.

**Optimization of the System.** Acetone was the initial solvent examined, as precedent for safe human exposure is provided by the common use of acetone as fingernail polish remover. A few additional solvents were examined. The time study depicted in Figure 5 was repeated with THF, acetonitrile, and isopropanol as solvent; the fluorescent response as a function of time was compared to that using acetone (Figure




**Figure 4.** Fluorescence trace of dansyl-TEMPO **7** (excitation in red, emission in purple) and *N*-alkoxyamine **8** (excitation in blue, emission in green),  $\lambda_{\text{ex}} = 350 \text{ nm}$ ,  $\lambda_{\text{em}} = 512 \text{ nm}$ .



**Figure 5.** Time study of fluorescence upon addition of  $3.0 \mu\text{mol}$  of catechol to a dilute acetone solution of  $1.5 \mu\text{mol}$  of *B*-*n*-butylboronic acid and  $3.0 \mu\text{mol}$  of dansyl-TEMPO **7** ( $\lambda_{\text{ex}} = 350 \text{ nm}$ ,  $\lambda_{\text{em}} = 512 \text{ nm}$ ).

6). Acetone gave a rapid response, followed by THF and then acetonitrile. Isopropanol gave the slowest increase in the




**Figure 6.** Effect of solvent on fluorescence intensity and reaction time in the reaction of catechol with profluorescent dansyl-TEMPO **7** and *B*-*n*-butyl boronic acid ( $\lambda_{\text{ex}} = 350 \text{ nm}$ ,  $\lambda_{\text{em}} = 512 \text{ nm}$ ).

fluorescent signal, presumably as the nucleophilic solvent may compete with the catechol for reaction with the *B*-*n*-alkyl boronic acid. The *B*-*n*-butyl boronic acid is insoluble in dichloromethane and chloroform. The fluorescence reached the greatest intensity most quickly with acetone compared to the other solvents. Thus, acetone continued to be the solvent of choice.

Alkylboronic acids are renowned for being marginally stable, dehydrating reversibly to the boroxine anhydrides,<sup>35</sup> which are

particularly sensitive to degradation by oxygen.<sup>36</sup> Arylboronic acids, which also reactive in this manner, are more robust:<sup>37</sup> thus the possibility of replacing the *n*-butyl with a phenyl group was examined. In Figure 7, it is clear that the *B*-*n*-butylboronic



**Figure 7.** Comparison between the use of *B*-*n*-butylboronic acid and *B*-phenylboronic acid on fluorescence intensity and reaction time in the reaction of catechol with profluorescent nitroxide **7** and *B*-(alkyl or aryl)boronic acid ( $\lambda_{\text{ex}} = 350 \text{ nm}$ ,  $\lambda_{\text{em}} = 512 \text{ nm}$ ).


acid is far more effective in generating a fluorescent signal than *B*-phenylboronic acid. This is not surprising, as the reaction mechanism for the formation of fluorescent *N*-alkoxyamine **8** entails generation of a carbon radical, following nitroxide addition to the catecholboronate ester **1**. The dissociation energy for a primary alkyl radical (for example,  $\text{Et-H}$  to  $\text{Et}\bullet$ ) has been measured at  $101.1 \pm 0.4 \text{ kcal/mol}$ , whereas for a phenyl radical ( $\text{Ph-H}$  to  $\text{Ph}\bullet$ ) at  $111.2 \pm 0.8 \text{ kcal/mol}$ ,<sup>38</sup> making the phenyl radical approximately  $10 \text{ kcal/mol}$  less stable than the butyl radical. Although anhydrous *B*-*n*-butylboronic acid tends to decompose upon storage, we found that a dilute acetone solution of *B*-*n*-butylboronic acid in the presence of water<sup>39</sup> was stable for at least several months. Thus the formulation of *B*-*n*-butylboronic acid as a component of a shelf-stable spray to detect urushiol should be viable. Although skin toxicity for  $10^{-6} \text{ M}$  *B*-*n*-butylboronic acid has not been determined, the use of the alkylboronic acid drug Velcade (Bortezomib)<sup>40</sup> in chemotherapy for multiple myeloma, the use of the arylboronic acid 4-dihydroxy-borylphenylalanine (BPA) in boron neutron capture therapy,<sup>41</sup> and the topical use of the closely related benzoxaboroles as a broad spectrum anti-



fungal,<sup>42,43</sup> make it likely that dilute solutions of an alkylboronic acid can be tolerated on the skin.

**Possible Competing Reactions.** Scaiano has shown that both catechols and phenols react slowly by transferring a phenoxy hydrogen to nitroxides,<sup>44,45,44</sup> and has demonstrated the use of profluorescent nitroxides to probe this antioxidant activity. In the urushiol detection system, reduction of the profluorescent nitroxide by phenols would form fluorescent *N*-hydroxyamines (second line, Scheme 8). In Figure 8,


### Scheme 8. Potential Competing Reactions



<sup>a</sup>reference 34; <sup>b</sup> references 44, 45; <sup>c</sup> reference 49; <sup>d</sup> reference 33; <sup>e</sup> reference 50.

replacement of catechol by phenol resulted in only a negligible increase of fluorescence after 2 h, indicating that phenols will not generate a significant fluorescent signal.


TEMPO 4 traps primary alkyl radicals at rate constants of approximately  $10^9 \text{ M}^{-1} \text{ s}^{-1}$  at room temperature<sup>34</sup> (this rate is slightly solvent dependent<sup>46</sup>). The rates of H-abstraction by TEMPO-type nitroxides from phenols such as Trolox and BHT in protic solvents have been measured to be  $10^{-1}$  to  $10^{-3} \text{ M}^{-1} \text{ s}^{-1}$  at room temperature.<sup>44,47</sup> Catechol has been shown to act as an effective hydrogen donor to carbon radicals.<sup>48,49</sup> A



**Figure 8.** Time study of fluorescence upon addition of  $3.0 \mu\text{mol}$  of phenol (red line) or  $3.0 \mu\text{mol}$  of catechol (blue line) to a solution of  $3.0 \mu\text{mol}$  dansyl-TEMPO and  $1.5 \mu\text{mol}$  of *B*-*n*-butylboronic acid in acetone.

possible concern in the presence of high concentrations of urushiol might be the reduction of the *n*-butyl radical by abstraction of a hydrogen from the catechol moiety of urushiol (third line of Scheme 8). The resulting oxygen-centered catechol radical would then be expected to add to a *B*-*n*-butylcatecholboronate ester **1**, generating another *n*-butyl radical and Meulenhoff's free acid **9**.<sup>48</sup> The rate of hydrogen atom abstraction from catechols by alkyl radicals has been recently determined by Renaud to be  $10^6 \text{ M}^{-1} \text{ s}^{-1}$  at  $80^\circ\text{C}$ . Thus in the presence of large amounts of catechol (for example,  $1 \text{ M}$ ), and very low concentrations of nitroxide (for example,  $3.0 \times 10^{-3} \text{ M}$ ), these two reactions could become competitive, and would lead to an underestimation of the urushiol concentration. For the detection of trace amounts of urushiol, hydrogen abstraction from the catechol by alkyl radical is expected to be a very minor reaction. The trapping of molecular oxygen by the *n*-butyl radical is much more likely. The resulting peroxy radical could participate in autoxidation, or abstract a  $\beta$  hydrogen from alkoxyamine **8** to reform the nitroxide, a pathway recently unveiled by ab initio calculations.<sup>50</sup>


As proof of concept for the detection of urushiol, a successful field test is demonstrated in Figure 9 with *Toxicodendron*



**Figure 9.** (a) Fresh triad of poison oak leaves; (b) print of the same leaves on a paper towel after treatment with a dilute acetone solution of dansyl-TEMPO **7** ( $3 \mu\text{molar}$ ) and *B*-*n*-butylboronic acid ( $1.5 \mu\text{molar}$ ), followed by illumination by a  $365 \text{ nm}$  TLC lamp.

*diversilobum* (western poison oak). A dilute acetone solution of dansyl-TEMPO **7** and *B*-*n*-butylboronic acid (2:1 molar ratio) was dripped onto a freshly harvested triad of poison oak leaves (Figure 9a). A new paper towel was laid over the wet leaves, and pressure was applied to make a "print." Immediate illumination with a long-wavelength UV TLC lamp resulted in the fluorescent image shown in Figure 9b. The particularly strong fluorescent spots are likely due to ruptures in the leaves while applying pressure, exposing the paper towel to localized areas containing high concentrations of urushiol.

Six months after creating the fluorescent leaf image, a small piece of the still-fluorescent print on the paper towel was analyzed by TOF-DART (Direct Analysis in Real Time) Mass Spectrometry:<sup>51</sup> the M+1 peak of the *N*-alkoxyamine product **8** was observed for both an authentic and paper towel-derived sample at 462.2808, confirming the formation and stability of the *N*-alkoxyamine **8** (Figure 10).




authentic alkoxyamine **8**

**M+1 = 462.2808**



alkoxyamine **8** from poison  
oak print from paper towel

**M+1 = 462.2808**



**Figure 10.** TOF-DART MS confirmation of the formation of *N*-alkoxyamine **8** from an authentic sample and from the paper towel 6 months after the leaf print was prepared. (Note: the 371 peak is plasticizer on the glass rod used for sampling.)

Catecholamines, including dopamine, epinephrine and norepinephrine, are biologically important neurotransmitters. They are involved in the regulation of both the central and peripheral nervous systems, and are implicated in a number of diseases, including Alzheimer's<sup>52</sup> and Parkinson's.<sup>53</sup> The ability to detect these species at very low concentration using this methodology based on *B*-alkylboronic acids and profluorescent nitroxides should lead to a number of biomedical applications. In preliminary experiments to detect L-DOPA in water, in analogy to urushiol in organic solvents, a fluorescent signal was observed. However, dansyl is a solvatochromic fluorophore;<sup>54,55</sup> dilute solutions of dansyl-TEMPO **7** with *B*-*n*-butylboronic acid in water gives a faint yet visible fluorescent signal *in the absence of catechol* (see Supporting Information). Use of the two-photon method<sup>56</sup> in a laboratory setting is an alternative strategy to avoid the fluorescence enhancement of dansyl fluorophore in water, but would not be applicable to a fast "in the field" test for urushiol. Thus, a less solvatochromic fluorophore is needed. This sensitivity to water also impacts the urushiol detection system. Efforts are now being directed at examining alternative nonsolvatochromic fluorophores with nonhydrolyzable linkages to the nitroxide as the profluorescent nitroxide partner, in order to avoid generation of a weak signal in the presence of water.

## CONCLUSION

Fluorescent detection to illuminate the location of urushiol contamination from contact with *Toxicodendron* species has been developed, utilizing the reversible reaction of catechols with a *B*-alkylboronic acids and a profluorescent nitroxide. The generation of the fluorescent signal is rapid: detection with a hand-held fluorescent lamp has been demonstrated, showing promise to identify the location of urushiol contamination in the home or field, without costly equipment, thus minimizing the severity of contact dermatitis for sensitive individuals. Investigation of the use of alternative fluorophores to dansyl is underway. The extension of this methodology to detect catecholamines for biomedical applications is under active development in our laboratory.

## EXPERIMENTAL SECTION

**4-Azido-2,2,6,6-tetramethyl-1-piperidine-1-oxyl Free Radical (4-Azido-TEMPO).** Following a modified procedure of Misharin et al.,<sup>57</sup> a solution of 4-methylsulfonyloxy-TEMPO (4.045 g, 16.16 mmol) and sodium azide (2.100 g, 32.32 mmol) in DMSO (54 mL, 0.3 M) was heated to 110 °C under N<sub>2</sub> overnight. Upon cooling to rt, ether (450 mL) was added and the solution was washed with a saturated solution of sodium bicarbonate (200 mL) followed by water (4 × 100 mL). The organic layer was dried over sodium sulfate, filtered and then concentrated. The resulting solid was recrystallized from hexanes to yield orange crystals (2.283 g, 72%).

TLC R<sub>f</sub> = 0.83 (1:1 hexanes/ethyl acetate, UV, PAA). IR (CDCl<sub>3</sub>) 2099 cm<sup>-1</sup> (-N<sub>3</sub> stretching). <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub> + phenylhydrazine): δ 3.60 (tt, *J* = 12.4, 4.2 Hz, 1H), 1.87 (dd, *J* = 12.4, 4.2 Hz, 2H) 1.57 (dd, *J* = 12.4, 12.4 Hz, 2H), 1.23 (s, 6H), 1.16 (s, 6H) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub> + phenylhydrazine): δ 59.6, 52.8, 44.1, 32.2, 20.2 ppm.

**4-Amino-2,2,6,6-tetramethyl-1-piperidine-1-oxyl Free Radical (4-Amino-TEMPO).** Following a modified procedure of Misharin et al.,<sup>57</sup> to a flame-dried round-bottom flask was added 4-azido-TEMPO (1.815 g, 9.201 mmol) and anh. THF (20 mL). Triphenylphosphine (4.827 g, 18.40 mmol) was added in portions over a period of three hours under N<sub>2</sub>. Effervescence was observed. Aqueous NH<sub>4</sub>OH (20 mL, 14.8 M) was added and the reaction mixture was stirred overnight. Most of the solvent was removed in vacuo using gentle heating. The concentrated reaction mixture was taken up in CHCl<sub>3</sub> (100 mL) before adding a solution of 10% acetic acid (50 mL). The phases were separated and 50 mL of 10% acetic acid solution was added. The combined aqueous layers were neutralized with solid sodium bicarbonate, followed by a small amount of 2N NaOH at the end of the neutralization to make the solution slightly basic as determined by pH paper. This solution was then extracted with CHCl<sub>3</sub> (3 × 100 mL). The organic layer was dried over sodium sulfate, filtered, and then concentrated to yield a viscous orange oil (1.071 g, 70%).

TLC R<sub>f</sub> = 0.58 (1:1 ethyl acetate/methanol, UV, PAA). <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub> + phenylhydrazine): δ 3.05 (tt, *J* = 15.2, 4.8 Hz, 1H), 1.74 (dd, *J* = 15.2, 4.8 Hz, 2H) 1.28 (dd, *J* = 15.2, 15.2 Hz, 2H), 1.16 (s, 6H), 1.14 (s, 6H) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub> + phenylhydrazine): δ 59.2, 49.7, 42.3, 32.6, 20.2 ppm.

***N*-Dansyl-TEMPO (7).** Following a modified procedure,<sup>58</sup> dry CH<sub>2</sub>Cl<sub>2</sub> (67.0 mL), 4-amino-TEMPO (1.439 g, 8.402 mmol), dansyl chloride (2.266 g, 28.402 mmol) and triethylamine (2.3 mL, 16.8 mmol, freshly distilled from calcium hydride) were added to a flame-dried round-bottom flask and placed under N<sub>2</sub>. The reaction mixture was allowed to stir for 16.5 h. The solvent was removed in vacuo and the crude product was purified by flash column chromatography (2:1 hexanes/ethyl acetate) to yield a fluffy orange solid (3.04 g, 89%).

TLC R<sub>f</sub> = 0.48 (1:1 hexanes:ethyl acetate, UV, I<sub>2</sub>, PAA). <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub> + phenylhydrazine): δ 8.54 (d, *J* = 10.2 Hz, 1H), 8.28–8.26 (m, 2H), 7.54–7.50 (m, 2H), 7.17 (d, *J* = 9 Hz, 1H), 3.51–3.45 (m, 1H), 2.88 (s, 6H), 1.52–1.40 (m, 4H), 1.10 (s, 6H), 1.00 (s, 6H) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub> + phenylhydrazine): δ 151.3,

135.9, 130.5, 129.9, 129.7, 129.4, 128.8, 123.2, 118.8, 115.2, 59.2, 46.2, 45.8, 45.4, 32.1, 19.5 ppm.

***B*-*n*-Butylcatecholboronate Ester (1).** Method A (Dean–Stark): Following a modified procedure from Green and Wagner,<sup>59</sup> to a flame-dried 25 mL round-bottom flask was added toluene (8 mL), catechol (324 mg, 2.94 mmol) and *B*-*n*-butylboronic acid (300 mg, 2.94 mmol). A Dean–Stark apparatus was attached and the solution was refluxed for 4 h. A separate water layer formed in the Dean–Stark trap. The solution was concentrated in vacuo to yield 516 mg of the title compound as a yellow oil (99.6% crude yield; <sup>1</sup>H NMR shows contamination with traces of toluene).

Method B (in pentane): Following a modified procedure of Brown et al.,<sup>60</sup> to a flame-dried 10 mL round-bottom flask was added catechol (216 mg, 1.96 mmol), *B*-*n*-butylboronic acid (200 mg, 1.96 mmol), and small scoop of sodium sulfate. The contents of the flask were dissolved in 1.6 mL of pentane and stirred under nitrogen at room temperature for 50 min. The solid was removed by filtration and the solvent removed in vacuo to give 287 mg (83.2%) of the desired product as a yellow oil.

Note: it has been previously reported<sup>61</sup> that the carbon attached to the boron is not visible by <sup>13</sup>C NMR due to the quadrupole relaxation of <sup>11</sup>B. In addition, <sup>1</sup>H–<sup>11</sup>B (<sup>11</sup>B, *I* = 3/2) coupling in the <sup>1</sup>H NMR spectrum of the butyl fragment is evident in the multiplet seen at 1.64–1.59 ppm.

<sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>): δ 7.21 (dd, *J* = 7.2, 4.2 Hz, 2H), 7.07 (dd, *J* = 7.2, 4.2 Hz, 2H), 1.64–1.59 (m, 2H), 1.45–1.39 (m, 2H), 1.32–1.29 (m, 2H), 0.96–0.93 (t, *J* = 9.9 Hz, 3H) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>): δ 148.3, 122.5, 112.2, 25.8, 25.2, 13.7 ppm.

**1-*n*-Butoxy-2,2,6,6-tetramethylpiperidine (5).** A. Preparation of authentic sample: Following the procedure of Whitesides,<sup>62</sup> a flame-dried 25 mL round-bottom flask was charged with TEMPO 4.28 (500 mg, 3.20 mmol) and dry toluene (17.8 mL). The solution was stirred and cooled to –78 °C under nitrogen before adding *n*-butyllithium (0.90 mL, 2.24 mmol, 2.5 M in hexanes) dropwise. The reaction mixture was allowed to warm to room temperature, and then washed with 18 mL of water. The solution was dried over magnesium sulfate, filtered and concentrated. The product was purified by flash column chromatography (4:1, hexanes/ethyl acetate) to give the desired product as a yellow oil (358 mg, >100%, slightly contaminated with the *N*-hydroxylamine).

TLC *R*<sub>f</sub> = 0.69 (4:1, hexanes/ethyl acetate, UV, I<sub>2</sub>, PAA). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 3.73 (t, *J* = 6 Hz, 2H), 1.51–1.31 (m, 10H), 1.15 (s, 6H), 1.09 (s, 6H), 0.92 (t, *J* = 6 Hz, 3H) ppm. <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>): δ 70.433.5, 32.0, 27.0, 24.8, 14.0, 13.6, 11.1, 8.2 ppm.

B. Formation by reaction of TEMPO 4 with *B*-*n*-butylboronic ester 1 in an NMR tube: an NMR tube was charged with *B*-*n*-butylboronic ester 1 (10.5 mg, 0.0600 mmol), CDCl<sub>3</sub> (700 μL) and TEMPO 4 (15.0 mg, 0.100 mmol). The tube was capped and <sup>1</sup>H NMR spectra were taken to monitor the reaction. Spectral data was consistent with that of the authentic sample prepared previously. In particular, the –OCH<sub>2</sub>– peak in the <sup>1</sup>H NMR at δ 3.73 (t) was diagnostic of the formation of *N*-alkoxyamine 5. Note: the product 5 is volatile, and is easily lost on the vacuum pump.

***O*-*n*-Butyl-dansyl-TEMPO (8).** A. Preparation of authentic sample: following the literature procedure,<sup>57</sup> a solution of dansyl-TEMPO 4.16 (500 mg, 1.24 mmol) in dry toluene (17.8 mL, 0.18M) was cooled to –78 °C before adding *n*-butyllithium (2.5 M in hexanes, 0.35 mL, 0.87 mmol) dropwise. The reaction mixture was allowed to warm to room temperature, and then was quenched with water (17 mL). The organic layer was separated and the solvent was removed in vacuo. The crude product was purified by flash column chromatography (2:1 hexanes/ethyl acetate) to give the title compound as a viscous oil (239 mg, 83.9%).

TLC *R*<sub>f</sub> = 0.27 (2:1 hexanes:ethyl acetate, UV, I<sub>2</sub>, PAA). IR (CDCl<sub>3</sub>): 1574 (N–O) 1373 (N–O) 1143 (S–O) cm<sup>–1</sup>. <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>): δ 8.54 (d, *J* = 8.4 Hz, 1H), 8.28–8.26 (m, 2H), 7.58–7.51 (m, 2H), 7.19 (d, *J* = 7.2 Hz, 1H), 3.61 (t, *J* = 6.6 Hz, 2H), 3.43–3.40 (m, 1H), 2.89 (s, 6H), 1.57 (s, NH), 2.05–1.20 (m, 8H), 1.00 (s, 6H), 0.94 (s, 6H), 0.89–0.85 (m, 3H) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>): δ 152.1, 135.8, 130.4, 129.9, 129.7, 129.2,

128.4, 123.2, 118.7, 115.2, 60.5, 59.6, 46.3, 45.9, 45.4, 32.6, 30.7, 20.3, 19.5, 14.0 ppm. HRMS (ESITOF) Calcd for C<sub>22</sub>H<sub>40</sub>N<sub>3</sub>O<sub>3</sub>S (M + 1) 462.2785; found 462.2785.

B. Formation of 8 by reaction of dansyl-TEMPO with *B*-*n*-butylboronic acid and catechol: catechol (136 mg, 1.24 mmol) was added to an acetone solution (24.7 mL) of dansyl-TEMPO (1.00 g, 2.47 mmol), PbO<sub>2</sub> (591 mg, 2.47 mmol), and *B*-*n*-butylboronic acid (126 mg, 1.24 mmol) under nitrogen and stirred overnight. The next day, the reaction mixture was filtered through a pad of Celite and the solvent removed in vacuo. The resulting oil was dissolved in 25 mL of CH<sub>2</sub>Cl<sub>2</sub> and washed with 2 M NaOH (2 × 25 mL) followed by 25 mL of deionized water. The solution was dried over MgSO<sub>4</sub>, filtered and then concentrated in vacuo. The crude product was purified by flash column chromatography (2:1, hexanes:ethyl acetate) to give the title compound as a viscous oil (64%). Spectral data was consistent with that of the authentic sample prepared with *n*-butyl lithium. (Note: Lead oxide was initially used to ensure the nitroxide was not reduced by the catechol to the *N*-hydroxylamine, however it was later found to be unnecessary, and was not used in subsequent experiments.)

*R*<sub>f</sub> = 0.27 (2:1 hexanes/ethyl acetate, UV, I<sub>2</sub>, PAA).

## ■ ASSOCIATED CONTENT

### ☉ Supporting Information

General experimental methods, spot tests of dansyl-TEMPO 7 and *B*-*n*-butylboronic acid on a paper towel with water, and with catechol, <sup>1</sup>H and <sup>13</sup>C NMR spectra of all intermediates and products, and the DEPT spectrum of *O*-*n*-butyl-dansyl-TEMPO 8. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [rbraslau@ucsc.edu](mailto:rbraslau@ucsc.edu). Tel: 831 459-3087.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

The University of California Santa Cruz Committee on Research is acknowledged for partial support of this research. We thank JEOL for obtaining the DART MS data. We thank a reviewer for pointing out the possible use of the two-photon method to circumvent the fluorescence enhancement of dansyl-TEMPO in water. We thank another reviewer for a discussion of side reactions in the presence of oxygen.

## ■ REFERENCES

- Gladman, A. C. *Wilderness Environ. Med.* **2006**, *17*, 120.
- Tanner, T. L. *Primary Care* **2000**, *27*, 493.
- Oltman, J.; Hensler, R. *Clin. Dermatol.* **1986**, *4*, 213.
- NIOSH Publication No. 89–136; Centers for Disease Control: Atlanta, GA, 1988; <http://www.cdc.gov/niosh/docs/89-136/> (accessed May 23, 2012).
- McGovern, T. W.; Barkley, T. M. *Int. J. Dermatol.* **1998**, *37*, 321.
- Kullavanijaya, P.; Ophaswongse, S. *Contact Dermatitis* **1997**, *36*, 244.
- Devich, K. B.; Lee, J. C.; Epstein, W. L.; Spittler, L. E.; Hopper, J., Jr. *Clin. Nephrol.* **1975**, *3*, 106.
- Gelber, C.; Gemmell, L.; McAteer, D.; Homola, M.; Swain, P.; Liu, A.; Wilson, K. J.; Gefter, M. *J. Immunol.* **1997**, *158*, 2425.
- Symes, W. F.; Dawson, C. R. *J. Am. Chem. Soc.* **1954**, *76*, 2959.
- Elshohly, M. A.; Adawadkar, P. D.; Ma, C. Y.; Turner, C. E. *J. Nat. Prod.* **1982**, *45*, 532.
- Stibich, A. S.; Yagan, M.; Sharma, V.; Herndon, B.; Montgomery, C. *Int. J. Dermatol.* **2000**, *39*, 515.
- Guin, J. D. *J. Am. Acad. Dermatol.* **1980**, *2*, 332.



- (13) Natelson, S.; Lugovoy, J. K.; Pincus, J. B. *Arch. Biochem.* **1949**, *23*, 157.
- (14) Pennington, J. P.; Schoneich, C.; Stobaugh, J. F. *Chromatographia* **2007**, *66*, 649.
- (15) Schaffner, A. P.; Renaud, P. *Eur. J. Org. Chem.* **2004**, 2291.
- (16) Renaud, P.; Beauseigneur, A.; Brecht-Forster, A.; Becattini, B.; Darmency, V.; Kandhasamy, S.; Montermini, F.; Ollivier, C.; Panchaud, P.; Pozzi, D.; Scanlan, E. M.; Schaffner, A. P.; Weber, V. *Pure Appl. Chem.* **2007**, *79*, 223.
- (17) Suzuki, A.; Nozawa, S.; Itoh, M.; Brown, H. C.; Negishi, E. I.; Gupta, S. K. *J. Chem. Soc. D* **1969**, 1009.
- (18) Brown, H. C.; Negishi, E. I. *J. Am. Chem. Soc.* **1971**, *93*, 3777.
- (19) Baban, J. A.; Goodchild, N. J.; Roberts, B. P. *J. Chem. Soc., Perkin Trans. 2* **1986**, 157.
- (20) Darmency, V.; Renaud, P. *Top. Curr. Chem.* **2006**, *263*, 71.
- (21) Cadot, C.; Dalko, P. I.; Cossy, J.; Ollivier, C.; Chuard, R.; Renaud, P. *J. Org. Chem.* **2002**, *67*, 7193.
- (22) Ollivier, C.; Chuard, R.; Renaud, P. *Synlett* **1999**, 807.
- (23) Pouliot, M.; Renaud, P.; Schenk, K.; Studer, A.; Vogler, T. *Angew. Chem., Int. Ed. Engl.* **2009**, *48*, 6037.
- (24) Braslau, R.; Anderson, M. O. In *Radicals in Organic Synthesis*, 1st ed.; Renaud, P., Sibi, M. P., Eds.; Wiley-VCH: Weinheim, NY, 2001; Vol. 2, p 127.
- (25) Stryer, L.; Griffith, O. H. *Proc. Nat. Acad. Sci. U.S.A.* **1965**, *54*, 1785.
- (26) Blough, N. V.; Simpson, D. J. *J. Am. Chem. Soc.* **1988**, *110*, 1915.
- (27) Blinco, J. P.; Fairfull-Smith, K. E.; Morrow, B. J.; Bottle, S. E. *Aust. J. Chem.* **2011**, *64*, 373.
- (28) Likhentein, G. I.; Ishii, K.; Nakatsuji, S. *Photochem. Photobiol.* **2007**, *83*, 871.
- (29) Ivan, M. G.; Scaiano, J. C. *Photochem. Photobiol.* **2003**, *78*, 416.
- (30) Brown, H. C.; Bhat, N. G.; Somayaji, V. *Organometallics* **1983**, *2*, 1311.
- (31) Whitesides, G. M.; Newirth, T. L. *J. Org. Chem.* **1975**, *40*, 3448.
- (32) Likhentein, G. I.; Bogatyrenko, V. R.; Kulikov, A. V.; Hideg, K.; Khankovskaya, G. O.; Lukoyanov, N. V.; Kotelnikov, A. I.; Tanaseichuk, B. S. *Dokl. Akad. Nauk SSSR* **1980**, *253*, 481.
- (33) Maillard, B.; Ingold, K. U.; Scaiano, J. C. *J. Am. Chem. Soc.* **1983**, *105*, 5095.
- (34) Bowry, V. W.; Ingold, K. U. *J. Am. Chem. Soc.* **1992**, *114*, 4992.
- (35) Korich, A. L.; Iovine, P. M. *Dalton Trans.* **2010**, *39*, 1423.
- (36) Davies, A. G.; Ingold, K. U.; Roberts, B. P.; Tudor, R. *J. Chem. Soc. B* **1971**, 698.
- (37) Snyder, H. R.; Kuck, J. A.; Johnson, J. R. *J. Am. Chem. Soc.* **1938**, *60*, 105.
- (38) Berkowitz, J.; Ellison, G. B.; Gutman, D. *J. Phys. Chem.* **1994**, *98*, 2744.
- (39) Hall, D. G. *Boronic Acids: Preparation and Applications in Organic Synthesis and Medicine*; Wiley-VCH Verlag GmbH: Weinheim, 2005; p 9.
- (40) Albanell, J.; Adams, J. *Drugs Future* **2002**, *27*, 1079.
- (41) Hawthorne, M. F.; Lee, M. W. *J. Neuro-Oncol.* **2003**, *62*, 33.
- (42) Adamczyk-Wozniak, A.; Cyranski, M. K.; Zubrowska, A.; Sporzynski, A. *J. Organomet. Chem.* **2009**, *694*, 3533.
- (43) Rock, F. L.; Mao, W. M.; Yaremchuk, A.; Tukalo, M.; Crepin, T.; Zhou, H. C.; Zhang, Y. K.; Hernandez, V.; Akama, T.; Baker, S. J.; Plattner, J. J.; Shapiro, L.; Martinis, S. A.; Benkovic, S. J.; Cusack, S.; Alley, M. R. *K. Science* **2007**, *316*, 1759.
- (44) Aliaga, C.; Aspee, A.; Scaiano, J. C. *Org. Lett.* **2003**, *5*, 4145.
- (45) Aliaga, C.; Rezende, M. C.; Tirapegui, C. *Tetrahedron* **2009**, *65*, 6025.
- (46) Beckwith, A. L. J.; Bowry, V. W.; Ingold, K. U. *J. Am. Chem. Soc.* **1992**, *114*, 4983.
- (47) Aliaga, C.; Juarez-Ruiz, J. M.; Scaiano, J. C.; Aspee, A. *Org. Lett.* **2008**, *10*, 2147.
- (48) Povie, G.; Villa, G.; Ford, L.; Pozzi, D.; Schiesser, C. H.; Renaud, P. *Chem. Commun.* **2010**, *46*, 803.
- (49) Villa, G.; Povie, G.; Renaud, P. *J. Am. Chem. Soc.* **2011**, *133*, 5913.
- (50) Gryn'ova, G.; Ingold, K. U.; Coote, M. L. *J. Am. Chem. Soc.* **2012**, *134*, 12979.
- (51) <http://www.jeolusa.com/PRODUCTS/AnalyticalInstruments/MassSpectrometers/AccuTOFDART/tabid/230/Default.aspx> (accessed May 23, 2012).
- (52) Liu, L. L.; Li, Q.; Li, N. J.; Ling, J. H.; Liu, R.; Wang, Y. X.; Sun, L. X.; Chen, X. H.; Bi, K. S. *J. Sep. Sci.* **2011**, *34*, 1198.
- (53) Bjorklund, T.; Hall, H.; Breyse, N.; Soneson, C.; Carlsson, T.; Mandel, R. J.; Carta, M.; Kirik, D. *J. Neurochem.* **2009**, *111*, 355.
- (54) Li, Y. H.; Chan, L. M.; Tyer, L.; Moody, R. T.; Himel, C. M.; Hercules, D. M. *J. Am. Chem. Soc.* **1975**, *97*, 3118.
- (55) Tewari, N.; Joshi, N. K.; Rautela, R.; Gahlaut, R.; Joshi, H. C.; Pant, S. *J. Mol. Liq.* **2011**, *160*, 150.
- (56) Ahn, H. Y.; Fairfull-Smith, K. E.; Morrow, B. J.; Lussini, V.; Kim, B.; Bondar, M. V.; Bottle, S. E.; Belfield, K. D. *J. Am. Chem. Soc.* **2012**, *134* (10), 4721–4730.
- (57) Bushmakina, N. G.; Misharin, A. Y. *Synthesis* **1986**, 966–966.
- (58) Zhao, Y.; Zhong, Z. *J. Am. Chem. Soc.* **2005**, *127*, 17894–17901.
- (59) Green, M. L.; Wagner, M. *J. Chem. Soc., Dalton Trans.* **1996**, 2467–2473.
- (60) Brown, H. C.; Bhat, N. G.; Somayaji, V. *Organometallics* **1983**, *2*, 1311–1316.
- (61) Hartman, J. S.; Kelusky, E. C. *Can. J. Chem.* **1981**, *59*, 1284–1289.
- (62) Whitesides, G. M.; Newirth, T. L. *J. Org. Chem.* **1975**, *40*, 3448–3450.