


Regioselective Polymethylation of α -(1 \rightarrow 4)-Linked Mannopyranose Oligosaccharides

Li Xia and Todd L. Lowary*

Alberta Glycomics Centre and Department of Chemistry, University of Alberta, Gunning–Lemieux Chemistry Centre, Edmonton, AB T6G 2G2, Canada

S Supporting Information



ABSTRACT: An approach has been developed of the regioselective methylation of α -(1 \rightarrow 4)-linked mannopyranose oligosaccharides via a four-step methodology. The key reaction involved n -Bu₂SnCl₂-mediated activation of *cis*-diols. By tuning protecting groups on the substrates, multiple *cis*-diols in the substrates were functionalized in a consistent and regioselective manner. Using optimized substrates and reaction conditions, regioselective methylation of di-, tri-, and tetrasaccharide substrates proceeded in isolated yields per *cis*-diol of 95, 88 and 79%, respectively. The methodology was also applied to functionalize *trans*-diols in α -cyclodextrin.

INTRODUCTION

Total synthesis plays an important role in providing compounds that can be used for understanding the biological role of complex organic molecules, both naturally occurring and synthetic. However, this approach is generally laborious and time-consuming. To overcome this problem, semisynthetic approaches can provide a more rapid solution by using readily available compound skeletons as templates.^{1,2} For example, a recent example of semisynthesis was the installation of a trifluoromethyl moiety onto the drug Lipitor, via direct C–H functionalization.³

Although appealing, direct and selective functionalization of carbohydrates, particularly glycans more complicated than monosaccharides, is difficult. Although it is often possible to exploit steric differences to selectively functionalize primary over secondary carbohydrate alcohols, differentiating secondary hydroxyl groups on oligosaccharides is challenging.^{4,5} Regioselective acylations have been successfully carried out directly on unprotected disaccharides, e.g., sucrose **1**⁶ and lactose **2**⁷ (Figure 1), using organotin reagents. However, further functionalization of these molecules is often difficult, as the introduced acyl groups are base-labile and thus prone to migrate under the basic conditions typically used in alkylation reactions.⁸ In another example, sulfate groups have been introduced onto unprotected oligosaccharides, however, in low yield, e.g., xylose trisaccharide **3** (Figure 1).⁹

On the other hand, regioselective alkylation reactions appear to be more challenging to carry out on oligosaccharides. Indeed, we were surprised to discover that there are only limited examples of regioselective alkylation of unprotected oligosaccharides, e.g., **4**,^{10,11} **5**,¹² **6**¹³ (Figure 1). The regioselectivities of the reactions leading to **4** and **5** were

achieved by deprotonating the most acidic hydroxyl groups with strong bases; lactoside **6** was obtained via the organotin-mediated selective activation of vicinal diols.

During studies of the α -(1 \rightarrow 4)-mannosyltransferase involved in the biosynthesis of 3-*O*-methyl-mannose polysaccharides (MMPs),¹⁴ we had the need to access a panel of α -(1 \rightarrow 4)-linked mannopyranose oligosaccharides with or without methyl groups (**A** and **B**, Figure 2). Although these analogues are structurally similar, traditional routes for their synthesis require different building blocks and/or multiple functional group interconversions on oligosaccharides. For example, to obtain tetrasaccharides of both analogues ($n = 2$ for both **A** and **B**), 15 steps are required for **A** and 24 steps for **B** in Figure 2. We were therefore interested to determine if it was possible to access **B** via functionalization of **A**. We report here our studies toward achieving this goal.

RESULTS AND DISCUSSION

Initial Attempts to Regioselectively Alkylate α -(1 \rightarrow 4)-Linked Mannopyranose Oligosaccharides. We first studied the selective methylation of α -(1 \rightarrow 4)-linked mannopyranose di- and trisaccharides. These substrates were synthesized from pyranone **7**, which was obtained in two steps by treatment of (*R*)-2-(benzyloxy)-1-(furan-2-yl)ethanol¹⁵ with *N*-bromosuccinimide followed by reaction with di-*tert*-butyl dicarbonate.¹⁶ With **7** in hand, use of the Pd-catalyzed iterative glycosylation methodology developed by O'Doherty and co-workers (Scheme 1)¹⁷ provided diene **10** and triene **11**. Exhaustive dihydroxylation of the double bonds in **10** and **11**

Received: January 20, 2013

Published: March 13, 2013




Figure 1. Examples of regioselective acylation, sulfation or alkylation of oligosaccharides.




Figure 2. Conventional methods to synthesize analogues A and B and the approach developed here.

Scheme 1. Synthesis of Disaccharide 12 and Trisaccharide 13






Table 1. Direct Regioselective Alkylation of Disaccharide **12**^a

entry	T (°C)	additive	products distribution (NMR yield % ^b)		recovered 12 (%)
			disubstituted	monosubstituted	
1	40	–	<5%	15% mixture	80
2	70	–	12a (19%), 12b (3%)	12c (35%), 15% other isomers	28
3	40	CsF	12a (29%), 12b (45%), 26% other isomers	–	–

^aReaction conditions: **12** (1 equiv), $n\text{-Bu}_2\text{SnO}$ (2 equiv), $n\text{-Bu}_4\text{NI}$ (2 equiv), MeI (20 equiv), additive (4 equiv), toluene (0.01 M). ^bYields were obtained by integration of anomeric protons of the products observed in crude ¹H NMR spectrum.

Scheme 2. Direct Regioselective Alkylation of **13**


provided the desired disaccharide **12** and trisaccharide **13** in excellent overall yields. The high diastereoselectivity observed in these reactions is consistent with a previous report.¹⁷

We began by investigating the possibility of direct introduction of two alkyl groups onto disaccharide **12**, at O-3 and O-3' positions. Because $n\text{-Bu}_2\text{SnO}$ is well-known to selectively activate the equatorial hydroxyl groups of *cis*-diols via a cyclic stannylene acetal,¹⁸ we anticipated that using 2 equiv of this reagent would result in formation of a bis-coordinated intermediate **14**¹⁹ from disaccharide **12** (Table 1).^{18–22} Subsequent addition of the alkylating reagent would lead to the formation of the disubstituted product **12a** (Table 1), with alkylation at both equatorial positions (O-3 and O-3').

To explore this possibility, disaccharide **12** was heated at reflux in toluene with 2 equiv of $n\text{-Bu}_2\text{SnO}$ and then treated with methyl iodide at various temperatures (Table 1). Formation of products was monitored by ¹H NMR spectroscopy of the crude reaction mixtures, and the yield was determined by integrating the anomeric protons in the newly formed products. Three major products were observed: two

disubstituted regioisomers (**12a** and **12b**) as well as a monosubstituted derivative, **12c**. When the alkylation was performed at 40 °C (entry 1, Table 1), 80% of starting material **12** remained. Raising the reaction temperature to 70 °C increased the amount of dimethylated products, **12a** (19%) and **12b** (3%). However, the reaction was incomplete, and a number of monomethylated products were produced (entry 2, Table 1). Addition of CsF drove the reaction to completion with the formation of exclusively dimethylated products (entry 3, Table 1).²³ However, the desired regioisomer **12a** was formed in only 29% yield, while the major product was the undesired isomer **12b** in 45% yield. The remaining mass balance was the other dimethylated isomers, based upon ¹H NMR analysis of the crude reaction mixtures. The structures of **12a**, **12b** and **12c** were assigned unambiguously after isolation of the corresponding products and characterization by ¹H NMR spectroscopy along with ¹H–¹H COSY experiments. Regioisomers **12a** and **12b** were distinguished by comparison of the ring protons adjacent to the alkylated hydroxyl groups. Chemical shifts of H-2, H-2', H-3 and H-3' of **12** are between

Scheme 3. Regioselective Acylation of 12

Figure 3. Regioselective acylation of 2,3-*cis*-diol of mannose residues.^{25,28–30}

3.94–3.59 ppm. After alkylation H-3 and H-3' of **12a** are significantly upfield-shifted to 3.46 and 3.33 ppm, respectively, while H-2 and H-2' slightly downfield-shifted to 4.00 and 4.08 ppm, respectively. This result indicated **12a** is O-3, O-3'-alkylated. In **12b**, H-2 and H-3' are upfield shifted to 3.37 and 3.33 ppm, indicating substitution taking place at 2-OH and 3'-OH positions. For the major monoalkylated adduct, **12c**, only the signal for H-3' was deshielded.

Despite the rather modest results obtained with disaccharide **12** as the substrate, we applied the *n*-Bu₂SnO-mediated alkylation conditions to trisaccharide **13** (Scheme 2). With this substrate, trisubstitution took place predominantly. No starting material was detected after the reaction; however, a minimum of five regioisomers were observed. The desired 3,3',3''-trisubstituted **13a** was isolated as the major product, but it accounted only for 41% of all the products. Two other identified isomers were O-3, O-2', O-3'' tri-*O*-methylated **13b** and O-2, O-3', O-3'' tri-*O*-methylated **13c**. The structures of the products were established as described above for the reactions with disaccharide **12**.

These studies suggest that the degree of alkylation of carbohydrate polyol systems can be controlled with *n*-Bu₂SnO-coordination. However, consistent regioselectivity was difficult to achieve with substrates such as disaccharide **12** and trisaccharide **13**. We therefore decided to explore if regioselectivity could be improved for acylation reactions. We first tested acylation of disaccharide **12**. Because *n*-Bu₂SnO-mediated acylation of *cis*-diols usually activates the equatorial hydroxyl group,¹⁸ we expected to see predominant O-3, O-3'-disubstitution of **12**.

Regioselective Acylation of Disaccharide 12. Regioselective acylation of **12** was explored using BzCl at room temperature after heating the substrate at reflux with a tin reagent (2 equiv of *n*-Bu₂SnO or 1.5 equiv of (*n*-Bu₃Sn)₂O) in toluene. Of the two reagents used, *n*-Bu₂SnO gave only trace amounts of disubstituted products. Isolation was not attempted because of the low yield. In contrast, when (*n*-Bu₃Sn)₂O was used, di-*O*-benzoylated products were obtained exclusively (method A, Scheme 3). In addition, only two of the four possible regioisomers were produced. Unexpectedly, the anticipated 3,3'-disubstituted **15a** was only formed as a minor product (21%, NMR yield). Instead, **15b**, in which O-2 and O-

3' were benzoylated, was the major product and was produced in a 79% yield as observed from the ¹H NMR spectrum of the crude reaction mixture. This regiochemistry was supported by the significant downfield shift of H-2 and H-3' in the ¹H NMR spectrum for **15b** compared to that for **12**. In addition to the method described above, which involved an excess of tin reagent, the reaction was also performed with a catalytic amount of *n*-Bu₂SnO (method B, Scheme 3).²⁴ However, this catalytic method also produced the 3,3'-disubstituted product **15a** in a low yield (25%, NMR yield). The major product of this reaction was 2,3'-disubstituted **15b**, which was obtained in 60% yield. The remaining 15% were other unidentified disubstituted isomers.

Before trying more acylation conditions, we considered the origin of the regioselectivity. Noticeably, the two diols of **12** (2',3'-*cis*-diol and 2,3-*cis*-diol) were alkylated with different regioselectivity to produce **15b** as the major product using either method A or method B (Scheme 3). The diol at the nonreducing moiety (2',3'-*cis*-diol) underwent substitution preferentially at the equatorial 3'-OH group. In contrast, the diol at the reducing end residue (2,3-*cis*-diol) showed the opposite selectivity, resulting in preferential functionalization of the axial 2-OH group.

This latter result is inconsistent with many reported tin-mediated reactions in which functionalization of the equatorial hydroxyl group of *cis*-diols on six-membered rings is preferential.¹⁸ For example, benzoylation of **16**, when carried out in the presence of *n*-Bu₂SnO, produced only **16b** with substitution at the equatorial hydroxyl group (Figure 3).²⁵ More recently developed methods, using either borinic acid derivatives²⁶ or Me₂SnCl₂,²⁷ also give the same selectivity with *cis*-diols. Nevertheless, it should be noted that examples giving different regioselectivity have been previously reported in the case of mannose. As examples, **16–19**, when reacted under the same conditions, resulted in different regioselectivities (Figure 3).^{25,28–30} These compounds differ only in the protecting groups, suggesting that these substituents play an important role in controlling the regioselectivity.

To rationalize the selectivity of the acylation reaction of **12**, we compared the steric environment of the two diols. We postulated that the selectivity of the diol at the reducing end (2,3-*cis*-diol) was influenced by the bulky mannose substituent





Figure 4. Rationale for the observed regioselectivity of the benzylation of 12 with *n*-Bu₂SnO.

at O-4, which rendered O-3 less accessible for reaction than O-2 (Figure 4A). On the other hand, the diol at the nonreducing end of 12 (2',3'-*cis*-diol) is not hindered, and the acylation occurs on the equatorial O-3' position preferentially. This rationale also explains the regioselectivity trends observed in 16–19 (Figure 3). With no substituent at O-4, benzylation occurred exclusively at O-3 to give 16b.²⁵ However, the presence of any substituent at O-4 (17,²⁸ 18,²⁹ or 19³⁰) decreased the regioselectivity, and significant amounts of O-2 substitution was also observed. Therefore, we hypothesized that introducing a bulky substituent at O-4' of disaccharide 12 would reverse the selectivity at the nonreducing residue diol and give an enhanced yield of the O-2 and O-2' di-*O*-acylated products (Figure 4B).

Substrate Modification and Optimization of Regioselective Acylation Conditions. To test this hypothesis, a disaccharide modified at O-4' with a *t*-butyldimethylsilyl (TBS) group (21, Scheme 4) was synthesized. The TBS group was

Scheme 4. Preparation of TBS-Modified Disaccharide 21



introduced by treating alcohol 10 with TBSOTf, and the resulting product, 20, was then dihydroxylated with OsO₄ to provide 21 in 91% overall yield. With 21 in hand, the regioselective acylation reaction was examined. We investigated three reported catalytic methods for regioselective acylation of diols, including two tin reagents, Me₂SnCl₂²⁷ and *n*-Bu₂SnO,²⁴ as well as a borinic acid derivative Ph₂BOCH₂CH₂NH₂.²⁶ All these methods were reported to efficiently catalyze the regioselective acylation of monosaccharides, but their application to oligosaccharide systems had not been investigated.

When 21 was treated with 2.4 equiv of BzCl (1.2 equiv per diol) in the presence of 20 mol % Me₂SnCl₂, *n*-Bu₂SnO or Ph₂BOCH₂CH₂NH₂ (10 mol % per diol), both tin reagents gave satisfactory conversions (entries 1 and 2, Table 2). However, the use of Ph₂BOCH₂CH₂NH₂ led only to ~10% conversion of 21 (entry 3, Table 2). Using of Me₂SnCl₂ gave 60% of a mixture of mono-*O*-benzoylated products and 40% of di-*O*-benzoylated products (entry 1). Of the disubstituted products, 22% was the expected 2,2'-di-*O*-benzoylated 21a and 18% was the 3,2'-di-*O*-benzoylated isomer 21d as determined by ¹H NMR spectroscopy. The other two isomers, 21b and 21c, were not detected. With *n*-Bu₂SnO, three di-*O*-benzoylated products were formed: 63% of the desired 21a, 9% of 21c and 16% of 21d (entry 2). These results support our


hypothesis that a bulky substituent at O-4' of 21 alters the regioselectivity of the nonreducing end diol to favor substitution at O-2'.

The regioselectivity employing *n*-Bu₂SnO (21a:21d = 3.9:1) was better than with Me₂SnCl₂ (21a:21d = 1.2:1), suggesting that the size of the alkyl group in the catalyst also influences acylation regioselectivity. Indeed, replacing Me₂SnCl₂ with the more hindered *n*-Bu₂SnCl₂ gave 21a as the only observed disubstituted isomer (entry 4, Table 2). However, 57% of the monobenzoylated products remained when this catalyst was used. When the *n*-Bu₂SnCl₂-catalyzed reaction of 21 was repeated with 10 equiv of BzCl, (5 equiv per diol), an 87% yield of 21a was produced (entry 5, Table 2). However, when the same amount of BzCl was employed in reactions using *n*-Bu₂SnO, mainly tribenzoylated products were produced (entry 6, Table 2).

Encouraged by the success of these preliminary studies, the regioselective benzylation of 21 catalyzed by *n*-Bu₂SnCl₂ was optimized using various solvents (Table 3). In all solvents, a similar trend in the regioselectivity (21a ≫ 21c, 21d > 21b) was observed, with the exception of pyridine, where the substrate formed a mixture of products containing more than two benzoyl groups. Among all the solvents tested, THF gave the least amount of undesired isomers (entry 1). Lowering the temperature to 0 °C did not improve the selectivity but, as expected, slowed the reaction (entry 6). Thus, future reactions employed THF as the solvent and were carried out at room temperature.

Next, we tested the optimized method with other acylating reagents (Table 4). The relatively unhindered acylation reagent AcCl gave no regioselectivity (entry 1). The bulky PivCl produced a single disubstituted isomer 22a with the expected 2,2'-disubstitution; however, 51% of the mono-2-*O*-pivaloylated product remained, as well as 36% of starting material 21 (entry 2). Replacing BzCl with *p*-TsCl gave the expected di-*O*-tosylated product 23a in excellent yield and regioselectivity (entry 3), although the equivalents of TsCl and reaction time had to be increased to drive the reaction to completion. When tosylation was performed with 15 equiv of TsCl for 24 h, 89% of 23a was observed (entry 4).

Having established that the introduction of a sterically demanding protecting group on O-4' of 21 successfully switched the regioselectivity of the nonreducing end residue, we next introduced a trityl substituent onto substrate instead of a TBS group (disaccharide 24, Table 4). This compound was found to be a better substrate than 21. Regioselective tosylation of 24 efficiently furnished a 91% isolated yield of the desired 25a (entry 5, Table 4). Only a trace amount of other isomers was produced from this reaction. On the basis of these results and for reasons discussed in greater detail below, we shifted our focus to tosylation reactions.

Table 2. Regioselective Benzoylation of **21** with Different Catalysts^a

entry	BzCl amount (equiv)	conditions	products distribution (NMR yield %)					monosubstituted	recovered 21
			disubstituted						
			21a	21b	21c	21d			
1	2.4	Me ₂ SnCl ₂ , DIPEA, THF	22	–	–	18	60	–	
2	2.4	<i>n</i> -Bu ₂ SnO, Et ₃ N, CH ₂ Cl ₂	63	–	9	16	–	–	
3	2.4	Ph ₂ BOCH ₂ CH ₂ NH ₂ , CH ₃ CN	–	–	–	–	<10	90	
4	2.4	<i>n</i> -Bu ₂ SnCl ₂ , DIPEA, THF	43	–	–	–	57	–	
5	10	<i>n</i> -Bu ₂ SnCl ₂ , DIPEA, THF	87	–	6	7	–	–	
6	10	<i>n</i> -Bu ₂ SnO, Et ₃ N, CH ₂ Cl ₂	tribenzoylated products observed					–	–

^aReaction conditions: **21** (1 equiv, 0.1 M), catalyst (20 mol %), base (4 equiv for entries 1–4, 10 equiv for entries 5 and 6), BzCl (2.4–10 equiv), room temperature, 3 h.

Table 3. Solvent Optimization for Regioselective Benzoylation of **21**^a

entry	solvent	<i>T</i> (°C)	product distribution (%)				
			disubstituted				
			21a	21b	21c	21d	
1	THF	rt	87	–	6	7	–
2	CH ₂ Cl ₂	rt	68	–	10	20	–
3	MeCN	rt	61	–	<4	28	–
4	Et ₂ O	rt	76	–	12	12	–
5	Pyridine	rt	overbenzylation				–
6	THF	0	66	–	8	7	19

^aReaction conditions: **21** (1 equiv, 0.1 M), *n*-Bu₂SnCl₂ (20 mol %), DIPEA (10 equiv), BzCl (10 equiv), 3 h.


Application to Tri- and Tetrasaccharides Substrates.

With a set of optimized conditions developed, the methodology was applied to more challenging substrates: trisaccharide **26** and tetrasaccharide **27**, which were obtained from their corresponding triene **11** and tetraene **28**, respectively (Scheme 5). Considering that regioselective acylation of each diol generates two possible regioisomers, trisaccharide **26**, with three diol pairs, would have eight possible tri-*O*-sulfonylated isomers and tetrasaccharide **27** would have 16 possible products. When **26** and **27** were subjected to the optimized regioselective sulfonation conditions, the desired trisubstituted product **26a** and tetrasubstituted product **27a** were obtained as the major regioisomers in both cases, in 69 and 38% isolated yields, respectively. In both cases, these products were the major regioisomers. Although the isolated yield of the desired regioisomers dropped when the substrates used went from the disaccharide to the tetrasaccharide, the efficiency of the regioselectivity is still remarkable. Tosylation of disaccharide **24** produced a 91% yield of **25a**, corresponding to a 95%

sulfonylation selectivity for each diol pair. In the case of the reaction with trisaccharide **26**, a 69% yield of **26a** corresponds to an 88% selectivity for each diol pair. For the reaction with tetrasaccharide **27**, a 38% yield is equivalent to sulfonylating each diol with 79% selectivity.

Subsequent Functionalization of Oligosaccharide by Alkylation. After successfully introducing multiple protecting groups onto the di-, tri- and tetrasaccharides with good regioselectivity, the ability to further functionalize these compounds was investigated. Attempts to methylate **21a** using NaH as base resulted in benzoyl group migration (condition a, Scheme 6A), a common problem during alkylation under basic conditions.⁸ The use of milder bases such as Ag₂O and Ag₂CO₃ failed to prevent acyl migration (condition b, Scheme 6A).^{31,32} In addition, methylation of **21a** using CH₂N₂ and MeOTf also proved unsuccessful (conditions c and d, Scheme 6A).^{33,34}


To overcome this acyl migration problem, the substrates with the sulfonyl groups were used.³⁵ As expected, the tosyl groups of **23a** were stable to alkylation with NaH and MeI. However, in addition to the desired product **29**, an unexpected byproduct, **30** (Scheme 6B), presumably resulting from intramolecular silyl group migration under the basic reaction conditions, was formed. The use of an alternative substrate with a base-stable trityl group on O-4' (**25a**) avoided this problem, giving the desired di-*O*-methyl disaccharide **31**. The trityl group in **31** was easily removed using HCl in methanol to give **32** in 83% yield over the two steps. Finally, the tosyl groups of **32** were cleaved with magnesium in methanol at reflux.³⁶ Subsequent hydrogenolysis removed the remaining benzyl groups, furnishing partially methylated disaccharide **33** (Scheme 6C).

Table 4. Protecting Group Scope of Regioselective Acylation^a

entry	substrate	RCl (amount, equiv)	products (NMR yield %)		
			disubstituted (%)	monosubstituted (%)	unreacted starting material (%)
1	21 (R ¹ = TBS)	AcCl (2.4)	trace	>80	10
2		PivCl (10)	22a (13)	51	36
3		TsCl (10)	23a (62)	31	trace
4		TsCl (15)	23a (89), 23d (11)	—	—
5	24 (R ¹ = Tr)	TsCl (10)	25a (96), 25d (4)	—	—

^aReaction conditions: substrate (1 equiv, 0.1 M), $n\text{-Bu}_2\text{SnCl}_2$ (20 mol %), DIPEA (10 equiv), RCl (2.4–15 equiv), room temperature, 24 h.

Scheme 5. Regioselective Modification of Trisaccharide 26 and Tetrasaccharide 27




Following this approach, the methylated trisaccharide 36 and tetrasaccharide 37 were obtained three steps from 26a and 27a, in 41 and 70% overall yields, respectively (Scheme 7).



Application to α -Cyclodextrin. To explore the generality of the methodology, it was applied to α -cyclodextrin. Chemical modifications of cyclodextrins are of significant interest and have been widely used to tune their physical properties.³⁷ However, methods to selectively functionalize the secondary face of these cyclic oligosaccharides are still limited. In particular, derivatives that are functionalized at O-3 positions are difficult to prepare.³⁸ Noticing of the similarity between the α -(1 → 4)-linked mannose oligosaccharides described above and the α -(1 → 4)-linked glucopyranose residues present in α -cyclodextrin, we envisioned our methodology might be able to

provide the 3-O-alkylated analogues. As expected, the TBS-protected α -cyclodextrin 38³⁹ underwent tosylation with high regioselectivity (Scheme 8). The expected fully 2-O-tosylated product 39 was obtained in 77% isolated yield. Of the other byproducts produced from this reaction, the major one, isolated in 5% yield, had five of the residues exclusively tosylated at O-2. After protection of all the O-2 positions of cyclodextrin, methyl groups were installed quantitatively onto O-3 of 39 to give 40. This substrate has *trans*- instead of *cis*-diols, which suggested the approach does not require a *cis*-diol.⁴⁰ This methodology thus provides a new method for the selective functionalization of cyclodextrins.

Scheme 6. Attempted Methylation with Disaccharides 21a, 23a and 25a



Scheme 7. Methylation of Trisaccharide 26a and Tetrasaccharide 27a

Scheme 8. Regioselective Functionalization of α -Cyclodextrin

CONCLUSION

In conclusion, we demonstrate here a method that can be used to regioselectively functionalize multiple hydroxyl groups in oligosaccharides. Key to the development of this approach was the realization that the regioselectivity of organotin-mediated acylation and sulfonation of diols can be altered by tuning the size of protecting groups on the substrate. Although such effects can, in retrospect, be identified in previous examples of such functionalizations,^{25,28–30} rational optimization of these steric effects appears not to have been carried out previously.

The focus of this paper has been on the selective methylation of *cis*-diols in α -(1 \rightarrow 4)-linked mannosopyranose oligomers, which are an interesting class of lipid-binding oligosaccharides.^{41–43} When coupled with O'Doherty's palladium-catalyzed pyranone glycosylation methodology,¹⁷ we consider this approach to be a viable alternative to more traditional approaches to these targets, which rely on the preparation of a number of selectively protected monosaccharides derivatives.^{14,44–50} We also provide examples showing that the

method is effective for selectively functionalizing not only *cis*-diols but also *trans*-diols (in α -cyclodextrin), and further work in applying this method to other oligosaccharide systems is ongoing. In particular, the stereochemistry at C-1 and C-4, and the identity of the protecting group at O-6, may influence the selectivity of the process. We note that during the preparation of this manuscript a similar method, also employing *n*- Bu_2SnCl_2 as the catalyst, was reported for the regioselective mono-sulfonylation of an unprotected disaccharide by Muramatsu.⁵¹ The major difference is that whereas Muramatsu aimed to protect only one of the hydroxyl groups of an oligosaccharide, our approach targets simultaneous functionalization of multiple hydroxyl groups.

EXPERIMENTAL SECTION

General Experimental Methods. All reagents were purchased from commercial sources and were used without further purification unless noted. All reactions were carried out under a positive pressure of argon or nitrogen at room temperature unless specified and were monitored by TLC on silica gel 60-F₂₅₄ (0.25 mm). Visualization of the reaction components was achieved using UV fluorescence (254 nm) and/or by charring with acidified anisaldehyde solution in ethanol. Organic solvents were evaporated under reduced pressure, and the products were purified by column chromatography on silica gel (230–400 mesh). Optical rotations were measured in a microcell (10 cm, 1 mL) at ambient temperature and are in units of degree-mL/(g-dm). ¹H NMR spectra were recorded at 500 or 600 MHz, and chemical shifts are referenced to residual CHCl_3 (7.26 ppm, CDCl_3), CHDCl_2 (5.32 ppm, CD_2Cl_2), or CHD_2OD (3.30 ppm, CD_3OD). ¹³C NMR spectra were recorded at 125 MHz, and chemical shifts are referenced to CDCl_3 (77.0 ppm) or CD_2Cl_2 (53.8 ppm). Reported splitting patterns are abbreviated as s = singlet, d = doublet, t = triplet, m = multiplet, br = broad, app = apparent. Assignments of NMR

spectra were based on two-dimensional experiments (^1H - ^1H COSY, HSQC, and HMBC) and stereochemistry of the anomeric centers of the pyranose rings were confirmed by measuring $^1J_{\text{C-1,H-1}}$ via coupled HSQC experiments. High resolution ESI-MS spectra (time-of-flight analyzer) were recorded on samples suspended in THF or CH_3OH and with added NaCl. Low resolution MALDI-MS (time-of-flight analyzer) used 2,5-dihydroxybenzoic acid (DHB) as matrix.

(2R,6R)-6-((Benzyloxy)methyl)-5-oxo-5,6-dihydro-2H-pyran-2-yl tert-butyl carbonate (7). A reported method was used for the synthesis of **7**.¹⁶ To a stirring ice-cold solution of (*R*)-2-(benzyloxy)-1-(furan-2-yl)ethanol¹⁵ (2.9 g, 13.2 mmol) in mixed THF and H_2O (21 mL, THF- H_2O 4:1) was added $\text{NaHCO}_3 \cdot 3\text{H}_2\text{O}$ (2.3 g, 26.5 mmol) and NaOAc (1.1 g, 13.2 mmol). *N*-Bromosuccinimide (2.4 g, 13.2 mmol) was added, and the resulting yellow solution was stirred at 0 °C for 1.5 h. The reaction mixture was concentrated to remove THF and then extracted with CH_2Cl_2 . After washing with saturated aqueous NaHCO_3 and brine, the organic layer was dried over Na_2SO_4 and concentrated to afford a yellow liquid (3.4 g). This yellow liquid (1.5 g, 5.7 mmol) was then dissolved in CH_2Cl_2 (8 mL) and cooled to -78 °C. Di-*t*-butyl dicarbonate (1.5 g, 6.8 mmol) was added into this solution followed by 4-dimethylaminopyridine (70.8 mg, 0.6 mmol). The solution was stirred for 1 h while warming to room temperature. The resulting black solution was concentrated, and the crude residue was purified by chromatography (hexane-EtOAc 9:1) to afford α isomer **7** (919.4 mg, 48% over two steps) as a pale yellow syrup. This reaction also produced the β isomer (212.1 mg, 11%) as yellow syrup. The following data are for α isomer **7** only: R_f 0.57 (hexane-EtOAc 3:1); $[\alpha]_{\text{D}} = -72.2$ (*c* 3.0, CHCl_3); IR 1751.6 (C=O), 1702.3 (Boc C=O); ^1H NMR (500 MHz, CDCl_3) δ 7.39–7.27 (m, 5H, Ar), 6.94 (dd, $J = 10.3, 3.7$ Hz, 1H, H-2), 6.48 (d, $J = 3.7$ Hz, 1H, H-1), 6.28 (d, $J = 10.3$ Hz, 1H, H-3), 4.72 (dd, $J = 4.4, 2.7$ Hz, 1H, H-5), 4.60 (s, 2H, $2 \times \text{OCH}_2\text{Ph}$), 3.97 (dd, $J = 10.9, 4.4$ Hz, 1H, H-6a), 3.92 (dd, $J = 10.9, 2.6$ Hz, 1H, H-6b), 1.54 (s, 9H, Boc); ^{13}C NMR (126 MHz, CDCl_3) δ 193.0 (C-4), 151.7 (Boc C=O), 141.4 (C-2), 137.8 (Ar), 129.1 (C-3), 128.4 (2C, Ar), 127.7 (2C, Ar), 127.7 (Ar), 89.2 (C-1), 83.7 (Boc *t*-Bu), 76.4 (C-5), 73.7 (OCH_2Ph), 68.5 (C-6), 27.7 (Boc *t*-Bu); HRMS (ESI) calcd $\text{C}_{18}\text{H}_{22}\text{O}_6$ $[\text{M} + \text{Na}]^+$ 357.1309, found 357.1311.

(2R,6S)-2-((Benzyloxy)methyl)-6-(octyloxy)-2H-pyran-3(6H)-one (8). A solution of CH_2Cl_2 (3 mL) containing **7** (352.9 mg, 1.1 mmol) and 1-octanol (0.5 mL, 3.2 mmol) with 4 Å molecular sieves was stirred at room temperature for 0.5 h before 2.5 mol % tris(dibenzylideneacetone)dipalladium(0) $\text{Pd}_2(\text{dba})_3$ (26.8 mg, 0.03 mmol) and 10 mol % triphenylphosphine PPh_3 (28.0 mg, 0.12 mmol) were added. After stirring for 1.5 h, the resulting purple solution was concentrated, and the resulting residue was purified by chromatography (hexane-EtOAc 15:1) to afford **8** (351.4 mg, 96%) as a colorless liquid: R_f 0.59 (hexane-EtOAc 3:1); $[\alpha]_{\text{D}} = -32.7$ (*c* 0.9, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ 7.39–7.26 (m, 5H, Ar), 6.91 (dd, $J = 10.2, 3.5$ Hz, 1H, H-2), 6.15 (d, $J = 10.3$ Hz, 1H, H-3), 5.33 (d, $J = 3.5$ Hz, 1H, H-1), 4.65–4.60 (m, 3H, H-5, $2 \times \text{OCH}_2\text{Ph}$), 3.95 (dd, $J = 10.8, 4.8$ Hz, 1H, H-6a), 3.92 (dd, $J = 10.9, 3.0$ Hz, 1H, H-6b), 3.88 (dt, $J = 9.6, 6.8$ Hz, 1H, octyl OCH_2), 3.62 (dt, $J = 9.6, 6.6$ Hz, 1H, octyl OCH_2), 1.70–1.59 (m, 2H, octyl OCH_2CH_2), 1.44–1.21 (m, 10H, octyl CH_2), 0.90 (t, $J = 7.0$ Hz, 3H, octyl CH_3); ^{13}C NMR (126 MHz, CDCl_3) δ 194.4 (C-4), 144.1 (C-2), 138.0 (Ar), 128.3 (2C, Ar), 127.8 (Ar), 127.6 (3C, Ar (2C), C-3), 93.2 (C-1), 74.5 (C-5), 73.7 (OCH_2Ph), 69.7 (octyl OCH_2), 68.7 (C-6), 31.8, 29.7, 29.4, 29.3, 26.2, 22.7 (6C, octyl CH_2), 14.1 (octyl CH_3); HRMS (ESI) calcd $\text{C}_{21}\text{H}_{30}\text{O}_4$ $[\text{M} + \text{Na}]^+$ 369.2036, found 369.2040.

(2R,3S,6S)-2-((Benzyloxy)methyl)-6-(octyloxy)-3,6-dihydro-2H-pyran-3-ol (9). To a solution of ketone **8** (343.0 mg, 1.1 mmol) in methanol (2.5 mL) at -78 °C was added NaBH_4 (42.3 mg, 1.1 mmol) and $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (392.7 mg, 1.1 mmol). The solution was stirred overnight while warming to room temperature. The mixture was concentrated to remove the methanol, and then the residue was redissolved in CH_2Cl_2 . After washing with water and brine, the organic layer was dried over Na_2SO_4 and then concentrated. The resulting residue was purified by chromatography (hexane-EtOAc 6:1) to afford alcohol **9** (307.3 mg, 88%) as a colorless syrup: R_f 0.35

(hexane-EtOAc 3:1); $[\alpha]_{\text{D}} = +20.0$ (*c* 0.8, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ 7.40–7.27 (m, 5H, Ar), 5.94 (d, $J = 10.2$ Hz, 1H, H-3), 5.76 (ddd, $J = 10.2, 2.7, 2.2$ Hz, 1H, H-2), 4.99 (d, $J = 2.6$ Hz, 1H, H-1), 4.65 (d, $J = 12.0$ Hz, 1H, OCH_2Ph), 4.61 (d, $J = 12.0$ Hz, 1H, OCH_2Ph), 4.23 (dddd, $J = 9.2, 5.6, 3.5, 1.8$ Hz, 1H, H-4), 3.89–3.83 (m, 1H, H-5), 3.82–3.75 (m, 2H, H-6a, octyl OCH_2), 3.72 (dd, $J = 10.0, 4.9$ Hz, 1H, H-6b), 3.50 (dt, $J = 9.6, 6.6$ Hz, 1H, octyl OCH_2), 2.67 (d, $J = 5.9$ Hz, 1H, OH-4), 1.67–1.55 (m, 2H, octyl OCH_2CH_2), 1.43–1.22 (m, 10H, octyl CH_2), 0.91 (t, $J = 7.0$ Hz, 3H, octyl CH_3); ^{13}C NMR (126 MHz, CDCl_3) δ 137.9 (Ar), 133.0 (C-3), 128.5 (2C, Ar), 127.8 (Ar), 127.7 (2C, Ar), 126.2 (C-2), 94.3 (C-1), 73.7 (OCH_2Ph), 70.7 (C-6), 70.0 (C-5), 68.8 (octyl OCH_2), 65.6 (C-4), 31.9, 29.8, 29.4, 29.3, 26.2, 22.7 (6C, octyl CH_2), 14.1 (octyl CH_3); HRMS (ESI) calcd $\text{C}_{21}\text{H}_{32}\text{O}_4$ $[\text{M} + \text{Na}]^+$ 371.2193, found 371.2189.

(2R,3S,6S)-2-((Benzyloxy)methyl)-6-(((2R,3S,6S)-2-((benzyloxy)methyl)-6-(octyloxy)-3,6-dihydro-2H-pyran-3-yl)-oxy)-3,6-dihydro-2H-pyran-3-ol (10). The reaction was performed as described for the synthesis of **8**, with alcohol **9** (611.8 mg, 1.8 mmol) and donor **7** (818.5 mg, 2.6 mmol) in the presence of $\text{Pd}_2(\text{dba})_3$ (41.8 mg, 0.05 mmol) and PPh_3 (60.0 mg, 0.23 mmol) in CH_2Cl_2 (6 mL). The crude residue was purified by chromatography (hexane-EtOAc 7:1) to afford a ketone (842.1 mg, 85%) as a colorless syrup. This ketone was then reduced as described for **9**, with NaBH_4 (68.6 mg, 1.8 mmol) and $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ (656.2 mg, 1.8 mmol) in methanol (6 mL). Chromatographic purification of the crude reaction mixture (hexane-EtOAc 6:1) furnished alcohol **10** (770.3 mg, 91%) as a colorless syrup: R_f 0.38 (hexane-EtOAc 2:1); $[\alpha]_{\text{D}} = +35.3$ (*c* 0.9, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ 7.40–7.24 (m, 10H, Ar), 6.05 (d, $J = 10.3$ Hz, 1H, H-3), 5.97 (d, $J = 10.2$ Hz, 1H, H-3'), 5.84 (ddd, $J = 10.3, 2.7, 1.8$ Hz, 1H, H-2), 5.68 (ddd, $J = 10.2, 2.4, 2.4$ Hz, 1H, H-2'), 5.20 (d, $J = 2.4$ Hz, 1H, H-1'), 5.03 (d, $J = 2.1$ Hz, 1H, H-1), 4.62–4.41 (m, 5H, H-4, $4 \times \text{OCH}_2\text{Ph}$), 4.26 (dddd, $J = 10.3, 3.4, 3.4, 1.7$ Hz, 1H, H-4'), 4.01 (ddd, $J = 9.4, 3.6, 3.6$ Hz, 1H, H-5), 3.81 (dt, $J = 9.5, 6.8$ Hz, 1H, octyl OCH_2), 3.77–3.73 (m, 2H, H-6a, H-6b), 3.73–3.68 (m, 1H, H-5'), 3.66 (dd, $J = 9.6, 4.3$ Hz, 1H, H-6a'), 3.55 (dd, $J = 9.6, 5.7$ Hz, 1H, H-6b'), 3.50 (dt, $J = 9.5, 6.6$ Hz, 1H, octyl OCH_2), 2.33 (d, $J = 4.9$ Hz, 1H, OH-4'), 1.67–1.57 (m, 2H, octyl OCH_2CH_2), 1.40–1.21 (m, 10H, octyl CH_2), 0.90 (t, $J = 7.0$ Hz, 3H, octyl CH_3); ^{13}C NMR (126 MHz, CDCl_3) δ 138.5, 137.6 (2C, Ar), 133.2 (C-3'), 129.3 (C-3), 128.5 (2C, Ar), 128.3 (2C, Ar), 127.9 (Ar), 127.8 (2C, Ar), 127.4 (3C, Ar), 127.3 (C-2), 125.8 (C-2'), 94.3 (C-1), 91.1 (C-1'), 73.7, 73.21 (2C, $2 \times \text{OCH}_2\text{Ph}$), 70.6 (C-6'), 69.9 (C-5'), 69.7 (C-6), 69.1 (C-5), 68.7 (octyl OCH_2), 67.3 (C-4), 66.0 (C-4'), 31.9, 29.8, 29.4, 29.3, 26.3, 22.7 (6C, octyl CH_2), 14.1 (octyl CH_3); HRMS (ESI) calcd $\text{C}_{34}\text{H}_{46}\text{O}_7$ $[\text{M} + \text{Na}]^+$ 589.3136, found 589.3136.

(2R,3S,6S)-2-((Benzyloxy)methyl)-6-(((2R,3S,6S)-2-((benzyloxy)methyl)-6-(octyloxy)-3,6-dihydro-2H-pyran-3-yl)-oxy)-3,6-dihydro-2H-pyran-3-ol (11). The reaction was performed as described for the synthesis of **8**, with alcohol **10** (978.0 mg, 1.73 mmol) and donor **7** (1.02 g, 3.05 mmol) in the presence of $\text{Pd}_2(\text{dba})_3$ (69.0 mg, 0.08 mmol) and PPh_3 (80.0 mg, 0.31 mmol) in CH_2Cl_2 (22 mL). The crude residue was purified by chromatography (hexane-EtOAc 9:1) to afford a ketone (1.28 g, 95%) as a pale yellow syrup: $[\alpha]_{\text{D}} = +5.1$ (*c* 0.6, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ 7.38–7.21 (m, 15H, Ar), 6.83 (dd, $J = 10.2, 3.5$ Hz, 1H, H-2''), 6.18 (d, $J = 10.3$ Hz, 1H, H-3''), 6.11 (d, $J = 10.3$ Hz, 1H, H-3'), 6.05 (d, $J = 10.3$ Hz, 1H, H-3), 5.85 (ddd, $J = 10.3, 2.7, 1.8$ Hz, 1H, H-2), 5.78 (ddd, $J = 10.3, 2.7, 1.9$ Hz, 1H, H-2'), 5.57 (d, $J = 3.5$ Hz, 1H, H-1''), 5.27 (d, $J = 2.1$ Hz, 1H, H-1'), 5.03 (d, $J = 2.2$ Hz, 1H, H-1), 4.63 (dd, $J = 9.3, 1.3$ Hz, 1H, H-4'), 4.60–4.37 (m, 8H, H-4, H-5'', $6 \times \text{OCH}_2\text{Ph}$), 4.02 (ddd, $J = 9.3, 5.6, 1.8$ Hz, 1H, H-5), 3.85–3.75 (m, 4H, H-5', H-6a, H-6a'', octyl OCH_2), 3.71 (dd, $J = 10.9, 5.7$ Hz, 1H, H-6b), 3.63–3.58 (m, 2H, H-6a', H-6b''), 3.54 (dd, $J = 10.8, 1.7$ Hz, 1H, H-6b'), 3.50 (dt, $J = 9.5, 6.6$ Hz, 1H, octyl OCH_2), 1.66–1.57 (m, 2H, octyl OCH_2CH_2), 1.41–1.22 (m, 10H, octyl CH_2), 0.90 (t, $J = 7.0$ Hz, 3H, octyl CH_3); ^{13}C NMR (126 MHz, CDCl_3) δ 194.0 (C-4''), 143.6 (C-2''), 138.5, 138.0, 137.8 (3C, Ar), 129.2 (2C, C-3, C-3'), 128.3 (2C, Ar), 128.3 (2C, Ar), 128.3 (2C, Ar), 128.2 (C-3''), 127.7 (2C, Ar), 127.6 (2C, Ar), 127.6 (2C, Ar), 127.4 (4C, $3 \times \text{Ar}$, C-

2), 127.3 (C-2'), 94.2 (C-1), 91.4 (C-1'), 90.0 (C-1''), 75.0 (C-5''), 73.7, 73.5, 73.2 (3C, 3 × OCH₂Ph), 69.9 (C-6), 69.7 (C-5'), 69.1 (C-5), 69.0 (C-6'), 68.7 (octyl OCH₂), 68.5 (C-6''), 67.5 (C-4), 67.4 (C-4'), 31.9, 29.8, 29.4, 29.3, 26.3, 22.7 (6C, octyl CH₂), 14.1 (octyl CH₃); HRMS (ESI) calcd C₄₇H₅₈O₁₀ [M + Na]⁺ 805.3922, found 805.3915. This ketone (208.9 mg, 0.21 mmol) was then reduced as described for **9**, with NaBH₄ (8.0 mg, 0.21 mmol) and CeCl₃·7H₂O (80.2 mg, 0.21 mmol) in methanol (2.5 mL). Chromatographic purification of the crude reaction mixture (hexane–EtOAc 3:1) furnished alcohol **11** (177.2 mg, 85%) as a colorless syrup: *R*_f 0.30 (hexane–EtOAc 3:1); [α]_D²⁰ = +35.6 (c 0.8, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.38–7.23 (m, 15H, Ar), 6.07 (d, *J* = 10.1 Hz, 1H, H-3'), 6.05 (d, *J* = 9.3 Hz, 1H, H-3), 5.99 (d, *J* = 10.2 Hz, 1H, H-3'), 5.84 (ddd, *J* = 10.3, 2.7, 1.8 Hz, 1H, H-2), 5.74 (ddd, *J* = 10.3, 2.7, 1.9 Hz, 1H, H-2'), 5.70 (ddd, *J* = 10.2, 2.7, 2.3 Hz, 1H, H-2''), 5.25 (d, *J* = 2.0 Hz, 1H, H-1'), 5.20 (d, *J* = 2.4 Hz, 1H, H-1''), 5.03 (d, *J* = 2.1 Hz, 1H, H-1), 4.60–4.39 (m, 8H, H-4, H-4', 3 × OCH₂Ph), 4.27 (ddd, *J* = 8.6, 4.4, 1.7 Hz, 1H, H-4''), 4.02 (ddd, *J* = 9.2, 6.0, 1.7 Hz, 1H, H-5), 3.88–3.76 (m, 3H, H-5', H-6a, octyl OCH₂), 3.74–3.62 (m, 4H, H-5'', H-6a, H-6a', H-6a''), 3.60 (dd, *J* = 10.8, 2.0 Hz, 1H, H-6b'), 3.55–3.46 (m, 2H, H-6b'', octyl OCH₂), 2.41 (d, *J* = 4.6 Hz, 1H, OH-4''), 1.67–1.55 (m, 2H, octyl OCH₂CH₂), 1.40–1.20 (m, 10H, octyl CH₂), 0.90 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (126 MHz, CDCl₃) δ 138.5, 138.4, 137.6 (3C, 3 × Ar), 133.3 (C-3''), 129.9, 129.3 (2C, C-3, C-3'), 128.5 (2C, Ar), 128.3 (2C, Ar), 128.3 (2C, Ar), 127.9 (Ar), 127.8 (2C, Ar), 127.4 (3C, Ar), 127.4 (2C, Ar), 127.4 (Ar), 127.3 (C-2), 126.7 (C-2'), 125.7 (C-2''), 94.2 (C-1), 91.3 (C-1'), 91.0 (C-1''), 73.7, 73.4, 73.2 (3C, 3 × OCH₂Ph), 70.7 (C-6''), 69.9 (C-6), 69.8 (2C, C-5', C-5''), 69.2 (C-6'), 69.1 (C-5), 68.7 (octyl OCH₂), 67.5 (C-4), 66.7 (C-4'), 66.1 (C-4''), 31.9, 29.8, 29.4, 29.3, 26.3, 22.7 (6C, octyl CH₂), 14.1 (octyl CH₃); HRMS (ESI) calcd C₄₇H₆₀O₁₀ [M + Na]⁺ 807.4079, found 807.4070.

Octyl 6-O-benzyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-α-D-mannopyranoside (12). To a solution of **10** (561.8 mg, 0.97 mmol) in *t*-butanol and acetone (4 mL, v/v 1:1) was added OsO₄ (2.5 wt.% in *t*-butanol, 250 μL, 0.02 mmol) and *N*-methyl-morpholine *N*-oxide (NMO, 50% w/v in water, 0.3 mL). After stirring at room temperature overnight, saturated aqueous Na₂SO₃ solution was added. The mixture was concentrated to remove *t*-butanol, and then the residue was extracted with CH₂Cl₂ three times. The combined organic layer was concentrated, and the resulting residue was purified by chromatography (CH₂Cl₂–methanol 15:1) to afford **12** (492.3 mg, 80%) as a pale yellow syrup: *R*_f 0.14 (CH₂Cl₂–methanol 15:1); [α]_D²⁰ = +59.5 (c 1.2, methanol); ¹H NMR (498 MHz, CD₃OD) δ 7.34–7.20 (m, 10H, Ar), 5.22 (d, *J* = 1.8 Hz, 1H, H-1'), 4.69 (d, *J* = 1.6 Hz, 1H, H-1), 4.53–4.41 (m, 4H, 4 × OCH₂Ph), 3.94 (dd, *J* = 3.1, 1.9 Hz, 1H, H-2'), 3.83–3.59 (m, 12H, H-2, H-3, H-3', H-4, H-4', H-5, H-5', 2 × H-6, 2 × H-6', octyl OCH₂), 3.39 (dt, *J* = 9.6, 6.4 Hz, 1H, octyl OCH₂), 1.61–1.52 (m, 2H, octyl OCH₂CH₂), 1.38–1.21 (m, 10H, octyl CH₂), 0.88 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (125 MHz, CD₃OD) δ 139.8, 139.7 (2C, Ar), 129.3 (2C, Ar), 129.3 (2C, Ar), 129.0 (2C, Ar), 128.9 (2C, Ar), 128.6 (Ar), 128.6 (Ar), 103.4 (C-1', ¹J_{C-1,H-1} = 171.4 Hz), 101.4 (C-1, ¹J_{C-1,H-1} = 167.0 Hz), 76.6 (C-4), 74.6, 74.4 (2C, 2 × OCH₂Ph), 74.4, 73.2, 72.7, 72.5, 72.2, 72.1 (6C, C-2, C-2', C-3, C-3', C-5, C-5'), 71.4, 71.4 (2C, C-6, C-6'), 68.8 (octyl OCH₂), 68.7 (C-4'), 33.0, 30.6, 30.4, 30.4, 27.3, 23.7 (6C, octyl CH₂), 14.5 (octyl CH₃); HRMS (ESI) calcd C₃₄H₅₀O₁₁ [M + Na]⁺ 657.3245, found 657.3239.

Synthesis of Octyl 6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-3-O-methyl-α-D-mannopyranoside (12a), Octyl 6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-2-O-methyl-α-D-mannopyranoside (12b) and Octyl 6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-α-D-mannopyranoside (12c). Compound **12** (0.01–0.04 mmol, 1 equiv) and *n*-Bu₄SnO (2 equiv) were heated at reflux in toluene (0.01 M) overnight. The resulting yellowish solution was cooled to room temperature before MeI (20 equiv), *t*-Bu₄NI (2 equiv) and CsF (none for entry 1 and 2, 4 equiv for entry 3) were added. The mixture was stirred at designated temperature (see Table 1) overnight before being concentrated. The crude products were then purified by

chromatography (CH₂Cl₂–methanol 30:1 to 20:1) to afford an inseparable mixture of **12a** and **12b** as a yellow syrup, as well as separable **12c** as a yellow syrup: *R*_f 0.51 for **12a** and **12b**, 0.43 for **12c** (CH₂Cl₂–methanol 12:1).

Octyl 6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-3-O-methyl-α-D-mannopyranoside (12a). ¹H NMR (600 MHz, CD₃OD) δ 7.39–7.18 (m, 10H, Ar), 5.16 (d, *J* = 1.9 Hz, 1H, H-1'), 4.76 (d, *J* = 1.8 Hz, 1H, H-1), 4.53–4.40 (m, 4H, 4 × OCH₂Ph), 4.08 (dd, *J* = 3.1, 2.1 Hz, 1H, H-2'), 4.00 (dd, *J* = 3.2, 1.9 Hz, 1H, H-2), 3.87–3.55 (m, 9H, H-4, H-4', H-5, H-5', 4 × H-6, octyl OCH₂), 3.46 (dd, *J* = 9.2, 3.3 Hz, 1H, H-3), 3.45 (s, 3H, OMe), 3.43–3.40 (m, 1H, octyl OCH₂), 3.41 (s, 3H, OMe), 3.33 (dd, *J* = 9.3, 3.1 Hz, 1H, H-3'), 1.62–1.55 (m, 2H, octyl OCH₂CH₂), 1.40–1.21 (m, 10H, octyl CH₂), 0.84 (t, *J* = 7.5 Hz, 3H, octyl CH₃); HRMS (ESI) calcd C₃₆H₅₄O₁₁ [M + Na]⁺ 685.3558, found 685.3555.

Octyl 6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-2-O-methyl-α-D-mannopyranoside (12b). ¹H NMR (600 MHz, CD₃OD) δ 7.39–7.18 (m, 10H, Ar), 5.24 (d, *J* = 1.9 Hz, 1H, H-1'), 4.87 (d, *J* = 1.6 Hz, 1H, H-1), 4.53–4.40 (m, 4H, 4 × OCH₂Ph), 4.16 (dd, *J* = 3.0, 2.1 Hz, 1H, H-2'), 3.87–3.55 (m, 10H, H-3, H-4, H-4', H-5, H-5', 4 × H-6, octyl OCH₂), 3.44 (s, 3H, OMe), 3.43–3.40 (m, 1H, octyl OCH₂), 3.37 (dd, *J* = 3.5, 1.7 Hz, 1H, H-2), 3.33 (dd, *J* = 9.3, 3.1 Hz, 1H, H-3'), 1.62–1.55 (m, 2H, octyl OCH₂CH₂), 1.40–1.21 (m, 10H, octyl CH₂), 0.84 (t, *J* = 7.5 Hz, 3H, octyl CH₃); HRMS (ESI) calcd C₃₆H₅₄O₁₁ [M + Na]⁺ 685.3558, found 685.3555.

Octyl 6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-α-D-mannopyranoside (12c). ¹H NMR (600 MHz, CD₃OD) δ 7.36–7.19 (m, 10H, Ar), 5.25 (d, *J* = 2.0 Hz, 1H, H-1'), 4.70 (d, *J* = 1.6 Hz, 1H, H-1), 4.54–4.40 (m, 4H, 2 × OCH₂Ph), 4.17 (dd, *J* = 3.0, 2.1 Hz, 1H, H-2'), 3.85–3.60 (m, 11H, H-2, H-3, H-4, H-4', H-5, H-5', 4 × H-6, octyl OCH₂), 3.45 (s, 3H, OMe), 3.42–3.38 (m, 1H, octyl OCH₂), 3.35 (dd, *J* = 9.3, 3.2 Hz, 1H, H-3'), 1.62–1.53 (m, 2H, octyl OCH₂CH₂), 1.38–1.23 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.1 Hz, 3H, octyl CH₃); HRMS (ESI) calcd C₃₅H₅₂O₁₁ [M + Na]⁺ 671.3402, found 671.3402.

Octyl 6-O-benzyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-α-D-mannopyranoside (13). The dihydroxylation reaction was performed as described for the synthesis of **12**, with **11** (243.3 mg, 0.31 mmol), OsO₄ (2.5 wt.% in *t*-butanol, 119 μL, 0.01 mmol), NMO (50% w/v in water, 0.9 mL) in *t*-butanol and acetone (1 mL, v/v 1:1). The crude residue was purified by chromatography (CH₂Cl₂–methanol 12:1) to afford trisaccharide **13** (205.1 mg, 75%) as a colorless foam: *R*_f 0.61 (CH₂Cl₂–methanol 6:1); [α]_D²⁰ = +64.8 (c 1.0, methanol); ¹H NMR (498 MHz, CD₃OD) δ 7.33–7.17 (m, 15H, Ar), 5.25 (d, *J* = 1.6 Hz, 1H, H-1'/H-1''), 5.23 (d, *J* = 1.7 Hz, 1H, H-1'/H-1''), 4.71 (d, *J* = 1.6 Hz, 1H, H-1), 4.52–4.33 (m, 6H, 6 × OCH₂Ph), 3.98 (dd, *J* = 3.0, 1.9 Hz, 1H, H-2'/H-2''), 3.90 (dd, *J* = 2.9, 2.2 Hz, 1H, H-2'/H-2''), 3.86–3.60 (m, 17H, H-2, 3 × H-3, 3 × H-4, 3 × H-5, 6 × H-6, octyl OCH₂), 3.38 (dt, *J* = 9.6, 6.3 Hz, 1H, octyl OCH₂), 1.61–1.52 (m, 2H, octyl OCH₂CH₂), 1.39–1.21 (m, 10H, OCH₂), 0.88 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (125 MHz, CD₃OD) δ 139.6 (Ar), 139.5 (2C, Ar), 129.3 (2C, Ar), 129.3 (2C, Ar), 129.2 (2C, Ar), 129.0 (2C, Ar), 129.0 (2C, Ar), 128.8 (2C, Ar), 128.6 (Ar), 128.6 (Ar), 128.5 (Ar), 103.3 (C-1'/C-1'', ¹J_{C-1,H-1} = 170 Hz), 103.0 (C-1'/C-1'', ¹J_{C-1,H-1} = 171 Hz), 101.3 (C-1, ¹J_{C-1,H-1} = 168.8 Hz), 76.4, 76.2 (2C, C-4, C-4'), 74.5, 74.4 (2C, 2 × OCH₂Ph), 74.3 (C-5'), 74.2 (OCH₂Ph), 73.2, 72.9, 72.8, 72.6, 72.6, 72.5, 72.1, 71.9 (9C, 3 × C-2, 3 × C-3, C-4'', C-5, C-5'), 71.3, 71.3, 71.1 (3C, 3 × C-6), 68.7 (octyl OCH₂), 32.9, 30.5, 30.3, 30.3, 27.2, 23.6 (6C, octyl CH₂), 14.5 (octyl CH₃); HRMS (ESI) calcd C₄₇H₆₆O₁₆ [M + Na]⁺ 909.4243, found 909.4244.

Synthesis of Octyl 6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-3-O-methyl-α-D-mannopyranoside (13a), Octyl 6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-3-O-methyl-α-D-mannopyranoside (13b) and Octyl 6-O-benzyl-3-O-methyl-α-D-mannopyranosyl-(1 → 4)-6-O-benzyl-2-O-methyl-α-D-mannopyranoside (13c). Compound **13** (33.3 mg,

0.04 mmol) and *n*-Bu₂SnO (31.8 mg, 0.12 mmol, 3 equiv) were heated at reflux in toluene (4 mL) for 14 h. The resulting yellowish solution was cooled to room temperature before MeI (70 μ L, 1.2 mmol), *t*-Bu₄NI (44.0 mg, 0.12 mmol, 3 equiv) were added. After stirring at 70 °C for 7 h, additional MeI (150 μ L, 2.4 mmol) was added, and the reaction was stirred overnight. The crude products were purified by chromatography (CH₂Cl₂–methanol 30:1 to 25:1) to afford three major products: an inseparable mixture (11.7 mg, 34%) of **13a** and **13c**, as well as **13b** (6.0 mg, 17%) both as colorless syrups: *R*_f 0.64 for **13b**, 0.61 for **13a** and **13c** (CH₂Cl₂–methanol 15:1); **13a** and **13c** were successfully separated after per-acetylation with acetic anhydride (0.1 mL) in pyridine (0.1 mL) and CH₂Cl₂ (1 mL). Chromatography purification gave acetylated **13a** (6.9 mg) and acetylated **13c** (3.5 mg) as colorless syrup.

Octyl 2,4-di-O-acetyl-6-O-benzyl-3-O-methyl- α -D-mannopyranosyl-(1 \rightarrow 4)-2-O-acetyl-6-O-benzyl-3-O-methyl- α -D-mannopyranoside (acetylated **13a).** ¹H NMR (600 MHz, CDCl₃) δ 7.39–7.16 (m, 15H, Ar), 5.39 (dd, *J* = 3.1, 2.0 Hz, 1H, H-2'), 5.38 (dd, *J* = 3.0, 2.1 Hz, 1H, H-2''), 5.28 (dd, *J* = 3.2, 1.8 Hz, 1H, H-2), 5.25 (d, *J* = 1.8 Hz, 1H, H-1'), 5.19 (app t, *J* = 10.0 Hz, 1H, H-4''), 5.19 (d, *J* = 1.8 Hz, 1H, H-1'), 4.80 (d, *J* = 1.7 Hz, 1H, H-1), 4.64–4.38 (m, 6H, 6 \times OCH₂Ph), 3.90 (app t, *J* = 10.0 Hz, 1H, H-4'), 3.89–3.79 (m, 5H, H-4, H-5, H-5', H-5''), 3.76–3.68 (m, 3H, H-6, H-6, octyl OCH₂), 3.67–3.62 (m, 2H, H-3, H-6, H-6), 3.61–3.56 (m, 2H, H-3', H-3''), 3.46–3.41 (m, 3H, H-6, H-6, octyl OCH₂), 3.41 (s, 3H, OMe), 3.40 (s, 3H, OMe), 3.36 (s, 3H, OMe), 2.11 (s, 3H, Ac), 2.08 (s, 3H, Ac), 2.06 (s, 3H, Ac), 1.99 (s, 3H, Ac), 1.66–1.58 (m, 2H, octyl OCH₂CH₂), 1.40–1.23 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.1 Hz, 3H, octyl CH₃); ¹³C NMR (126 MHz, CDCl₃) δ 170.3, 170.1, 170.0, 169.9 (4C, 4 \times OAc), 138.5 (2C, Ar), 138.2 (Ar), 128.3 (2C, Ar), 128.3 (2C, Ar), 128.2 (2C, Ar), 127.8 (2C, Ar), 127.6 (Ar), 127.4 (3C, Ar), 127.4 (2C, Ar), 127.4 (Ar), 99.7 (C-1''), 99.4 (C-1'), 97.5 (C-1), 80.0 (C-3), 79.9 (C-3'), 76.5 (C-3''), 74.5 (C-4), 73.6 (C-4'), 73.6, 73.4, 73.3 (3 \times OCH₂Ph), 71.9, 71.0, 70.8 (3C, C-5, C-5', C-5''), 70.0, 69.9, 69.5 (3C, C-6, C-6', C-6''), 68.4 (C-4''), 68.3 (octyl OCH₂), 67.7 (2C, C-2, C-2'), 67.4 (C-2''), 57.6, 57.2, 57.1 (3C, 3 \times OMe), 31.8, 29.4, 29.4, 29.2, 26.1, 22.7 (6C, octyl CH₂), 21.0, 21.0, 21.0, 21.0 (4C, 4 \times Ac), 14.1 (octyl CH₃); HRMS (ESI) calcd C₅₈H₈₀O₂₀ [M + Na]⁺ 1119.5135, found 1119.5114.

Octyl 6-O-benzyl-3-O-methyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-2-O-methyl- α -D-mannopyranoside (13b). ¹H NMR (600 MHz, CD₃OD) δ 7.41–7.13 (m, 15H, Ar), 5.24 (d, *J* = 2.0 Hz, 1H, H-1''), 5.24 (d, *J* = 1.7 Hz, 1H, H-1'), 4.76 (d, *J* = 1.8 Hz, 1H, H-1), 4.53–4.32 (m, 6H, 6 \times OCH₂Ph), 4.17 (dd, *J* = 3.0, 2.1 Hz, 1H, H-2''), 4.01 (dd, *J* = 3.2, 1.9 Hz, 1H, H-2), 3.86–3.56 (m, 14H, H-3', H-4, H-4', H-4'', H-5, H-5', H-5'', 6 \times H-6, 1 \times octyl OCH₂), 3.48–3.40 (m, 3H, H-2', H-3, octyl OCH₂), 3.45 (s, 3H, OMe), 3.43 (s, 3H, OMe), 3.42 (s, 3H, OMe), 3.36 (dd, *J* = 9.2, 3.2 Hz, 1H, H-3'), 1.65–1.52 (m, 2H, octyl OCH₂CH₂), 1.42–1.20 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.1 Hz, 3H, octyl CH₃); HRMS (ESI) calcd C₅₀H₇₂O₁₆ [M + Na]⁺ 951.4713, found 951.4708.

Octyl 2,4-di-O-acetyl-6-O-benzyl-3-O-methyl- α -D-mannopyranosyl-(1 \rightarrow 4)-2-O-acetyl-6-O-benzyl-3-O-methyl- α -D-mannopyranoside (acetylated **13c).** ¹H NMR (600 MHz, CDCl₃) δ 7.39–7.13 (m, 15H, Ar), 5.37 (dd, *J* = 3.1, 2.1 Hz, 1H, H-2''), 5.22 (dd, *J* = 9.6, 3.3 Hz, 1H, H-3), 5.21 (d, *J* = 1.9 Hz, 1H, H-1'), 5.18 (app t, *J* = 9.8 Hz, 1H, H-4''), 5.16 (dd, *J* = 2.9, 2.1 Hz, 1H, H-2'), 4.99 (d, *J* = 1.9 Hz, 1H, H-1'), 4.86 (d, *J* = 1.7 Hz, 1H, H-1), 4.61–4.39 (m, 6H, 6 \times OCH₂Ph), 4.01 (app t, *J* = 9.6 Hz, 1H, H-4), 3.91–3.60 (m, 10H, H-2, H-4', H-5, H-5', H-5'', 4 \times H-6, 1 \times octyl OCH₂), 3.59 (dd, *J* = 9.8, 3.2 Hz, 1H, H-3''), 3.56 (dd, *J* = 9.0, 3.1 Hz, 1H, H-3'), 3.43 (s, 3H, OMe), 3.47–3.39 (m, 3H, 2 \times H-6, 1 \times octyl OCH₂), 3.37 (s, 3H, OMe), 3.36 (s, 3H, OMe), 2.20 (s, 3H, OAc), 2.10 (s, 3H, OAc), 2.06 (s, 3H, OAc), 1.99 (s, 3H, OAc), 1.66–1.58 (m, 2H, octyl OCH₂CH₂), 1.40–1.21 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (126 MHz, CDCl₃) δ 170.7 (OAc), 170.1 (OAc), 170.0 (OAc), 169.9 (OAc), 138.7, 138.4, 138.0, 128.3, 128.2, 128.2, 127.8, 127.6, 127.5, 127.4, 127.4, 127.3 (18C, Ar),

100.2 (C-1'), 99.5 (C-1''), 97.0 (C-1), 79.5 (C-3'), 78.4 (C-2), 76.5 (C-3''), 75.7 (C-4), 73.8 (2C, C-3, C-4'), 73.6, 73.3, 73.3 (3C, 3 \times OCH₂Ph), 71.9 (C-5''), 71.2 (C-5'), 70.8 (C-5), 69.9, 69.8, 69.5 (3C, C-6, C-6', C-6''), 68.4 (C-4''), 68.2 (octyl OCH₂), 67.7 (C-2'), 67.6 (C-2''), 59.3, 57.6, 57.2 (3C, 3 \times OMe), 31.8, 29.5, 29.4, 29.2, 26.1, 22.7 (6C, octyl CH₂), 21.2, 21.0, 20.9, 20.9 (4C, 4 \times Ac), 14.1 (octyl CH₃); HRMS (ESI) calcd C₅₈H₈₀O₂₀ [M + Na]⁺ 1119.5135, found 1119.5117.

Synthesis of Octyl 3-O-benzoyl-6-O-benzyl- α -D-mannopyranosyl-(1 \rightarrow 4)-3-O-benzoyl-6-O-benzyl- α -D-mannopyranoside (15a) and Octyl 3-O-benzoyl-6-O-benzyl- α -D-mannopyranosyl-(1 \rightarrow 4)-2-O-benzoyl-6-O-benzyl- α -D-mannopyranoside (15b). Method A: Disaccharide **12** (13.3 mg, 0.02 mmol) and (*n*-Bu₂Sn)₂O (175 μ L, 0.03 mmol) were heated at reflux in toluene (1.2 mL) for 15 h. The resulting yellowish solution was cooled to room temperature before BzCl (48 μ L, 0.4 mmol) was added. After stirring at room temperature for 6 h, the reaction mixture was concentrated, and the crude products were confirmed by ¹H NMR spectroscopy to be a mixture of **15a** (21% NMR yield) and **15b** (79% NMR yield). Method B: To a solution of **12** (14.1 mg, 0.02 mmol) in CH₂Cl₂ (0.5 mL) was added *n*-Bu₂SnO (1.3 mg, 0.004 mmol), Et₃N (9 μ L, 0.06 mmol) followed by BzCl (6.5 μ L, 0.05 mmol). The reaction mixture was stirred at room temperature overnight. The crude NMR spectrum indicated a mixture of **15a** (25% NMR) and **15b** (60% NMR) and 15% other regioisomers.

Octyl 3-O-benzoyl-6-O-benzyl- α -D-mannopyranosyl-(1 \rightarrow 4)-3-O-benzoyl-6-O-benzyl- α -D-mannopyranoside (15a). ¹H NMR (600 MHz, CDCl₃) δ 8.23–6.99 (m, 20H, Ar), 5.49 (dd, *J* = 9.6, 3.0 Hz, 1H, H-3), 5.23 (dd, *J* = 9.8, 3.2 Hz, 1H, H-3'), 5.22 (d, *J* = 1.4 Hz, 1H, H-1'), 4.87 (d, *J* = 1.9 Hz, 1H, H-1), 4.61–4.49 (m, 4H, 4 \times OCH₂Ph), 4.38 (app t, *J* = 9.5 Hz, 1H, H-4), 4.20–4.18 (m, 1H, H-2), 4.10 (app td, *J* = 9.6, 4.3 Hz, 1H, H-4'), 4.01–3.71 (m, 6H, H-2', H-5, H-5', H-6a, H-6b, octyl OCH₂), 3.69 (dd, *J* = 10.2, 3.6 Hz, 1H, H-6a'), 3.61 (dd, *J* = 10.2, 4.2 Hz, 1H, H-6b'), 3.50–3.43 (m, 1H, octyl OCH₂), 2.71 (d, *J* = 4.3 Hz, 1H, OH-4'), 2.09 (d, *J* = 6.6 Hz, 1H, OH-2), 1.89 (d, *J* = 5.2 Hz, 1H, OH-2'), 1.78–1.58 (m, 2H, octyl OCH₂CH₂), 1.55–1.18 (m, 10H, octyl CH₂), 0.90 (t, *J* = 7.1 Hz, 3H, octyl CH₃); HRMS (ESI) calcd C₄₈H₅₈O₁₃ [M + Na]⁺ 865.3770, found 865.3776.

Octyl 3-O-benzoyl-6-O-benzyl- α -D-mannopyranosyl-(1 \rightarrow 4)-2-O-benzoyl-6-O-benzyl- α -D-mannopyranoside (15b). ¹H NMR (600 MHz, CD₃OD) δ 8.23–6.99 (m, 20H, Ar), 5.38 (d, *J* = 2.0 Hz, 1H, H-1'), 5.28 (dd, *J* = 9.7, 3.2 Hz, 1H, H-3'), 5.22 (dd, *J* = 2.8, 1.8 Hz, 1H, H-2), 4.87 (d, *J* = 1.8 Hz, 1H, H-1), 4.61–4.49 (m, 4H, 4 \times OCH₂Ph), 4.25 (dd, *J* = 3.1, 2.0 Hz, 1H, H-2'), 4.23–4.19 (m, 1H, H-3), 4.07 (app t, *J* = 9.9 Hz, 1H, H-4'), 4.01–3.92 (m, 3H, H-5, H-5', H-6), 3.90–3.83 (m, 2H, H-4, H-6), 3.79–3.67 (m, 3H, 2 \times H-6, octyl OCH₂), 3.49 (app dt, *J* = 9.7, 6.4 Hz, 1H, octyl OCH₂), 1.78–1.58 (m, 2H, octyl OCH₂CH₂), 1.55–1.18 (m, 10H, octyl CH₂), 0.90 (t, *J* = 7.1 Hz, 3H, octyl CH₃); HRMS (ESI) calcd C₄₈H₅₈O₁₃ [M + Na]⁺ 865.3770, found 865.3776.

(((2R,3S,6S)-2-((Benzyloxy)methyl)-6-(((2R,3S,6S)-2-((benzyloxy)methyl)-6-(octyloxy)-3,6-dihydro-2H-pyran-3-yl)-oxy)-3,6-dihydro-2H-pyran-3-yl)oxy)(tert-butyl)dimethylsilane (20). To a solution of alcohol **10** (302.3 mg, 0.53 mmol) in CH₂Cl₂ (6 mL) at –10 °C was added 2,6-lutidine (280 μ L, 2.40 mmol) followed by TBSOTf (380 μ L, 1.60 mmol). The resulting solution was stirred for 0.5 h before being concentrated. The crude product was purified by chromatography (hexane–EtOAc 12:1) to afford trisaccharide **20** (360.4 mg, 99%) as a colorless syrup: *R*_f 0.65 (hexane–EtOAc 3:1); [α]_D = +58.8 (c 1.1, CHCl₃); ¹H NMR (498 MHz, CDCl₃) δ 7.36–7.20 (m, 10H, Ar), 6.02 (d, *J* = 10.3 Hz, 1H, H-3), 5.87 (d, *J* = 10.2 Hz, 1H, H-3'), 5.81 (ddd, *J* = 10.3, 2.6, 1.9 Hz, 1H, H-2), 5.60 (ddd, *J* = 10.2, 2.7, 2.1 Hz, 1H, H-2'), 5.21 (d, *J* = 2.8 Hz, 1H, H-1'), 5.01 (d, *J* = 2.0 Hz, 1H, H-1), 4.58 (d, *J* = 12.2 Hz, 1H, OCH₂Ph), 4.53 (d, *J* = 12.2 Hz, 1H, OCH₂Ph), 4.47 (s, 2H, OCH₂Ph), 4.38 (dd, *J* = 9.5, 1.2 Hz, 1H, H-4), 4.34 (dd, *J* = 9.1, 1.4 Hz, 1H, H-4'), 4.02 (ddd, *J* = 9.0, 6.0, 1.7 Hz, 1H, H-5), 3.84–3.73 (m, 2H, H-6a, octyl OCH₂), 3.71–3.62 (m, 2H, H-5', H-6b), 3.55 (dd, *J* = 10.5, 3.7 Hz, 1H, H-6a'), 3.51–3.44 (m, 2H, H-6b', octyl OCH₂), 1.68–1.52 (m, 2H, octyl OCH₂CH₂), 1.38–1.18 (m, 10H, octyl CH₂), 0.97–0.81 (m, 12H,

octyl CH₃, TBS(*t*-Bu)), 0.08 (s, 3H, TBS(Me)), 0.02 (s, 3H, TBS(Me)); ¹³C NMR (125 MHz, CDCl₃) δ 138.5, 138.1 (2C, Ar), 134.8 (C-3'), 129.4 (C-3), 128.3 (2C, Ar), 128.2 (2C, Ar), 127.7 (2C, Ar), 127.5 (Ar), 127.4 (2C, Ar), 127.3 (Ar), 127.2 (C-2), 125.1 (C-2'), 94.2 (C-1), 91.6 (C-1'), 73.5, 73.2 (2C, 2 × OCH₂Ph), 71.6 (C-5'), 69.9 (C-6), 69.1 (C-5), 68.6, 68.6 (2C, C-6', octyl OCH₂), 67.5 (C-4), 63.8 (C-4'), 31.8, 29.8, 29.4, 29.3, 26.3 (5C, octyl CH₂), 25.7 (TBS(*t*-Bu)), 22.7 (octyl CH₂), 17.9 (TBS(*t*-Bu)), 14.1 (octyl CH₃), -4.2, -4.9 (2C, 2 × TBS(Me)); HRMS (ESI) calcd C₄₀H₆₀O₇Si [M + Na]⁺ 703.4001, found 703.3994.

Octyl 6-O-benzyl-4-O-*t*-butyldimethylsilyl- α -D-mannopyranosyl-(1 → 4)-6-O-benzyl- α -D-mannopyranoside (21). The dihydroxylation reaction was performed as described for the synthesis of **12**, with **20** (588.6 mg, 0.86 mmol), OsO₄ (2.5 wt.% in *t*-butanol, 213 μ L, 0.017 mmol), NMO (50% w/v in water, 1.3 mL) in *t*-butanol and acetone (9 mL, v/v 1:1). The crude residue was purified by chromatography (CH₂Cl₂–methanol 30:1) to afford **21** (68.4 mg, 91%) as a colorless syrup: *R*_f 0.49 (CH₂Cl₂–methanol 15:1); [α]_D = +58.7 (c 0.7, CHCl₃); ¹H NMR (500 MHz, CD₃OD) δ 7.33–7.22 (m, 10H, Ar), 5.24 (d, *J* = 1.9 Hz, 1H, H-1'), 4.71 (d, *J* = 1.5 Hz, 1H, H-1), 4.57–4.41 (m, 4H, 4 × OCH₂Ph), 3.89 (dd, *J* = 3.1, 2.2 Hz, 1H, H-2), 3.84–3.77 (m, 3H, H-4, H-6a, H-6b), 3.77–3.67 (m, 6H, H-2', H-3, H-4', H-5, H-5', octyl OCH₂), 3.60 (dd, *J* = 8.5, 3.3 Hz, 1H, H-3'), 3.59–3.55 (m, 2H, H-6a', H-6b'), 3.40 (dt, *J* = 9.6, 6.3 Hz, 1H, octyl OCH₂), 1.65–1.52 (m, 2H, octyl OCH₂CH₂), 1.41–1.23 (m, 10H, octyl CH₂), 0.89 (t, *J* = 6.9 Hz, 3H, octyl CH₃), 0.86 (s, 9H, TBS(*t*-Bu)), 0.14 (s, 3H, TBS(Me)), 0.05 (s, 3H, TBS(Me)); ¹³C NMR (126 MHz, CD₃OD) δ 139.7 (Ar), 139.5 (Ar), 129.4 (2C, Ar), 129.4 (2C, Ar), 129.2 (2C, Ar), 128.8 (2C, Ar), 128.7 (Ar), 128.6 (Ar), 103.1 (C-1', ¹J_{C-1,H-1} = 171.7 Hz), 101.5 (C-1, ¹J_{C-1,H-1} = 167.3 Hz), 76.1 (C-4), 74.6 (C-5'), 74.5, 74.4 (2C, 2 × OCH₂Ph), 73.3 (C-4'), 72.9, 72.7, 72.7, 72.1 (4C, C-2, C-2', C-3, C-5), 71.4 (C-6), 71.0 (C-6'), 70.5 (C-3'), 68.9 (octyl OCH₂), 33.0, 30.6, 30.4, 27.4 (5C, octyl CH₂), 26.7 (TBS(*t*-Bu)), 23.7 (octyl CH₂), 19.2 (TBS(*t*-Bu)), 14.5 (octyl CH₃), -3.4, -4.7 (2C, 2 × TBS(Me)); HRMS (ESI) calcd C₄₀H₆₄O₁₁Si [M + Na]⁺ 771.4110, found 771.4106.

General Procedures for Reactions in Table 2. To a solution of **21** (1 equiv, 0.1 M) in the indicated solvent (0.1 M) was added catalyst (20 mol %) and base (4 equiv for entry 1–4, 10 equiv for entry 5–6) followed by BzCl (2.4–10 equiv). The resulting mixture was stirred at room temperature for 3 h before being concentrated. The crude products **21a**, **21c** and **21d** were purified by chromatography as indicated below.

Octyl 2-O-benzoyl-6-O-benzyl-4-O-*t*-butyldimethylsilyl- α -D-mannopyranosyl-(1 → 4)-2-O-benzoyl-6-O-benzyl- α -D-mannopyranoside (21a). The reaction was performed as the general procedure described above. Chromatographic purification (hexane–EtOAc 7:1) gave **21a** as a colorless film: *R*_f 0.66 (toluene–EtOAc 5:1); [α]_D = +13.9 (c 0.4, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 8.05 (dd, *J* = 8.4, 1.3 Hz, 2H, Ar), 8.01 (dd, *J* = 8.4, 1.3 Hz, 2H, Ar), 7.60–7.52 (m, 2H, Ar), 7.44–7.26 (m, 14H, Ar), 5.49 (d, *J* = 1.8 Hz, 1H, H-1'), 5.36 (dd, *J* = 3.4, 1.8 Hz, 2H, H-2, H-2'), 4.97 (d, *J* = 1.8 Hz, 1H, H-1), 4.73–4.54 (m, 4H, 4 × OCH₂Ph), 4.42 (dd, *J* = 9.6, 3.5 Hz, 1H, H-3), 4.27 (app t, *J* = 9.6 Hz, 1H, H-4), 4.12–4.03 (m, 1H, H-3'), 4.01–3.90 (m, 4H, H-4', H-5, H-5', H-6a), 3.88 (dd, *J* = 10.4, 1.5 Hz, 1H, H-6b), 3.76 (dt, *J* = 9.6, 6.8 Hz, 1H, octyl OCH₂), 3.67 (dd, *J* = 10.3, 1.0 Hz, 1H, H-6a'), 3.63 (dd, *J* = 10.3, 5.1 Hz, 1H, H-6b'), 3.49 (dt, *J* = 9.6, 6.6 Hz, 1H, octyl OCH₂), 1.68–1.57 (m, 2H, octyl OCH₂CH₂), 1.44–1.22 (m, 10H, octyl CH₂), 0.91 (t, *J* = 7.0 Hz, 3H, octyl CH₃), 0.87 (s, 9H, TBS(*t*-Bu)), 0.13 (s, 3H, TBS(Me)), 0.05 (s, 3H, TBS(Me)); ¹³C NMR (126 MHz, CDCl₃) δ 166.3 (C=O), 66.2 (C=O), 138.6 (Ar), 138.1 (Ar), 133.3 (Ar), 133.2 (Ar), 129.9 (2C, Ar), 129.8 (2C, Ar), 129.7 (Ar), 129.6 (Ar), 128.5 (2C, Ar), 128.4 (2C, Ar), 128.3 (2C, Ar), 128.3 (2C, Ar), 127.5 (3C, Ar), 127.4 (3C, Ar), 97.3 (C-1), 96.9 (C-1'), 75.2 (C-4), 73.5, 73.5 (2C, C-2, C-2'), 73.4, 73.3 (2C, 2 × OCH₂Ph), 73.2 (C-5'), 70.8 (C-3'), 69.9, 69.6 (2C, C-4', C-5), 69.5 (C-6), 69.3 (C-6'), 69.2 (C-3), 68.3 (octyl OCH₂), 31.8, 29.4, 29.4, 29.2, 26.2 (5C, octyl CH₂), 25.9 (TBS(*t*-Bu)), 22.7 (octyl CH₂), 18.3 (TBS(Me)), 14.1 (octyl CH₃), -4.0,

-4.9 (2C, 2 × TBS(Me)); HRMS (ESI) calcd C₅₄H₇₂O₁₃Si [M + Na]⁺ 979.4634, found 979.4628.

Octyl 3-O-benzoyl-6-O-benzyl-4-O-*t*-butyldimethylsilyl- α -D-mannopyranosyl-(1 → 4)-2-O-benzoyl-6-O-benzyl- α -D-mannopyranoside (21c). The reaction was performed as the general procedure described above. Chromatographic purification (hexane–EtOAc 2:1) gave **21c** as a colorless film: *R*_f 0.14 (toluene–EtOAc 5:1); [α]_D = +17.2 (c 0.2, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 8.15–7.97 (m, 4H, Ar), 7.62–7.17 (m, 16H, Ar), 5.37 (dd, *J* = 8.0, 3.0 Hz, 1H, H-3'), 5.35 (dd, *J* = 3.4, 1.8 Hz, 1H, H-2), 5.32 (d, *J* = 2.7 Hz, 1H, H-1'), 4.97 (d, *J* = 1.7 Hz, 1H, H-1), 4.70–4.49 (m, 4H, 4 × OCH₂Ph), 4.40 (dd, *J* = 9.5, 3.4 Hz, 1H, H-3), 4.24–4.14 (m, 3H, H-2', H-4, H-4'), 4.02–3.92 (m, 3H, H-5, H-5', H-6a), 3.91–3.85 (m, 1H, H-6b), 3.76 (dt, *J* = 9.6, 6.7 Hz, 1H, octyl OCH₂), 3.59 (d, *J* = 3.7 Hz, 2H, H-6a', H-6b'), 3.48 (dt, *J* = 9.6, 6.7 Hz, 1H, octyl OCH₂), 1.74–1.57 (m, 2H, octyl OCH₂CH₂), 1.47–1.23 (m, 10H, octyl CH₂), 0.92 (t, *J* = 6.9 Hz, 3H, octyl CH₃), 0.76 (s, 9H, TBS(*t*-Bu)), -0.01 (s, 3H, TBS(Me)), -0.05 (s, 3H, TBS(Me)); ¹³C NMR (126 MHz, CDCl₃) δ 166.3 (C=O), 166.0 (C=O), 138.6 (Ar), 137.9 (Ar), 133.2 (2C, Ar), 130.0 (Ar), 129.9 (2C, Ar), 129.9 (2C, Ar), 129.7 (Ar), 128.4 (2C, Ar), 128.4 (2C, Ar), 128.4 (2C, Ar), 128.3 (2C, Ar), 127.7 (2C, Ar), 127.6 (Ar), 127.3 (3C, Ar), 100.5 (C-1'), 97.4 (C-1), 75.9 (C-4), 75.4 (C-3'), 73.7 (C-5'), 73.4 (OCH₂Ph), 73.4 (C-2), 73.3 (OCH₂Ph), 70.3 (C-5), 69.8, 69.7 (2C, C-3, C-4'), 69.6 (C-6), 69.1 (C-6'), 68.3 (octyl OCH₂), 66.6 (C-2'), 31.9, 29.5, 29.4, 29.3, 26.2 (5C, octyl CH₂), 25.7 (TBS(*t*-Bu)), 22.7 (octyl CH₂), 18.0 (TBS(*t*-Bu)), 14.1 (octyl CH₃), -4.2, -4.9 (2C, 2 × TBS(Me)); HRMS (ESI) calcd C₅₄H₇₂O₁₃Si [M + Na]⁺ 979.4634, found 979.4629.

Octyl 2-O-benzoyl-6-O-benzyl-4-O-*t*-butyldimethylsilyl- α -D-mannopyranosyl-(1 → 4)-3-O-benzoyl-6-O-benzyl- α -D-mannopyranoside (21d). The reaction was performed as the general procedure described above. Chromatographic purification (hexane–EtOAc 4:1) gave **21d** as a colorless film: *R*_f 0.43 (toluene–EtOAc 5:1); [α]_D = +42.9 (c 0.2, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 8.11 (dd, *J* = 8.4, 1.3 Hz, 2H, Ar), 7.81 (dd, *J* = 8.4, 1.3 Hz, 2H, Ar), 7.60–7.22 (m, 16H, Ar), 5.53 (dd, *J* = 9.6, 3.2 Hz, 1H, H-3), 5.25 (d, *J* = 1.8 Hz, 1H, H-1'), 5.14 (dd, *J* = 3.1, 2.0 Hz, 1H, H-2'), 4.89 (d, *J* = 1.8 Hz, 1H, H-1), 4.70–4.53 (m, 4H, 4 × OCH₂Ph), 4.36 (app t, *J* = 9.6 Hz, 1H, H-4), 4.21 (dd, *J* = 3.1, 2.0 Hz, 1H, H-2), 4.02–3.95 (m, 2H, H-4', H-5), 3.94–3.89 (m, 2H, H-3', H-6a), 3.85 (dd, *J* = 10.8, 1.8 Hz, 1H, H-6b), 3.83–3.74 (m, 2H, H-5', octyl OCH₂), 3.73 (dd, *J* = 10.7, 4.5 Hz, 1H, H-6a'), 3.62 (dd, *J* = 10.6, 1.7 Hz, 1H, H-6b'), 3.49 (dt, *J* = 9.6, 6.7 Hz, 1H, octyl OCH₂), 1.72–1.60 (m, 2H, octyl OCH₂CH₂), 1.49–1.24 (m, 10H, octyl CH₂), 0.92 (t, *J* = 6.8 Hz, 3H, octyl CH₃), 0.87 (s, 9H, TBS(*t*-Bu)), 0.07 (s, 3H, TBS(Me)), 0.03 (s, 3H, TBS(Me)); ¹³C NMR (126 MHz, CDCl₃) δ 165.6 (C=O), 165.6 (C=O), 138.6 (Ar), 138.3 (Ar), 133.3 (Ar), 133.1 (Ar), 129.8 (2C, Ar), 129.7 (2C, Ar), 129.5 (Ar), 129.4 (Ar), 128.5 (Ar), 128.4 (2C, Ar), 128.30 (2C, Ar), 128.28 (2C, Ar), 128.2 (2C, Ar), 127.6 (2C, Ar), 127.4 (Ar), 127.3 (2C, Ar), 99.5 (C-1), 99.4 (C-1'), 74.9 (C-3), 73.9 (C-5'), 73.5 (OCH₂Ph), 73.4 (C-4), 73.3 (OCH₂Ph), 73.0 (C-2'), 70.9, 70.6, 69.6 (3C, C-3', C-4', C-5), 69.3 (C-6), 69.2 (C-6'), 69.0 (C-2), 68.3 (octyl OCH₂), 31.9, 29.4, 29.4, 29.3, 26.2 (5C, octyl CH₂), 26.0 (TBS(*t*-Bu)), 22.7 (octyl CH₂), 18.3 (TBS(*t*-Bu)), 14.1 (octyl CH₃), -4.1, -5.0 (2C, 2 × TBS(Me)); HRMS (ESI) calcd C₅₄H₇₂O₁₃Si [M + Na]⁺ 979.4634, found 979.4628.

Octyl 6-O-benzyl-4-O-*t*-butyldimethylsilyl-2-O-pivaloyl- α -D-mannopyranosyl-(1 → 4)-6-O-benzyl-2-O-pivaloyl- α -D-mannopyranoside (22a). The reaction was performed as the general procedure for Table 2 described above. The amounts for each reagent was listed in Table 4. The reaction mixture were concentrated and examined by crude ¹H NMR: ¹H NMR (600 MHz, CDCl₃) δ 7.34–7.23 (m, 10H, Ar), 5.27 (d, *J* = 1.8 Hz, 1H, H-1'), 5.04 (dd, *J* = 3.2, 1.7 Hz, 1H, H-2'), 4.98 (dd, *J* = 3.7, 1.8 Hz, 1H, H-2), 4.78 (d, *J* = 2.0 Hz, 1H, H-1), 4.62–4.44 (m, 4H, 4 × OCH₂Ph), 4.22 (dd, *J* = 9.5, 3.5 Hz, 1H, H-3), 4.06 (dd, *J* = 9.5, 3.2 Hz, 1H, H-3'), 4.00–3.35 (m, 10H, H-4, H-4', H-5, H-5', 4 × H-6, 2 × octyl OCH₂), 1.72–1.60 (m, 2H, octyl OCH₂CH₂), 1.35–1.24 (m, 10H, octyl CH₂), 0.87 (t, *J* = 6.8 Hz, 3H, octyl CH₃), 1.26 (s, 9H, Piv), 0.83 (s, 9H, TBS(*t*-Bu)), 0.09 (s, 3H, TBS(Me)), 0.00 (s, 3H, TBS(Me)).

Synthesis of Octyl 6-O-benzyl-4-O-*t*-butyldimethylsilyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-2-O-toluenesulfonyl- α -D-mannopyranoside (23a) and Octyl 6-O-benzyl-4-O-*t*-butyldimethylsilyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-toluenesulfonyl- α -D-mannopyranoside (23d). To a solution of **23** (50.5 mg, 0.067 mmol) in THF (0.7 mL) was added *n*-Bu₂SnCl₂ (20 mol %, 4.2 mg, 0.0013 mmol) and DIPEA (240 μ L, 1.34 mmol) followed by TsCl (190.7 mg, 0.013 mmol). The reaction mixture was stirred at room temperature for 24 h before being concentrated. The crude products **23a** and **23d** were purified by chromatography as described below.

Octyl 6-O-benzyl-4-O-*t*-butyldimethylsilyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-2-O-toluenesulfonyl- α -D-mannopyranoside (23a). Chromatographic purification (hexane–EtOAc 5:1) to give **23a** (47.7 mg, 67%) as a colorless syrup: *R*_f 0.66 (CH₂Cl₂–methanol 30:1); [α]_D = +27.2 (c 0.9, CHCl₃); ¹H NMR (600 MHz, CDCl₃) δ 7.82 (d, *J* = 8.3 Hz, 2H, Ar), 7.77 (d, *J* = 8.3 Hz, 2H, Ar), 7.35–7.22 (m, 14H, Ar), 5.09 (d, *J* = 1.8 Hz, 1H, H-1'), 4.80 (d, *J* = 1.7 Hz, 1H, H-1), 4.68 (dd, *J* = 3.1, 2.0 Hz, 1H, C-2'), 4.65 (dd, *J* = 3.4, 1.8 Hz, 1H, H-2), 4.53–4.40 (m, 4H, 4 \times OCH₂Ph), 4.02 (dd, *J* = 8.8, 3.2 Hz, 1H, H-3), 3.78–3.73 (m, 2H, H-3', H-5'), 3.72–3.65 (m, 3H, H-4, H-5, H-6a), 3.65–3.58 (m, 3H, H-4', H-6b, octyl OCH₂), 3.56 (dd, *J* = 10.3, 1.7 Hz, 1H, H-6a'), 3.44 (dd, *J* = 10.3, 6.8 Hz, 1H, H-6b'), 3.36 (dt, *J* = 9.6, 6.6 Hz, 1H, octyl OCH₂), 2.43 (s, 3H, ArCH₃), 2.40 (s, 3H, ArCH₃), 1.58–1.48 (m, 2H, octyl OCH₂CH₂), 1.33–1.22 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.1 Hz, 3H, octyl CH₃), 0.79 (s, 9H, TBS(*t*-Bu)), 0.06 (s, 3H, TBS(Me)), –0.04 (s, 3H, TBS(Me)); ¹³C NMR (125 MHz, CDCl₃) δ 145.3, 145.1, 138.4, 137.8, 133.2, 133.0, 129.9, 129.8, 128.3, 128.2, 128.1, 127.6, 127.5, 127.4 (24C, Ar), 97.4 (C-1'), 97.1 (C-1), 79.4 (C-2'), 79.2 (C-2), 75.8 (C-4), 73.3 (2C, C-5', OCH₂Ph), 73.2 (OCH₂Ph), 70.0, 69.9 (2C, C-3', C-5), 69.4 (C-4'), 69.3 (C-6), 69.1 (C-6'), 68.5 (C-3), 68.2 (octyl OCH₂), 31.8, 29.3, 29.2, 26.0 (5C, octyl CH₂), 25.8 (TBS(*t*-Bu)), 22.6 (octyl CH₂), 21.7 (ArCH₃), 21.6 (ArCH₃), 18.1 (TBS(*t*-Bu)), 14.1 (octyl CH₃), –4.0, –5.0 (2C, TBS(Me)); HRMS (ESI) calcd C₅₄H₇₆O₁₅S₂Si [M + Na]⁺ 1079.4287, found 1079.4294.

Octyl 6-O-benzyl-4-O-*t*-butyldimethylsilyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-toluenesulfonyl- α -D-mannopyranoside (23d). Chromatographic purification to give **23d** as a colorless syrup: *R*_f 0.37 (CH₂Cl₂–methanol 30:1); [α]_D = +27.5 (c 1.5, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.92 (d, *J* = 8.2 Hz, 2H, Ar), 7.76 (d, *J* = 8.2 Hz, 2H, Ar), 7.42 (d, *J* = 8.2 Hz, 2H, Ar), 7.39–7.23 (m, 12H, Ar), 7.20 (d, *J* = 8.1 Hz, 2H, Ar), 4.76 (d, *J* = 1.7 Hz, 1H, H-1), 4.72 (dd, *J* = 9.5, 3.1 Hz, 1H, H-3), 4.64 (dd, *J* = 3.3, 1.7 Hz, 1H, H-2'), 4.61 (d, *J* = 1.4 Hz, 1H, H-1'), 4.56–4.36 (m, 4H, 4 \times OCH₂Ph), 4.04 (br s, 1H, H-2), 3.90 (dd, *J* = 11.0, 3.6 Hz, 1H, H-6a), 3.76 (t, *J* = 9.6 Hz, 1H, H-4), 3.73–3.53 (m, 8H, H-3', H-4', H-5', H-6b, H-6a', octyl OCH₂), 3.38 (dt, *J* = 9.6, 6.8 Hz, 1H, octyl OCH₂), 2.48 (s, 3H, ArCH₃), 2.39 (s, 3H, ArCH₃), 2.17 (br d, *J* = 5.0 Hz, 1H, OH-2), 1.91 (d, *J* = 6.8 Hz, 1H, OH-3'), 1.64–1.53 (m, 2H, octyl OCH₂CH₂), 1.40–1.24 (m, 10H, octyl CH₂), 0.92 (t, *J* = 6.7 Hz, 3H, octyl CH₃), 0.86 (s, 9H, TBS(*t*-Bu)), 0.12 (s, 3H, TBS(Me)), 0.03 (s, 3H, TBS(Me)); ¹³C NMR (126 MHz, CDCl₃) δ 145.5, 145.1, 138.3, 138.2, 133.3, 132.8, 130.3, 129.9 (8C, Ar), 128.3 (3C, Ar), 128.3 (3C, Ar), 128.2 (2C, Ar), 128.0 (2C, Ar), 127.9 (2C, Ar), 127.6 (2C, Ar), 127.6 (Ar), 127.5 (Ar), 100.6 (C-1'), 99.2 (C-1), 81.0 (C-3), 79.1 (C-2'), 75.2 (C-4), 73.9 (C-5'), 73.3, 73.2 (2C, 2 \times OCH₂Ph), 71.2 (C-5), 69.6 (C-6'), 69.6, 69.5, 69.2 (3C, C-2, C-3', C-4'), 68.6 (C-6), 68.2 (octyl OCH₂), 31.8, 29.4, 29.3, 29.3, 26.1 (5C, octyl CH₂), 26.0 (TBS(*t*-Bu)), 22.7 (octyl CH₂), 21.8, 21.6 (2C, 2 \times ArCH₃), 18.3 (TBS(*t*-Bu)), 14.1 (octyl CH₃), –3.9, –5.0 (2C, 2 \times TBS(Me)); HRMS (ESI) calcd C₅₄H₇₆O₁₅S₂Si [M + Na]⁺ 1079.4287, found 1079.4291.

Octyl 6-O-benzyl-4-O-triphenylmethyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl- α -D-mannopyranoside (24). TrOTf was prepared in situ by adding a CH₂Cl₂ solution of TMSOTf (10% v/v, 1 mL, 0.53 mmol) into stirring ice-cold TfOH (137 mg, 0.53 mmol) in CH₂Cl₂ (4 mL). The bright yellow solution of TrOTf was formed after stirring at 0 °C for 5 min. This fresh-made TrOTf solution was slowly added into a stirring ice-cold solution of alcohol **10** (107 mg, 0.19

mmol) and 2,4,6-collidine (0.11 mL, 0.79 mmol) in CH₂Cl₂ (2 mL) over 0.5 h. Methanol (0.5 mL) was then added, and then the reaction mixture was concentrated, and the crude residue was purified by chromatography (hexane–EtOAc 8:1) to afford the expected trityl ether (150.4 mg, 98%) as a colorless syrup: *R*_f 0.69 (hexane–EtOAc 3:1); [α]_D = +96.8 (c 0.8, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.55–7.10 (m, 25H, Ar), 6.01 (d, *J* = 10.3 Hz, 1H, H-3), 5.83 (ddd, *J* = 10.3, 2.7, 1.8 Hz, 1H, H-2), 5.75 (d, *J* = 10.5 Hz, 1H, H-3'), 5.44 (ddd, *J* = 10.5, 2.7, 1.9 Hz, 1H, H-2'), 5.10 (d, *J* = 2.3 Hz, 1H, H-1'), 5.04 (d, *J* = 2.3 Hz, 1H, H-1), 4.69 (d, *J* = 12.2 Hz, 1H, OCH₂Ph), 4.61 (d, *J* = 12.2 Hz, 1H, OCH₂Ph), 4.38 (ddd, *J* = 9.5, 2.9, 1.6 Hz, 1H, H-4), 4.34 (s, 2H, OCH₂Ph), 4.16–4.08 (m, 2H, H-5, H-5'), 3.93–3.88 (m, 2H, H-4', H-6a), 3.86 (dt, *J* = 9.5, 6.9 Hz, 1H, octyl OCH₂), 3.77 (dd, *J* = 10.9, 6.4 Hz, 1H, H-6b), 3.58–3.47 (m, 3H, H-6a, H-6b, octyl OCH₂), 1.73–1.56 (m, 2H, octyl OCH₂CH₂), 1.45–1.19 (m, 10H, octyl CH₂), 0.92 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (126 MHz, CDCl₃) δ 144.8 (3C, Ar), 138.6 (Ar), 138.2 (Ar), 133.0 (C-3'), 129.5 (C-3), 129.0 (6C, Ar), 128.3 (2C, Ar), 128.2 (2C, Ar), 127.7 (6C, Ar), 127.6 (2C, Ar), 127.5 (2C, Ar), 127.4 (Ar), 127.4 (Ar), 127.2 (C-2), 127.2 (3C, Ar), 124.7 (C-2'), 94.2 (C-1), 91.8 (C-1'), 86.7 (Ph₃C), 73.4, 73.2 (2C, 2 \times OCH₂Ph), 70.5 (C-5'), 70.0, 69.3 (2C, C-6, C-6'), 69.2 (C-5), 68.7 (octyl OCH₂), 67.9 (C-4), 66.1 (C-4'), 31.9, 29.9, 29.5, 29.3, 26.3, 22.7 (6C, octyl CH₂), 14.2 (octyl CH₃); HRMS (ESI) calcd C₅₃H₆₀O₇ [M + Na]⁺ 831.4231, found 831.4223. The dihydroxylation of the trityl ether (165.3 mg, 0.20 mmol) was carried out with OsO₄ (2.5 wt% in *t*-butanol, 40 μ L, 0.004 mmol) and NMO (50% w/v in water, 0.3 mL) in *t*-butanol and acetone (2 mL, v/v 1:1). After stirring at room temperature overnight, the resulting saturated aqueous Na₂SO₃ solution was added. The mixture was concentrated to remove *t*-butanol, and then the residue was extracted with CH₂Cl₂ three times. The combined organic layer was concentrated, and the residue was purified by chromatography (CH₂Cl₂–methanol 30:1) to afford **24** (187.3 mg, 92%) as a white foam: *R*_f 0.26 (CH₂Cl₂–methanol 20:1); [α]_D = +33.5 (c 0.6, methanol); ¹H NMR (600 MHz, CDCl₃) δ 7.43–7.14 (m, 25H, Ar), 4.96 (d, *J* = 4.8 Hz, 1H, H-1'), 4.83 (d, *J* = 1.4 Hz, 1H, H-1), 4.63–4.58 (m, 2H, OCH₂Ph), 4.37 (d, *J* = 4.8 Hz, 1H, OH-3), 4.33 (d, *J* = 12.0 Hz, 1H, OCH₂Ph), 4.27 (d, *J* = 12.0 Hz, 1H, OCH₂Ph), 4.11–4.03 (m, 1H, H-3), 3.95 (app t, *J* = 9.8 Hz, 1H, H-4), 3.92–3.88 (m, 2H, H-5', H-2), 3.88–3.83 (m, 3H, H-5, H-2', H-3'), 3.81–3.77 (m, 2H, H-6a, H-6b), 3.69 (dt, *J* = 9.6, 6.8 Hz, 1H, octyl OCH₂), 3.50 (app t, *J* = 5.1 Hz, 1H, H-4'), 3.41 (dt, *J* = 9.6, 6.6 Hz, 1H, octyl OCH₂), 3.36 (dd, *J* = 10.3, 2.8 Hz, 1H, H-6a'), 3.16 (s, 1H, OH-3'), 3.06 (dd, *J* = 10.3, 7.5 Hz, 1H, H-6b'), 2.66 (d, *J* = 2.6 Hz, 1H, OH-2), 2.51 (d, *J* = 3.7 Hz, 1H, OH-2'), 1.59–1.53 (m, 2H, octyl OCH₂CH₂), 1.37–1.22 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.1 Hz, 3H, octyl CH₃); ¹³C NMR (126 MHz, CD₃OD) δ 146.0 (3C, Ar), 139.9 (Ar), 139.5 (Ar), 130.3 (3C, Ar), 129.4 (Ar), 129.3 (Ar), 128.87 (3C, Ar), 128.86 (3C, Ar), 128.6 (Ar), 128.5 (Ar), 128.4 (2C, Ar), 101.48, 101.47 (2C, C-1', ¹J_{C-1,H-1} = 169.0 Hz; C-1, ¹J_{C-1,H-1} = 167.6 Hz), 89.0 (Ph₃C), 77.5 (C-4), 76.2 (C-5'), 74.4 (OCH₂Ph), 73.9 (OCH₂Ph), 73.6, 72.9, 72.4, 72.3, 72.2 (5C, C-2, C-3, C-3', C-4', C-5), 72.0 (C-2'), 71.3 (C-6), 70.5 (C-6'), 68.7 (octyl OCH₂), 33.0, 30.6, 30.5, 30.4, 27.4, 23.8 (6C, octyl CH₂), 14.5 (octyl CH₃); HRMS (ESI) calcd C₅₃H₆₄O₁₁ [M + Na]⁺ 899.4341, found 899.4334.

Octyl 6-O-benzyl-2-O-toluenesulfonyl-4-O-triphenylmethyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-2-O-toluenesulfonyl- α -D-mannopyranoside (25a). To a solution of **24** (36.5 mg, 0.04 mmol) in THF (0.4 mL) was added *n*-Bu₂SnCl₂ (20 mol %, 2.5 mg, 0.008 mmol) and DIPEA (85 μ L, 0.48 mmol) followed by TsCl (80.9 mg, 0.42 mmol). The reaction mixture was stirred at room temperature for 24 h before being concentrated. The crude product was purified by chromatography (hexane–EtOAc 7:2) to give **25a** (45.0 mg, 91%) as a colorless film: *R*_f 0.43 (toluene–EtOAc 5:1); ¹H NMR (500 MHz, acetone) δ 7.91 (d, *J* = 8.3 Hz, 2H, Ar), 7.87 (d, *J* = 8.3 Hz, 2H, Ar), 7.58–7.24 (m, 29H), 5.32 (d, *J* = 5.4 Hz, 1H, H-1'), 4.92 (d, *J* = 1.7 Hz, 1H, H-1), 4.79 (dd, *J* = 5.4, 2.9 Hz, 1H, H-2'), 4.65 (dd, *J* = 3.5, 1.7 Hz, 1H, H-2), 4.63–4.55 (m, 2H, OCH₂Ph), 4.40–4.33 (m, 2H, OCH₂Ph), 4.10 (dd, *J* = 5.0, 2.9 Hz, 1H, H-3'), 4.02 (dd, *J* = 9.5, 3.4 Hz, 1H, H-3), 3.96 (dd, *J* = 9.8, 6.6 Hz, 1H, H-5'), 3.90 (dd, *J* = 11.0, 1.6 Hz, 1H, H-6a), 3.87–3.79 (m, 2H, H-4, octyl

OCH₂), 3.73 (dd, *J* = 10.9, 6.1 Hz, 1H, H-6b), 3.70–3.62 (m, 2H, H-5, H-4'), 3.55 (dt, *J* = 9.8, 6.7 Hz, 1H, octyl OCH₂), 3.41 (dd, *J* = 10.5, 3.0 Hz, 1H, H-6a'), 3.37 (dd, *J* = 10.8, 6.9 Hz, 1H, H-6b'), 2.53 (s, 3H, ArCH₃), 2.51 (s, 3H, ArCH₃), 1.78–1.67 (m, 2H, octyl OCH₂CH₂), 1.56–1.36 (m, 10H, octyl CH₂), 0.99 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (101 MHz, CDCl₃) δ 144.9, 144.9, 144.1, 138.6, 137.6, 133.5, 133.4, 130.0, 129.9, 129.7, 129.6, 129.0, 128.5, 128.4, 128.4, 128.3, 128.2, 128.0, 128.0, 128.0, 127.8, 127.7, 127.7, 127.5, 127.5, 127.4, 127.3.0 (42C, Ar), 97.3 (C-1), 97.0 (C-1'), 88.0 (Ph₃P), 79.4 (C-2), 79.3 (C-2'), 75.3 (C-4), 73.3 (OCH₂Ph), 73.2 (OCH₂Ph), 73.1 (C-5'), 71.6 (C-4'), 70.1 (C-5), 69.6 (C-6'), 69.5 (C-6), 69.4 (C-3'), 68.4 (C-3), 68.3 (octyl OCH₂), 31.9, 29.4, 29.3, 29.3, 26.1, 22.7 (6C, octyl CH₂), 21.7 (ArCH₃), 21.7 (ArCH₃), 14.2 (octyl CH₃) (this compound is not stable in CDCl₃; when running the ¹³C NMR spectrum, a minor byproduct appeared over time, because of decomposition); HRMS (ESI) calcd C₆₇H₇₆O₁₅S₂ [M + Na]⁺ 1207.4518, found 1207.4530.

Octyl 6-O-benzyl-4-O-triphenylmethyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl- α -D-mannopyranoside (26). The synthesis was performed as described for 24. Installation of the trityl group was carried out using 11 (72.5 mg, 0.09 mmol), 2,4,6-collidine (75 μ L, 0.56 mmol), TrOH (73.2 mg, 0.28 mmol) and TMSOTf (10% v/v in CH₂Cl₂, 0.5 mL, 0.28 mmol) in CH₂Cl₂ (2 mL). The crude residue was purified by chromatography (hexane–EtOAc 7:1) to afford trityl ether (100 mg, quantitative) as a pale yellow syrup: [α]_D = +78.6 (c 0.9, CH₂Cl₂); ¹H NMR (500 MHz, CD₂Cl₂) δ 7.47–7.10 (m, 30H, Ar), 6.02 (dd, *J* = 10.3, 3.8 Hz, 2H, H-3, H-3'), 5.80 (ddd, *J* = 10.2, 2.8, 1.8 Hz, 1H, H-2), 5.72 (ddd, *J* = 10.4, 2.7, 1.9 Hz, 1H, H-2'), 5.71 (d, *J* = 10.3 Hz, 1H, H-3), 5.45 (ddd, *J* = 10.4, 2.6, 1.9 Hz, 1H, H-2''), 5.21 (d, *J* = 2.3 Hz, 1H, H-1'), 5.05 (d, *J* = 2.4 Hz, 1H, H-1''), 4.97 (d, *J* = 2.5 Hz, 1H, H-1), 4.54–4.45 (m, 4H, 4 \times OCH₂Ph), 4.36–4.30 (m, 2H, H-4, H-4'), 4.27–4.22 (d, *J* = 11.9 Hz, 1H, OCH₂Ph), 4.24 (d, *J* = 11.9 Hz, 1H, OCH₂Ph), 4.11–4.07 (m, 1H, H-5''), 3.95 (ddd, *J* = 8.5, 6.4, 1.6 Hz, 1H, H-5), 3.88 (ddd, *J* = 7.6, 5.5, 1.6 Hz, 1H, H-5'), 3.84–3.72 (m, 4H, H-4'', H-6a, H-6a', octyl OCH₂), 3.70–3.63 (m, 2H, H-6b, H-6b'), 3.54–3.48 (m, 2H, H-6a'', H-6b''), 3.46 (dt, *J* = 9.5, 6.6 Hz, 1H, octyl OCH₂), 1.64–1.48 (m, 2H, octyl OCH₂CH₂), 1.38–1.21 (m, 10H, octyl CH₂), 0.88 (t, *J* = 7.0 Hz, 1H, octyl CH₃); ¹³C NMR (126 MHz, CD₂Cl₂) δ 145.3 (3C; Tr), 139.3, 139.1, 138.8 (3C; Ar), 133.0 (C-3''), 130.5 (C-3'), 129.5 (C-3), 129.3, 128.6, 128.6, 128.5, 128.1, 127.9, 127.9, 127.8, 127.7, 127.7, 127.6 (31C; 30 \times Ar, C-2), 127.0 (C-2'), 125.3 (C-2''), 94.5 (C-1), 92.1 (C-1''), 91.7 (C-1'), 87.0 (Tr), 73.7, 73.5, 73.4 (3C; 3 \times OCH₂Ph), 71.0 (C-5''), 70.7 (C-6), 70.3 (C-5'), 70.3 (C-6), 69.8 (C-6), 69.6 (C-5), 68.9 (octyl OCH₂), 68.1 (C-4'), 67.8 (C-4), 66.5 (C-4''), 32.2, 30.2, 29.8, 29.7, 26.65, 23.1 (6C; octyl CH₂), 14.3 (octyl CH₃); HRMS (ESI) calcd C₆₆H₇₄O₁₀ [M + Na]⁺ 1049.5174, found 1049.5168. The dihydroxylation of the trityl ether was carried out with OsO₄ (2.5 wt.% in *t*-butanol, 75 μ L, 0.006 mmol) and NMO (50% w/v in water, 0.3 mL) in mixed *t*-butanol and acetone (3 mL, v/v 1:1). The crude residue was purified by chromatography (CH₂Cl₂–MeOH 15:1) to afford trisaccharide 26 (79.5 mg, 76%) as a white foam: *R*_f 0.26 (CH₂Cl₂–methanol 15:1); [α]_D = +63.2 (c 0.1, CH₂Cl₂); ¹H NMR (500 MHz, CD₃OD) δ 7.46–7.11 (m, 30H, Ar), 5.27 (d, *J* = 1.6 Hz, 1H, H-1'), 5.03 (d, *J* = 5.7 Hz, 1H, H-1''), 4.71 (d, *J* = 1.3 Hz, 1H, H-1), 4.58–4.17 (m, 6H, 6 \times OCH₂Ph), 3.98–3.91 (m, 2H, H-2', H-2''), 3.91–3.77 (m, 9H, H-3, H-3', H-3'', H-4, H-4', H-5, H-5', H-5'', H-6), 3.76–3.63 (m, 5H, H-2, 3 \times H-6, octyl OCH₂), 3.55 (app t, *J* = 4.9 Hz, 1H, H-4'), 3.44–3.37 (m, 2H, H-6a'', octyl OCH₂), 3.30–3.24 (m, 1H, H-6b''), 1.63–1.48 (m, 2H, octyl OCH₂CH₂), 1.43–1.18 (m, 10H, octyl CH₂), 0.90 (t, *J* = 6.9 Hz, 3H, octyl CH₃); ¹³C NMR (126 MHz, CD₃OD) δ 146.0 (3C, Ar), 139.9, 139.6, 139.5, 130.4, 129.5, 129.4, 129.3, 129.1, 129.0, 128.9, 128.87, 128.7, 128.6, 128.5, 128.4 (33C, Ar), 102.9 (C-1'), 101.7 (C-1''), 101.5 (C-1), 89.1 (Ph₃C), 77.5 (C-4), 76.2, 76.0 (2 \times C-5), 74.5, 74.4, 73.9 (3C, 3 \times OCH₂Ph), 73.6, 73.4, 73.0, 72.8, 72.7, 72.4, 72.3, 72.1, 72.0 (9C, 3 \times C-2, 3 \times C-3, 2 \times C-4, C-5), 71.3 (C-6'), 70.6 (C-6/C-6'), 68.7 (octyl OCH₂), 67.4 (C-6'/C-6), 33.0, 30.6, 30.5, 30.5, 27.4, 23.8 (6C, octyl CH₂), 14.6 (octyl CH₃); HRMS (ESI) calcd C₆₆H₈₀O₁₆ [M + Na]⁺ 1151.5339, found 1151.5333.

Octyl 6-O-benzyl-2-O-toluenesulfonyl-4-O-triphenylmethyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-2-O-toluenesulfonyl- α -D-mannopyranoside (26a). To a solution of 26 (29.3 mg, 0.026 mmol) in THF (0.3 mL) was added *n*-Bu₂SnCl₂ (30 mol %, 2.0 mg, 0.008 mmol) and DIPEA (83 μ L, 0.47 mmol) followed by TsCl (75.2 mg, 0.39 mmol). The reaction mixture was stirred at room temperature for 24 h before being concentrated. Chromatographic purification (hexane–EtOAc 5:2) gave 26a (28.3 mg, 69%) as a colorless film: [α]_D = +36.2 (c 0.1, CH₂Cl₂); ¹H NMR (500 MHz, CD₂Cl₂) δ 7.83–7.62 (m, 6H, Ar), 7.37–7.08 (m, 36H, Ar), 5.13 (d, *J* = 1.7 Hz, 1H, H-1'), 5.09 (d, *J* = 3.9 Hz, 1H, H-1''), 4.75 (d, *J* = 1.7 Hz, 1H, H-1), 4.69 (dd, *J* = 3.8, 3.2 Hz, 1H, H-2''), 4.64–4.60 (m, 2H, H-2, H-2'), 4.55–4.39 (m, 4H, 2 \times OCH₂Ph), 4.23 (s, 2H, OCH₂Ph), 4.06 (app dt, *J* = 8.4, 3.4 Hz, 1H, H-3), 3.99–3.91 (m, 2H, H-3', H-5''), 3.86 (ddd, *J* = 9.9, 6.1, 1.6 Hz, 1H, H-5'), 3.78–3.59 (m, 8H, H-3'', H-4, H-4', H-5, H-6a, H-6b, H-6a', octyl OCH₂), 3.56 (dd, *J* = 10.7, 6.2 Hz, 1H, H-6b'), 3.43–3.34 (m, 2H, H-4'', octyl OCH₂), 3.33 (dd, *J* = 10.7, 2.2 Hz, 1H, H-6a''), 3.00 (dd, *J* = 10.4, 6.2 Hz, 1H, H-6b''), 2.96 (d, *J* = 7.2 Hz, 1H, OH-3'), 2.71 (d, *J* = 8.0 Hz, 1H, OH-3), 2.42 (s, 3H, ArCH₃), 2.40 (s, 3H, ArCH₃), 2.40 (s, 3H, ArCH₃), 1.59–1.49 (m, 2H, octyl OCH₂CH₂), 1.35–1.22 (m, 10H, octyl CH₂), 0.89 (t, *J* = 6.9 Hz, 1H, octyl CH₃); ¹³C NMR (126 MHz, CD₂Cl₂) δ 145.8, 145.8, 145.7, 144.6, 138.9, 138.6, 138.1, 133.7, 133.5, 133.4, 130.3, 130.2, 130.1, 129.3, 128.7, 128.6, 128.4, 128.4, 128.3, 128.2, 128.1, 128.0, 128.0, 127.9, 127.8 (54C, Ar), 97.8, 97.6, 97.5 (3C, C-1, C-1', C-1''), 88.3 (Tr), 80.0, 79.9, 79.8 (3C, C-2, C-2', C-2''), 75.9 (C-4), 75.2 (C-4'), 73.7 (OCH₂Ph), 73.6 (OCH₂Ph), 73.6 (C-5''), 73.5 (OCH₂Ph), 72.1 (C-4''), 71.3 (C-5'), 70.2 (C-5), 69.9 (C-6''), 69.8 (C-3''), 69.7 (C-6'), 69.4 (C-6), 68.9 (C-3'), 68.8 (C-3), 68.6 (octyl OCH₂), 32.2, 29.7, 29.6, 29.6, 26.4, 23.0 (6C, octyl CH₂), 21.79 (2C, 2 \times ArCH₃), 21.78 (ArCH₃), 14.3 (octyl CH₃); HRMS (ESI) calcd C₈₇H₉₈O₂₂S₃ [M + Na]⁺ 1613.5604, found 1613.5602.

Octyl 6-O-benzyl-4-O-triphenylmethyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl- α -D-mannopyranoside (27). Synthesis was performed as described for 24. Installation of the trityl group was carried out with 28 (24 mg, 0.02 mmol), 2,4,6-collidine (25 μ L, 0.19 mmol), TrOH (23.4 mg, 0.09 mmol) and TMSOTf (10% v/v in CH₂Cl₂, 160 μ L, 0.09 mmol) in CH₂Cl₂ (0.6 mL). The crude residue from CH₂Cl₂ extraction was used for the next step without further purification. The dihydroxylation of the trityl ether was carried out with OsO₄ (2.5 wt.% in *t*-butanol, 30 μ L, 0.002 mmol) and NMO (50% w/v in water, 0.12 mL) in mixed *t*-butanol and acetone (1 mL, v/v 1:1). The crude residue was purified by chromatography (CH₂Cl₂–MeOH 12:1) to afford tetrasaccharide 27 (29.7 mg, 90% over two steps) as a white foam: *R*_f 0.40 (CH₂Cl₂–methanol 9:1); [α]_D = +63.2 (c 0.2, methanol); ¹H NMR (600 MHz, CD₃OD) δ 7.45–7.13 (m, 35H, Ar), 5.26 (d, *J* = 1.8 Hz, 1H, H-1'/H-1''), 5.25 (d, *J* = 1.8 Hz, 1H, H-1'/H-1''), 5.04 (d, *J* = 5.5 Hz, 1H, H-1''), 4.72 (d, *J* = 1.7 Hz, 1H, H-1), 4.58–4.18 (m, 8H, 8 \times OCH₂Ph), 3.99 (dd, *J* = 5.5, 3.1 Hz, 1H, H-2''), 3.96 (dd, *J* = 3.2, 2.0 Hz, 1H, H-2'/H-2''), 3.95–3.90 (m, 2H, H-5''), H-5'/H-5''), 3.89–3.75 (m, 11H, H-2'/H-2'', 4 \times H-3, 3 \times H-4, H-5, 2 \times H-6), 3.75–3.67 (m, 5H, H-2, H-5'/H-5'', 2 \times H-6, octyl OCH₂), 3.58 (d, *J* = 3.4 Hz, 2H, 2 \times H-6), 3.54 (app t, *J* = 5.3 Hz, 1H, H-4''), 3.41 (dt, *J* = 9.7, 6.3 Hz, 1H, octyl OCH₂), 3.38 (dd, *J* = 11.3, 8.2 Hz, 1H, H-6''), 3.32–3.30 (m, 1H, H-6''), 1.64–1.54 (m, 2H, octyl OCH₂CH₂), 1.41–1.23 (m, 10H, octyl CH₂), 0.90 (t, *J* = 7.1 Hz, 1H, octyl CH₃); ¹³C NMR (126 MHz, CD₃OD) δ 146.0, 139.8, 139.7, 139.6, 139.5, 130.4, 129.5, 129.4, 129.4, 129.3, 129.3, 129.1, 128.9, 128.8, 128.7, 128.6, 128.5, 128.4 (42C, Ar), 103.0, 102.9 (2C, C-1', C-1'', average ¹J_{C-1,H-1} = 174.6 Hz), 102.1 (C-1'''), ¹J_{C-1,H-1} = 169.4 Hz), 101.5 (C-1, ¹J_{C-1,H-1} = 169.3 Hz), 89.1 (Ph₃P), 77.9 (C-4), 76.0, 76.0, 75.9 (3C), 74.5 (2C, 2 \times OCH₂Ph), 74.3 (OCH₂Ph), 74.0 (OCH₂Ph), 73.6, 73.4, 73.1, 73.0, 72.8, 72.7, 72.7, 72.4, 72.3, 72.2, 72.0 (15C), 71.4, 71.4, 71.2, 70.8 (4C, 4 \times C-6), 68.7 (octyl OCH₂), 33.0, 30.6, 30.4, 30.4, 27.4, 23.8 (6C, octyl CH₂), 14.6 (octyl CH₃); HRMS (ESI) calcd C₇₉H₉₆O₂₁ [M + Na]⁺ 1403.6336, found 1403.6340.

Octyl 6-O-benzyl-2-O-toluenesulfonyl-4-O-triphenylmethyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-2-O-toluenesulfonyl- α -D-mannopyranoside (27a). To a solution of 27 (21.1 mg, 0.015 mmol) in THF (0.3 mL) was added *n*-Bu₂SnCl₂ (40 mol %, 2.0 mg, 0.006 mmol) and DIPEA (65 μ L, 0.36 mmol) followed by TsCl (56.2 mg, 0.30 mmol). The reaction mixture was stirred at room temperature for 22 h before being concentrated. Chromatographic purification (hexane–EtOAc 3:2) gave 27a (11.7 mg, 38%) as a colorless film: $[\alpha]_D^{25} = +35.8$ (c 0.6, methanol); ¹H NMR (600 MHz, CD₃OD) δ 7.88–7.72 (m, 8H, Ar), 7.45–7.07 (m, 43H, Ar), 5.20 (d, *J* = 5.5 Hz, 1H, H-1''), 5.01 (d, *J* = 1.9 Hz, 1H, H-1'), 4.95 (d, *J* = 1.7 Hz, 1H, H-1''), 4.78 (dd, *J* = 3.2, 2.1 Hz, 1H, H-2'), 4.73 (d, *J* = 1.8 Hz, 1H, H-1), 4.71–4.67 (m, 2H, H-2'', H-2'''), 4.53 (dd, *J* = 3.4, 1.8 Hz, 1H, H-2), 4.46–4.30 (m, 4H, 2 \times OCH₂Ph), 4.21 (d, *J* = 12.0 Hz, 1H, OCH₂Ph), 4.16 (d, *J* = 12.0 Hz, 1H, OCH₂Ph), 3.91 (dd, *J* = 9.8, 2.9 Hz, 1H, H-3'), 3.89 (dd, *J* = 9.5, 3.0 Hz, 1H, H-3), 3.79 (brs, 1H, H-3'''), 3.76 (dd, *J* = 9.3, 3.2 Hz, 1H, H-3''), 3.75–3.72 (m, 1H, H-5'''), 3.72–3.68 (m, 1H, H-5'), 3.68 (app t, *J* = 9.5 Hz, 1H, H-4''), 3.65–3.56 (m, 7H, H-4, H-4', H-5, H-6a, H-6b, H-6a'', octyl OCH₂), 3.57–3.45 (m, 5H, H-4'', H-5'', H-6b'', H-6a'', H-6b''), 3.35 (dt, *J* = 9.9, 6.3 Hz, 1H, octyl OCH₂), 3.27–3.21 (m, 1H, H-6a'''), 3.20–3.14 (m, 1H, H-6b'''), 2.43 (s, 3H, ArCH₃), 2.39 (s, 3H, ArCH₃), 2.36 (s, 3H, ArCH₃), 2.34 (s, 3H, ArCH₃), 1.55–1.46 (m, 2H, octyl OCH₂CH₂), 1.35–1.17 (m, 10H, octyl CH₂), 0.86 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (125 MHz, CD₃OD) δ 146.6, 146.5, 146.3, 145.6, 139.8, 139.7, 139.7, 139.4, 135.2, 135.1, 135.0, 131.1, 131.0, 131.0, 130.9, 130.3, 129.5, 129.4, 129.4, 129.3, 129.3, 129.3, 129.0, 128.9, 128.9, 128.8, 128.8, 128.7, 128.6, 128.5, 128.5 (66C, Ar), 100.2 (C-1'), 100.0 (C-1''), 98.5 (C-1), 97.4 (C-1'''), (octyl OCH₂), 33.0, 30.4, 30.4, 30.89.2 (Ph₃C), 81.4, 81.23, 81.20, 81.17 (4C, C-2, C-2', C-2'', C-2'''), 76.72 (C-4), 76.68 (C-4'), 75.7 (C-5'''), 74.3 (3 \times OCH₂Ph), 74.0 (OCH₂Ph), 73.1 (C-5'), 73.0 (C-4''), 72.3 (C-5''), 71.0 (C-3'''), 70.9 (C-6''), 70.5 (C-6), 70.4 (2C, C-6', C-6'''), 70.2 (C-3), 69.9 (C-3''), 69.7 (C-3'), 69.3 4, 27.2, 23.7 (6C, octyl CH₂), 22.1 (ArCH₃), 21.7 (ArCH₃), 21.6 (2C, 2 \times ArCH₃), 14.5 (octyl CH₃); HRMS (ESI) calcd C₁₀₇H₁₂₀O₂₉S₄ [M + Na]⁺ 2019.6690, found 2019.6684.

(2R,3S,6S)-2-((benzyloxy)methyl)-6-(((2R,3S,6S)-2-((benzyloxy)methyl)-6-(((2R,3S,6S)-2-((benzyloxy)methyl)-6-((octyloxy)-3,6-dihydro-2H-pyran-3-yl)oxy)-3,6-dihydro-2H-pyran-3-yl)oxy)-3,6-dihydro-2H-pyran-3-yl)oxy)-3,6-dihydro-2H-pyran-3-yl)oxy)-3,6-dihydro-2H-pyran-3-ol (28). The coupling step was performed as described for the synthesis of 8, with alcohol 11 (219.2 mg, 0.28 mmol) and donor 7 (184.3 mg, 0.55 mmol) in the presence of Pd₂(dba)₃ (12.9 mg, 0.01 mmol) and PPh₃ (16.0 mg, 0.06 mmol) in CH₂Cl₂ (5 mL). The crude residue was purified by chromatography (hexane–EtOAc 3.5:1) to afford a ketone (208.9 mg, 82%) as pale yellow syrup: $[\alpha]_D^{25} = +7.7$ (c 2.0, CHCl₃); ¹H NMR (600 MHz, CDCl₃) δ 7.36–7.18 (m, 20H, Ar), 6.82 (dd, *J* = 10.2, 3.5 Hz, 1H, H-2''), 6.17 (d, *J* = 10.3 Hz, 1H, H-3'''), 6.12 (d, *J* = 10.3 Hz, 1H, H-3''), 6.06 (d, *J* = 10.4 Hz, 1H, H-3'), 6.03 (d, *J* = 10.4 Hz, 1H, H-3), 5.82 (ddd, *J* = 10.3, 2.6, 1.9 Hz, 1H, H-2), 5.78 (ddd, *J* = 10.3, 2.6, 2.0 Hz, 1H, H-2''), 5.73 (ddd, *J* = 10.3, 2.6, 2.0 Hz, 1H, H-2'), 5.56 (d, *J* = 3.5 Hz, 1H, H-1'''), 5.25 (d, *J* = 2.3 Hz, 1H, H-1''), 5.23 (d, *J* = 2.3 Hz, 1H, H-1'), 5.00 (d, *J* = 2.3 Hz, 1H, H-1), 4.64 (dd, *J* = 9.3, 1.3 Hz, 1H, H-4''), 4.46 (dd, *J* = 9.3, 1.3 Hz, 1H, H-4'), 4.56–4.32 (m, 8H, 8 \times OCH₂Ph), 4.38 (dd, *J* = 9.4, 1.2 Hz, 1H, H-4), 4.01 (ddd, *J* = 9.1, 6.2, 1.7 Hz, 1H, H-5), 3.85 (ddd, *J* = 9.3, 3.4, 3.4 Hz, 1H, H-5'), 3.82–3.76 (m, 4H, H-5'', H-6a, H-6a'', octyl OCH₂), 3.68 (dd, *J* = 11.0, 6.2 Hz, 1H, H-6b), 3.63–3.55 (m, 4H, H-6a', H-6b', H-6a'', H-6b'''), 3.50 (dd, *J* = 10.7, 1.8 Hz, 1H, H-6b''), 3.48 (dt, *J* = 9.5, 6.6 Hz, 1H, octyl OCH₂), 1.64–1.54 (m, 2H, octyl OCH₂CH₂), 1.38–1.20 (m, 10H, octyl CH₂), 0.88 (t, *J* = 7.1 Hz, 3H, octyl CH₃); ¹³C NMR (151 MHz, CDCl₃) δ 194.0 (C=O), 143.6 (C-2''), 138.6, 138.3, 137.9, 137.8 (4C, Ar), 129.8, 129.3, 129.3 (3C; C-3, C-3', C-3''), 128.3, 128.3, 128.3, 128.2, 127.6, 127.6, 127.6, 127.4, 127.4, 127.3, 127.2, 126.8 (24C; 20 \times Ar, C-2, C-2', C-2'', C-3''), 94.1 (C-1), 91.3, 91.3 (2C; C-1', C-1''), 90.0 (C-1'''), 75.0 (C-5'''), 73.7, 73.5, 73.4, 73.1 (4C; 4 \times OCH₂Ph), 70.0 (C-6), 69.8, 69.7 (2C; C-5', C-5''), 69.4

(C-6'), 69.1 (C-5), 68.9 (C-6''), 68.7 (octyl OCH₂), 68.4 (C-6'''), 67.6 (C-4), 67.2 (C-4''), 67.0 (C-4'), 31.8, 29.8, 29.4, 29.3, 26.2, 22.7 (octyl CH₂), 14.1 (octyl CH₃); HRMS (ESI) calcd C₆₀H₇₂O₁₃ [M + Na]⁺ 1023.4865, found 1023.4856. This ketone (208.9 mg, 0.21 mmol) was then reduced as described for 9, with NaBH₄ (8.0 mg, 0.21 mmol) and CeCl₃·7H₂O (80.2 mg, 0.21 mmol) in MeOH (2.5 mL). Chromatography purification (hexane–EtOAc 3:1) furnished alcohol 28 (177.2 mg, 85%) as colorless syrup: *R*_f 0.37 (hexane–EtOAc 2:1); $[\alpha]_D^{25} = +32.2$ (c 1.0, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.44–7.17 (m, 20H, Ar), 6.12–6.03 (m, 3H, H-3, H-3', H-3''), 6.00 (d, *J* = 10.2 Hz, 1H, H-3'''), 5.84 (ddd, *J* = 10.3, 2.7, 1.8 Hz, 1H, H-2), 5.79–5.69 (m, 3H, H-2', H-2'', H-2'''), 5.29–5.24 (m, 2H, H-1', H-1''), 5.22 (d, *J* = 2.6 Hz, 1H, H-1'''), 5.02 (d, *J* = 2.2 Hz, 1H, H-1), 4.58–4.36 (m, 11H, 4 \times OCH₂Ph, H-4, H-4', H-4''), 4.28 (dddd, *J* = 6.2, 3.6, 1.8, 1.8 Hz, 1H, H-4'''), 4.03 (ddd, *J* = 9.3, 6.3, 1.7 Hz, 1H, H-5), 3.91–3.78 (m, 4H, H-5', H-5'', H-6a, octyl OCH₂), 3.76–3.56 (m, 7H, H-5'', H-6b, H-6a', H-6b', H-6a'', H-6b'''), 3.55–3.46 (m, 2H, H-6b''', octyl OCH₂), 2.46 (d, *J* = 4.5 Hz, 1H, OH-4''), 1.66–1.56 (m, 2H, octyl OCH₂CH₂), 1.40–1.21 (m, 10H, octyl CH₂), 0.90 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (126 MHz, CDCl₃) δ 138.6, 138.4, 138.4, 137.6 (4C; Ar), 133.3, 130.0, 129.9, 129.3, 128.5, 128.3, 128.2, 128.0, 127.8, 127.5, 127.4, 127.4, 127.3, 127.3 (20C; Ar), 126.7 (C-2), 126.6 (2C; C-2', C-2''), 125.7 (C-2''), 94.1 (C-1), 91.3, 91.2 (C-1', C-1''), 91.0 (C-1'''), 73.7, 73.4, 73.4, 73.1 (4C; 4 \times OCH₂Ph), 70.7 (C-6), 70.1 (C-6), 69.8 (3C; C-5', C-5'', C-5'''), 69.5 (C-6), 69.2 (C-6), 69.1 (C-5), 68.7 (octyl OCH₂), 67.6 (C-4), 67.0, 66.6 (2C; C-4', C-4''), 66.0 (C-4'''), 31.9, 29.8, 29.4, 29.3, 26.3, 22.7 (6C; octyl CH₂), 14.1 (octyl CH₃); HRMS (ESI) calcd C₆₀H₇₄O₁₃ [M + Na]⁺ 1025.5022, found 1025.5027.

Synthesis of Octyl 6-O-benzyl-4-O-*t*-butyldimethylsilyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranoside (29) and Octyl 6-O-benzyl-3-O-*t*-butyldimethylsilyl-4-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranoside (30). To a stirring ice-cold solution of alcohol 23a (9.3 mg, 0.01 mmol) and MeI (10 μ L, 0.16 mmol) in DMF (0.7 mL) was added NaH (60% in mineral oil, 6.3 mg, 0.15 mmol). The resulting solution was stirred at 0 °C for 15 min before methanol (0.1 mL) was added. The resulting reaction mixture was diluted with CH₂Cl₂ and washed with brine. The separated organic layer was dried over Na₂SO₄ and concentrated, and the resulting residue was purified by chromatography (hexane–EtOAc 8:1) to afford an inseparable mixture of 29 and 30 (8.7 mg, 91%, 29/30 = 1.8:1) as a colorless syrups. *R*_f 0.63 (hexane–EtOAc 4:1). Distinguishing between 29 and 30 was assisted by both ¹H–¹H COSY and 1D-TOCSY NMR spectroscopy.

Octyl 6-O-benzyl-4-O-*t*-butyldimethylsilyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranoside (29). ¹H NMR (600 MHz, CDCl₃) δ 7.87–7.78 (m, 4H, Ar), 7.36–7.21 (m, 14H, Ar), 5.00 (d, *J* = 2.1 Hz, 1H, H-1'), 4.95 (dd, *J* = 2.9, 2.2 Hz, 1H, H-2'), 4.91 (d, *J* = 1.9 Hz, 1H, H-1), 4.79 (dd, *J* = 3.2, 1.9 Hz, 1H, H-2), 4.55–4.38 (m, 4H, 4 \times OCH₂Ph), 3.75–3.53 (m, 9H); 3.51 (dd, *J* = 9.4, 3.3 Hz, 1H, H-3), 3.41–3.38 (m, 1H, octyl OCH₂); 3.25 (dd, *J* = 8.7, 3.2 Hz, 1H, H-3'), 3.19 (s, 3H, OMe), 3.14 (s, 3H, OMe), 2.44 (s, 3H, ArCH₃), 2.40 (s, 3H, ArCH₃), 1.60–1.53 (m, 2H, octyl OCH₂CH₂), 1.34–1.24 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.0 Hz, 3H, octyl CH₃), 0.80 (s, 9H, TBS(*t*-Bu)), –0.02 (s, 3H, TBS(Me)), –0.06 (s, 3H, TBS(Me)); HRMS (ESI) calcd C₅₆H₈₀O₁₅S₂Si [M + Na]⁺ 1107.4600, found 1107.4594.

Octyl 6-O-benzyl-3-O-*t*-butyldimethylsilyl-4-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranoside (30). ¹H NMR (600 MHz, CDCl₃) δ 7.87–7.78 (m, 4H, Ar), 7.36–7.21 (m, 14H, Ar), 4.85 (br d, *J* = 1.9 Hz, 2H, H-1, H-1'), 4.78–4.76 (m, 2H, H-2, H-2'), 4.55–4.38 (m, 4H, 4 \times OCH₂Ph), 3.94 (dd, *J* = 8.9, 2.7 Hz, 1H, H-3'), 3.75–3.53 (m, 5H); 3.47–3.42 (m, 2H, H-3, H-6a'); 3.41–3.38 (m, 1H, octyl OCH₂); 3.39 (s, 3H, OMe-4'), 3.35 (app t, *J* = 8.7 Hz, 1H, H-4'), 3.20 (s, 3H, OMe-3), 2.43 (s, 3H, ArCH₃), 2.40 (s, 3H, ArCH₃), 1.60–1.53 (m, 2H, octyl OCH₂CH₂), 1.34–1.24 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.0 Hz, 3H, octyl CH₃), 0.94 (s, 9H,

TBS(*t*-Bu)), 0.13 (s, 3H, TBS(Me)), 0.12 (s, 3H, TBS(Me)); HRMS (ESI) calcd C₅₆H₈₀O₁₅S₂Si [M + Na]⁺ 1107.4600, found 1107.4594.

Octyl 6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranoside (32). To a stirring ice-cold solution of alcohol 25a (56.3 mg, 0.05 mmol) and MeI (24 μ L, 0.38 mmol) in THF (0.6 mL) was added NaH (60% in mineral oil, 24 mg, 0.6 mmol). The resulting solution was stirred at 0 °C for 20 min before methanol (1 mL) was added. Then a methanolic solution of HCl (0.1 mL, 10% v/v) was added, and the reaction mixture was stirred for 0.5 h. The yellowish solution was concentrated and redissolved in CH₂Cl₂, washed with saturated aqueous NaHCO₃, followed by saturated Na₂SO₃ and brine. The separated organic layer was dried over Na₂SO₄ and concentrated, and the resulting residue was purified by chromatography (hexane–EtOAc 2.5:1) to afford 32 (38.2 mg, 83% over two steps) as a colorless syrup; *R*_f 0.31 (hexane–EtOAc 2:1); [α]_D = +6.2 (c 0.4, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.88 (d, *J* = 8.3 Hz, 2H, Ar), 7.82 (d, *J* = 8.3 Hz, 2H, Ar), 7.38–7.23 (m, 14H, Ar), 5.15 (d, *J* = 1.7 Hz, 1H, H-1'), 4.96–4.92 (m, 2H, H-1, H-2'), 4.81 (dd, *J* = 3.0, 2.0 Hz, 1H, H-2), 4.54–4.44 (m, 4H, 4 \times OCH₂Ph), 3.82–3.64 (m, 7H, H-4', H-4, H-5, H-5', H-6, H-6, octyl OCH₂), 3.62–3.52 (m, 3H, H-3, H-6, H-6), 3.42 (dt, *J* = 9.7, 6.7 Hz, 1H, octyl OCH₂), 3.39 (dd, *J* = 9.5, 3.0 Hz, 1H, H-3'), 3.25 (s, 3H, OMe), 3.21 (s, 3H, OMe), 2.63 (brs, 1H, OH-4'), 2.46 (s, 3H, ArCH₃), 2.44 (s, 3H, ArCH₃), 1.65–1.54 (m, 2H, octyl OCH₂CH₂), 1.37–1.25 (m, 10H, octyl CH₂), 0.91 (t, *J* = 6.9 Hz, 3H, octyl CH₃); ¹³C NMR (126 MHz, CDCl₃) δ 144.9, 144.8, 138.4, 138.0, 134.0, 133.7, 129.8, 129.7, 128.4, 128.3, 128.1, 127.9, 127.7, 127.5, 127.4 (24C, Ar), 99.6 (C-1'), 97.3 (C-1), 79.4 (C-3), 78.4 (C-3'), 74.2 (C-2), 73.9 (C-2'), 73.6 (OCH₂Ph), 73.5 (C-4), 73.4 (OCH₂Ph), 72.0 (C-5), 71.0 (C-5'), 70.2, 69.4 (2C, C-6, C-6'), 68.3 (octyl OCH₂), 67.4 (C-4'), 57.1, 56.8 (2C, 2 \times OMe), 31.8, 29.4, 29.3, 29.2, 26.1, 22.7 (6C, octyl CH₂), 21.7 (ArCH₃), 21.6 (ArCH₃), 14.1 (octyl CH₃); HRMS (ESI) calcd C₅₀H₆₆O₁₅S₂ [M + Na]⁺ 993.3735, found 993.3736.

Octyl 3-O-methyl- α -D-mannopyranosyl-(1 \rightarrow 4)-3-O-methyl- α -D-mannopyranoside (33). A solution of 32 (25.5 mg, 0.03 mmol) and Mg (39.9 mg, 1.66 mmol) was heated in dry methanol (1 mL) at reflux overnight. The solution was cooled to room temperature, and then 1 M HCl (5 mL) was added. The resulting mixture was extracted with CH₂Cl₂ and washed with brine. The organic layer was then concentrated and purified by chromatography (CH₂Cl₂–methanol 30:1) to afford a partially deprotected disaccharide (12.5 mg, 72%) as a colorless syrup; *R*_f 0.30 (CH₂Cl₂–methanol 20:1); ¹H NMR (500 MHz, CD₃OD) δ 7.36–7.11 (m, 10H, Ar), 5.15 (d, *J* = 1.6 Hz, 1H, H-1'), 4.75 (d, *J* = 1.5 Hz, 1H, H-1), 4.54–4.38 (m, 4H, 4 \times OCH₂Ph), 4.08 (dd, *J* = 2.9, 2.1 Hz, 1H, H-2'), 4.00 (dd, *J* = 3.1, 1.9 Hz, 1H, H-2), 3.85–3.78 (m, 2H, H-4, H-6), 3.78–3.65 (m, 6H, H-4', H-5, H-5', H-6, H-6, octyl OCH₂), 3.62 (dd, *J* = 10.7, 5.5 Hz, 1H, H-6), 3.48–3.44 (m, 1H, H-3), 3.45 (s, 3H, OMe), 3.44–3.39 (m, 1H, octyl OCH₂), 3.41 (s, 3H, OMe), 3.33 (dd, *J* = 9.1, 3.0 Hz, 1H, H-3'), 1.64–1.55 (m, 2H, octyl OCH₂CH₂), 1.41–1.24 (m, 10H, octyl CH₂), 0.89 (t, *J* = 6.9 Hz, 3H, octyl OCH₃); ¹³C NMR (126 MHz, CD₃OD) δ 139.8 (Ar), 139.7 (Ar), 129.34 (2C, Ar), 129.30 (2C, Ar), 129.0 (2C, Ar), 128.9 (2C, Ar), 128.6 (Ar), 128.6 (Ar), 103.7 (C-1'), 101.4 (C-1), 83.3 (C-3), 82.2 (C-3'), 75.3 (C-4), 74.6 (C-5), 74.6 (OCH₂Ph), 74.4 (OCH₂Ph), 72.3 (C-5'), 71.4, 71.3 (2C, C-6, C-6'), 68.9 (octyl OCH₂), 68.1 (C-2'), 67.7 (C-2), 67.5 (C-4'), 57.3, 56.7 (2C, 2 \times OMe), 33.0, 30.5, 30.4, 30.4, 27.4, 23.7 (6C, octyl CH₂), 14.5 (octyl CH₃); HRMS (ESI) calcd C₃₆H₅₄O₁₁ [M + Na]⁺ 685.3558, found 685.3551. Hydrogenolysis of this disaccharide (9.8 mg, 0.01 mmol) was performed in with Pd–C (5 wt.%, 10 mg) in methanol (0.5 mL) under H₂ atmosphere for overnight. The catalyst was removed by filtration through Celite. The filtrate was then concentrated, and the residue was purified by chromatography (CH₂Cl₂–methanol 9:1) to afford 33 (6.9 mg, 97%) as a colorless syrup; *R*_f 0.11 (CH₂Cl₂–methanol 9:1); [α]_D = +82.5 (c 0.04, methanol); the NMR spectra were identical to those previously reported.¹⁴

Octyl 6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-methyl-2-

O-toluenesulfonyl- α -D-mannopyranoside (34). Installation of the methyl groups was performed as described for the synthesis of 32, with alcohol 26a (16.5 mg, 0.01 mmol), NaH (60% in mineral oil, 7.8 mg, 0.20 mmol) and MeI (8 μ L, 0.12 mmol) in THF (0.5 mL), at 0 °C for 1h. The trityl group was removed by addition of a methanolic solution of HCl (0.1 mL, 10% v/v). The crude product was purified by chromatography to afford 34 (10.2 mg, 73%) as colorless syrup; [α]_D = +9.4 (c 0.3, CHCl₃); ¹H NMR (600 MHz, CDCl₃) δ 7.89–7.77 (m, 6H, Ar), 7.36–7.19 (m, 21H, Ar), 5.10 (d, *J* = 1.7 Hz, 1H, H-1'), 5.06 (d, *J* = 1.9 Hz, 1H, H-1'), 4.95–4.91 (m, 2H, H-2', H-2''), 4.90 (d, *J* = 1.8 Hz, 1H, H-1), 4.80 (dd, *J* = 3.0, 2.0 Hz, 1H, H-2), 4.51–4.36 (m, 6H, 6 \times OCH₂Ph), 3.77 (app t, *J* = 9.7 Hz, 1H, H-4'), 3.72 (app t, *J* = 9.7 Hz, 1H, H-4'), 3.71–3.61 (m, 6H, H-4, H-5, H-5', H-5'', H-5''', octyl OCH₂), 3.61–3.49 (m, 6H, H-3, 5 \times H-6), 3.46 (dd, *J* = 9.3, 3.0 Hz, 1H, H-3'), 3.40 (dt, *J* = 9.8, 6.6 Hz, 1H, octyl OCH₂), 3.36 (dd, *J* = 9.5, 3.1 Hz, 1H, H-3''), 3.25 (s, 3H, OMe), 3.24 (s, 3H, OMe), 3.20 (s, 3H, OMe), 2.63 (d, *J* = 1.6 Hz, 1H, OH-4''), 2.45 (s, 3H, ArCH₃), 2.41 (s, 3H, ArCH₃), 2.41 (s, 3H, ArCH₃), 1.62–1.51 (m, 2H, octyl OCH₂CH₂), 1.35–1.23 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (126 MHz, CDCl₃) δ 145.0, 144.9, 144.8, 138.3, 138.3, 138.0, 134.0, 134.0, 133.6, 129.9, 129.8, 129.8, 128.4, 128.3, 128.2, 128.1, 127.9, 127.7, 127.6, 127.5, 127.5, 127.4 (36C, Ar), 99.5 (2C, C-1', C-1''), 97.3 (C-1), 79.2 (C-3'), 79.0 (C-3), 78.4 (C-3'), 74.4 (C-4), 74.1, 74.1, 73.9 (3C, C-2, C-2', C-2''), 73.7, 73.4, 73.4 (3C, 3 \times OCH₂Ph), 73.1 (C-4'), 72.2, 71.9, 71.2 (3C, C-5, C-5', C-5''), 70.2, 69.6, 69.3 (3C, C-6, C-6', C-6''), 68.4 (octyl OCH₂), 67.4 (C-4''), 57.1, 56.8, 56.7 (3C, 3 \times OMe), 31.8, 29.4, 29.3, 29.2, 26.0, 22.7 (6C, octyl CH₂), 21.7, 21.6, 21.6 (3C, 3 \times ArCH₃), 14.1 (octyl CH₃); HRMS (ESI) calcd C₇₁H₉₀O₂₂S₃ [M + Na]⁺ 1413.4978, found 1413.4964.

Octyl 6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranosyl-(1 \rightarrow 4)-6-O-benzyl-3-O-methyl-2-O-toluenesulfonyl- α -D-mannopyranoside (35). Installation of the methyl groups was performed as described for the synthesis of 32, with alcohol 27a (10.5 mg, 5.3 μ mol), NaH (60% in mineral oil, 8.2 mg, 0.21 mmol) and MeI (14 μ L, 0.22 mmol) in THF (0.4 mL), at 0 °C for 0.5h. The trityl group was removed by addition of a methanolic solution of HCl (0.1 mL, 10% v/v). The crude product was purified by chromatography to afford 35 (8.3 mg, 87%) as colorless syrup; [α]_D = +14.5 (c 0.6, CHCl₃); ¹H NMR (600 MHz, CDCl₃) δ 7.88–7.77 (m, 8H, Ar), 7.38–7.17 (m, 28H, Ar), 5.11 (d, *J* = 1.4 Hz, 1H, H-1''), 5.05 (d, *J* = 1.7 Hz, 1H, H-1'), 5.02 (d, *J* = 1.5 Hz, 1H, H-1''), 4.95–4.92 (m, 3H, H-2', H-2'', H-2'''), 4.90 (d, *J* = 1.7 Hz, 1H, H-1), 4.80 (dd, *J* = 2.9, 2.0 Hz, 1H, H-2), 4.52–4.31 (m, 8H, 8 \times OCH₂Ph), 3.77 (br app t, *J* = 9.6 Hz, 1H, H-4''), 3.73 (app t, *J* = 9.8 Hz, 1H, H-4''), 3.73–3.56 (m, 10H, H-4, H-4', H-5, H-5', H-5'', 3 \times H-6, octyl OCH₂), 3.57–3.43 (m, 8H, H-3, H-3', H-3'', 5 \times H-6), 3.40 (dt, *J* = 9.8, 6.9 Hz, 1H, octyl OCH₂), 3.37 (dd, *J* = 9.5, 3.1 Hz, 1H, H-3'''), 3.27 (s, 6H, 2 \times OMe), 3.22 (s, 3H, OMe), 3.20 (s, 3H, OMe), 2.63 (br s, 1H, OH-4'''), 2.45 (s, 3H, ArCH₃), 2.42 (s, 3H, ArCH₃), 2.41 (s, 6H, 2 \times ArCH₃), 1.61–1.52 (m, 2H, octyl OCH₂CH₂), 1.36–1.23 (m, 10H, octyl CH₂), 0.89 (t, *J* = 7.0 Hz, 3H, octyl CH₃); ¹³C NMR (151 MHz, CDCl₃) δ 145.0, 144.9, 144.8, 138.3, 138.3, 138.2, 137.9, 134.0, 134.0, 133.9, 133.6, 129.9, 129.8, 129.7, 128.3, 128.2, 128.1, 127.9, 127.6, 127.6, 127.5, 127.4, 127.4, 127.4 (48C, Ar), 99.5 (2C, C-1', C-1''), 99.4 (C-1'), 97.2 (C-1), 79.1 (C-3), 78.9 (C-3''), 78.8 (C-3'), 78.4 (C-3''), 74.6 (C-4), 74.0, 74.0, 74.0, 73.9 (4C, C-2, C-2', C-2''), 73.8 (C-4'), 73.6 (OCH₂Ph), 73.5 (2C, 2 \times OCH₂Ph), 73.4 (OCH₂Ph), 73.1 (C-4''), 72.2 (C-5'), 72.1 (C-5''), 71.9 (C-5'''), 71.2 (C-5), 70.2, 69.7, 69.5, 69.2 (4C, C-6, C-6', C-6'', C-6'''), 68.4 (octyl OCH₂), 67.4 (C-4'''), 57.0 (OMe), 56.7 (OMe), 56.62 (OMe), 56.61 (OMe), 31.8, 29.3, 29.3, 29.2, 26.0, 22.6 (6C, octyl CH₂), 21.7 (ArCH₃), 21.6 (3C, 3 \times ArCH₃), 14.1 (octyl CH₃); HRMS (ESI) calcd C₉₂H₁₁₄O₂₉S₄ [M + Na]⁺ 1833.6221, found 1833.6199.

Synthesis of Octyl 3-O-methyl- α -D-mannopyranosyl-(1 \rightarrow 4)-3-O-methyl- α -D-mannopyranosyl-(1 \rightarrow 4)-3-O-methyl- α -D-mannopyranoside (36). Deprotection was performed as for 33.

Removal of tosyl group was carried out with Mg (30 mg) in MeOH (2 mL) for 2 overnights. The crude product was purified by chromatography (CH₂Cl₂–methanol 15:1) to afford a partially deprotected trisaccharide (4.9 mg, 64%) as a colorless syrup. The benzyl groups of this tetrasaccharide (4.5 mg) were then removed by hydrogenolysis with Pd–C (5 wt.%, 5 mg) in methanol (1.5 mL) under H₂ atmosphere for 3 overnights. The catalyst was removed by filtration through Celite. The filtrate was then concentrated and purified by chromatography to afford **36** (2.8 mg, 88%) as a colorless film: [α]_D = +57.8 (c 0.2, MeOH); HRMS (ESI) calcd C₂₉H₃₄O₁₆ [M + Na]⁺ 681.3304, found 681.3298. The ¹H NMR spectra was identical to the previously reported.¹⁴

Octyl 3-O-methyl- α -D-mannopyranosyl-(1 → 4)-3-O-methyl- α -D-mannopyranosyl-(1 → 4)-3-O-methyl- α -D-mannopyranosyl-(1 → 4)-3-O-methyl- α -D-mannopyranoside (37). Deprotection was performed as for **33**. Removal of tosyl group was carried out with Mg (20 mg) in MeOH (1 mL) for overnight. The crude product was purified by chromatography (CH₂Cl₂–methanol 15:1) to afford a partially deprotected tetrasaccharide (5.3 mg, quantitative) as a colorless syrup. The benzyl groups of this tetrasaccharide were then removed by hydrogenolysis with Pd–C (5 wt.%, 13 mg) in methanol (1.5 mL) under H₂ atmosphere for 3 overnights. The catalyst was removed by filtration through Celite. The filtrate was then concentrated to afford **37** (2.7 mg, 80%) as a colorless film: [α]_D = +35.1 (c 0.3, MeOH); HRMS (ESI) calcd C₃₆H₆₆O₂₁ [M + Na]⁺ 857.3989, found 857.3993. The ¹H NMR spectra was identical to the previously reported.¹⁴

Hexakis(6-O-*t*-butyldimethylsilyl-2-O-toluenesulfonyl)- α -cyclodextrin (39). To a solution of **38**³⁹ (50.7 mg, 0.03 mmol) in THF (0.3 mL) was added *n*-Bu₂SnCl₂ (6.4 mg, 0.02 mmol) and *N,N*-diisopropylethylamine (190 μ L, 1.08 mmol) followed by toluenesulfonyl chloride (177.0 mg, 0.90 mmol). The reaction mixture was stirred at room temperature for six days and then purified by chromatography (hexane–EtOAc 3:1 to 1:1) to afford **39** (60.6 mg, 77%) as a colorless film: *R*_f 0.77 (CH₂Cl₂–methanol 30:1); [α]_D = +54.0 (c 0.6, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ 7.82 (d, *J* = 8.3 Hz, 2H, Ar), 7.36 (d, *J* = 8.1 Hz, 2H, Ar), 5.13 (d, *J* = 3.5 Hz, 1H, H-1), 4.28 (dd, *J* = 9.9, 3.5 Hz, 1H, H-2), 3.95 (ddd, *J* = 9.9, 8.5, 3.3 Hz, 1H, H-3), 3.91 (dd, *J* = 11.5, 2.9 Hz, 1H, H-6a), 3.71 (app t, *J* = 9.0 Hz, 1H, H-4), 3.64 (d, *J* = 11.0 Hz, 1H, H-6b), 3.56 (dd, *J* = 9.3, 2.7 Hz, 1H, H-5), 3.07 (d, *J* = 3.3 Hz, OH-3), 2.49 (s, 3H, ArCH₃), 0.89 (s, 9H, TBS(*t*-Bu)), 0.03 (s, 6H, 2 × TBS(Me)); ¹³C NMR (126 MHz, CDCl₃) δ 145.1 (1C, Ar), 133.0 (1C, Ar), 129.6 (2C, Ar), 128.4 (2C, Ar), 99.3 (C-1), 81.2 (C-4), 79.7 (C-2), 71.7 (C-5), 70.1 (C-3), 61.8 (C-6), 25.9 (TBS(*t*-Bu)), 21.8 (ArCH₃), 18.3 (TBS(*t*-Bu)), –5.1, –5.2 (2C, 2 × TBS(Me)); HRMS (ESI) calcd C₁₁₄H₁₈₀O₄₂S₆Si₆ [M + 2(NH₄)]²⁺ 1308.4783, found 1308.4761.

Hexakis(6-O-*t*-butyldimethylsilyl-3-O-methyl-2-O-toluenesulfonyl)- α -cyclodextrin (40). To a stirring ice-cold solution of alcohol **39** (30.6 mg, 0.01 mmol) and MeI (15 μ L, 0.22 mmol) in THF (0.5 mL) was added NaH (60% in mineral oil, 17 mg, 0.43 mmol). The resulting solution was stirred at 0 °C for 1 h before methanol (1 mL) was added. The reaction mixture was then concentrated, and the residue was purified by chromatography (hexane–EtOAc 3:1) to afford **40** (34.3 mg, quantitative) as a pale yellow syrup: *R*_f 0.66 (hexane–EtOAc 2:1); ¹H NMR (500 MHz, CDCl₃) δ 7.86 (d, *J* = 8.3 Hz, 2H, Ar), 7.35 (d, *J* = 8.1 Hz, 2H, Ar), 4.83 (d, *J* = 3.2 Hz, 1H, H-1), 4.07 (dd, *J* = 9.6, 3.2 Hz, 1H, H-2), 3.89 (dd, *J* = 11.4, 2.5 Hz, 1H, H-6a), 3.65–3.49 (m, 4H, H-3, H-4, H-5, H-6b), 3.45 (s, 3H, OMe), 2.48 (s, 3H, ArCH₃), 0.84 (s, 9H, TBS(*t*-Bu)), –0.02 (s, 3H, TBS(Me)), –0.03 (s, 3H, TBS(Me)); ¹³C NMR (126 MHz, CDCl₃) δ 145.0 (Ar), 133.5 (Ar), 129.6 (2C, Ar), 128.5 (Ar), 100.1 (C-1), 80.9 (C-3), 79.2 (C-4), 78.7 (C-2), 72.8 (C-5), 61.9 (OMe), 61.8 (C-6), 25.8 (TBS(*t*-Bu)), 21.8 (ArCH₃), 18.2 (TBS(*t*-Bu)), –5.0, –5.2 (2C, 2 × TBS(Me)); HRMS (ESI) calcd C₁₂₀H₁₉₂O₄₂S₆Si₆ [M + NH₄]⁺ 2683.0166, found 2683.0076.

■ ASSOCIATED CONTENT

📄 Supporting Information

NMR spectra for compounds **7–40**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: tlowary@ualberta.ca.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the Centre for Oil Sands Innovation, the Natural Sciences and Engineering Research Council of Canada, and the Department of Chemistry, University of Alberta. We are grateful to Professor George A. O'Doherty from the Department of Chemistry and Chemical Biology at Northeastern University for helpful discussions and providing information regarding the preparation of **7**.

■ REFERENCES

- (1) Cordell, G. A. *Phytochemistry* **1995**, *40*, 1585–1612.
- (2) Wessjohann, L. A. *Curr. Opin. Chem. Biol.* **2000**, *4*, 303–309.
- (3) Nagib, D. A.; MacMillan, D. W. C. *Nature* **2011**, *480*, 224–228.
- (4) Codee, J. D. C.; Ali, A.; Overkleef, H. S.; van der Marel, G. A. C. *R. Chim.* **2011**, *14*, 178–193.
- (5) Smoot, J. T.; Demchenko, A. V. *Adv. Carbohydr. Chem. Biochem.* **2009**, *62*, 161–250.
- (6) Ogawa, T.; Matsui, M. *Carbohydr. Res.* **1977**, *56*, C1–C6.
- (7) Ogawa, T.; Matsui, M. *Tetrahedron* **1981**, *37*, 2363–2369.
- (8) Angyal, S. J.; Melrose, G. J. H. *J. Chem. Soc.* **1965**, 6494–6500.
- (9) Abad-Romero, B.; Mereiter, K.; Sixta, H.; Hofinger, A.; Kosma, P. *Carbohydr. Res.* **2009**, *344*, 21–28.
- (10) Jackowski, O.; Bussiere, A.; Vanhaverbeke, C.; Baussanne, I.; Peyrin, E.; Mingot-Leclercq, M. P.; Decout, J. L. *Tetrahedron* **2012**, *68*, 737–746.
- (11) Although **4** is not an oligosaccharide, it is classified as such here given its structural similarity to a disaccharide.
- (12) Bergeron, R. J. M., M. P.; Machida, Y. *Bioorg. Chem.* **1976**, *5*, 121–126.
- (13) Wang, Y.; Huang, X.; Zhang, L. H.; Ye, X. S. *Org. Lett.* **2004**, *6*, 4415–4417.
- (14) Xia, L.; Zheng, R. B.; Lowary, T. L. *ChemBioChem* **2012**, *13*, 1139–1151.
- (15) Achmatowicz, O., Jr.; Bielski, R. *Carbohydr. Res.* **1977**, *55*, 165–176.
- (16) Li, M. S.; Scott, J.; O'Doherty, G. A. *Tetrahedron Lett.* **2004**, *45*, 1005–1009.
- (17) Babu, R. S.; Zhou, M.; O'Doherty, G. A. *J. Am. Chem. Soc.* **2004**, *126*, 3428–3429.
- (18) David, S.; Hanessian, S. *Tetrahedron* **1985**, *41*, 643–663.
- (19) It should be appreciated that stannylene acetals of carbohydrate diols typically adopt structure-dependent higher-order oligomers (e.g., dimers, pentamers) in solution.^{18–22} The depiction of **14** as a monomer is a simplified representation of the intermediate, because, if indeed it is formed, its oligomeric structure in solution has not been investigated and is therefore unknown.
- (20) Grindley, T. B.; Wasylishen, R. E.; Thangarasa, R.; Power, W. P.; Curtis, R. D. *Can. J. Chem.* **1992**, *70*, 205–217.
- (21) Grindley, T. B. *Adv. Carbohydr. Chem. Biochem.* **1998**, *53*, 17–142.
- (22) Holzapfel, C. W.; Koekemoer, J. M.; Marais, C. F.; Kruger, G. J.; Pretorius, J. A. S. *Afr. J. Chem.* **1982**, *35*, 80–88.
- (23) Nagashima, N.; Ohno, M. *Chem. Lett.* **1987**, 141–144.

- (24) Martinelli, M. J.; Vaidyanathan, R.; Pawlak, J. M.; Nayyar, N. K.; Dhokte, U. P.; Doecke, C. W.; Zollars, L. M.; Moher, E. D.; Khau, V. V.; Kosmrlj, B. *J. Am. Chem. Soc.* **2002**, *124*, 3578–3585.
- (25) Peri, F.; Cipolla, L.; Nicotra, F. *Tetrahedron Lett.* **2000**, *41*, 8587–8590.
- (26) Lee, D.; Taylor, M. S. *J. Am. Chem. Soc.* **2011**, *133*, 3724–3727.
- (27) Demizu, Y.; Kubo, Y.; Miyoshi, H.; Maki, T.; Matsumura, Y.; Moriyama, N.; Onomura, O. *Org. Lett.* **2008**, *10*, 5075–5077.
- (28) Nashed, M. A.; Anderson, L. *Tetrahedron Lett.* **1976**, 3503–3506.
- (29) Wu, X. F.; Kong, F. Z. *Carbohydr. Res.* **1987**, *162*, 166–169.
- (30) Tam, P. H.; Lowary, T. L. *Carbohydr. Res.* **2007**, *342*, 1741–1772.
- (31) Malik, S.; Dixit, V. A.; Bharatam, P. V.; Kartha, K. P. R. *Carbohydr. Res.* **2010**, *345*, 559–564.
- (32) Purdie, T.; Irvine, J. C. *J. Chem. Soc.* **1903**, *83*, 1021–1037.
- (33) Neeman, M.; Caserio, M. C.; Roberts, J. D.; Johnson, W. S. *Tetrahedron* **1959**, *6*, 36–47.
- (34) Evans, D. A.; Ratz, A. M.; Huff, B. E.; Sheppard, G. S. *Tetrahedron Lett.* **1994**, *35*, 7171–7172.
- (35) Sureshan, K. M.; Shashidhar, M. S.; Praveen, T.; Gonnade, R. G.; Bhadbhade, M. M. *Carbohydr. Res.* **2002**, *337*, 2399–2410.
- (36) Sridhar, M.; Kumar, B. A.; Narender, R. *Tetrahedron Lett.* **1998**, *39*, 2847–2850.
- (37) Khan, A. R.; Forgo, P.; Stine, K. J.; D'Souza, V. T. *Chem. Rev.* **1998**, *98*, 1977–1996.
- (38) Chiu, S. H.; Myles, D. C.; Garrell, R. L.; Stoddart, J. F. *J. Org. Chem.* **2000**, *65*, 2792–2796.
- (39) Grachev, M. K.; Edunov, A. V.; Kurochkina, G. I.; Levina, I. I.; Nifant'ev, E. *Russ. J. Gen. Chem.* **2011**, *81*, 322–329.
- (40) Stannylene-acetal intermediates of *trans*-diols, in particular α -glucopyranosides, have been described.^{18–22}
- (41) Ballou, C. E. *Pure Appl. Chem.* **1981**, *53*, 107–112.
- (42) Jackson, M.; Brennan, P. J. *J. Biol. Chem.* **2009**, *284*, 1949–1953.
- (43) Mendes, V.; Maranhã, A.; Alarico, S.; Empadinhas, N. *Nat. Prod. Rep.* **2012**, *29*, 834–844.
- (44) Liu, L.; Bai, Y.; Sun, N.; Xia, L.; Lowary, T. L.; Klassen, J. S. *Chem.—Eur. J.* **2012**, *18*, 12059–12067.
- (45) Hsu, M. C.; Lee, J.; Kishi, Y. *J. Org. Chem.* **2007**, *72*, 1931–1940.
- (46) Cheon, H. S.; Lian, Y.; Kishi, Y. *Org. Lett.* **2007**, *9*, 3323–3326.
- (47) Hirooka, M.; Terayama, M.; Mitani, E.; Koto, S.; Miura, A.; Chiba, K.; Takabatake, A.; Tashiro, T. *Bull. Chem. Soc. Jpn.* **2002**, *75*, 1301–1309.
- (48) Liao, W. S.; Lu, D. P.; Li, A. H.; Kong, F. Z. *J. Carbohydr. Chem.* **1997**, *16*, 877–890.
- (49) Liao, W. S.; Lu, D. P. *Carbohydr. Res.* **1997**, *300*, 347–349.
- (50) Liao, W. S.; Lu, D. P. *Carbohydr. Res.* **1996**, *296*, 171–182.
- (51) Muramatsu, W. *J. Org. Chem.* **2012**, *77*, 8083–8091.