


Developing Ligands for Palladium(II)-Catalyzed C–H Functionalization: Intimate Dialogue between Ligand and Substrate

Keary M. Engle and Jin-Quan Yu*

Department of Chemistry, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037, United States

ABSTRACT: Homogeneous transition-metal-catalyzed reactions are indispensable to all facets of modern chemical synthesis. It is thus difficult to imagine that for much of the early 20th century, the reactivity and selectivity of all known homogeneous metal catalysts paled in comparison to their heterogeneous and biological counterparts. In the intervening decades, advances in ligand design bridged this divide, such that today some of the most demanding bond-forming events are mediated by ligand-supported homogeneous metal species. While ligand design has propelled many areas of homogeneous catalysis, in the field of Pd(II)-catalyzed C–H functionalization, suitable ligand scaffolds are lacking, which has hampered the development of broadly practical transformations based on C–H functionalization logic. In this Perspective, we offer an account of our research employing three ligand scaffolds, mono-*N*-protected amino acids, 2,6-disubstituted pyridines, and 2,2'-bipyridines, to address challenges posed by several synthetically versatile substrate classes. Drawing on this work, we discuss principles of ligand design, such as the need to match a ligand to a particular substrate class, and how ligand traits such as tunability and modularity can be advantageous in reaction discovery.



1. INTRODUCTION

1.1. The Role of Ligand Design in Transition-Metal Catalysis. The importance of ligands in modern homogeneous transition-metal catalysis cannot be overstated. Because ligand coordination changes the structure and reactivity of a metal catalyst, it inherently changes the activation energy of elementary steps in a given catalytic process. This change is manifested in the kinetic reactivity,¹ which can broaden the effective substrate scope of the reaction. Moreover, ligands can influence the selectivity (i.e., enantioselectivity, diastereoselectivity, regioselectivity, and chemoselectivity) in transformations where more than one product is produced, improve the solubility of metal catalysts in organic solvents, and extend catalyst lifetimes by suppressing metal catalyst degradation pathways. For instance, in cases where precipitation of metal nanoparticles is an irreversible catalyst deactivation pathway, strongly binding organic ligands can protect two molecules of catalyst from contacting one another and precipitating. Finally, as a consequence of these previous points, the ligand can affect the operational properties of the reaction: the compatibility with air and/or moisture, the reaction temperature, pressure, etc. In essence, ligands are chemists' "hands" for influencing the bond-making and -breaking processes that occur at metal centers, and arguably, in the past several decades, ligand design has been the principal force driving the improvement of known catalytic reactions and the discovery of new ones.²

During the past century, simple, commonly available metal salts (e.g., CuCl, AuCl₃, and Pd(OAc)₂) have been applied extensively in catalysis, leading to the discovery of a myriad of different reactions, several of which have become mainstays in

organic synthesis. This flurry of research in catalytic organometallic chemistry set the stage for the design of catalysts with improved reactivity and selectivity, such that desired products could be synthesized with high levels of purity and minimal amounts of waste. Because improved catalytic performance hinges on the properties of ancillary ligands, new ligand scaffolds have been vigorously pursued in all subfields of homogeneous metal catalysis.

Much of the search for new ligands has taken place in the context of research into catalytic *asymmetric* reactions, where in the absence of chiral information the reaction will proceed via one of two pathways of equal energy to give a 1:1 mixture of enantiomeric products. By using a chiral nonracemic ligand, however, the reaction can be rendered enantioselective, leading to the predominant production of one enantiomer. Prominent early examples include the use of dialkyl tartrate ligands in the Sharpless asymmetric epoxidation^{3,4} and the development of chiral diphosphine ligands for asymmetric hydrogenation, such as DIOP,⁵ DiPAMP,⁶ chiraphos,⁷ and BINAP⁸ by Kagan, Knowles, Bosnich, and Noyori, respectively. Ligand design has played an equally important role in achieving *enhanced kinetic reactivity* for catalytic transformations that otherwise exhibit prohibitively slow reaction rates (or a complete lack of reactivity all together). Notable examples include the popularization of triphenylphosphine as a ligand in the 1960s, particularly in Wilkinson's seminal applications to rhodium-catalyzed hydroformylation⁹ and hydrogenation,¹⁰ and more

Received: February 12, 2013

Published: April 8, 2013




Figure 1. "Privileged" chiral ligands for asymmetric transition-metal catalysis.²⁰




Figure 2. "Privileged" achiral ligands for high kinetic reactivity in transition metal catalysis.⁴⁴

recently, the advent of dialkylaryl- and trialkylphosphines and *N*-heterocyclic carbenes as ligands in ruthenium-catalyzed olefin metathesis^{11,12} and palladium-catalyzed cross-coupling reactions.^{13–17} Moreover, there are also important cases where the search for ligands for stereoselection led to the identification of more reactive catalysts and vice versa.^{18,19}


1.2. "Privileged" Ligands in Transition-Metal Catalysis. During the past half-century of research in asymmetric transition-metal catalysis, a small collection of "privileged" ligand scaffolds have emerged, a term first coined by Yoon and Jacobsen.²⁰ These ligand scaffolds have been found to be useful in many different types of asymmetric transformations, even among those with few or no mechanistic similarities (1–9, Figure 1). Privileged chiral ligands have been used extensively for reaction development in organic chemistry, and their application accounts for a high percentage of new asymmetric reactions developed each year.

As discussed above, in parallel to the progress in asymmetric catalysis, homogeneous catalytic transformations that do not induce chirality have been extensively studied. These reactions have required catalysts with high kinetic reactivity to ensure broad substrate scope, short reaction times, and favorable product distributions, and here too, ligand development has played a central role. In nonstereoselective reactions, there exist achiral "privileged" ligands that have been utilized ubiquitously


to enable a variety of different catalytic reactions, all with a myriad of distinct elementary steps (Figure 2). These include triaryl- and trialkylphosphines (10–13),^{21–24} biaryl(dialkyl)-phosphines (14–19),^{25,26} diphosphines (20–25),^{27,28} "pincer" ligands (26–28),^{29,30} *N*-heterocyclic carbenes (29–33),^{16,31,32} cyclopentadienides (34–36),^{33–38} acetylacetonates (37–39),³⁹ diamines (40–42),^{39–41} bipyridines (43–45),^{39,41–43} and pyridines (46–48).⁴¹

1.3. Matching a Substrate Class to a Metal Catalyst and Identifying a Suitable Ligand Scaffold. Though privileged ligand scaffolds are remarkably general in their utility across reaction types and substrate classes, they are still not universally applicable. One common feature of ligand-controlled metal-catalyzed reactions is that the catalytic system must be designed to match the properties of the substrate. Each individual class of substrates possesses a characteristic set of properties that will affect its approach to and interaction with the metal center. The course of the reaction is dictated by the intrinsic properties of the metal center (oxidation state, electron count, coordination geometry, etc.) and its established redox activity patterns. Together the properties of the substrate and metal then inform the selection and/or design of appropriate ligand scaffolds. Important substrate considerations include the steric and electronic properties (particularly those proximate to the reactive functional group), any existing

Scheme 1. Application of a Diverse Array of Catalyst/Ligand Structures to Asymmetric Transformations of Olefins: (a) Asymmetric Hydrogenation of Olefins by Noyori^{47,54} and Pfaltz;⁴⁹ (b) Asymmetric Epoxidation of Olefins by Sharpless^{3,4} and Jacobsen;⁵³ (c) Ligand Structures



Scheme 2. Application of Sterically Bulky, Electron-Rich Phosphine Ligands in the Suzuki–Miyaura Cross-Coupling of Unactivated Aryl Chlorides with Boronic Acids, As Reported Independently by the Groups of (a) Buchwald²⁵ and (b) Fu²³



stereochemistry, the presence or absence of other potentially reactive functional groups, and the presence or absence of Lewis basic chelating functional groups (directing groups).^{45,46} For example, consider asymmetric olefin hydrogenation (Scheme 1a). For substrates containing a chelating functional group, such as α,β -unsaturated carboxylic acid 49, the presence of an open binding site on the metal for carboxylate coordination is desirable. With the C₂-symmetric Noyori [Ru(BINAP)(OAc)₂] catalyst, the substrate can smoothly displace the acetate, without compromising catalyst activity, and subsequent directed asymmetric hydrogenation proceeds with excellent levels of stereoselection.⁴⁷ In contrast, in the Pfaltz nondirected asymmetric hydrogenation of olefins (51), a Crabtree-type catalyst⁴⁸ bearing a non-C₂-symmetric P,N-phosphite–pyridine ligand 52 and a noncoordinating counterion, BARF, is highly effective in the absence of a chelating functional group, relying on weak coordinative interactions for orienting the substrate.^{49,50} Similarly, for asymmetric olefin epoxidation (Scheme 1b), the Sharpless system^{3,4,51} is highly effective for allylic alcohols (54), and the Jacobsen system works well for internal (Z)-olefins not containing directing

groups (57).^{52,53} Although the net transformations appear similar in both cases, the properties of the substrate dictate the selection of a metal, the mechanistic manifold, and ultimately the ligand used for stereoselection.

In nonstereoselective catalysis, tailoring ligand properties to match the substrate is similarly important. For instance, in Pd-catalyzed Suzuki–Miyaura⁵⁵ cross-coupling reactions, unactivated aryl chlorides were traditionally an unreactive class of substrates. Ligands that are commonly used with aryl iodides, -bromides, and -triflates, such as triphenylphosphine, BINAP, and dppf, are generally unreactive with electron-neutral or -rich aryl chlorides, such as 59,¹³ presumably because oxidative addition of a C(sp²)–Cl bond to a Pd(0) center is kinetically unfavorable.⁵⁶ This challenge can be surmounted, however, through the application of sterically demanding, electron-rich phosphine ligands such as DavePhos (16)²⁵ and P(*t*-Bu)₃ (12) (Scheme 2).²³ The σ -donating character of these ligands increases the nucleophilicity of the Pd(0) center which lowers the activation energy for oxidative addition. The steric bulk is thought to facilitate reductive elimination and promote formation of the catalytically active 1:1 Pd(0)/ligand complex




Figure 3. Selected examples of commercially available BINOL derivatives.




Figure 4. (a) Modular synthesis of chiral PHOX-type ligands.⁷⁰ (b) Representative PHOX ligands.⁷¹

in solution.¹³ Through the selection and design of appropriate ligands, a new class of widely available substrates was rendered reactive in this and other related cross-coupling chemistries.

1.4. Modifiability and Modularity of Ligand Scaffolds.

Because it is difficult to predict a priori the potentially dramatic influence that subtle changes to a ligand's structure will have on the metal center as it cycles through many elementary steps of a catalytic process,^{57,58} one key to the success of a ligand scaffold is that it can be readily modified. Multiple sites of variation both near and distal to the metal allow for the steric and electronic properties to be finely tuned to meet the energetic requirements of the reaction. For instance, the BINAP/BINOL scaffold has been adapted through modification at nearly every conceivable position since its introduction in asymmetric catalysis^{8,59–62} and has continued to play a central role in reaction discovery (Figure 3). In the early stages of reaction discovery and development, catalyst modifiability is particularly important because many of the crucial reaction parameters are typically poorly understood. As evidence of the importance of modifiability, it is becoming increasingly common for commercial suppliers to offer ligand kits, containing multiple structural variants on a single ligand scaffold.

For similar reasons, many widely used ligand scaffolds are modular, meaning that they can be synthesized in a small number of steps from readily available components, allowing for a library of ligand candidates to be synthesized expediently.⁶³ For instance, chiral phosphine–oxazoline (PHOX, **66**) ligands pioneered by Pfaltz,⁶⁴ Helmchen,⁶⁵ and Williams⁶⁶ possess a nonsymmetrical P,N-donor system⁶⁷ with one soft atom (P) and one hard atom (N) (Figure 4). Their unique structure has proven to be critical for challenging asymmetric metal-catalyzed transformations, particularly asymmetric allylic substitution. A major practical aspect that makes this class of ligands convenient to use is the modular synthesis,

where an entire family of ligand candidates can be synthesized from widely available commercial chiral building blocks.^{68,69}

Tailoring ligand properties for a desired metal-catalyzed transformation requires careful consideration of the key features of the substrates class as well as the mechanism of each elementary step in the would-be catalytic process. Ligands that have a rich history of success in metal-catalyzed reactions (those that are considered privileged) are often useful starting points in reaction discovery and development. In this vein, modifiability and modularity are key in providing libraries of ligands for medium- and high-throughput screening campaigns. As discussed below, many of the key advancements in our own laboratory have hinged upon ligand design, so the considerations outlined above are ones that we have returned to repeatedly as our research program in Pd(II)-catalyzed C–H functionalization has matured.

In the ensuing section, we discuss some of our general goals and sources of inspiration and outline how ligand development plays a key role in the realization of our objectives. We then go on to describe the challenges in identifying effective ligand scaffolds for C(sp²)-H and C(sp³)-H functionalization using catalytic Pd(II). Drawing primarily from our own results, we present three stories about three different ligand scaffolds and the substrate classes with which they were designed to promote selective C–H functionalization. We show how ligand design can enable enantioselective C–H activation and also serve to enhance reactivity with achiral substrates, thereby dramatically broadening substrate scope. We further discuss the need to match a target substrate and ligand to enable a productive interplay at the metal center as a prerequisite to efficient catalysis. In recounting these examples, we illustrate the principles articulated above and reflect on the lessons learned from our work.

2. OVERVIEW OF OUR RESEARCH PROGRAM IN Pd(II)-CATALYZED C–H FUNCTIONALIZATION

2.1. Philosophy and Guidelines for Reaction Development. While there remain numerous challenges in organic synthesis that can be addressed through transition-metal catalysis, a primary focus of our own research program has been the creation of novel and strategic retrosynthetic disconnections^{72–75} that allow C–H bonds to be viewed as dormant functional groups that can be converted to a desired C–C or C–X bond at any stage of a synthesis.^{76,77} To make this approach broadly useful for synthetic applications, we have adhered to the following guidelines in developing C–H functionalization reactions:

(1) *Substrates.* We utilize substrate classes that are readily available in both bulk quantity and structural diversity and that can lead to a variety of complex synthetic targets upon C–H functionalization (Figure 5). Newly uncovered modes of




Figure 5. Aryl- and alkylcarboxylic acids, representative starting material classes that are abundant in quantity and rich in structural diversity.

reactivity can then be extended to feedstock chemicals, including those that contain weakly coordinating functional groups that can be used to facilitate C–H cleavage.⁷⁷ Overall, appropriate selection of target substrate classes can further improve both the economy⁷⁸ and sustainability⁷⁹ of organic synthesis.

(2) *Products.* Equally important is that C–H activation and subsequent functionalization take place at a strategic position and install a desirable functional group so that the product represents a commonly encountered structural motif, such as a β -functionalized carboxylic acid, amino acid, or biaryl. Establishing reliable routes to these substructures facilitates synthesis planning that incorporates C–H activation transformations as key disconnections. When combined with the selection of simple starting materials, drastic increases in molecular complexity can be achieved using this approach (Scheme 3).


(3) *Reactivity.* We preferentially target catalytic manifolds that allow flexibility in terms of the scope of nucleophiles, electrophiles, and/or radical species that can serve as reacting partners with C–H bonds. Versatility in the functionalization step allows the [C–M] bond resulting from C–H cleavage to be efficiently mapped into any carbon–carbon or carbon–heteroatom bond (Scheme 4).

(4) *Catalysts/Ligands.* For the reasons outlined above, ligand-controlled C–H functionalization reactions represent an especially fertile ground for discovering new modes of reactivity and for gleaning fundamental insights into chemical reactivity (Scheme 5). We view ligand development as the long-term solution for improving catalyst performance: reactivity (TON, TOF, and substrate scope), selectivity (site-selectivity, chemo-selectivity, and stereoselectivity), and operating conditions (reaction time, temperature, and tolerance for air and moisture).


(5) *Operational Convenience.* In order for novel C–H functionalization reactions to be embraced by the synthetic community, the reactions must be operationally simple and robust. In this respect, substrates, coupling partners, catalysts, and ligands that are commercially available or easily accessible are preferable. Reactions that do not require special equipment and can be set up on the benchtop without rigorous exclusion of air or moisture are similarly desirable.

In line with these goals, much of our research has focused on C–H functionalization reactions catalyzed by Pd(II).^{77,91–95}


Scheme 3. Novel Retrosynthetic Disconnections Enabled by Our Methodology: Routes to (a) the Kinamycins⁸⁰ and (b) (+)-Lithospermic Acid⁸¹



Scheme 4. Diverse Pd(II)-Catalyzed C–H Functionalization of (a) Benzoic Acids^{82–86} and (b) Benzoic Acid Derived N-Arylamides^{87–90} Reported by Our Group



Scheme 5. Enantioselective Pd(II)-Catalyzed C–H Activation Using a Chiral Ligand



One of the attractive aspects of this approach is the established, versatile reactivity of [R–Pd(II)] (R = aryl or alkyl) intermediates. In the context of Pd(0)-catalyzed aryl halide functionalization, it has been well-established that [R–Pd(II)] intermediates can effectively couple with a range of carbon- and heteroatom-based nucleophiles via Pd(0)/Pd(II) catalysis. Thus, C–H cleavage with Pd(II), which is generally believed to be redox neutral with concomitant loss of HX (X = halide, acetate, etc.), can be viewed as an alternative and complementary entry point to this same collection of bond-forming reactions. Key to the achieving this type of general reactivity is the development of effective ligands to promote C–H cleavage,⁷⁶ which is typically the rate-limiting and selectivity-determining step in Pd(II)-catalyzed C–H functionalization reactions. It is also our intention that fundamental studies on the interplay between Pd(II), substrate, and ligand could lend useful information to the development of C–H functionalization reactions catalyzed by other metals, especially Cu(II),⁹⁶ Rh(III),⁹⁷ and Ru(II),⁹⁸ which can potentially perform redox chemistry similar to that of Pd(II).⁹⁹

2.2. The Need for Effective Ligand Scaffolds. During the past 10 years that we have worked in this area, an

impediment to our research has been the lack of ligand scaffolds for accelerating C–H cleavage with Pd(II).¹⁰⁰ We have consistently resorted to the simplest Pd(II) salts available (Pd(OAc)₂, Pd(TFA)₂, PdI₂, etc.), and with these basic catalysts, many of our goals, including the development of methodology for functionalizing bulk chemicals and for achieving site- and stereoselectivity in the C–H cleavage step, remained elusive. Thus, the lack of suitable ligand scaffolds represents a major roadblock in harnessing the full potential of Pd(II)-catalyzed C–H functionalization. Indeed, this need for ligands is at the heart of commonly cited problems in the field, like the inability to control positional selectivity of functionalized arenes¹⁰¹ or more generally, the inability to selectively functionalize one of the many inequivalent C–H bonds present in organic small molecules.

We became keenly aware of these shortcomings when developing methodology for stereoselective C–H activation which would require an efficient chiral [Pd(II)–ligand]* catalyst (section 3.1).⁹³ As we began to move from a chiral auxiliary approach to a chiral catalyst approach, we confronted a long-standing challenge in the field: uncovering a chiral metal catalyst for enantioselective C–H activation. Our motivation for investigating enantioselectivity was rooted in the hypothesis that a chiral ligand effective for stereoselection is likely to be involved in the C–H cleavage transition state. Involvement in the C–H cleavage step suggests that this ligand scaffold, or a closely related scaffold, could ultimately be used for the interrelated problems of rate acceleration and site-selectivity.

3. IDENTIFICATION OF LIGAND SCAFFOLDS FOR Pd(II)-CATALYZED C–H FUNCTIONALIZATION

3.1. Considerations for Ligand Design. When considering potential ancillary ligand scaffolds for Pd(II)-catalyzed C–H functionalization, there are many interrelated challenges and complications to keep in mind:

(1) *Assembling a 1:1 Ligand/Substrate Coordination Structure.* The ligand could outcompete the substrate for coordination to the Pd(II) center or vice versa. Ligands by their very nature contain coordinating functional groups (either in the form of nonbonding lone pair electrons or bonding π -systems). By contrast, unactivated C–H bonds are paraffin in nature, meaning that they are nonpolar and have inherently low affinity for cationic metal centers. Achieving high effective molarity of the substrate in the presence of a ligand is thus difficult, particularly with simple hydrocarbon substrates, like toluene or hexane. In directed C–H activation, on the other hand, many traditional directing groups (pyridines, oxazolines, oximes, etc.),^{92,102,103} could inhibit ligand coordination prior to C–H cleavage or could render the C–H cleavage step so facile that the ligand would not play a dominant role in influencing reactivity and/or selectivity.

(2) *Promoting the C–H Cleavage Step.* Under different mechanistic paradigms for C–H cleavage, the electronic requirements of the Pd(II) center, and thus the supporting ligand, are different. For instance, in C(aryl)–H cleavage along a electrophilic palladation mechanism¹⁰⁴ there is a need to maintain relatively electrophilic Pd(II) center. In a concerted metalation/deprotonation (CMD) mechanism,^{105–108} a more nucleophilic Pd(II) center can be effective. The operative mechanism for C–H cleavage typically depends on a combination of factors, including the substrate, the reaction conditions, and also the ligand(s).⁷⁶

(3) *Compatibility with the Other Steps in the Catalytic Cycle and the Reaction Conditions.* Different Pd(II)-mediated C–H functionalization reactions proceed via different elementary steps, and each poses challenges in ligand engineering. For example, some steps may require ligand dissociation to provide a vacant coordination site, while others could require the ligand to be resistant to reductive elimination from high-valent intermediates. Additionally, under the reaction conditions, the ligand could react with the substrate, the coupling partner, or the Pd(II) catalyst along any of a number of possible degradation pathways.

(4) *Modifiability and Modularity.* Because it is unlikely that a single ligand will be adequate for each combination of substrate and reaction partner, it is advantageous to pursue multiple distinct ligand scaffolds in parallel. In addition to developing multiple complementary ligand scaffolds, utilizing modifiable and modular ligand scaffolds can be advantageous for fine-tuning a ligand to a specific combination of substrate and reaction type.

These general considerations and design guidelines provide a conceptual framework for ligand development in Pd(II)-catalyzed C–H functionalization. Naturally, as our research program has progressed, our theoretical understanding of ligand design principles has evolved as a result of new insights from experimental work. Owing to the inherent unpredictability of ligand discovery, there is also an element of empiricism and serendipity to this research area, which is why it is important to think carefully but still be adventurous in exploring old and new ligand architectures.

Below we present three case studies from our research group about three ligand scaffolds, mono-*N*-protected amino acids, 2,6-disubstituted pyridines, and 2,2'-bipyridines, which we have found to be effective for Pd(II)-catalyzed C–H functionalization (Figure 6). Of the three ligand scaffolds, we have studied




Figure 6. Ligand scaffolds discussed in this Perspective.

mono-*N*-protected amino acids in the most depth to date; we thus provide a more comprehensive discussion of this work and shorter synopses of studies done with the other two ligand scaffolds. We describe previous findings that led to identification of these ligand scaffolds and examine how the principles outlined above are at play in each case.

3.2. Mono-*N*-Protected Amino Acid Ligands.

3.2.1. Background. Drawn to the richness of mechanistic questions and potential synthetic applications of C–H activation chemistry, we established our research program in 2002 at the University of Cambridge to investigate stereoselective induction in the palladation of prochiral C–H bonds, which we viewed as fundamentally important to the understanding of C–H cleavage by Pd(II) centers.⁹³ To gain structural information concerning the transition state of directed palladation reactions, we focused on reactions that would proceed through well-defined intermediates that could be characterized. Inspired by the important role that removable chiral auxiliaries play in modern organic synthesis, our first approach centered on the

use of a chiral oxazoline group^{109–111} that could be readily installed onto aliphatic carboxylic acids. Following this strategy, we found high levels of stereoselective induction (up to >99:1 dr) could be achieved in the diastereoselective C–H iodination and acetoxylation of prochiral α,α -gem-dimethyl groups, α,α -gem-diphenyl groups, and cyclopropanes along a Pd(II)/Pd(IV) catalytic cycle (Scheme 6a).^{91,112–114}


tert-Butyl-substituted oxazoline **110** proved to be remarkably effective in directing C(sp³)-H activation, even under mild conditions (including temperatures as low as 0 °C). We have sought to understand the root cause of this reactivity through structural characterization both in solution and in the solid state. Further studies to elucidate the transition state structure via computational analysis have also been performed in collaboration with the Houk group (Scheme 6b).¹¹⁵ We have characterized several trinuclear palladium intermediates (e.g., **119b**), performed calculations that revealed the involvement of monomeric precursor **115** as the reactive intermediate, and modeled various possible transition states for the diastereoselective C–H cleavage step (**116**). This work has confirmed the importance of conformation and geometry in the square planar pretransition-state coordination structure **115**, in which the target C–H bond is approximately coplanar to the acetate anion. This arrangement minimizes the dihedral angle between the C–H bond and the Pd(II)–OAc, which is critical during the CMD transition state.

3.2.2. Enantioselective C–H Activation. As this work was progressing, we took the next conceptual step forward and questioned whether we could develop a chiral ligand capable of achieving similar levels of stereoselective induction in the absence of a chiral auxiliary. To that end, we used diphenyl-2-pyridylmethane (**120**) as our pilot substrate and developed efficient nonstereoselective conditions to achieve Pd(II)-catalyzed C–H/R–BX_n cross-coupling (Schemes 7 and 8).⁹² In the long term, we hoped to use this reaction system as a model for developing a broad range of enantioselective carbon–carbon and carbon–heteroatom bond-forming reactions, in which C–H cleavage could be directed by a diverse collection of functional groups.


Following the isolation and characterization of cyclopalladated dimer **122**, we reasoned that prior to the C–H cleavage step, the putative pretransition-state coordination structure **124** offered the opportunity to replace the achiral nonbridging acetate ligands on the Pd(II) center with chiral carboxylates. In this manner, we hypothesized, stereoselective C–H cleavage could take place at one of the two prochiral phenyl groups. Although this line of thinking is not entirely accurate on the basis of our current understanding, it nonetheless served as an invaluable starting point to probe enantioselective C–H activation.

To develop a catalytic enantioselective reaction, we first surveyed a variety of chiral acid ligands in an attempt to induce stereoselectivity in the Pd(II)-catalyzed enantioselective C–H activation of **125** (Table 1).¹¹⁷ We were encouraged by modest enantioselectivity (<25%) using this approach, and we reasoned that a more rigid backbone structure would limit the conformational degrees of freedom that the ligand possessed. For this reason, we prepared chiral cyclopropane amino acids **136–139** and found that the ee values could be improved to >40%, while maintaining good yield (Table 2). Consistent with expectations, enantiomers **136** and **138** led to product mixtures with opposite enantioselectivity values. Surprisingly, pseudo-C₂-symmetric ligand **139** led to a nearly racemic mixture. These

Scheme 6. (a) Diastereoselective C(sp³)-H Iodination of *gem*-Dimethyl Groups Using a Removable Chiral Oxazoline Auxiliary. (b) General Catalytic Cycle for C-H Iodination via Pd(II)/Pd(IV) Catalysis. (c) Model for Diastereocontrol, Including Proposed Transition-State Structure



Scheme 7. General Catalytic Cycle for C-H/R-BX_n Cross-Coupling via Pd(II)/Pd(0) Catalysis




data, in conjunction with previous work in the literature characterizing amino acid-bound palladacycles,¹¹⁸ suggested that during the C-H cleavage step the ligand was bound in a bidentate fashion with the N-H bond possibly remaining intact, inducing chirality at the nitrogen atom. In this way, chiral information could be effectively geared from the α -carbon atom of the ligand to the nitrogen atom, which would rationalize why **139**, which has two α -substituents of similar steric bulk, was ineffective.

With a refined mechanistic understanding, we returned to optimizing the ligand scaffold by maximizing the steric

difference between the two α -substituents. To our delight, we found that simple, commercially available Boc-protected amino acids¹¹⁹ gave high levels of stereoselection (entries 1–7, Table 3). Among the Boc-protected amino acids examined, leucine (entry 7) proved to be the optimal backbone structure. In an effort to improve the yield, we then fine-tuned the alkyl substituent on the carbamoyl protecting group. By introducing a (–)-menthyl group, we were able to improve the yield while maintaining the stereoselectivity (entry 11).

Importantly, we were also able to extend this concept to enantioselective C(sp³)-H activation using pyridyl substrates containing *gem*-dimethyl groups (Scheme 9). In this case, one of the rigid cyclopropane amino acid ligands from above, **136**, was found to be optimal. Although, the alkylated product was only obtained in modest yield and ee, it served as crucial proof of concept that through future advances in ligand design, synthetically useful levels of stereoselection could be achieved in enantioselective C(sp³)-H activation.

These experiments led us to elucidate how the chiral mono-*N*-protected amino acid ligand coordinates with the Pd(II) center and to propose a stereomodel as a basis for further development of amino acid-based ligands to control C-H activation (Scheme 10).^{116,120} Through NMR studies and inorganic synthesis of putative coordinating structures, we believe that the mono-*N*-protected amino acid ligand coordinates to Pd(II) with high affinity via a bidentate binding

Scheme 8. (a) Racemic Route to Dimeric Palladacycle 122 from Prochiral Starting Material 120. (b) Initial Hypothesis for Achieving Stereinduction in the C–H Cleavage Step through Use of a Chiral Carboxylate Ligand¹¹⁶Table 1. Early Attempts to Use Chiral Carboxylate Ligands in the Catalytic Enantioselective C(sp²)-H Activation of 125¹¹⁷





Entry	Ligand	Yield (%) ^a	ee (%) ^b	Entry	Ligand	Yield (%) ^a	ee (%)
1		62	5	6		78	20
2		75	20	7		63	15
3		65	20	8 ^{b,e}		9	21
4		87	6				
5 ^{c,d}		25	0				

^aIsolated yield. ^bee determined by HPLC using a chiral stationary phase. ^cYield determined by ¹H NMR. ^d10 mol % of ligand, 85 °C. ^e15 mol % of ligand, 85 °C.

mode (153). Upon complexation, the nitrogen atom becomes stereogenic, and the sterically encumbered protecting group is forced *trans* to the bulky alkyl substituent through a “gearing” effect. The pyridine of the substrate coordinates *trans* to the carbonyl moiety of the ligand, situating the substrate’s *ortho*-C–H bonds in close proximity to the acetate group.


Computational studies performed in collaboration with the Musaeu group suggest that at this stage, activation of the N–H bond by internal acetate takes place, opening a coordination site at the Pd(II) center.¹²⁰ Concerted metalation/deprotonation via external base delivers the palladacycle with concomitant formation of a new stereocenter at the methine carbon atom.












Table 2. Catalytic Enantioselective C(sp²)-H Activation of 125 with Chiral Cyclopropane Amino Acid Ligands¹¹⁶

Entry	Ligand	Yield (%) ^a	ee (%) ^b	Entry	Ligand	Yield (%) ^a	ee (%) ^b
1		46	46 (-)	3		63	42 (+)
2		71	41 (+)	4		58	<5 (-)


^aIsolated yield. ^bee determined by HPLC using a chiral stationary phase.

Table 3. Catalytic Enantioselective C(sp²)-H Activation of 125 with Mono-N-protected Amino Acid Ligands¹¹⁶




Entry	Ligand	Yield (%) ^a	ee (%) ^{b,c}	Entry	Ligand	Yield (%) ^a	ee (%) ^{b,c}
1		60	52	7		63	90
2		69	70	8		88	79
3		47	85	9		89	85
4		60	80	10		87	85
5		65	88	11		91	87
6		65	88				

^aIsolated yield. ^bee determined by HPLC using a chiral stationary phase. ^cAbsolute configuration of 127 determined by single-crystal X-ray diffraction.¹²⁰


Scheme 9. Catalytic Enantioselective C(sp³)-H Activation of 151¹¹⁶

Consistent with our experimental results, pretransition-state coordination structure 155 and transition state 156, which lead to 157, containing a newly formed (*R*)-stereocenter, are computed to be lower in energy than 159 and 160, which lead to 161, containing a newly formed (*S*) stereocenter. The precise steric and electronic effects of the amino acid ligand and its influence on the relative energies of key intermediates 154, 155, 158, and 159 and transition states 156 and 160 are topics of continuing investigation. After enantioselective C-H activation, palladacycle 157 then reacts with the boronic acid via a transmetalation/reductive elimination sequence, generat-

Scheme 10. Working Stereomodel, As Supported by Computational Studies.^{116,120–122}

ing Pd(0), which is reoxidized by Ag(I) and re-enters the catalytic cycle (Scheme 7).


Our next objectives were (1) to broaden this chemistry to substrate- and product classes more relevant to organic synthesis and (2) to test the viability of other catalytic manifolds for C–C and C–heteroatom bond formation. Drawing on our extensive experience using carboxylic acid-containing substrates (see Scheme 4a),⁷⁷ we questioned whether we could employ mono-*N*-protected amino acid ligands to induce enantioselectivity in a Pd(II)-catalyzed aerobic C–H olefination reaction of phenylacetic acids that we had recently discovered in our laboratory.¹²³ Consistent with our philosophy for reaction development, we envisioned at the outset that the carboxylate directing group^{82,86,124–127} would be versatile for further functionalization, and in this way, we would have a unique method of constructing enantio-enriched chiral building blocks containing densely substituted stereocenters. Gratifyingly, exposing sodium 2,2-diphenylpropanoate (**162**) to our parent reaction conditions in the presence of Boc-Ile-OH (**164**) led to the product in high yield and with high ee (Scheme 11).¹²³ Interestingly, the combination of the preformed sodium salt and KHCO₃ as the base were critical; alternative combinations led to lower yield and/or ee.

Scheme 11. Enantioselective C(sp²)-H Olefination of Diphenylacetic Acid Derivative **162**¹²¹

We were able to confirm the absolute configuration of one product through single-crystal X-ray diffraction and assign the other products by analogy, including (*R*)-**165**. The observation that this set of products has the same absolute configuration as (*R*)-**127** in Table 3 is consistent with the notion that the two mechanisms of stereoinduction are analogous, with the sodium carboxylate acting like the 2-pyridyl group as an L-type neutral donor during the enantioselective cyclometalation step.

Importantly, this work also established the compatibility of the amino acid ligand scaffold with other elementary steps in catalysis: namely, olefin association, 1,2-migratory insertion, and β -hydride elimination (Scheme 12).

Scheme 12. General Catalytic Cycle for C–H Olefination via Pd(II)/Pd(0) Catalysis



More recently, we targeted another important and underutilized class of starting materials, cyclopropanecarboxylic acids. Encouraged by the high levels of diastereoselectivity that we previously observed in cyclopropane C–H iodination using our chiral oxazoline auxiliary¹¹² and by our preliminary results using amino acid ligands for enantioselective C(sp³)-H activation of *gem*-dimethyl groups (Scheme 9),¹¹⁶ we reasoned that we could develop a catalytic enantioselective method for cyclopropane C–H functionalization.

We viewed this as a potentially powerful and direct means of synthesizing enantiopure *cis* chiral cyclopropanes, which are challenging to access using other methods. Because of the partial π -character of the C–C bonding system in cyclopropanes (Walsh orbital analysis), the cleavage of cyclopropyl C–H bonds is known to be more facile than that of completely unactivated C(sp³)-H bonds, and one would expect the pretransition state coordination structure to be rigid, conceivably allowing C–H activation to take place under mild conditions. To facilitate C(sp³)-H cleavage we installed an acidic *N*-arylamide directing group (see Scheme 4b),^{128,129} enabling Pd(II)-catalyzed C–H/R–BX_n cross-coupling at temperatures as low as 40 °C.¹³⁰ Following optimization of a nonstereoselective version of the reaction, we systematically examined a structurally diverse array of amino acid ligands for stereoinduction (Table 4). In our early screening studies, we found that of a wide range of protecting groups examined, the

Table 4. Catalytic Enantioselective C(sp³)-H Activation of Cyclopropanecarboxylic Acid Derivative 166 with Mono-N-protected Amino Acid Ligands¹¹⁶

Entry	Ligand	Yield (%) ^a	ee (%) ^b	Entry	Ligand	Yield (%) ^a	ee (%) ^b
1		55	80	7		51	88
2		42	75	8		44	91
3		47	78	9		50	90
4		47	55	10		48	90
5		31	33	11		40	88
6		47	85	12		47	93

^aYield determined by ¹H NMR analysis of the crude reaction mixture using CH₂Br₂ as an internal standard. ^bee determined by HPLC using a chiral stationary phase.

TcBoc group was most successful, likely due to the large steric bulk and moderate electron-withdrawing character of the Me₂(Cl₃C)- moiety. Of the backbones tested, those that had aromatic rings like phenylalanine, tyrosine, and tryptophan were superior. Within the TcBoc-protected series, TcBoc-Phe-OH (173) proved to be optimal (entry 6). Increasing the steric demands of the alkyl substituents on the TcBoc group and tuning the electronic properties on the arene led to the identification of ligand 179 (entry 12). Though the yield was low, this problem could be remedied by adding the reagents consecutively in two batches, giving up to 81% yield, 91% ee for the model reaction of 166 to 168. This reaction was found to be compatible with aryl-, vinyl-, and alkylboron reagents and could tolerate a number of different substituents at the α -position.

3.2.3. Position-Selective C-H Activation. In light of mounting evidence that the mono-N-protected amino acid ligands were intimately involved in the C-H cleavage step of the reactions described above, we then examined the use of these ligands for a conceptually related challenge, controlling the positional selectivity in the C-H functionalization of substituted arenes.

While exploring the substrate scope of nonstereoselective Pd(II)-catalyzed *ortho*-C-H olefination of phenylacetic acids¹²³ we became interested in the functionalization of 2-(3-methoxy-5-methylphenyl)acetic acid 180 because of its potential as a synthon in natural products synthesis (Table 5). When we attempted the reaction under our parent reaction conditions, we observed an intractable 1.4:1 mixture of the two possible products, 182-A and 182-B, in 68% conversion. Upon surveying a collection of ligands, we ultimately found that the application of For-Ile-OH (189) furnished 182-A and 183-B in a 20:1 ratio without a substantial decrease in conversion (45%) (entry 12). Simply increasing the catalyst loading to 7% lifted the conversion to 75% with identical levels of selectivity.

Though a precise mechanistic rationalization of this change in selectivity remains to be determined, one plausible explanation is that the ligand causes a change in the mechanism of C-H cleavage. In the absence of ligand, electron-rich and electron-neutral phenylacetic acids are highly reactive, while substrates with electron-withdrawing groups are unreactive, a pattern consistent with an electrophilic palladation mechanism for C-H cleavage.¹²³ In the presence of certain amino acid ligands, electron-deficient phenylacetic acids are preferentially

Table 5. Position-Selective C–H Olefination of **180** Using Mono-*N*-protected Amino Ligands

Entry	Ligand	Conv. (%) ^a	A:B ^a	Entry	Ligand	Conv. (%) ^a	A:B ^a
1	---	68	1.4:1	7		50	3:1
2		17	3:1	8		16	6:1
3		17	5:1	9		24	13:1
4		23	6:1	10		16	5:1
5		24	7:1	11		23	10:1
6		27	8:1	12		45 (75) ^b	20:1

^aThe conversion and A:B ratio were determined by ¹H NMR of the crude reaction mixture. ^bValue in parentheses represents the conversion using 7 mol % of Pd(OAc)₂ and 14 mol % of For-Ile-OH under otherwise identical conditions.

reactive, which is more consistent with a CMD mechanism (vide infra).⁷⁶ It is possible that the drastic change in product distribution is a reflection of the difference in relative rates under the two mechanistic regimes. An alternative explanation is that the ligand could merely be enhancing the steric demands around the metal center, favoring reaction away from the bulkier methyl group.^{131,132}

3.2.4. Ligand-Accelerated C–H Activation. The ability of mono-*N*-protected amino acids to control enantioselectivity and positional selectivity serves as evidence that the ligand is coordinated to Pd(II) during the C–H cleavage event and is influencing the corresponding transition state energies. Accordingly, we wondered whether these ligands could promote C–H activation for otherwise unreactive substrates.^{76,123,133} In our C–H olefination of phenylacetic acids,¹²³ we had found that electron-poor substrates gave low yields even after extended reaction times (up to 48 h). For instance, 2-(trifluoromethyl)phenylacetic acid (**190**) was found to give <15% yield of **191** after 48 h under our standard reaction conditions. We began investigating the effect of mono-*N*-protected amino acids (Table 6), and in order to measure the kinetic reactivity of the various ligands, we assayed the conversion at 20 min and 2 h. Of the Boc-protected amino acids tested (entries 2–6), valine was found to be the best backbone (entry 6), giving 46% conversion after 20 min and

nearly quantitative conversion after 2 h. Optimizing the structure of the protecting group revealed that acetyl was the most reactive (entries 7–10). We then reoptimized the backbone structure using acetyl as the protecting group, leading to the discovery that Ac-Ile-OH (**188**) gave the highest kinetic reactivity for this transformation (entry 12).


With this new ligand-accelerated catalytic system, the substrate scope could be dramatically expanded to include electron-poor phenylacetic acids bearing CF₃ or NO₂ substituents, and it gave improved activity with electron-rich and -neutral substrates.^{76,133} By employing amino acid ligands, hydrocinnamic acids, which contain a more distal carboxylate directing group (via a seven-membered palladacycle) were also found to be reactive in C–H olefination for the first time (Scheme 13).^{123,133} Using Ac-Ile-OH (**188**) as the ligand, exceptionally high catalytic activity was observed with phenylacetic acid substrates. With **190**, for example, catalyst loadings as low as 0.2 mol % could be used to give >90% conversion (>450 TON), which is among the most efficient aerobic C–H olefination reactions reported to date.^{134,135} Because of the stability of the catalyst, the reaction can also be run with air (1 atm), rather than O₂ (1 atm), as the sole terminal oxidant, albeit with extended reaction times (87% conv. of **190** to **191** after 48 h with 1 mol % of catalyst). This transformation can readily be applied in practical settings, for instance, in the direct

Table 6. Ligand-Accelerated C–H Olefination of 2-(Trifluoromethyl)phenylacetic Acid (190) Using Mono-*N*-protected Amino Ligands⁷⁶

Conv. (%) ^{a,b}				Conv. (%) ^{a,b}			
Entry	Ligand	20 min	2 h	Entry	Ligand	20 min	2 h
1	---	<5 ^c	7 ^c	7		21	<i>n.d.</i> ^d
2		28	88	8		2	<i>n.d.</i> ^d
3		38	98	9		31	<i>n.d.</i> ^d
4		37	98	10		57	>99
5		37	92	11		55	>99
6		46 ^c	98 ^c (96)	12		72 ^c	>99 (96)

^aThe conversion was determined by ¹H NMR of the crude reaction mixture. ^bIsolated yield shown in parentheses. ^cAverage of three trials. ^d*n.d.* = not determined.

Scheme 13. Ligand-Promoted Diolefination of Hydrocinnamic Acid (197)^a



^aThe mono-olefinated product (35% conversion) was also observed by ¹H NMR but was not isolated.¹³³

functionalization of NSAIDs (e.g., ibuprofen, ketoprofen, and naproxen). Moreover, it has been used as the key convergent step in the synthesis of (+)-lithospermic acid (see Scheme 3b for retrosynthesis).⁸¹

Mechanistic studies performed in our laboratory⁷⁶ and in collaboration with the Blackmond group¹³⁶ suggest that the observed rate enhancement in the presence of amino acid ligand is due to acceleration of the C–H cleavage step, possibly due to a change in the mechanism of C–H cleavage from electrophilic palladation (in the absence of ligand) to a CMD


pathway (with ligand), as described previously.¹²⁰ Through kinetic studies, off-cycle catalyst reservoirs from the olefin and the product binding to the active Pd(II) catalyst have been identified, which decrease the reaction rate but may protect against catalyst deactivation.¹³⁶

In recent work, we were able to transpose the ligand-enhanced reactivity of phenylacetic acids to the first example of ligand-accelerated Pd(II)-catalyzed C(sp²)-H/Ar–BX_n cross-coupling. By using mono-*N*-protected amino acid ligands in conjunction with Ag₂CO₃ as a stoichiometric reoxidant, we developed a new protocol for the functionalization of phenylacetic acid substrates that offers shorter reaction times, improved substrate scope, and higher yields compared to our previous method.¹³⁷ Importantly, this work established that mono-*N*-protected amino acids are compatible with the transmetalation, reductive elimination, and reoxidation steps in ligand-accelerated C–H/R–M cross-coupling.

In addition to carboxylic acid substrates, we have found that other key classes of widely available, synthetically useful starting materials can undergo Pd(II)-mediated C–H functionalization promoted by amino acid ligands. This catalytic system has proven to be instrumental for substrates containing weakly coordinating functional groups, where in the absence of a highly reactive catalyst, cyclopalladation is sluggish. Contemporaneous to our initial studies in ligand-accelerated catalysis

with phenylacetic acids, we also began exploring the effects of amino acid ligands in alcohol-directed reactions.^{138,139} We viewed aryl and alkyl alcohols as ideal starting materials in C–H functionalization because of their abundance and prevalence in desirable product motifs.^{140–143} With the exception of examples using phenol-type directing groups,^{144,145} hydroxy-directed C–H functionalization using Pd catalysts had not been reported prior to our work. Through careful optimization of the reaction conditions, we were able to achieve *ortho*-C–H olefination¹³⁸ and carbonylation¹³⁹ of phenethyl alcohols (Scheme 14). The

Scheme 14. Ligand-Promoted *ortho*-C–H Functionalization of Phenethyl Alcohols: (a) Olefination¹³⁸ and (b) Carbonylation.¹³⁹ (c) General Catalytic Cycle for C–H Carbonylation via Pd(II)/Pd(0) Catalysis; Nu = Generic Nucleophile



use of (+)-Men-Leu-OH as a supporting ligand was critical for obtaining synthetically useful yields; in both transformations depicted in Scheme 14, <40% yield was observed in the absence of ligand. For both reactions, selection of a noncoordinating


solvent was key due to the weak coordinative affinity of alcohols for Pd(II) centers in comparison to traditional directing groups. As part of a collaboration with Pfizer, we applied amino acid ligands to achieve high reactivity in *ortho*-C–H olefination directed by sulfonamides,¹⁴⁶ privileged pharmacophores in medicinal chemistry (Scheme 15a).¹⁴⁷ With this result as the entry point, we developed a collection of categorically distinct, divergent C–H functionalization reactions for synthesizing analogue libraries of drugs like Celebrex. Using ether directing groups,¹⁴⁸ another class of prevalent, weakly coordinating functional groups, we were able to develop an *ortho*-C–H olefination reactions in the presence of Ac-Gly-OH and hexafluoro-2-propanol (HFIP) (Scheme 15b).¹⁴⁹




Finally, we again turned to amino acid ligands during our efforts to develop a *meta*-selective Pd(II)-catalyzed C–H olefination reaction of hydrocinnamic acid derivatives (Scheme 16).¹⁵⁰ Our reaction design centered on the use of an end-on template, which was inspired by gas-phase studies of Schwarz in which metal centers were observed to adopt a linear coordination geometry with nitriles, enabling activation of remote C–H bonds.^{151,152} To reduce this idea to practice, a simple nitrile-containing amide auxiliary was installed on a hydrocinnamic acid. End-on coordination of Pd(II) to the nitrile through the nitrogen atom situates the catalyst directly over the desired C–H bond. In this system, the combination of Ac-Gly-OH (207) and HFIP were found to be critical for high activity in C–H activation.

3.2.5. Outlook. This collection of results, which we have gathered during the past five years, demonstrates that mono-*N*-protected amino acid ligands effect enantioselectivity, positional selectivity, and rate enhancement in Pd(II)-catalyzed C–H functionalization. It establishes that amino acid ligands are bound to the Pd(II) center during C–H cleavage and are capable of controlling the mechanism (and thus activation energy) of C–H cleavage. Amino acid ligands have been proven to control reactivity with many different substrate classes containing diverse directing groups, and they are compatible with a variety of elementary steps found in the catalytic cycles of Pd(II)-catalyzed C–H/R–BX_n cross-coupling, C–H olefination, and C–H carbonylation.

An advantage of mono-*N*-protected amino acid ligands for *directed* Pd(II)-catalyzed C–H functionalization is the non-symmetrical N,O-mixed donor system, which desymmetrizes the remaining two sites on the square planar Pd(II) center,

Scheme 15. Ligand-Promoted *ortho*-C–H Olefination of (a) Benzylsulfonamides (204)¹⁴⁷ and (b) Phenethyl Ethers (206)¹⁴⁹



Scheme 16. Ligand-Promoted *meta*-C–H Olefination of Hydrocinnamic Acid Derivative 209 Using an End-On Template Approach¹⁵⁰Scheme 17. (a) Initial Discovery of Pd(II)-Mediated C–H Olefination Using Stoichiometric Complex 211 by Fujiwara and Moritani in 1967.¹⁶⁰ (b) Distribution of Positional Isomers (*o*-, *m*- and *p*-215) in the Stoichiometric Pd(II)-Mediated C–H Olefination of Toluene (214)¹⁷⁰Scheme 18. Representative Examples of Tactics for Controlling Positional Selectivity in C–H Olefination: (a) Intermolecular *ortho*-C–H Olefination of *N*-(*p*-Tolyl)acetamide (216) by de Vries and van Leeuwen;¹⁷⁴ (b) Intramolecular C–H Olefination of *N*-Methylindole Substrate 219 by Stoltz¹⁷³

such that the directing group can favorably bind at one and the C–H bond can react at the other. Another benefit is their practical convenience; many are commercially available and others can easily be prepared through modular synthesis. They are air- and moisture-stable and simple to employ in screening libraries. Furthermore, the mono-*N*-protected amino acid ligand scaffold offers several sites for tuning, which was pivotal for us in adapting the steric and electronic properties of the ligand to the unique demands of each of the transformations depicted above. Indeed, much to our delight, other groups have already found success employing them to develop new Pd(II)-catalyzed C–H activation reactions.^{153–159} We further anticipate the use of amino acid ligands for metal catalysis will be applied to develop hybrid transition metal/enzyme catalysts for C–H activation.

3.3. 2,6-Disubstituted Pyridine Ligands. **3.3.1. Background.** Simple, monofunctionalized arenes constitute a class of abundant chemicals with rich synthetic potential. An attractive approach to map these substrates onto more complex multisubstituted arene building blocks would be position-selective C–H functionalization to form new C–C bonds. Since the early work of Fujiwara and Moritani in the late 1960s (Scheme 17),^{160,161} C–H olefination of simple arenes has been extensively studied.^{134,135,162–168} Despite these efforts, several problems have persisted: (a) the need for high molar excess of arene substrates, which are often used as solvent (b) limited positional selectivity, (c) the need for high catalyst loadings and stoichiometric metal reoxidants, and (d) low reactivity with electron-poor substrates.^{169,170} Through extensive efforts during the past several years, methods for position-selective C–H olefination of electron-rich heterocycles^{171–173} and

Table 7. Ligand-Promoted C–H Olefination of 1,3-Bis(trifluoromethyl)benzene (**222**)¹⁷⁵

Entry	Ligand	Yield (%) ^a	Entry	Ligand	Yield (%) ^b
1	---	<2	6		<5
2		<5	7		10
3		<2	8		24 (<i>E</i> : <i>Z</i> = 7.8:1) ^b
4		<5	9		52 (<i>E</i> : <i>Z</i> = 5.9:1) ^b
5		<2			

^aThe yield was determined by ¹H NMR of the crude reaction mixture using CH₂Br₂ as an internal standard. ^bThe *E*/*Z* ratio was determined by ¹H NMR of the crude reaction mixture.

substrates containing *ortho*-directing groups^{125,174} have been established (Scheme 18). To realize the full synthetic utility of this approach, however, new strategies to promote reactivity and control selectivity are needed.

We sought to address these problems through the design of an ancillary ligand that could accomplish the following goals: (1) enhance reactivity with electron-poor arenes, ideally using only 1 equiv of substrate, (2) favor selectivity at the *meta* position to offer complementary selectivity to directed methods, and (3) promote reoxidation with molecular O₂^{134,135} at ambient pressure.









In order to realize these goals, there were several obstacles that warranted consideration at the outset. The first potential problem is that the π -systems of electron-deficient arenes are comparatively poor σ -donors, meaning that they may be unable to compete with the majority of L-type ligands for binding to Pd(II). An ideal ligand would be capable of tightly binding to Pd(II) yet would still allow the substrate to bind, even in low concentration. The second challenge, as alluded to above, is that it is generally believed that under standard reaction conditions, Pd(II)-mediated C–H olefination proceeds through an electrophilic palladation mechanism for C–H cleavage, in a similar manner to that of a Friedel–Crafts reactions. To achieve high reactivity with electron-poor arenes, the supporting ligand would need to increase greatly the electrophilicity of the Pd(II) catalyst or bias the C–H cleavage step toward a CMD mechanism. Third, to achieve *meta* selectivity, the catalyst would need to be able to recognize minor differences in the steric and electronic environments of the different C–H bonds, promoting selective C–H cleavage according to these subtle cues. Lastly, the ligand would need to

be compatible with (and ideally would promote) the reoxidation step of Pd(0) to Pd(II) with O₂.


3.3.2. *meta*-Selective C–H Activation of Electron-Deficient Arenes. To begin our investigation, we first attempted to develop conditions to effect Pd(II)-catalyzed C–H olefination of 1,3-bis(trifluoromethyl)benzene (**222**), a highly electron-deficient arene.¹⁷⁵ Due to steric effects, we presumed that this substrate would give a single product in C–H olefination, which would allow us to focus on reactivity during optimization. Under standard aerobic Fujiwara–Moritani conditions, using Pd(OAc)₂ as the catalyst and **222** as the solvent, we found the reaction gave only trace quantities of the desired product (entry 1, Table 7).

To overcome the low reactivity and lack of catalytic turnover, we first turned to pyridine-based ligands, which have long been known to be beneficial in oxidative Pd(II)-catalyzed transformations, especially those under aerobic conditions.^{176–183} Seminal work by Ferreira and Stoltz established that an electron-deficient pyridine ligand, ethyl nicotinate, greatly improved the yield in an intramolecular C–H olefination reaction of an electron-rich nitrogen heterocycles under aerobic conditions (Scheme 18b).¹⁷³ Other groups have also successfully employed pyridine ligands in C–H functionalization reactions along a Pd(II)/Pd(0) catalytic cycle.^{168,182,184–188} In our study, we first tested pyridine (py, **46**) and 2,6-lutidine (**47**), but these ligands gave low yields (entries 2 and 3). At this stage, we speculated that the pyridine ligands were bis-ligated to the Pd(II) center (e.g., Pd(II)-(py)₂OAc)₂ and that displacement of a ligand with the electron-poor arene **222** was unfavorable, owing to the strength of the Pd(II)–N bond. We hypothesized that the Pd(II)–N bond

Table 8. Substrate Scope of Ligand-Promoted C–H Olefination of Electron-Poor Arenes (Selected Examples)¹⁷⁵

Entry ^a	Product	Yield (%) ^b	<i>m</i> : <i>p</i> ^c	Time (h)	Entry ^a	Product	Yield (%) ^b	<i>m</i> : <i>p</i> ^c	Time (h)
1 ^d		62	---	36	5		72	3.5:1	20
2		68	---	24	6		71	4.9:1	5
3		77	---	24	7		80	3.5:1	56
4 ^d		70	4.0:1	2	8		71	3.3:1	36

^aUnless otherwise specified, the reaction conditions were as follows: alkene (0.6 mmol), arene (2 mL, 20–30 equiv), Pd(OAc)₂ (10 mol %), **229** (20 mol %), Ac₂O (1.0 equiv), O₂ (1 atm), 90 °C. ^bIsolated yield. ^cThe *meta/para* ratio was determined by GC. Reference samples of the pure *meta*- and *para*-isomers were prepared independently via Mizoroki–Heck chemistry. ^dAc₂O (1.5 equiv).

Scheme 19. Sequential C–H Functionalization Route to Tetrasubstituted Arene **240**.¹⁷⁵

could be destabilized by introducing steric bulk at the 2- and 6-positions of the pyridine ring or by substituting the pyridine ring with an electron-withdrawing group; however, 2,6-diisopropylpyridine (**225**), 2,6-di-*tert*-butylpyridine (**226**), and ethyl nicotinate (**220**) all gave low yields (entries 5–7). With these ligands, formation of the corresponding Wacker olefin oxidation product and accompanying precipitation of palladium black (Pd(0)_n) suggested that ligand binding was too weak.

We reasoned that what was needed was a pyridine-based ligand that would be strongly binding yet preferentially monoligating. Our breakthrough came in recognizing that we could achieve this combination of properties by minimizing steric bulk at the carbon atoms immediately adjacent to the nitrogen atom and introducing steric bulk at a more distal position. In this manner, one molecule of ligand would be free to bind Pd(II), but coordination of a second ligand molecule *trans* to the first would be unfavorable because of steric clashing away from the metal center. In other words, the ligands would experience “mutual repulsion” when both bound to the Pd(II) center. To reduce this idea to practice, we prepared and tested ligands **227**, **228**, and **229** and found, much to our delight, that the yield could be improved to 10–52% (entries 7–9). Longer and more branched side chains led to improved yield, with **229** proving to be optimal.



The substrate scope with electron-deficient arenes proved to be quite broad (selected examples shown in Table 8). The reaction was compatible with a variety of electron-withdrawing groups; trifluoromethyl and nitro substituents were well tolerated, as were esters and ketones. 1,3-Disubstituted arenes

bearing two electron-withdrawing groups, reacted selectively at the 5 position (entry 1). 1,4-Disubstituted arenes with one electron-releasing group and one electron-withdrawing group reacted *meta* to the electron-withdrawing group (entry 2). Monosubstituted electron-poor arenes reacted at the *meta*- and *para*-positions with an *m*:*p* ratio typically around 4:1 (entries 4–8). A variety of olefin coupling partners could be used, including those with substitution as the β -position.

As mentioned above, one limitation of traditional non-directed Fujiwara–Moritani reactions is the need to use the substrate as the solvent. This requirement is highly undesirable and inherently limits the substrate scope to simple, low-melting arene starting materials. The high levels of reactivity promoted by **229** allowed us to reduce the arene substrate loading to 5 equiv relative to olefin using EtOAc as the solvent (see Scheme 19 for an example). These findings suggest that with future advances in ligand design, a catalytic system could be devised such that 1 equiv of arene starting material and exclusive selectivity is obtained for functionalization of a single C–H bond.

In an effort to showcase the synthetic utility of this *meta*-selective transformation, we attempted to use it as the first step in a sequential C–H functionalization route¹³³ to build up molecular complexity from *tert*-butyl 4-methylbenzoate (**238**), a commercially available benzoic acid derivative (Scheme 19). Following ligand-promoted C–H olefination, Pd/C-mediated hydrogenation, and *tert*-butyl ester deprotection, free acid **239** was obtained in 66% yield over the three steps. Next, we found that **239** could undergo directed *ortho*-C–H iodination⁸² to

Scheme 20. Proposed Equilibrium Species Based on Structural and Spectroscopic Studies

Scheme 21. (a) Ligand-Promoted Methylene C(sp³)-H Arylation of Butanoic Acid Derivative **244**.¹⁹¹ (b) General Catalytic Cycle for C-H Arylation with Aryl Iodides via Pd(II)/Pd(IV) Catalysis

give differentially functionalized 1,2,4,5-tetrasubstituted arene **240** in 67% yield following conversion to the methyl ester to simplify purification.

To understand the mechanism of this reaction and the role of the 2,6-dialkylpyridine ligand **229**, we studied relevant Pd(II) intermediates by ¹H NMR and X-ray crystallography (Scheme 20). By stirring 2 equiv of ligand **229** and Pd(OAc)₂ in hexanes, we were able to prepare Pd(**229**)₂(OAc)₂ complex **241**, which we characterized by single-crystal X-ray diffraction. The structure shows that the two ligand molecules **229** are approximately coplanar and coordinated *trans* to one another. Consistent with our mutual repulsion hypothesis, the two sets of branching arms lie on opposite sides of the Pd–N–C₂ plane (as expressed in the relative stereochemistry drawn for **241**), and the Pd–N bond length is 0.05 Å longer than the Pd–N bond in Pd(Py)₂(OAc)₂·H₂O.¹⁸⁹ When we dissolved complex **241** in CDCl₃ and monitored the solution by ¹H NMR at 2 and 12 h, we observed formation of a new species consistent with **243** along with free ligand **229**. In contrast, when we performed the same experiments with the pyridine and 2,6-lutidine complexes, Pd(py)₂(OAc)₂ and Pd(**47**)(OAc)₂, we observed that they were stable in CDCl₃ for several days.

On the basis of these data, we propose that the exceptional reactivity observed with ligand **229** is due to its mutually repulsive properties. It is energetically unfavorable for two

molecules of **229** to be bound at Pd(II) simultaneously, which creates an opportunity for the substrate π -system to coordinate to the metal. The high reactivity with electron-poor substrates is consistent with a CMD mechanism for the C–H cleavage step. Furthermore, in line with the precedents mentioned above, the pyridine-type **229** ligand could be effective in part because it stabilizes reduced Pd(0) species and facilitates reoxidation with O₂. In terms of selectivity, the observed product distribution remains somewhat unclear. The lack of reactivity at the *ortho* position with monosubstituted arenes can be attributed to the steric bulk of the catalyst, which would force substituents on the arene away upon association to the catalyst. The precise origin of *meta*-selectivity over *para*-selectivity is possibly due to subtle differences in electronic density or C–H bond acidity between the two positions. Differences between the thermodynamic stability of the developing Pd(II)–C bonds formed via C–H cleavage at the different positions could also contribute to the observed selectivity patterns.

Computational studies performed by Zhang and co-workers found that a CMD mechanism was favorable for the C–H cleavage step in this system and supported the notion that steric interactions between the pyridine ligand and the substrate are dominant in influencing the positional selectivity, with electronic effects playing a minor effect.¹⁹⁰ Their model also

suggests that the ligand dissociates from the metal upon olefin binding, a step that would also be facilitated by a ligand with weaker Pd–N bond strength.

We have recently gone on to apply the mutually repulsive ligand design concept to address a different problem, methylene C(sp³)–H functionalization of aliphatic acid derivatives (Scheme 21a).¹⁹¹ Our goal was to expand the diverse reactivity that we have developed using our weakly coordinating *N*-arylamide auxiliary to methylene C(sp³)–H functionalization. However, methylene C(sp³)–H cleavage is generally sluggish with Pd(II) catalysts in the absence of a strongly coordinating (typically bidentate directing group).^{192,193} Drawing on the lessons from above, we discovered an electron-rich, mutually repulsive 2-alkoxy quinoline ligand **246**, which dramatically promoted methylene C(sp³)–H arylation of **244** along a Pd(II)/Pd(IV) catalytic cycle (Scheme 21b).

3.3.3. Outlook. Generally speaking, mutually repulsive 2,6-disubstituted pyridines represent a ligand scaffold that is carefully tailored for the steric and electronic demands of electron-poor arene substrates that do not contain directing groups. This class of substrates poses unique challenges in Pd(II)-mediated C–H cleavage that informed our ligand design criteria. A benefit of the ligand scaffold is that it has multiple tuning sites. In our case, we adjusted the steric properties to minimize steric crowding near nitrogen and maximizing steric clashing away from nitrogen. The ability to tune both adjacent to and remote from the binding atom were critical. In addition, modular synthesis allowed each of the ligand candidates to be synthesized in a single step, which was vital in the reaction discovery stage. The compatibility of this ligand scaffold with other elementary steps in other catalytic manifolds remains to be examined.


3.4. 2,2'-Bipyridine Ligands. **3.4.1. Background.** Ubiquitous in natural products, pharmaceutical agents, agrochemicals, organic materials, and ligands for coordination chemistry, pyridines (and related azines) represent one of the most common heterocyclic motifs. Owing to the potential synthetic utility of functionalized pyridine derivatives in many different realms of chemistry, metal-catalyzed position-selective pyridine C–H functionalization has been the topic of considerable interest in the past few years.¹⁹⁴ However, owing to its tendency to adopt an *N*-bound coordination mode and its electron-poor nature, pyridine has proven to be a challenging substrate to use in these transformations.

Nonetheless, with metals other than Pd(II), some success has been achieved in position-selective C–H functionalization of pyridines.¹⁹⁴ Representative strategies include the use of low-valent metals (e.g., Ru(0) and Rh(I)) for C2 functionalization facilitated by *N*-chelation,^{195–199} C3/C4-selective borylation using Ir(III) species,^{200–206} Ni(0)/Lewis acid catalysis for functionalization of the C2 or C4 positions,^{207–209} and utilization of *ortho*-directing groups on the pyridine ring with Ru(0) or Pd(0) catalysts.^{210–212}

We were intrigued with the idea of developing position-selective pyridine C–H functionalization methodology using Pd(II) catalysts due to the potential versatility in the functionalization step. In terms of precedents, previously published approaches relied upon minimizing or eliminating the chelation strength of the nitrogen atom through substitution of the pyridine ring. For example, by protecting the nitrogen atom as an *N*-oxide^{213–217} or *N*-iminopyridinium ylide,^{218,219} C2 selectivity has been achieved with Pd(II),

Pd(0), and Cu(I) catalysts (Scheme 22a). Additionally, the introduction of steric blocking groups and/or electron-

Scheme 22. Literature Precedents for Pyridine C–H Functionalization with Pd(II) Catalysts: (a) C2-Selective C–H Olefination of Pyridine *N*-Oxide (248**);²¹⁴ (b) C4-Selective C–H/R–B(OH)₂ Cross-Coupling of 2,3,5,6-Tetrafluoropyridine (**251**)²²⁰**




withdrawing groups has been demonstrated to suppress *N*-chelation; for instance, C4-selective C–H functionalization of 2,3,5,6-tetrafluoropyridine has been achieved using a Pd(II) catalyst (Scheme 22b).^{220,221}

Based on our success in achieving reactivity with electron-deficient arenes, we hoped that we could draw on the lessons learned about the interplay between substrate and ligand at the Pd(II) center to design a ligand scaffold that would accomplish the following: (1) promote pyridine C–H bond cleavage, (2) functionalize the C3 position to offer complementary selectivity to other methodologies, and (3) maintain compatibility with a variety of coupling partners when applied to categorically distinct transformations.

In order to achieve these goals through ligand design, several factors were considered. The first major problem is that pyridine is a strongly binding ligand, tending to adopt an *N*-bound coordination mode with the catalyst, which situates the metal center distal from the target C–H bonds (Scheme 23).

Scheme 23. Postulated Coordination Equilibrium of Pyridine with Pd(II): Unproductive *N*-Bound (Left) and Productive π -Bound (Right)



Pyridine substrate molecules could potentially outcompete the ancillary ligand for binding to the metal center. Second, the pyridine ring is electron-poor, which makes initial coordination of the π -system unfavorable and renders C–H cleavage via an electrophilic palladation mechanism untenable (similar to the situation described with electron-poor arenes in section 3.2). Thus, the ligand, in addition to disfavoring the *N*-bound coordination mode, would need to promote C–H cleavage along a CMD pathway. Third, achieving C3 selectivity would

Table 9. Ligand-Promoted C3-Selective C–H Olefination of Pyridine (py, **46**)²⁴⁶

C1=CC=NC=C1 (**46**, 16 equiv.) + CC(=O)C=C (**181**) $\xrightarrow[\text{air, 12 h}]{\text{DMF, 140 }^\circ\text{C, 10 mol\% Pd(OAc)}_2, \text{13 mol\% Ligand, 0.5 equiv. Ag}_2\text{CO}_3}$ C1=CC=NC=C1C=C(C)C(=O)OCC (**256**)

Entry	Ligand	Yield (%) ^a	C3:C2:C4 ^b	Entry	Ligand	Yield (%) ^a	C3:C2:C4 ^b
1	---	21	5:1:1	7	pyrox, 261	11	6:4:1
2	bipy, 43	50	14:1:2	8	quinox, 45	7	5:1:1
3	257	29	5:1:1	9	phen, 44	87	12:1:1
4	258	55	11:1:1	10	bphen, 262	86	12:1:1
5	DAF, 259	16	3:1:1	11	263	6	1:1:1
6	DAF, 260	27	4:1:1	12	bcp, 264	9	4:4:1

^aThe yield was determined by ¹H NMR of the crude reaction mixture using CH₂Br₂ as an internal standard. ^bThe C3:C2:C4 ratio was determined by ¹H NMR of the crude reaction mixture.

require that the catalyst be sensitive to the steric or electronic properties of the three different positions and would need to be resistant to *N*-coordination-assisted C2 functionalization. Fourth, to accommodate many catalytic transformations along Pd(II)/Pd(0) and Pd(II)/Pd(IV) redox couples, the ligand would need to be compatible with a range of different elementary steps. For example in C–H arylation along a Pd(II)/Pd(IV) catalytic cycle, it should stabilize high-energy Pd(IV) intermediates and be recalcitrant to reductive elimination.

3.4.2. C3-Selective Pyridine C–H Functionalization. At the outset, we hypothesized that a strongly chelating bidentate ligand would be required to compete effectively for binding at the Pd(II) center and to disfavor *N*-bound coordination of the substrate. In particular, we surmised that *N,N*-bidentate ligands might prove to be effective in these pursuits. Bidentate N(sp²),N(sp²)-ligands, such as 2,2'-bipyridines, 2,2'-bis(2-oxazoline)s, 2-(2'-pyridyl)oxazoline, and 2-(2-quinolinyl)-oxazolines, have a rich history in oxidative transformations with Pd(II), both in asymmetric and nonstereoselective reactions.^{42,181,222–229} In the context of Pd(II)-catalyzed C–H functionalization, Eberson probed the effects of pyridine- and 2,2'-bipyridine-type ligands in C(aryl)–H acetoxylation along a Pd(II)/Pd(IV) catalytic cycle, and found that use of 2,2'-bipyridine (bipy, **43**) improved the reactivity and positional selectivity with substituted arenes.^{230–233} Fujiwara employed

1,10-phenanthroline (phen, **44**) as a ligand in the C–H carbonylation of naphthalene along a Pd(II)/Pd(0) catalytic cycle, finding that it gave both improved conversion and better selectivity for the β-position.²³⁴ More recently, bidentate N(sp²),N(sp²)-ligands have been found to be beneficial in a variety of Pd(II)-catalyzed C–H functionalization transformations involving Pd(II)/Pd(0) and Pd(II)/Pd(IV) catalysis.^{43,142,148,235–245}


We envisioned that by using a bidentate *N,N*-ligand, we could enhance dissociation of the *N*-bound substrate both sterically and electronically (Scheme 23). The ligand would potentially increase the steric demands of *N*-bound coordination due to unfavorable edge-to-edge or edge-to-face interactions and could also exert a kinetic *trans*-labilizing effect through electronic induction (vide infra). By promoting dissociation of the *N*-bound substrate, the catalyst would facilitate high effective concentration of reactive pyridine molecules around the Pd(II) center. The dissociated substrate would possess various orientations with respect to the Pd(II) catalyst, and under these conditions, C3-selective C–H cleavage could take place upon assembly of an intermediate with the appropriate orientation between the pyridine π-system and the Pd(II) center.

We began by studying Pd(II)-catalyzed C3-selective C–H olefination of pyridine (py, **46**) (Table 9)²⁴⁶ with the long-term aim of developing a wide range of C–C and C–heteroatom

Table 10. Substrate Scope for Ligand-Promoted C3-Selective C–H Olefination of Pyridines (Selected Examples)²⁴⁶

Entry ^a	Product	Yield (%) ^b	C3:C2:C4 ^c	Entry ^a	Product	Yield (%) ^b	C3:C2:C4 ^c
1	 265	54	---	5	 C3/C2/C4-256	73	12:1:1
2	 C3/C2/C4-266	61	10:2:1	6	 C2/C3/C4-269	68	7:1:1
3	 C3/C2/C4-267	63	30:1:1	7	 C2/C3/C4-270	57	14:1:1
4	 C3/C2-268	44	3:1:0	8	 C2/C3/C4-271	45	30:3:1

^aUnless otherwise specified, the reaction conditions were as follows: alkene (0.5 mmol), pyridine substrate (16 equiv), Pd(OAc)₂ (10 mol %), phen (**44**) (13 mol %), Ag₂CO₃ (0.5 equiv), DMF (1 mL), air (1 atm), 140 °C, 12 h. ^bIsolated yield of C3-substituted product. ^cThe C3:C2:C4 ratio was determined by ¹H NMR of the crude product mixture prior to purification.

Table 11. Substrate Scope for Ligand-Promoted C3-Selective C–H Arylation of Pyridines (Selected Examples)²⁴⁸

Entry ^a	Product	Yield (%) ^b	C3:C2:C4 ^c	Entry ^a	Product	Yield (%) ^b	C3:C2:C4 ^c
1	 C3/C2-275	58	15:1:0	5	 C3/C2/C4-279	82	28:1:1
2	 C3/C2-276	65	3:1:0	6 ^f	 C2/C4-280	90	15:0:1
3 ^e	 277	88	---	7	 C2/C3/C4-281	80	---
4 ^e	 278	70	---	8 ^f	 C2/C3-282	73	9:1:0

^aUnless otherwise specified the reaction conditions were as follows: aryl iodide (0.5 mmol), pyridine (3 mL), Pd(OAc)₂ (5 mol %), phen (**44**) (15 mol %), Cs₂CO₃ (3.0 equiv), 140 °C, 48 h. ^bIsolated yield of C3-substituted product. ^cThe C3:C2:C4 ratio was calculated from the isolated yields for all product isomers. ^dPyridine substrate (60 equiv) and DMF (mL). ^eThe corresponding ArBr was used in place of the ArI.

bond-forming reactions. We elected to use an excess of substrate to ensure high concentration around the Pd(II) catalyst. In the absence of an ancillary ligand, using ethyl acrylate (**181**) as the limiting reagent in the presence of Ag₂CO₃ (0.5 equiv) and Pd(OAc)₂ (10 mol %) in DMF, we observed 21% yield of **256** in a 5:1:1 C3:C2:C4 mixture (entry

1). We reasoned that at such high concentration of pyridine in solution, it was likely that during the C–H cleavage step, at least two pyridine molecules were bound through nitrogen and were serving as ligands, suggesting that the pyridine moiety had approximately the appropriate donor strength for promoting the reaction. In accordance with our rationale for ligand design,

Scheme 24. Expedient Synthesis of (±)-Preclamol Using C3-Selective C–H Arylation





Figure 7. Relevant examples of Pd(II)(Py)_nL_m complexes that have been characterized by X-ray diffraction.^{189,250–252}

we turned our attention to bidentate N,N-ligand scaffolds, surveying a broad range of different structures. We found that 2,2'-bipyridine ligands had a modestly positive effect on yield but gave substantially higher selectivity (entries 2–6). Mixed N,N-donors like pyrox (entry 7) and quinox (entry 8) were ineffective, potentially because the oxazoline moiety is too electron-donating. 1,10-Phenanthroline-based ligands, which are less conformationally flexible and thus stronger binding, proved to be optimal in terms of combining high yield and selectivity (entries 9 and 10).

Generally speaking, other methodology for C3- and C4-selective pyridine C–H functionalization has suffered from restricted substrate scope. We were thus delighted to find that under our optimized reaction conditions for C3-selective C–H olefination, a wide range of azines could be efficiently olefinated, generally with high selectivity for the C3 position (selected examples shown in Table 10). Both electron-releasing (entries 2 and 3) and electron-withdrawing substituents (entries 1 and 3) were tolerated on the pyridine ring, and quinoline was also found to be compatible (entry 4). Several different functional groups were accommodated on the olefin as well (entries 5–8). Under the optimal conditions, a large excess of substrate was required for high yield; however, the compatibility of the reaction with DMF as a solvent nonetheless allowed for the use of substrates that were solids at room temperature.

Having established high reactivity and C3-selectivity in C–H activation using the [Pd(II)-phen]-catalyst,²⁴⁷ we subsequently sought to test other coupling partners and catalytic manifolds with this system. In particular, owing to the importance of heterobiaryls in medicinal chemistry and agrochemistry, our next target was a C3-selective C–H arylation reaction with aryl iodides via Pd(II)/Pd(IV) catalysis. Starting with conditions similar to those we had developed for C–H olefination, we were ultimately able to devise a protocol for C–H arylation with aryl iodides and bromides.²⁴⁸ Through the use of Cs₂CO₃, which presumably regenerates a catalytically active Pd(II) species from PdI₂ or PdI(OAc), we were able to avoid Ag(I) salts, which are often employed as halide scavengers in arylation reactions. To achieve high yields, we found that a large excess

of pyridine substrate was required (generally used as solvent). The scope of the reaction was quite broad, tolerating a vast array of substituted aryl bromides and iodides,²⁴⁹ as well as a variety of azines (Table 11). Importantly, these results established the viability of this catalyst/substrate combination in supporting multiple coupling partners and elementary steps during the catalytic cycle.

To demonstrate the practicality of this transformation, we developed a concise route to the preclinical drug (±)-preclamol (286), a partial dopamine receptor agonist, using our C3-selective C–H arylation on gram-scale as the first step (Scheme 24). Following construction of the carbon skeleton, a three-step sequence of N-alkylation, hydrogenation, and deprotection yielded the final product.

While a detailed mechanistic model for these reactions remains the subject of active investigation in our laboratory, there are nonetheless several pertinent observations that we would like to highlight here to help guide future analysis. First, in both the pyridine C–H olefination and arylation reactions, large primary kinetic isotope effects were observed $k_H/k_D \geq 4.0$, suggesting that a CMD mechanism, rather than an electrophilic palladation mechanism, may be operative. (At this stage, however, we cannot rule out the possibility of electrophilic palladation with rate-limiting deprotonation.) Moreover, with the C–H arylation reaction, we have observed that pyridine reacts at a higher rate than benzene, despite being far less electron-rich, which is consistent with a CMD mechanism. This observation is also potentially consistent with the hypothesis that initial N-coordination of pyridine increases the molarity of reactive substrate around Pd(II) to facilitate C–H cleavage.

Several relevant solid-state structures had also been characterized by other groups prior to our work (Figure 7),^{189,250–255} analysis of which sheds some light on the role of the ligand in this reaction. Qualitatively, there seems to be more strain in the Pd(phen)(py)₂(PF₆)₂ (289) and Pd(bipy)(py)₂(PF₆)₂ (290) structures compared to the Pd(py)₄(BF₄)₂ structure (288). In the Pd(py)₄(BF₄)₂ structure, adjacent pyridine rings are all oriented face-to-face to one another and perpendicular to the Pd(II) square plane. In contrast, in the bipy- and phen-ligated structures, there are two enforced and

sterically unfavorable edge-to-face interactions between the ligand and the pyridine substrate, which would potentially make pyridine dissociation more favorable. It should be noted, however, that across structures 287–290, the [Pd(II)–py] bond lengths remain nearly constant (all within 0.011 Å).

Alongside steric considerations, the electronic influence of the ligand also warrants discussion. Generally speaking N- and N,N-ligands are known to be relatively weak *trans*-labilizing ligands (though still stronger than OAc[−]).^{41,256,257} In the case of our reaction system, however, because dissociation of the bidentate ligand will be prohibitively slow compared to dissociation of pyridine, the effective *trans* influence of the phen ligand on N-bound pyridine could still be substantial.

The solution dynamics of complexes of the type shown in Figure 7 have also been studied, and pyridine-dissociation mechanisms mediated by coordinating solvents and acetate anion have been reported.^{252,255} It would seem likely that these mechanisms act in concert with the steric and electronic influences of the ligand in promoting the dissociation of N-bound pyridine to allow for population of the π -bound state at elevated temperatures (Scheme 23). Beyond its role in assisting the assembly of the active pretransition state coordination structure, the ligand could have other beneficial roles in the catalytic cycle, such as promoting reoxidation of reduced Pd(0) species in the C–H olefination reaction or stabilizing Pd(IV) species in the C–H arylation reaction.

The strong preference for C–H cleavage at the C3 position is not entirely clear at this time. The C3 position is expected to be the most electron-rich, so one possibility is that initial π -coordination and subsequent CMD is consequently more favorable. There are other explanations that are also consistent with the observed selectivity trends. Recent computational studies of C(heteroaryl)–H cleavage selectivity patterns via CMD with a related complex, (Ph)(PMe₃)Pd(OAc), have been performed by Gorelsky using distortion–interaction analysis, and with pyridine, C3 selectivity was found to be favorable on the basis of the proposed CMD mechanism.²⁵⁸ Ess and co-workers have studied the same system using density functional calculations and have correlated the CMD transition state energy to the thermodynamic stability of the developing Pd(II)–Ar bond, and have found that this too is in agreement with selectivity at the C3 position for pyridine.²⁵⁹ While these findings are intriguing, caution should be taken in extrapolating the data from this more electron-rich (Ph)(PMe₃)Pd(OAc) system to the [Pd(II)–phen] catalyst used in our studies; further studies are needed to evaluate whether the same trends hold in both cases.

Recently, our group and others have gone on to apply this [Pd(II)–phen] catalytic system to effect selective C–H arylation of other classes of biologically active azaheterocycles, including pyrazoles and 1*H*-indazoles.²⁶⁰ Importantly, we were also able to demonstrate the practical utility of this procedure in a concise total synthesis of the natural product nigellidine hydrobromide.

3.4.3. Outlook. 2,2'-Bipyridines, and N(sp²),N(sp²)-ligands more generally, have a rich history in Pd(II) catalysis and are now beginning to be used to address problems in Pd(II)-catalyzed C–H functionalization. One of the strengths of this ligand scaffold is that there are multiple tuning sites to adjust the steric and electron properties proximate and distal to the nitrogen donor atoms. Additionally, the ground-state torsion angle (and thus the binding strength) of the biaryl backbone can further be adjusted. Each of the two donor sites can be

tuned individually, offering the opportunity to desymmetrize the two remaining sites on the Pd(II) square plane. Many ligands within this class are commercially available and others can be prepared in a small number of steps through modular synthesis, which facilitates ligand screening and reaction optimization. This ligand scaffold was demonstrated to be compatible with the elementary steps of C–H olefination and C–H arylation along Pd(II)/Pd(0) and Pd(II)/Pd(IV) catalytic manifolds, respectively.

Despite being a strong chelator to Pd(II), this ligand scaffold still allows substrate coordination and subsequent C–H activation. In our case, the strong bidentate coordination seems to be key for facilitating the dissociation of N-bound pyridine, leading to the productive π -bound assembly. Through this interplay, high effective molarity of pyridine can be maintained around the catalyst, allowing high reactivity and selectivity to be achieved, despite the electron-poor nature of the arene. Further optimization of the catalyst to achieve high yield with 1 equiv of pyridine substrate relative to olefin is the next challenge that our group is pursuing.

4. CONCLUSIONS

Ligand development has been a principal driving force in the rapid maturation of the field of homogeneous catalysis during the past several decades. In both asymmetric and non-stereoselective reactions, privileged ligand scaffolds have played leading roles in the discovery of a wide range of catalytic reactions with little to no mechanistic similarity. Nonetheless, each combination of substrate, reaction type, and catalyst poses a unique set of requirements for the ligand (steric and electronic properties, denticity, etc.) that are difficult to predict from first principles. For this reason, ligand scaffolds that are tunable and modular are particularly useful because the ligand can rapidly be adjusted to meet the challenge at hand.

In our own research in Pd(II)-catalyzed C–H functionalization, we have made the discovery, development, and application of suitable ligands for controlling C–H cleavage a major priority. We view ligand development as the key to meeting our long-term goals of engineering reactions to convert feedstock chemicals into value-added intermediates through selective and efficient C–H functionalization to form any desired C–C or C–heteroatom bond. In this review, we have discussed the challenges that we faced in trying to develop catalytic systems with high reactivity and selectivity with several distinct substrate classes, and we have examined the ligand design considerations at play in each reaction. Using mono-*N*-protected amino acid ligands, we were able to achieve enantioselectivity, positional selectivity, and rate enhancement with substrates containing directing groups. With 2,6-disubstituted pyridine ligands we were able to achieve *meta*-selective C–H olefination of electron-poor arenes under aerobic conditions and promote methylene C(sp³)–H functionalization of aliphatic acid derivatives. Finally, with 2,2'-bipyridine ligands, we successfully developed C3-selective C–H olefination and arylation of pyridines and related azines. All three ligand classes have proven to be compatible with both Pd(II)/Pd(0) and Pd(II)/Pd(IV) catalysis. Across these case studies, our success in achieving unprecedented levels of reactivity and selectivity came by tailoring the ligand structure to the substrate properties, enabling a productive interplay at the Pd(II) center. In this work, identifying an optimal ligand typically stemmed from a dialogue between theory and empiricism, requiring

robust ligand scaffolds that are compatible with a multitude of catalytic transformations and possess several sites for tuning.

Moving forward, ligand development will play an increasingly important role in realizing the untapped synthetic potential of Pd(II)-mediated C–H functionalization, and as a central challenge in the field, it continues to stimulate and inspire us.

AUTHOR INFORMATION

Corresponding Author

*E-mail: yu200@scripps.edu.

Notes

The authors declare no competing financial interest.

Biography



Jin-Quan Yu received his B.Sc. in Chemistry from East China Normal University, M.Sc. from the Guangzhou Institute of Chemistry, and Ph.D. from the University of Cambridge (with Prof. J. B. Spencer). Following time as a Junior Research Fellow at Cambridge, he joined the laboratory of Prof. E. J. Corey at Harvard University as a postdoctoral fellow. He then began his independent career at Cambridge (2003–2004) before moving to Brandeis University (2004–2007) and finally to TSRI, where he is currently the Frank and Bertha Hupp Professor of Chemistry.

ACKNOWLEDGMENTS

This work was made possible by financial support from TSRI, the NSF (CHE-1011898), the NIH (NIGMS, 1 R01 GM084019-04; NIGMS 1 R01 GM102265-01), NSF under the CCI Center for Stereoselective C–H Functionalization (CHE-1205646), Bristol-Myers Squibb, Syngenta, Novartis, Pfizer, Amgen, and Eli Lilly. We are grateful to TSRI, the NSF GRFP, the NDSEG Fellowship program, and the Skaggs–Oxford Scholarship program for predoctoral fellowships to K.M.E. We acknowledge Peter S. Thuy-Boun (Yu laboratory, TSRI), Will R. Gutekunst (Baran laboratory, TSRI), Jae H. Chu (Mauro laboratory, TSRI), and Matthew Tredwell (Gouverneur laboratory, Oxford) for generous assistance in editing this review and for thoughtful advice during its preparation. We also wish to recognize our colleagues at University of Cambridge, Brandeis University, and TSRI for encouragement and inspiration over the years. Finally, and most importantly, we thank all present and former group members, whose tenacity and creativity have been at the heart of the research described here. TSRI Manuscript no. 21887.

REFERENCES

(1) Berrisford, D. J.; Bolm, C.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **1995**, *34*, 1059–1070.

(2) As a point of clarification, “ligand” can refer to any molecule that is coordinated to a metal catalyst (anionic donor, X-type; or neutral donor, L-type). Throughout the text, we generally focus on ligands that possesses an element of design (i.e., a molecule that was independently synthesized for the express purpose of coordinating it to the metal to influence the reactivity) rather than common counterions (OAc[−], OTf[−], Cl[−], etc.) or solvent molecules (MeCN, THF, etc.), but those too can have a dramatic impact on the reactivity and/or selectivity of catalytic transformations.

(3) Katsuki, T.; Sharpless, K. B. *J. Am. Chem. Soc.* **1980**, *102*, 5974–5976.

(4) Gao, Y.; Hanson, R. M.; Klunder, J. M.; Ko, S. Y.; Masamune, H.; Sharpless, K. B. *J. Am. Chem. Soc.* **1987**, *109*, 5765–5780.

(5) Dang, T. P.; Kagan, H. B. *J. Chem. Soc. D* **1971**, 481.

(6) Vineyard, B. D.; Knowles, W. S.; Sabacky, M. J.; Bachman, G. L.; Weinkauff, D. J. *J. Am. Chem. Soc.* **1977**, *99*, 5946–5952.

(7) Fryzuk, M. D.; Bosnich, B. *J. Am. Chem. Soc.* **1977**, *99*, 6262–6267.

(8) Miyashita, A.; Yasuda, A.; Takaya, H.; Toriumi, K.; Ito, T.; Souchi, T.; Noyori, R. *J. Am. Chem. Soc.* **1980**, *102*, 7932–7934.

(9) Osborn, J. A.; Wilkinson, G.; Young, J. F. *Chem. Commun.* **1965**, 17.

(10) Osborn, J. A.; Jardine, F. H.; Young, J. F.; Wilkinson, G. *J. Chem. Soc. A* **1966**, 1711–1732.

(11) Trnka, T. M.; Grubbs, R. H. *Acc. Chem. Res.* **2001**, *34*, 18–29.

(12) Nolan, S. P.; Clavier, H. *Chem. Soc. Rev.* **2010**, *39*, 3305–3316.

(13) Littke, A. F.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 4176–4211.

(14) Hartwig, J. F. *Acc. Chem. Res.* **2008**, *41*, 1534–1544.

(15) Martin, R.; Buchwald, S. L. *Acc. Chem. Res.* **2008**, *41*, 1461–1473.

(16) Herrmann, W. A. *Angew. Chem., Int. Ed.* **2002**, *41*, 1290–1309.

(17) Hillier, A. C.; Grasa, G. A.; Viciu, M. S.; Lee, H. M.; Yang, C.; Nolan, S. P. *J. Organomet. Chem.* **2002**, *653*, 69–82.

(18) Jacobsen, E. N.; Markó, I.; Mungall, W. S.; Schröder, G.; Sharpless, K. B. *J. Am. Chem. Soc.* **1988**, *110*, 1968–1970.

(19) Andersson, M. A.; Epple, R.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 472–475.

(20) Yoon, T. P.; Jacobsen, E. N. *Science* **2003**, *299*, 1691–1693.

(21) Tolman, C. A. *Chem. Rev.* **1977**, *77*, 313–348.

(22) Nishiyama, M.; Yamamoto, T.; Koie, Y. *Tetrahedron Lett.* **1998**, *39*, 617–620.

(23) Littke, A. F.; Fu, G. C. *Angew. Chem., Int. Ed.* **1998**, *37*, 3387–3388.

(24) Hartwig, J. F.; Kawatsura, M.; Hauck, S. I.; Shaughnessy, K. H.; Alcazar-Roman, L. M. *J. Org. Chem.* **1999**, *64*, 5575–5580.

(25) Old, D. W.; Wolfe, J. P.; Buchwald, S. L. *J. Am. Chem. Soc.* **1998**, *120*, 9722–9723.

(26) Nieto-Oberhuber, C.; López, S.; Echavarren, A. M. *J. Am. Chem. Soc.* **2005**, *127*, 6178–6179.

(27) Hughes, O. R.; Unruh, J. D. *J. Mol. Catal.* **1981**, *12*, 71–83.

(28) Kranenburg, M.; van der Burgt, Y. E. M.; Kamer, P. C. J.; van Leeuwen, P. W. N. M.; Goubitz, K.; Fraanje, J. *Organometallics* **1995**, *14*, 3081–3089.

(29) Gupta, M.; Hagen, C.; Flesher, R. J.; Kaska, W. C.; Jensen, C. M. *Chem. Commun.* **1996**, 2083–2084.

(30) Liu, F.; Pak, E. B.; Singh, B.; Jensen, C. M.; Goldman, A. S. *J. Am. Chem. Soc.* **1999**, *121*, 4086–4087.

(31) Nolan, S. P. *Acc. Chem. Res.* **2011**, *44*, 91–100.

(32) Scholl, M.; Ding, S.; Lee, C. W.; Grubbs, R. H. *Org. Lett.* **1999**, *1*, 953–956.

(33) Tajima, Y.; Kunioka, E. *J. Org. Chem.* **1968**, *33*, 1689–1690.

(34) Stern, R.; Sajus, L. *Tetrahedron Lett.* **1968**, *9*, 6313–6314.

(35) Miyake, A.; Kondo, H. *Angew. Chem., Int. Ed.* **1968**, *7*, 631–632.

(36) Funk, R. L.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1977**, *99*, 5483–5484.

(37) Uto, T.; Shimizu, M.; Ueura, K.; Tsurugi, H.; Satoh, T.; Miura, M. *J. Org. Chem.* **2008**, *73*, 298–300.

(38) Hyster, T. K.; Rovis, T. *Chem. Sci.* **2011**, *2*, 1606–1610.

- (39) Evano, G.; Blanchard, N.; Toumi, M. *Chem. Rev.* **2008**, *108*, 3054–3131.
- (40) Surry, D. S.; Buchwald, S. L. *Chem. Sci.* **2010**, *1*, 13–31.
- (41) Togni, A.; Venanzi, L. M. *Angew. Chem., Int. Ed.* **1994**, *33*, 497–526.
- (42) Michel, B. W.; Camelio, A. M.; Cornell, C. N.; Sigman, M. S. *J. Am. Chem. Soc.* **2009**, *131*, 6076–6077.
- (43) Campbell, A. N.; White, P. B.; Guzei, I. A.; Stahl, S. S. *J. Am. Chem. Soc.* **2010**, *132*, 15116–15119.
- (44) The examples included in this figure are not intended to constitute a comprehensive list. Additionally, it should also be noted that many of the ligands highlighted in Figure 1 are also known to give enhanced kinetic reactivity in asymmetric and nonstereoselective transformations.
- (45) Brown, J. M. *Angew. Chem., Int. Ed.* **1987**, *26*, 190–203.
- (46) Hoveyda, A. H.; Evans, D. A.; Fu, G. C. *Chem. Rev.* **1993**, *93*, 1307–1370.
- (47) Ohta, T.; Takaya, H.; Kitamura, M.; Nagai, K.; Noyori, R. *J. Org. Chem.* **1987**, *52*, 3174–3176.
- (48) Crabtree, R. H.; Felkin, H.; Morris, G. E. *J. Organomet. Chem.* **1977**, *141*, 205–215.
- (49) Bell, S.; Wüstenberg, B.; Kaiser, S.; Menges, F.; Netscher, T.; Pfaltz, A. *Science* **2006**, *311*, 642–644.
- (50) Kaiser, S.; Smidt, S. P.; Pfaltz, A. *Angew. Chem., Int. Ed.* **2006**, *45*, 5194–5197.
- (51) Hanson, R. M.; Sharpless, K. B. *J. Org. Chem.* **1986**, *51*, 1922–1925.
- (52) Zhang, W.; Loebach, J. L.; Wilson, S. R.; Jacobsen, E. N. *J. Am. Chem. Soc.* **1990**, *112*, 2801–2803.
- (53) Jacobsen, E. N.; Zhang, W.; Muci, A. R.; Ecker, J. R.; Deng, L. *J. Am. Chem. Soc.* **1991**, *113*, 7063–7064.
- (54) Noyori, R. *Angew. Chem., Int. Ed.* **2002**, *41*, 2008–2022.
- (55) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457–2483.
- (56) Grushin, V. V.; Alper, H. *Chem. Rev.* **1994**, *94*, 1047–1062.
- (57) Knowles, W. S. *Acc. Chem. Res.* **1983**, *16*, 106–112.
- (58) Reetz, M. T. *Angew. Chem., Int. Ed.* **2008**, *47*, 2556–2588.
- (59) Sakane, S.; Maruoka, K.; Yamamoto, H. *Tetrahedron Lett.* **1985**, *26*, 5535–5538.
- (60) Sasai, H.; Suzuki, T.; Arai, S.; Arai, T.; Shibasaki, M. *J. Am. Chem. Soc.* **1992**, *114*, 4418–4420.
- (61) Keck, G. E.; Tarbet, K. H.; Geraci, L. S. *J. Am. Chem. Soc.* **1993**, *115*, 8467–8468.
- (62) Hoveyda, A. H.; Schrock, R. R. *Chem.—Eur. J.* **2001**, *7*, 945–950.
- (63) Harper, K. C.; Sigman, M. S. *Science* **2011**, *333*, 1875–1878.
- (64) von Matt, P.; Pfaltz, A. *Angew. Chem., Int. Ed.* **1993**, *32*, 566–568.
- (65) Sprinz, J.; Helmchen, G. *Tetrahedron Lett.* **1993**, *34*, 1769–1772.
- (66) Dawson, G. J.; Frost, C. G.; Williams, J. M. J.; Coote, S. J. *Tetrahedron Lett.* **1993**, *34*, 3149–3150.
- (67) Brown, J. M.; Hulmes, D. I.; Guiry, P. J. *Tetrahedron* **1994**, *50*, 4493–4506.
- (68) Helmchen, G.; Pfaltz, A. *Acc. Chem. Res.* **2000**, *33*, 336–345.
- (69) Williams, J. M. J. *Synlett* **1996**, 705–710.
- (70) Pfaltz, A.; Drury, W. J., III. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 5723–5726.
- (71) Tani, K.; Behenna, D. C.; McFadden, R. M.; Stoltz, B. M. *Org. Lett.* **2007**, *9*, 2529–2531.
- (72) Corey, E. J.; Cheng, X.-M. *The Logic of Chemical Synthesis*; John Wiley & Sons: New York, 1995.
- (73) McMurray, L.; O'Hara, F.; Gaunt, M. J. *Chem. Soc. Rev.* **2011**, *40*, 1885–1898.
- (74) Gutekunst, W. R.; Baran, P. S. *Chem. Soc. Rev.* **2011**, *40*, 1976–1991.
- (75) Yamaguchi, J.; Yamaguchi, A. D.; Itami, K. *Angew. Chem., Int. Ed.* **2012**, *51*, 8960–9009.
- (76) Engle, K. M.; Wang, D.-H.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 14137–14151.
- (77) Engle, K. M.; Mei, T.-S.; Wasa, M.; Yu, J.-Q. *Acc. Chem. Res.* **2012**, *45*, 788–802.
- (78) Newhouse, T.; Baran, P. S.; Hoffmann, R. W. *Chem. Soc. Rev.* **2009**, *38*, 3010–3021.
- (79) Li, C.-J.; Trost, B. M. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 13197–13202.
- (80) Kimura, S.; Kobayashi, S.; Kumamoto, T.; Akagi, A.; Sato, N.; Ishikawa, T. *Helv. Chim. Acta* **2011**, *94*, 578–591.
- (81) Wang, D.-H.; Yu, J.-Q. *J. Am. Chem. Soc.* **2011**, *133*, 5767–5769.
- (82) Mei, T.-S.; Giri, R.; Mangel, N.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2008**, *47*, 5215–5219.
- (83) Zhang, Y.-H.; Yu, J.-Q. *J. Am. Chem. Soc.* **2009**, *131*, 14654–14655.
- (84) Giri, R.; Yu, J.-Q. *J. Am. Chem. Soc.* **2008**, *130*, 14082–14083.
- (85) Zhang, Y.-H.; Shi, B.-F.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2009**, *48*, 6097–6100.
- (86) Giri, R.; Mangel, N.; Li, J.-J.; Wang, D.-H.; Breazzano, S. P.; Saunders, L. B.; Yu, J.-Q. *J. Am. Chem. Soc.* **2007**, *129*, 3510–3511.
- (87) Yoo, E. J.; Ma, S.; Mei, T.-S.; Chan, K. S. L.; Yu, J.-Q. *J. Am. Chem. Soc.* **2011**, *133*, 7652–7655.
- (88) Chan, K. S. L.; Wasa, M.; Wang, X.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2011**, *50*, 9081–9084.
- (89) Dai, H.-X.; Yu, J.-Q. *J. Am. Chem. Soc.* **2012**, *134*, 134–137.
- (90) Zhang, X.-G.; Dai, H.-X.; Wasa, M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2012**, *134*, 11948–11951.
- (91) Yu, J.-Q.; Giri, R.; Chen, X. *Org. Biomol. Chem.* **2006**, *4*, 4041–4047.
- (92) Chen, X.; Engle, K. M.; Wang, D.-H.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2009**, *48*, 5094–5115.
- (93) Giri, R.; Shi, B.-F.; Engle, K. M.; Mangel, N.; Yu, J.-Q. *Chem. Soc. Rev.* **2009**, *38*, 3242–3272.
- (94) Wasa, M.; Engle, K. M.; Yu, J.-Q. *Isr. J. Chem.* **2010**, *50*, 605–616.
- (95) Mei, T.-S.; Kou, L.; Ma, S.; Engle, K. M.; Yu, J.-Q. *Synthesis* **2012**, *44*, 1778–1791.
- (96) Because Cu is known to readily participate in single-electron transfer events, it has historically been challenging to elucidate the mechanism for Cu-catalyzed cross-coupling reactions. This has also proven to be the case with Cu-catalyzed C–H functionalization reactions, and the mechanism of these transformations is the topic of ongoing investigation. It is possible that these reactions involve a Cu(II)/Cu(III)/Cu(I) or Cu(III)/Cu(I) redox couple, the latter of which would be isoelectronic to the Pd(II)/Pd(0) redox couple. It is difficult, however, to generalize from individual mechanistic studies because there could be fundamental differences from one reaction to the next. For a detailed discussion of the mechanism of these transformations and related Cu-catalyzed cross-coupling reactions, see: (a) King, A. E.; Huffman, L. M.; Casitas, A.; Costas, M.; Ribas, X.; Stahl, S. S. *J. Am. Chem. Soc.* **2010**, *132*, 12068–12073. For selected examples of C–H functionalization using Cu(II) precatalysts, see: (b) Chen, X.; Hao, X.-S.; Goodhue, C. E.; Yu, J.-Q. *J. Am. Chem. Soc.* **2006**, *128*, 6790–6791. (c) Brasche, G.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2008**, *47*, 1932–1934. (d) Ueda, S.; Nagasawa, H. *Angew. Chem., Int. Ed.* **2008**, *47*, 6411–6413. (e) Do, H.-Q.; Daugulis, O. *J. Am. Chem. Soc.* **2009**, *131*, 17052–17053.
- (97) For selected examples of Rh(III)-catalyzed C–H functionalization, see: (a) Ueura, K.; Satoh, T.; Miura, M. *Org. Lett.* **2007**, *9*, 1407–1409. (b) Stuart, D. R.; Bertrand-Laperle, M.; Burgess, K. M. N.; Fagnou, K. *J. Am. Chem. Soc.* **2008**, *130*, 16474–16475. (c) Patureau, F. W.; Glorius, F. *J. Am. Chem. Soc.* **2010**, *132*, 9982–9983. (d) Hyster, T. K.; Rovis, T. *J. Am. Chem. Soc.* **2010**, *132*, 10565–10569. (e) Pesset, M.; Oehrlrich, D.; Rombouts, F.; Molander, G. A. *Org. Lett.* **2013**, *15*, 1528–1531.
- (98) For selected examples of Ru(II)-catalyzed C–H functionalization, see: (a) Weissman, H.; Song, X.; Milstein, D. *J. Am. Chem. Soc.* **2001**, *123*, 337–338. (b) Oi, S.; Fukita, S.; Hirata, N.; Watanuki, N.; Miyano, S.; Inoue, Y. *Org. Lett.* **2001**, *3*, 2579–2581. (c) Ackermann, L. *Org. Lett.* **2005**, *7*, 3123–3125. (d) Özdemir, I.; Demir, S.; Çetinkaya, B.; Gourlaouen, C.; Maseras, F.; Bruneau, C.; Dixneuf, P.

H. *J. Am. Chem. Soc.* **2008**, *130*, 1156–1157. (e) Padala, K.; Jegannmohan, M. *Org. Lett.* **2011**, *13*, 6144–6147.

(99) For early examples of an alternative approach to effect C–H functionalization using low-valent Ru(0) and Rh(I) catalysts, see: (a) Murai, S.; Kakiuchi, F.; Sekine, S.; Tanaka, Y.; Kamatani, A.; Sonoda, M.; Chatani, N. *Nature* **1993**, *366*, 529–531. (b) Trost, B. M.; Imi, K.; Davies, I. W. *J. Am. Chem. Soc.* **1995**, *117*, 5371–5372. (c) Lenges, C. P.; Brookhart, M. *J. Am. Chem. Soc.* **1999**, *121*, 6616–6623. (d) Jun, C.-H.; Hong, J.-B.; Kim, Y.-H.; Chung, K.-Y. *Angew. Chem., Int. Ed.* **2000**, *39*, 3440–3442. (e) Tan, K. L.; Bergman, R. G.; Ellman, J. A. *J. Am. Chem. Soc.* **2001**, *123*, 2685–2686.

(100) It is important to clarify that along a Pd(0)/Pd(II) catalytic cycle, the cross-coupling of C(aryl)–H bonds with aryl halides can be promoted with phosphine and NHC ligands. Analogous to traditional Pd(0)-catalyzed cross-coupling reactions, this type of reaction begins with oxidative addition of the aryl halide as the first elementary step in the catalytic cycle. Though Pd(II) salts are often used as precatalysts in these transformations, in situ reduction generates a [Pd(0)L_n] species, which is the active catalyst. This is a distinct mechanistic manifold that is outside of the scope of this Perspective. For reviews discussing this mode of catalysis, see ref 92 and the following: (a) Campeau, L.-C.; Stuart, D. R.; Fagnou, K. *Aldrichimica Acta* **2007**, *40*, 35–41. (b) Alberico, D.; Scott, M. E.; Lautens, M. *Chem. Rev.* **2007**, *107*, 174–238. For lead references using chiral [Pd(0)–L*] catalysts in C–H functionalization, see: (c) Albicker, M. R.; Cramer, N. *Angew. Chem., Int. Ed.* **2009**, *48*, 9139–9142. (d) Renaudat, A.; Jean-Gérard, L.; Jazzar, R.; Kefalidis, C. E.; Clot, E.; Baudoin, O. *Angew. Chem., Int. Ed.* **2010**, *49*, 7261–7265. (e) Nakanishi, M.; Katayev, D.; Besnard, C.; Kündig, E. P. *Angew. Chem., Int. Ed.* **2011**, *50*, 7438–7441. (f) Anas, S.; Cordi, A.; Kagan, H. B. *Chem. Commun.* **2011**, *47*, 11483–11485. (g) Saget, T.; Lemouzy, S. J.; Cramer, N. *Angew. Chem., Int. Ed.* **2012**, *51*, 2238–2242. (h) Martin, N.; Pierre, C.; Davi, M.; Jazzar, R.; Baudoin, O. *Chem.–Eur. J.* **2012**, *18*, 4480–4484. (i) Saget, T.; Cramer, N. *Angew. Chem., Int. Ed.* **2012**, *51*, 12842–12845.

(101) Neufeldt, S. R.; Sanford, M. S. *Acc. Chem. Res.* **2012**, *45*, 936–946.

(102) Daugulis, O.; Do, H.-Q.; Shabashov, D. *Acc. Chem. Res.* **2009**, *42*, 1074–1086.

(103) Lyons, T. W.; Sanford, M. S. *Chem. Rev.* **2010**, *110*, 1147–1169.

(104) Ryabov, A. D.; Sakodinskaya, I. K.; Yatsimirsky, A. K. *J. Chem. Soc., Dalton Trans.* **1985**, 2629–2638.

(105) Gómez, M.; Granell, J.; Martinez, M. *Organometallics* **1997**, *16*, 2539–2546.

(106) Davies, D. L.; Donald, S. M. A.; Macgregor, S. A. *J. Am. Chem. Soc.* **2005**, *127*, 13754–13755.

(107) The mechanisms and associated nomenclature of metal-mediated C–H cleavage are topics of ongoing discussion in the literature. With Pd(II), oxidative addition, electrophilic palladation, and concerted metalation/deprotonation (CMD) have all been proposed, and based on our current knowledge the latter two seem to be operative in the preponderance of cases. It is important to consider, however, that in actuality these broad categories likely represent extremes on a continuum of possible mechanistic scenarios. For example, one could reasonably argue that the “CMD” category actually encompasses nearly all base-assisted, redox-neutral C–H activation mechanisms. If proton removal at the transition state is not advanced compared to Pd–C bonding, the CMD pathway would possess electrophilic substitution character. If, on the other hand, proton removal is advanced compared to Pd–C bonding, the CMD pathway would have deprotonative character. As the community gains a deeper understanding of the true mechanism at play in each unique reaction, the adoption more precise nomenclature, such “CMD with deprotonative character” or “CMD with electrophilic substitution character” may be in order. To simplify discussion in the present review, we use “electrophilic palladation” and “CMD” in their traditional sense. The former we use to describe Pd(II)-mediated C(sp²)–H activation that likely involves a Wheland (arenium) intermediate. The latter we use to describe Pd(II)-mediated

C(sp³)–H activation and C(sp²)–H activation with strongly deprotonative character (involving minimal charge build-up in the transition state).

(108) We kindly acknowledge a reviewer for pointing out the different manifestations of CMD discussed in ref 107.

(109) Gant, T. G.; Meyers, A. I. *Tetrahedron* **1994**, *50*, 2297–2360.

(110) Xia, J.-B.; You, S.-L. *Organometallics* **2007**, *26*, 4869–4871.

(111) Takebayashi, S.; Shizuno, T.; Otani, T.; Shibata, T. *Beilstein J. Org. Chem.* **2012**, *8*, 1844–1848.

(112) Giri, R.; Chen, X.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2005**, *44*, 2112–2115.

(113) Giri, R.; Liang, J.; Lei, J.-G.; Li, J.-J.; Wang, D.-H.; Chen, X.; Naggar, I. C.; Guo, C.; Foxman, B. M.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2005**, *44*, 7420–7424.

(114) Giri, R.; Chen, X.; Hao, X.-S.; Li, J.-J.; Liang, J.; Fan, Z.-P.; Yu, J.-Q. *Tetrahedron: Asymmetry* **2005**, *16*, 3502–3505.

(115) Giri, R.; Lan, Y.; Liu, P.; Houk, K. N.; Yu, J.-Q. *J. Am. Chem. Soc.* **2012**, *134*, 14118–14126.

(116) Shi, B.-F.; Mangel, N.; Zhang, Y.-H.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2008**, *47*, 4882–4886.

(117) For original data corresponding to entries 5 and 8, see: Mangel, N. L. Palladium-Catalyzed Alkylation, Arylation and Dehydrogenation of Unactivated C–H bonds. Syntheses of the Tetracyclic Aminoquinone Moiety of Marmycin A, (±)-Nosyberkol (Isotuberculosinol, Revised Structure of Edaxadiene), and (±)-Tuberculosinol. Ph.D. Thesis, Brandeis University, Waltham, MA, February 2011. For all other entries, see the Supporting Information of ref 116.

(118) Navarro, R.; García, J.; Urriolabeitia, E. P.; Cativiela, C.; Diaz-de-Villegas, M. D. *J. Organomet. Chem.* **1995**, *490*, 35–43.

(119) Mono-*N*-protected amino acids have been used to control the stereoselectivity in stoichiometric cyclopalladation reactions. For examples, see: (a) Sokolov, V. I.; Troitskaya, L. L. *Chimia* **1978**, *32*, 122–123. (b) Sokolov, V. I.; Troitskaya, L. L.; Reutov, O. A. *J. Organomet. Chem.* **1979**, *182*, 537–546. (c) Sokolov, V. I.; Troitskaya, L. L.; Khrushchova, N. S. *J. Organomet. Chem.* **1983**, *250*, 439–446. (d) Günay, M. E.; Richards, C. J. *Organometallics* **2009**, *28*, 5833–5836. (e) Dendele, N.; Bisaro, F.; Gaumont, A.-C.; Perrio, S.; Richards, C. J. *Chem. Commun.* **2012**, *48*, 1991–1993.

(120) Musaeov, D. G.; Kaledin, A.; Shi, B.-F.; Yu, J.-Q. *J. Am. Chem. Soc.* **2012**, *134*, 1690–1698.

(121) Shi, B.-F.; Zhang, Y.-H.; Lam, J. K.; Wang, D.-H.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 460–461.

(122) Prior to determining the absolute configuration of product (–)-**127** as (*R*) (see ref 120), we put forward a working model for stereoselection in ref 116. However, the model actually predicted preferential formation of the opposite enantiomer. That model has since been abandoned, and a revised model was put forward in refs 120 and 121. The discussion herein is based on the revised model.

(123) Wang, D.-H.; Engle, K. M.; Shi, B.-F.; Yu, J.-Q. *Science* **2010**, *327*, 315–319.

(124) Kao, L.-C.; Sen, A. *J. Chem. Soc., Chem. Commun.* **1991**, 1242–1243.

(125) Miura, M.; Tsuda, T.; Satoh, T.; Pivsa-Art, S.; Nomura, M. *J. Org. Chem.* **1998**, *63*, 5211–5215.

(126) Dangel, B. D.; Johnson, J. A.; Sames, D. *J. Am. Chem. Soc.* **2001**, *123*, 8149–8150.

(127) Lee, J. M.; Chang, S. *Tetrahedron Lett.* **2006**, *47*, 1375–1379.

(128) Wasa, M.; Engle, K. M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2009**, *131*, 9886–9887.

(129) Wasa, M.; Engle, K. M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 3680–3681.

(130) Wasa, M.; Engle, K. M.; Lin, D. W.; Yoo, E. J.; Yu, J.-Q. *J. Am. Chem. Soc.* **2011**, *133*, 19598–19601.

(131) Kutter, E.; Hansch, C. *J. Med. Chem.* **1969**, *12*, 647–652.

(132) Sotomatsu, T.; Fujita, T. *J. Org. Chem.* **1989**, *54*, 4443–4448.

(133) Engle, K. M.; Wang, D.-H.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2010**, *49*, 6169–6173.

(134) Shue, R. S. *J. Chem. Soc. D* **1971**, 1510–1511.

- (135) Dams, M.; De Vos, D. E.; Celen, S.; Jacobs, P. A. *Angew. Chem., Int. Ed.* **2003**, *42*, 3512–3515.
- (136) Baxter, R. D.; Sale, D.; Engle, K. M.; Yu, J.-Q.; Blackmond, D. G. *J. Am. Chem. Soc.* **2012**, *134*, 4600–4606.
- (137) Engle, K. M.; Thuy-Boun, P. S.; Dang, M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2011**, *133*, 18183–18193.
- (138) Lu, Y.; Wang, D.-H.; Engle, K. M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 5916–5921.
- (139) Lu, Y.; Leow, D.; Wang, X.; Engle, K. M.; Yu, J.-Q. *Chem. Sci.* **2011**, *2*, 967–971.
- (140) Simmons, E. M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2010**, *132*, 17092–17095.
- (141) Simmons, E. M.; Hartwig, J. F. *Nature* **2012**, *483*, 70–73.
- (142) Xiao, B.; Gong, T.-J.; Liu, Z.-J.; Liu, J.-H.; Luo, D.-F.; Xu, J.; Liu, L. *J. Am. Chem. Soc.* **2011**, *133*, 9250–9253.
- (143) Wei, Y.; Yoshikai, N. *Org. Lett.* **2011**, *13*, 5504–5507.
- (144) Miura, M.; Tsuda, T.; Satoh, T.; Nomura, M. *Chem. Lett.* **1997**, *26*, 1103–1104.
- (145) Hennings, D. D.; Iwasa, S.; Rawal, V. H. *J. Org. Chem.* **1997**, *62*, 2–3.
- (146) Pham, M. V.; Ye, B.; Cramer, N. *Angew. Chem., Int. Ed.* **2012**, *51*, 10610–10614.
- (147) Dai, H.-X.; Stepan, A. F.; Plummer, M. S.; Zhang, Y.-H.; Yu, J.-Q. *J. Am. Chem. Soc.* **2011**, *133*, 7222–7228.
- (148) Iglesias, Á.; Álvarez, R.; de Lera, Á. R.; Muñiz, K. *Angew. Chem., Int. Ed.* **2012**, *51*, 2225–2228.
- (149) Li, G.; Leow, D.; Wan, L.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2013**, *52*, 1245–1247.
- (150) Leow, D.; Li, G.; Mei, T.-S.; Yu, J.-Q. *Nature* **2012**, *486*, 518–522.
- (151) Schwarz, H. *Acc. Chem. Res.* **1989**, *22*, 282–287.
- (152) Dietl, N.; Schlangen, M.; Schwarz, H. *Chem.—Eur. J.* **2011**, *17*, 1783–1788.
- (153) Huang, C.; Chattopadhyay, B.; Gevorgyan, V. *J. Am. Chem. Soc.* **2011**, *133*, 12406–12409.
- (154) Novák, P.; Correa, A.; Gallardo-Donaire, J.; Martin, R. *Angew. Chem., Int. Ed.* **2011**, *50*, 12236–12239.
- (155) Wang, Y.-N.; Guo, X.-Q.; Zhu, X.-H.; Zhong, R.; Cai, L.-H.; Hou, X.-F. *Chem. Commun.* **2012**, *48*, 10437–10439.
- (156) Gao, D.-W.; Shi, Y.-C.; Gu, Q.; Zhao, Z.-L.; You, S.-L. *J. Am. Chem. Soc.* **2013**, *135*, 86–89.
- (157) Cheng, X.-F.; Li, Y.; Su, Y.-M.; Yin, F.; Wang, J.-Y.; Sheng, J.; Vora, H. U.; Wang, X.-S.; Yu, J.-Q. *J. Am. Chem. Soc.* **2013**, *135*, 1236–1239.
- (158) Cong, X.; You, J.; Gao, G.; Lan, J. *Chem. Commun.* **2013**, *49*, 662–664.
- (159) For a recent example in which mono-*N*-protected amino acid ligands were used in Rh(III)-catalyzed C–H functionalization, see ref 97e.
- (160) Moritani, I.; Fujiwara, Y. *Tetrahedron Lett.* **1967**, *8*, 1119–1122.
- (161) Fujiwara, Y.; Moritani, I.; Matsuda, M.; Teranishi, S. *Tetrahedron Lett.* **1968**, *9*, 3863–3865.
- (162) Fujiwara, Y.; Moritani, I.; Danno, S.; Asano, R.; Teranishi, S. *J. Am. Chem. Soc.* **1969**, *91*, 7166–7169.
- (163) Asano, R.; Moritani, I.; Fujiwara, Y.; Teranishi, S. *J. Chem. Soc. D* **1970**, 1293.
- (164) Tsuji, J.; Nagashima, H. *Tetrahedron* **1984**, *40*, 2699–2702.
- (165) Jia, C.; Lu, W.; Kitamura, T.; Fujiwara, Y. *Org. Lett.* **1999**, *1*, 2097–2100.
- (166) Jia, C.; Piao, D.; Oyamada, J.; Lu, W.; Kitamura, T.; Fujiwara, Y. *Science* **2000**, *287*, 1992–1995.
- (167) Yokota, T.; Tani, M.; Sakaguchi, S.; Ishii, Y. *J. Am. Chem. Soc.* **2003**, *125*, 1476–1477.
- (168) Kubota, A.; Emmert, M. H.; Sanford, M. S. *Org. Lett.* **2012**, *14*, 1760–1763.
- (169) Fujiwara, Y.; Moritani, I.; Asano, R.; Tanaka, H.; Teranishi, S. *Tetrahedron* **1969**, *25*, 4815–4818.
- (170) Fujiwara, Y.; Asano, R.; Moritani, I.; Teranishi, S. *J. Org. Chem.* **1976**, *41*, 1681–1683.
- (171) Fujiwara, Y.; Maruyama, O.; Yoshidomi, M.; Taniguchi, H. *J. Org. Chem.* **1981**, *46*, 851–855.
- (172) Itahara, T.; Ikeda, M.; Sakakibara, T. *J. Chem. Soc., Perkin Trans. 1* **1983**, 1361–1363.
- (173) Ferreira, E. M.; Stoltz, B. M. *J. Am. Chem. Soc.* **2003**, *125*, 9578–9579.
- (174) Boele, M. D. K.; van Strijdonck, G. P. F.; de Vries, A. H. M.; Kamer, P. C. J.; de Vries, J. G.; van Leeuwen, P. W. N. M. *J. Am. Chem. Soc.* **2002**, *124*, 1586–1587.
- (175) Zhang, Y.-H.; Shi, B.-F.; Yu, J.-Q. *J. Am. Chem. Soc.* **2009**, *131*, 5072–5074.
- (176) Nishimura, T.; Onoue, T.; Ohe, K.; Uemura, S. *Tetrahedron Lett.* **1998**, *39*, 6011–6014.
- (177) Nishimura, T.; Onoue, T.; Ohe, K.; Uemura, S. *J. Org. Chem.* **1999**, *64*, 6750–6755.
- (178) Nishimura, T.; Ohe, K.; Uemura, S. *J. Am. Chem. Soc.* **1999**, *121*, 2645–2646.
- (179) Fix, S. R.; Brice, J. L.; Stahl, S. S. *Angew. Chem., Int. Ed.* **2002**, *41*, 164–166.
- (180) Trend, R. M.; Ramtohl, Y. K.; Ferreira, E. M.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2003**, *42*, 2892–2895.
- (181) Trend, R. M.; Ramtohl, Y. K.; Stoltz, B. M. *J. Am. Chem. Soc.* **2005**, *127*, 17778–17788.
- (182) Schiffner, J. A.; Oestreich, M. *Eur. J. Org. Chem.* **2011**, 1148–1154.
- (183) Kandukuri, S. R.; Oestreich, M. *J. Org. Chem.* **2012**, *77*, 8750–8755.
- (184) Stuart, D. R.; Fagnou, K. *Science* **2007**, *316*, 1172–1175.
- (185) Wang, C.; Rakshit, S.; Glorius, F. *J. Am. Chem. Soc.* **2010**, *132*, 14006–14008.
- (186) Izawa, Y.; Stahl, S. S. *Adv. Synth. Catal.* **2010**, *352*, 3223–3229.
- (187) Izawa, Y.; Pun, D.; Stahl, S. S. *Science* **2011**, *333*, 209–213.
- (188) Rentner, J.; Breinbauer, R. *Chem. Commun.* **2012**, *48*, 10343–10345.
- (189) Kravtsova, S. V.; Romm, I. P.; Stash, A. I.; Belsky, V. K. *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.* **1996**, *52*, 2201–2204.
- (190) Zhang, S.; Shi, L.; Ding, Y. *J. Am. Chem. Soc.* **2011**, *133*, 20218–20229.
- (191) Wasa, M.; Chan, K. S. L.; Zhang, X.-G.; He, J.; Miura, M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2012**, *134*, 18570–18572.
- (192) Zaitsev, V. G.; Shabashov, D.; Daugulis, O. *J. Am. Chem. Soc.* **2005**, *127*, 13154–13155.
- (193) Giri, R.; Mangel, N.; Foxman, B. M.; Yu, J.-Q. *Organometallics* **2008**, *27*, 1667–1670.
- (194) Nakao, Y. *Synthesis* **2011**, 3209–3219.
- (195) Moore, E. J.; Pretzer, W. R.; O’Connell, T. J.; Harris, J.; LaBounty, L.; Chou, L.; Grimmer, S. S. *J. Am. Chem. Soc.* **1992**, *114*, 5888–5890.
- (196) Godula, K.; Sezen, B.; Sames, D. *J. Am. Chem. Soc.* **2005**, *127*, 3648–3649.
- (197) Kawashima, T.; Takao, T.; Suzuki, H. *J. Am. Chem. Soc.* **2007**, *129*, 11006–11007.
- (198) Lewis, J. C.; Bergman, R. G.; Ellman, J. A. *J. Am. Chem. Soc.* **2007**, *129*, 5332–5333.
- (199) Berman, A. M.; Lewis, J. C.; Bergman, R. G.; Ellman, J. A. *J. Am. Chem. Soc.* **2008**, *130*, 14926–14927.
- (200) Takagi, J.; Sato, K.; Hartwig, J. F.; Ishiyama, T.; Miyaura, N. *Tetrahedron Lett.* **2002**, *43*, 5649–5651.
- (201) Mkhaliid, I. A. I.; Coventry, D. N.; Albesa-Jove, D.; Batsanov, A. S.; Howard, J. A. K.; Perutz, R. N.; Marder, T. B. *Angew. Chem., Int. Ed.* **2006**, *45*, 489–491.
- (202) Murphy, J. M.; Liao, X.; Hartwig, J. F. *J. Am. Chem. Soc.* **2007**, *129*, 15434–15435.
- (203) Fischer, D. F.; Sarpong, R. *J. Am. Chem. Soc.* **2010**, *132*, 5926–5927.

- (204) Hurst, T. E.; Macklin, T. K.; Becker, M.; Hartmann, E.; Kügel, W.; Parisienne-La Salle, J.-C.; Batsanov, A. S.; Marder, T. B.; Snieckus, V. *Chem.—Eur. J.* **2010**, *16*, 8155–8161.
- (205) Li, B.-J.; Shi, Z.-J. *Chem. Sci.* **2011**, *2*, 488–493.
- (206) Cho, J.-Y.; Iverson, C. N.; Smith, M. R., III. *J. Am. Chem. Soc.* **2000**, *122*, 12868–12869.
- (207) Nakao, Y.; Kanyiva, K. S.; Hiyama, T. *J. Am. Chem. Soc.* **2008**, *130*, 2448–2449.
- (208) Tsai, C.-C.; Shih, W.-C.; Fang, C.-H.; Li, C.-Y.; Ong, T.-G.; Yap, G. P. A. *J. Am. Chem. Soc.* **2010**, *132*, 11887–11889.
- (209) Nakao, Y.; Yamada, Y.; Kashihara, N.; Hiyama, T. *J. Am. Chem. Soc.* **2010**, *132*, 13666–13668.
- (210) Grigg, R.; Savic, V. *Tetrahedron Lett.* **1997**, *38*, 5737–5740.
- (211) Gürbüz, N.; Özdemir, I.; Çetinkaya, B. *Tetrahedron Lett.* **2005**, *46*, 2273–2277.
- (212) Wasa, M.; Worrell, B. T.; Yu, J.-Q. *Angew. Chem., Int. Ed.* **2010**, *49*, 1275–1277.
- (213) Campeau, L.-C.; Rousseaux, S.; Fagnou, K. *J. Am. Chem. Soc.* **2005**, *127*, 18020–18021.
- (214) Cho, S. H.; Hwang, S. J.; Chang, S. *J. Am. Chem. Soc.* **2008**, *130*, 9254–9256.
- (215) Wu, J.; Cui, X.; Chen, L.; Jiang, G.; Wu, Y. *J. Am. Chem. Soc.* **2009**, *131*, 13888–13889.
- (216) Xi, P.; Yang, F.; Qin, S.; Zhao, D.; Lan, J.; Gao, G.; Hu, C.; You, J. *J. Am. Chem. Soc.* **2010**, *132*, 1822–1824.
- (217) Tan, Y.; Barrios-Landeros, F.; Hartwig, J. F. *J. Am. Chem. Soc.* **2012**, *134*, 3683–3686.
- (218) Larivée, A.; Mousseau, J. J.; Charette, A. B. *J. Am. Chem. Soc.* **2008**, *130*, 52–54.
- (219) Mousseau, J. J.; Bull, J. A.; Charette, A. B. *Angew. Chem., Int. Ed.* **2010**, *49*, 1115–1118.
- (220) Wei, Y.; Kan, J.; Wang, M.; Su, W.; Hong, M. *Org. Lett.* **2009**, *11*, 3346–3349.
- (221) Wei, Y.; Su, W. *J. Am. Chem. Soc.* **2010**, *132*, 16377–16379.
- (222) Lin, B.-L.; Labinger, J. A.; Bercaw, J. E. *Can. J. Chem.* **2009**, *87*, 264–271.
- (223) McDonald, R. I.; White, P. B.; Weinstein, A. B.; Tam, C. P.; Stahl, S. S. *Org. Lett.* **2011**, *13*, 2830–2833.
- (224) Perch, N. S.; Widenhofer, R. A. *J. Am. Chem. Soc.* **1999**, *121*, 6960–6961.
- (225) Tsujihara, T.; Takenaka, K.; Onitsuka, K.; Hatanaka, M.; Sasai, H. *J. Am. Chem. Soc.* **2009**, *131*, 3452–3453.
- (226) Takenaka, K.; Akita, M.; Tanigaki, Y.; Takizawa, S.; Sasai, H. *Org. Lett.* **2011**, *13*, 3506–3509.
- (227) He, W.; Yip, K.-T.; Zhu, N.-Y.; Yang, D. *Org. Lett.* **2009**, *11*, 5626–5628.
- (228) Jiang, F.; Wu, Z.; Zhang, W. *Tetrahedron Lett.* **2010**, *51*, 5124–5126.
- (229) Yang, G.; Shen, C.; Zhang, W. *Angew. Chem., Int. Ed.* **2012**, *51*, 9141–9145.
- (230) Ebersson, L.; Jönsson, E. *Acta Chem. Scand. Ser. B* **1974**, *28*, 771–776.
- (231) Ebersson, L.; Jönsson, L. *J. Chem. Soc., Chem. Commun.* **1974**, 885–886.
- (232) Ebersson, L.; Jönsson, L. *Acta Chem. Scand. Ser. B* **1976**, *30*, 361–364.
- (233) Ebersson, L.; Jönsson, L. *Liebigs Ann. Chem.* **1977**, 233–241.
- (234) Jintoku, T.; Taniguchi, H.; Fujiwara, Y. *Chem. Lett.* **1987**, *16*, 1159–1162.
- (235) Campbell, A. N.; Meyer, E. B.; Stahl, S. S. *Chem. Commun.* **2011**, *47*, 10257–10259.
- (236) Diao, T.; Wadzinski, T. J.; Stahl, S. S. *Chem. Sci.* **2012**, *3*, 887–891.
- (237) Hickman, A. J.; Sanford, M. S. *ACS Catal.* **2011**, *1*, 170–174.
- (238) Zhu, C.; Falck, J. R. *Org. Lett.* **2011**, *13*, 1214–1217.
- (239) Kirchberg, S.; Tani, S.; Ueda, K.; Yamaguchi, J.; Studer, A.; Itami, K. *Angew. Chem., Int. Ed.* **2011**, *50*, 2387–2391.
- (240) Steinmetz, M.; Ueda, K.; Grimme, S.; Yamaguchi, J.; Kirchberg, S.; Itami, K.; Studer, A. *Chem.—Asian J.* **2012**, *7*, 1256–1260.
- (241) Yamaguchi, K.; Yamaguchi, J.; Studer, A.; Itami, K. *Chem. Sci.* **2012**, *3*, 2165–2169.
- (242) Shibahara, F.; Yamauchi, T.; Yamaguchi, E.; Murai, T. *J. Org. Chem.* **2012**, *77*, 8815–8820.
- (243) Ma, Y.; You, J.; Song, F. *Chem.—Eur. J.* **2013**, *19*, 1189–1193.
- (244) Jiang, T.-S.; Wang, G.-W. *Org. Lett.* **2013**, *15*, 788–791.
- (245) Izawa, Y.; Zheng, C.; Stahl, S. S. *Angew. Chem., Int. Ed.* **2013**, *52*, 3672–3675.
- (246) Ye, M.; Gao, G.-L.; Yu, J.-Q. *J. Am. Chem. Soc.* **2011**, *133*, 6964–6967.
- (247) Subsequent to our work, an example of Pd(II)-catalyzed C2-selective pyridine C–H olefination was reported: Wen, P.; Li, Y.; Zhou, K.; Ma, C.; Lan, X.; Ma, C.; Huang, G. *Adv. Synth. Catal.* **2012**, *354*, 2135–2140. In the absence of an ancillary ligand, N-coordination appears to facilitate C–H cleavage at the proximal C2 position.
- (248) Ye, M.; Gao, G.-L.; Edmunds, A. J. F.; Worthington, P. A.; Morris, J. A.; Yu, J.-Q. *J. Am. Chem. Soc.* **2011**, *133*, 19090–19093.
- (249) In a recent report, aryl tosylates were also found to be effective coupling partners in this transformation: Dai, F.; Gui, Q.; Liu, J.; Yang, Z.; Chen, X.; Guo, R.; Tan, Z. *Chem. Commun.* **2013**, *49*, 4634–4636.
- (250) De León, A.; Pons, J.; Solans, X.; Font-Bardia, M. *Acta Crystallogr., Sect. E: Struct. Rep. Online* **2007**, *63*, m2164.
- (251) Milani, B.; Scarel, A.; Zangrando, E.; Mestroni, G.; Carfagna, C.; Binotti, B. *Inorg. Chim. Acta* **2003**, *350*, 592–602.
- (252) Rotondo, A.; Bruno, G.; Cusumano, M.; Rotondo, E. *Inorg. Chim. Acta* **2009**, *362*, 4767–4773.
- (253) Holzbock, J.; Sawodny, W.; Thewalt, U. Z. *Anorg. Allg. Chem.* **2000**, *626*, 2563–2568.
- (254) Tebbe, K.-F.; Gräfe-Kavoosian, A.; Freckmann, B. Z. *Naturforsch., B: J. Chem. Sci.* **1996**, *51*, 999–1006.
- (255) Ma, L.; Smith, R. C.; Protasiewicz, J. D. *Inorg. Chim. Acta* **2005**, *358*, 3478–3482.
- (256) Quagliano, J. V.; Schubert, L. *Chem. Rev.* **1952**, *50*, 201–260.
- (257) Coe, B. J.; Glenwright, S. J. *Coord. Chem. Rev.* **2000**, *203*, 5–80.
- (258) Gorelsky, S. I. *Organometallics* **2012**, *31*, 794–797.
- (259) Petit, A.; Flygare, J.; Miller, A. T.; Winkel, G.; Ess, D. H. *Org. Lett.* **2012**, *14*, 3680–3683.
- (260) (a) Ye, M.; Edmunds, A. J. F.; Morris, J. A.; Sale, D.; Zhang, Y.; Yu, J.-Q. *Chem. Sci.* **2013**, DOI: 10.1039/C3SC50184A. (b) Hattori, K.; Yamaguchi, K.; Yamaguchi, J.; Itami, K. *Tetrahedron* **2012**, *68*, 7605–7612. (c) Ben-Yahia, A.; Naas, M.; El Kazzouli, S.; Essassi, E. M.; Guillaumet, G. *Eur. J. Org. Chem.* **2012**, 7075–7081.

NOTE ADDED AFTER ASAP PUBLICATION

Figure 1 contained errors in the version published ASAP on May 3, 2013; the correct version reposted on June 19, 2013.