


Regioselective Synthesis of 1,2-Dihydroquinolines by a Solvent-Free MgBr₂-Catalyzed Multicomponent Reaction

Rsuini U. Gutiérrez, Hans C. Correa,[†] Rafael Bautista, José Luis Vargas, Alberto V. Jerezano, Francisco Delgado, and Joaquín Tamariz*

Departamento de Química Orgánica, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, Prol Carpio y Plan de Ayala, 11340 México, D.F., Mexico

S Supporting Information

ABSTRACT: A highly efficient and regioselective synthesis of 1,2-dihydroquinolines via a multicomponent reaction between an aniline and two ketones is described. This reaction was catalyzed by magnesium bromide and carried out under solvent-free conditions. When the reaction was performed by using 3-substituted anilines and nonsymmetrically substituted ketones, principally a single product was found among the four expected regioisomers. A variety of anilines and ketones, including cyclic ketones, were evaluated providing a series of 1,2-dihydroquinolines with diverse substitution patterns. A study of the mechanism is discussed. There is evidence of the in situ formation of the imine as a result of the reaction between the aniline and one of the ketones, before annulation to the heterocyclic ring.



INTRODUCTION

1,2-Dihydroquinolines (1,2-DHQ) represent a privileged heterocyclic scaffold to build pharmacologically and biologically active molecules, as well as compounds for industrial uses. Examples are antibacterial,¹ antitrypanosomal,² antioxidant³ and antidiabetic agents,⁴ antijuvenile hormone insecticides,⁵ and natural products.⁶ Furthermore, 1,2-DHQ are antioxidants used as preservatives in animal nutriment and in vegetable and animal oils,⁷ as rubber antioxidants,⁸ as linkers for solid-phase organic synthesis,⁹ and as the core in ferromagnetic compounds.¹⁰ Despite their biological and industrial importance, only in the few past decades has 1,2-DHQ received considerable attention as synthetic targets. In the 1880s, they were prepared as intermediates in the quinoline synthesis carried out by Skraup, Riehm, and Doebner, and von Miller.¹¹ This multicomponent reaction (MCR)¹² was performed by Reddelien,¹³ Craig,¹⁴ and Vaughan¹⁵ years later for the synthesis of acetone–anil by using aniline (**1a**) and acetone (**2a**), catalyzed by hydrochloric acid or iodine (Scheme 1).

The latter methodology was recently employed with more versatile reaction conditions, which led to greater yield¹⁶ and suggested a mechanistic pathway.¹⁷ Analogous MCR catalyzed by transition metals used terminal acetylenes instead of ketones.¹⁸ Recently, an Au/Ag-catalyzed cyclization of *o*-


(hydroxyallyl)anilines under mild conditions was reported.¹⁹ Other methodologies include transition-metal catalysts,²⁰ zeolite,²¹ Lewis^{16,22} and Brønsted²³ acids, a cascade Michael/aldol reaction,²⁴ and an Ir-catalyzed allylic amination/ring-closing metathesis reaction,²⁵ which were mainly designed on the basis of heterocyclic moiety formation, starting from the respective functionalized anilines. However, most of these methods have severe limitations, such as the difficulty in properly introducing the A-ring substituents and the generation of undesirable byproducts, or compounds with low regioselectivity. Additionally, many use expensive catalysts, complex multistep pathways, harmful solvents, or harsh reaction conditions.

Owing to the relevance of 1,2-DHQ, and within the context of our ongoing studies oriented toward the design of new approaches for the preparation of pharmacologically key heterocyclic frameworks,²⁶ we herein disclose an efficient, versatile and regioselective method for the synthesis of these heterocycles, via a solvent-free MgBr₂-catalyzed MCR process. Due to the controversial nature of the mechanism for their formation,¹⁷ the mechanistic pathway was also explored in an attempt to establish the possible species involved in such cascade reaction.

RESULTS AND DISCUSSION


With a new and more efficient catalyst for the synthesis of 1,2-DHQ **3**, the reaction was carried out on the basis of the MCR method depicted in Scheme 1, and with the use of solvent-free conditions. Preliminary results showed that compound **3b** could be obtained, albeit in low to modest yields (Table 1,

Scheme 1. Synthesis of Acetone–Anil by Reaction of Aniline (1a) and Acetone (2a)



Received: May 11, 2013


Published: September 17, 2013

Table 1. Yields of 1,2-DHQ 3a–d/4a, Obtained by Reaction of 3-Methoxyaniline (1b) and Ketones 2a–d^a


entry	2 (R)	cat. ^b	base	solvent ^c	T (°C)	time (h)	products ^d (%)
1	2b (Me)	Br ₂	K ₂ CO ₃		20	6	3b (6)
2	2b (Me)	Br ₂	K ₂ CO ₃		40	6	3b (19)
3	2b (Me)	Br ₂	K ₂ CO ₃		80	6	3b (58)
4	2b (Me)	NBS	K ₂ CO ₃		60	6	^e
5	2b (Me)	HBr	K ₂ CO ₃		60	6	^e
6	2b (Me)	<i>n</i> -Bu ₄ NBr	K ₂ CO ₃		60	6	3b (25)
7	2b (Me)	CuBr ₂	Li ₂ CO ₃		60	12	3b (<3)
8	2b (Me)	KBr	Li ₂ CO ₃		60	12	3b (8)
9	2b (Me)	LiBr	Li ₂ CO ₃		60	12	3b (5)
10	2b (Me)	I ₂	Li ₂ CO ₃		60	12	3b (13)
11	2b (Me)	BBr ₃	K ₂ CO ₃		60	6	3b (50)
12	2b (Me)	PyHBr ₃	K ₂ CO ₃		70	6	3b (62)
13	2b (Me)	PyHBr ₃	Li ₂ CO ₃		45	12	3b (75)
14	2b (Me)	PyHBr ₃	Li ₂ CO ₃		60	12	3b (85)
15	2b (Me)	MgBr ₂	Li ₂ CO ₃		20	12	3b (13)
16	2b (Me)	MgBr ₂	Li ₂ CO ₃		60	12	3b (93)
17	2b (Me)	MgCO ₃	Li ₂ CO ₃		60	12	[^e]
18	2b (Me)	PyHBr ₃	Li ₂ CO ₃	MeCN	60	12	3b (10)
19	2b (Me)	MgBr ₂	Li ₂ CO ₃	THF	60	12	3b (19)
20	2b (Me)	MgBr ₂	Li ₂ CO ₃	(CH ₂) ₂ Cl ₂	60	12	3b (8)
21	2a (H)	PyHBr ₃	Li ₂ CO ₃ ^f		60	1	3a/4a (98:2) (77)
22	2a (H)	MgBr ₂ ^g	Li ₂ CO ₃ ^g		60	1	3a/4a (98:2) (89)
23	2b (Me)	MgBr ₂	Li ₂ CO ₃ ^g		60	1	3b (98)
24	2c (Et)	MgBr ₂	Li ₂ CO ₃ ^g		60	1	3c (80)
25	2d (Pr)	MgBr ₂	Li ₂ CO ₃ ^g		60	1	3d (62)

^aReaction conditions: 1b (1.0 molar equiv), 2 (5.0 molar equiv), catalyst (0.5–1.0 molar equiv), and M₂CO₃ (0.5–2.0 molar equiv). ^bAll the catalysts in 1.0 molar equiv, except PyHBr₃ (0.5 molar equiv). ^c3.0 mL. ^dAfter column chromatography. ^eNo reaction. ^f2.0 molar equiv. ^g0.5 molar equiv.

Scheme 2. Possible Regioisomeric Products from the Reaction of 1b with 2b



entries 1–3), by using bromine along with potassium carbonate at 20–80 °C for 6 h. Interestingly, among the four possible isomeric products, 3b–6b, only 3b was detected in the crude mixture (Scheme 2).

Whereas the desired product was yielded with bromine (Table 1, entries 4 and 5), which has both bromonium and bromide species, this was not the case with agents that provide only one of these species, such as *N*-bromosuccinimide (NBS) or hydrobromic acid. Other bromide salt catalysts or even iodine provided 3b in low yields (Table 1, entries 6–10). However, when the reaction was carried out in the presence of BBr₃ and PyHBr₃, reagents that can generate both kinds of species,²⁷ derivative 3b was obtained in higher yields (Table 1,

entries 11 and 12). The yields from the latter catalyst were improved by using Li₂CO₃ as the base, decreasing the temperature, and increasing the reaction time (Table 1, entries 13 and 14).


Unlike with copper, potassium, or lithium bromides, magnesium bromide proved to be highly efficient and regioselective (Table 1, entries 7–9 and 16). The latter and other magnesium salts have been used as Lewis acid promoters in diverse procedures, where both the magnesium and bromide ions play an important role in transformations.²⁸ It seems that both ions are also important in the present case, since with magnesium carbonate the reaction did not occur (Table 1, entry 17). Moreover, the highest efficacy was obtained under

solvent-free conditions (Table 1, entries 16 and 18–20). Although the regioselectivity is similar between MgBr_2 and PyHBr_3 , the efficiency of the former catalyst was an advantage, as shown by the reaction between aniline **1b** and acetone (**2a**) (Table 1, entries 21 and 22). It was necessary to use twice the amount of Li_2CO_3 for the PyHBr_3 -catalyzed process, as otherwise the reaction was too slow and the mixture of products was obtained in only 40% yield after 1 h of reaction. Contrarily, with the use of MgBr_2 gave a total conversion of the starting material and a higher isolated yield of **3a/4a**. The amount of PyHBr_3 cannot be increased to improve the conversion because the product decomposes.

Upon applying the optimal reaction conditions, the series of 1,2-DHQ **3b–d** was prepared by condensing the corresponding alkyl methyl ketones **2b–d** with aniline **1b** (Table 1, entries 23–25). It is noteworthy that the bigger the substituent in the ketone, the lower the yield. Actually, the reaction of diethyl ketone (**2e**) with **1b** leads to a very low yield (6%) of product **3e**, which can be attributed to the low reactivity of the hindered ketone.^{22b}

Consequently, we investigated the behavior of cyclic ketones **2f,g** under the same reaction conditions. In contrast to **2e**, the procedure with **2f,g** was highly efficient, leading to the desired products **3f,g** in quantitative yields (Scheme 3). The yield was

Scheme 3. Highly Efficient Synthesis of Polycyclic 1,2-DHQ **3f,g**




low (45%) when using PyHBr_3 as the catalyst in the reaction with **2g** to give **3g**. These results are in contrast with a recent report for an analogous iodine-catalyzed reaction,¹⁷ in which the main products were the 1,2-dihydroquinolines substituted not only by the expected 2-spirocycloalkyl 4-cycloalkenyl-fused 1,2-dihydroquinoline core, similar to products **3f,g** (Scheme 3), but also by derivatives bearing cycloalkenyl groups at the C-6 and/or C-8 position of the heterocycle. This polyalkenylation was not observed in our reactions in spite of the relatively large amount of cycloalkanones **2f,g** employed (5 molar equiv) to upgrade the conversion into **3f,g**. The greater quantity of **2f–g** was necessary in order to compensate for the formation of dimers or oligomers of these cyclic ketones under the reaction conditions.

Therefore, the highly efficient reactions obtained with cyclic ketones **2f–g**, in comparison with the poor yield observed with diethyl ketone (**2e**), indicates that the process is indeed sterically dependent on the substituents in the ketone.

With the aim of evaluating the versatility of this methodology and of standardizing a method of synthesis for a broad series of 1,2-DHQ **3**, we performed an MgBr_2 -catalyzed condensation with acetophenones **2h–k** using similar conditions. The series of heterocycles **3h–k** were obtained in low to moderate yields (10–50%) as stable solids. To enhance the yields, reactions were carried out with different temperatures and periods of time, finding that compounds **3h–k** could be isolated in better yields and in high regioselectivity at 90 °C and for 5 h (Scheme 4). It is worth mentioning that the presence of electron-

Scheme 4. Synthesis of 2,4-Diaryl-1,2-DHQ **3h–k**




withdrawing groups in the benzene ring of the acetophenone (i.e., **2k**) led to a less efficient conversion, which suggests that the reaction mechanism is sensitive to the electron-demand in the carbonyl group of the ketone (vide infra).

Considering that the electronic effect of the substituent is involved in the reactivity of the ketone, methyl pyruvate (**2l**) was submitted to the same conditions as those used for alkyl ketones, limiting the reaction time to 1 h in order to compare their reactivity and conversion rate (Scheme 5).²⁹ Whereas PyHBr_3 was unable to promote the reaction, MgBr_2 catalyzed the conversion into a mixture of regioisomers **3l/4l** (79:21) in modest yields. It was found that this reaction is very fast, and after 1 h of heating the starting material was consumed, though with the presence of side products. Indeed, ketone **2l** is a much more reactive substrate probably due to the presence of the vicinal electron-withdrawing methoxycarbonyl group.^{22b} Since the latter group can also be activated by the catalyst, a synergic activation effect is promoted.

The course and rate conversion of the reaction seems to be dictated not only by the electron demand or the hindrance of substituents in the ketone but also by the electron demand of the aniline substituents. Taking this into account, the reactivity of aniline **1b** should depend on the electron-donating effect of the methoxy group at the *meta* position of the amino group. To test this hypothesis, we evaluated the reactivity of a series of *meta*-substituted anilines **1c,d**, whose substituents have a lesser electron-donating effect than the methoxy group (Table 2). Thus, *m*-chloroaniline (**1c**) reacted with **2b** under the same conditions to cleanly give 1,2-dihydroquinoline **3m** with high regioselectivity, albeit in moderate yield, and recovery of the starting material (Table 2, entry 1). In the case of *m*-toluidine (**1d**), when using a reaction time of 1 h the conversion was less than that observed for **1b**, as evidenced by the fact that **3n** was isolated in low yield, though with excellent regioselectivity (Table 2, entry 2). It took 7 h of reaction to consume all the starting material, furnishing an almost quantitative yield of **3n** (Table 2, entry 3). When the ketone was changed to **2f**, a similar behavior was observed for 1 and 7 h of heating at 60 °C (Table 2, entries 4 and 5). With the latter period of time, **3o** was obtained in a quantitative yield. Acetophenones reacted in an analogous way, affording 1,2-dihydroquinoline **3p** in excellent yield with a temperature of 90 °C and longer reaction times (Table 2, entries 6 and 7).

Interestingly, and in contrast with **1b**, aniline **1e** regioselectively reacted with methyl pyruvate (**2l**) to form the single isomer of the tricyclic 1,2-DHQ **3q** in almost quantitative yield (Scheme 6). Although **1e** should not be as reactive as **1b**, the conversion efficiency of the former aniline into **3q** suggests the efficacy of the catalyst when the ketone is activated with the ester group.^{22b} This activation of the ketone seems to be relevant, considering that with dialkyl ketones, such as **2a** and


Scheme 5. Formation of Regioisomers 3l/4l by Condensation of 1b with Methyl Pyruvate (2l)

Table 2. Yields of 1,2-DHQ 3m–p, Obtained by Reaction of *Meta*-Substituted Anilines 1c,d and Ketones 2b,f,h^a

Entry	1 (R ¹)	2 (R ² , R ³)	T (°C)	t (h)	3 (%) ^b
1	1c (3-Cl)	2b (Et, Me)	60	6	3m (45)
2	1d (3-Me)	2b (Et, Me)	60	1	3n (39)
3	1d (3-Me)	2b (Et, Me)	60	7	3n (98)
4	1d (3-Me)	2f (-(CH ₂) ₄ -)	60	1	3o (26)
5	1d (3-Me)	2f (-(CH ₂) ₄ -)	60	7	3o (99)
6	1d (3-Me)	2h (C ₆ H ₅ , Me)	90	5	3p (17)
7	1d (3-Me)	2h (C ₆ H ₅ , Me)	90	19	3p (98)

^aReaction conditions: 1 (1.0 molar equiv), 2 (5.0 molar equiv), MgBr₂ (1.0 molar equiv), and Li₂CO₃ (0.5 molar equiv). ^bAfter column chromatography.

Scheme 6. Synthesis of Tricyclic 1,2-DHQ 3q



2b, the reaction with 1e did not furnish the expected 1,2-DHQ, but instead complex mixtures of products after heating at 90 °C for a longer reaction time than that employed with 2l.

We investigated the reactivity of anilines substituted by electron-donating and electron-withdrawing groups in the *para* position to the amino group. Thus, the reaction of 1f with ketone 2l under milder reaction conditions resulted in the desired 1,2-DHQ 3r in high yield (Table 3, entry 1). Low reactive anilines, such as 1g–i, were assessed with ketones 2b and 2l. These reactions furnished the corresponding 1,2-DHQ 3s–v in low to high yields (Table 3, entries 2–6). As expected, anilines 1h,i bearing electron-withdrawing groups also reacted

with 2b or 2l, albeit more slowly, with recovery of the starting material in some cases (Table 3, entries 5 and 6). Unexpectedly, the reaction between aniline 1i or 3-nitroaniline (1j) with ketone 2l resulted in a complex mixture of products. These results support the idea that the reaction process depends on the electron demand of both anilines and ketones. That is, the reaction is favored with electron-donating substituted anilines and electron-withdrawing substituted ketones.

o-Anisidine (1k) was also evaluated with ketones 2b and 2l. The former ketone reacted under harder conditions (90 °C, 15 h) than the latter (60 °C, 1 h) to give 1,2-dihydroquinolines 3w,x, respectively, in modest to high yields (Scheme 7). These results could be explained by a steric effect of the *ortho*-substituent in the arylamine, which possibly prevents full conjugation of the amino group with the phenyl ring.³⁰ Consequently, this effect could diminish the efficiency of the reaction with 2b, but is not significant enough to affect the cyclization reaction with ketone 2l.


Table 3. Yields of 1,2-DHQ 3r–v, Obtained by Reaction of Substituted Anilines 1f–i and Ketones 2b and 2l^a

1 + 2 **2b**, R² = Et
2l, R² = CO₂Me → **3**

Entry	1 (R ¹)	2 (R ²)	T (°C)	t (h)	3 (%) ^b
1	1f (4-OMe)	2l (CO ₂ Me)	60	1	 3r (82)
2	1g (4-Me)	2l (CO ₂ Me)	60	1	 3s (78)
3	1h (4-CN)	2l (CO ₂ Me)	90	3	 3t (94)
4	1h (4-CN)	2b (Et)	90	15	 3u (26) ^c
5	1i (4-NO ₂)	2b (Et)	90	15	 3v (31) ^c


^aReaction conditions: **1** (1.0 molar equiv), **2** (5.0 molar equiv), MgBr₂ (1.0 molar equiv), and Li₂CO₃ (0.5 molar equiv). ^bAfter column chromatography. ^cStarting aniline is recovered.

Scheme 7. Reaction of Hindered Aniline **1k** with **2b** and **2l** To Yield 1,2-DHQ **3w** and **3x**



Compound **3x** crystallized and its structure was established by X-ray diffraction crystallography (see the Supporting Information). The crystal structure shows a half-chair conformation for the dihydroheterocyclic ring, where the C-2 methoxycarbonyl group adopts an axial conformation, and consequently the methyl group assumes the equatorial conformation. Interestingly, the C-4 methoxycarbonyl group is not completely coplanar to the C-3/C-4 double bond of the heterocycle (torsion angle C(3)–C(4)–C(16)–O(17) = –149.52(15)°).

Scheme 8. Proposed Reaction Mechanisms for the Synthesis 1,2-DHQ **3**, Starting from anilines **1** and Ketones **2**




Two proposed mechanisms for the formation of 1,2-dihydroquinolines are useful to consider^{17,22b,23,31,32} in regard to our conditions and catalysts. Mechanism (1): The previous formation of enone **III**, as the Mg^{2+} -catalyzed self-condensation product of ketone **2** species **I** and **II**, is followed by the conjugate addition of the aniline **1** to yield the β -anilino ketone intermediate **IV**.^{31,32a} This in turn undergoes consecutive cyclization and dehydration reactions,^{22c} again catalyzed by the Mg^{2+} species, to give precursor **V** and the final product **3**.^{32a} (Scheme 8). Mechanism (2): The previous formation of imine species **VI**, as the product of the Mg^{2+} -catalyzed reaction between species **I** and aniline **1**, is followed by the addition of enolate **II** of ketone **2** to yield the β -anilino ketone intermediate **VII**,^{32a} which in turn is converted into the observed 1,2-dihydroquinoline **3** via precursor **V**.²³ (Scheme 8). Of course, both mechanisms can also involve the formation of the corresponding imine and 4-amino species **VIII** and **IX**, respectively, as possible precursors of **3**.^{32a}

The solvent-free conditions of this methodology avoid unfavorable coordinating interactions between the Mg^{2+} ions and the solvent, maintaining the wanted coordinating interactions with the oxygen of the ketone carbonyl group that enhance the electrostatic catalytic effect.³³ This would explain the inefficiency of the assays when a solvent is used (Table 1, entries 18–20). Furthermore, the stronger coordinating ability of Mg^{2+} or $BrMg^+$ ($[Mg]$) ions compared to Li^+ ions prevents a competitive effect between both species for the catalytic sites.³³ The large number of these sites accounts for the amount of catalyst required in the process. It is likely that the bromide ion assists the generation of the base species, such as $LiCO_3^-$, in the middle of the solvent-free reaction. We monitored several reactions attempting to detect or isolate some intermediates under typical reaction conditions. However, we were unable to isolate any of these intermediates or to detect them by 1H NMR analysis. Therefore, in order to test the feasibility of these two mechanisms, some of the plausible stable intermediates were independently prepared or purchased and submitted to the same reaction conditions.

To test the first mechanism, mesityl oxide (**7**) (Mg -free complex of **III**) was treated with *m*-anisidine (**1b**) for 1 h, providing a mixture of 1,2-dihydroquinolines **3a/4a** (98:2). This was purified to give 48% and 1% yields, respectively, along with recovery of the starting material (Scheme 9). Although the


Scheme 9. Reaction of Aniline 1b with Mesityl Oxide (7)



regioselectivity was identical to that obtained with **1b** and acetone (**2a**) as starting materials (Table 1, entry 22), the conversion rate was less, evidenced by the fact that only half of the aniline was consumed in the former reaction.

To test the second mechanism, we examined the in situ generation of the imine Mg complex **VI** as by performing the $Hg(I)$ -catalyzed reaction of aniline **1b** with phenylacetylene (**9**), followed by treatment with sodium borohydride, giving imine **8** in good yield (Scheme 10).³⁴ In contrast with the process depicted in Scheme 9, that of **8** with ketone **2h** turned out to efficiently produce the desired 1,2-dihydroquinoline **3h**

Scheme 10. Formation of Imine 8 and Its Conversion into 1,2-Dihydroquinoline 3h



in quantitative yield. This result provides evidence that imine is the most probable intermediate in the formation of the heterocycles, which is in agreement with other reports.^{22b,23,32,35} Although the formation of the imine can be favored by the presence of electron-withdrawing groups in the benzene ring of the acetophenone (i.e., **2k**), the enhanced stability of the conjugate base species **II** of the latter compound may lead to a decrease in its reactivity and performance in the condensation process toward the heterocycle. This could be the reason for the modest yield of **3k** (Scheme 4). On the other hand, it is expected that the formation of the imine should not be favored by the presence of electron-withdrawing groups in the aniline, due to a decrease in nucleophilicity. This idea is in accordance with the observed depletion of reactivity when anilines **1h,i** reacted with ketones **2b** and **2l** (Table 3).

However, the reaction mechanism may be much more complex than either of those illustrated in Scheme 8, since diverse intermediate species can be associated with the mechanistic steps,³⁶ mainly depending on the starting materials and reaction conditions.^{32a} For instance, recent mechanistic studies were conducted to rationalize the formation of spiro-1,2-dihydroquinoline analogues of **3f,g**,¹⁷ proposing that an isolated *o*-alkenylimino intermediate is involved at the final annulation step toward the observed products. This finding is supported by the fact that the acid- or iodine-catalyzed reactions of *o*-alkenylanilines with cyclic ketones yielded 2-spiro-1,2-dihydroquinoline derivatives.^{32c,d,37} However, in our transformations this kind of intermediate was not detected or isolated, nor were alkenyl-substituted 1,2-dihydroquinoline products¹⁷ α,β -unsaturated ketones^{32a} or 4-amino tetrahydroquinolines.³⁸

It is noteworthy that most of the reported quinoline and 1,2-dihydroquinoline syntheses have been carried out in the presence of solvents, such as benzene, toluene or acetonitrile, which may stabilize some of the proposed intermediates and therefore lead not only to the expected quinoline products but also to a variety of side-products. In contrast, under our solvent-free conditions only anilines and ketones are interacting, of course with the assistance of the catalyst, to give the 1,2-DHQ with high regioselectivity and efficiency, even if some deactivated anilines are used. Consequently, compared to methods involving a condensing phase, it seems that our approach follows a simpler pathway, probably limiting side pathways by the strong magnesium ion coordination. Nevertheless, the first proposal involving the enone **7** cannot be

completely ruled out as a competitive mechanism in view of the complexity of the quinoline formation.^{32a}

The high regioselectivity observed in the cyclization step, affording 1,2-dihydroquinoline **3** as the major and **4** as the minor regioisomer, could be rationalized on the basis of two factors,^{36,39} the electronic effects of the substituents in the aniline ring and the steric hindrance generated by both the aniline and ketone.⁴⁰ The electrophilic aromatic substitution, which is presumably involved in the 1,2-dihydroquinoline heterocyclic ring formation, is usually kinetically controlled during the reaction step leading to the arenium ion intermediate (Wheland σ -complex).⁴¹ Although the *ortho*–*para*-directing groups in anilines such as **1b** reinforce each other to orient the position of the incoming group, the stability of the arenium ion in a *para*-quinoid canonical structure with respect to the methoxy group (attack at the C-6 position) is greater than that in the *ortho*-quinoid structure (attack at the C-2 position).³⁰ Therefore, the fact that **3** is obtained as the major regioisomer in detriment to **4** is expected.^{39,42}

Furthermore, an explanation for this preference is offered on the basis of hard and soft interaction arguments.⁴³ Since the *ortho* position (C-2) is harder than the *para* position (C-6) in a benzene ring monosubstituted by an electron-donating group, then the softer electrophiles give more *para* substitution (C-6). Considering that in the proposed mechanisms (Scheme 8) the soft metal complex species **VII** or **VIII** (rather than harder small positive-charged species) are formed as precursors in the cyclization step, it is likely that the process takes place rather at the C-6 position (*ortho*–*para* orientation), which would favor regioisomer **3**, rather than at the harder C-2 position (*ortho*–*ortho* orientation), which would favor regioisomer **4**. Molecular orbital calculations agree with this prediction indicating that the attack preferentially occurs at the *para* position in anisole, among other electron-donating groups, which has the highest π -electron density and the lowest energy of the π -molecular orbital of the σ -complex.^{43,44} Among other theoretical constructs,⁴⁵ density-functional theory (DFT)-based reactivity criteria and descriptors have been used to clarify this question,⁴⁶ predicting the preferential *para* approach of the electrophile as a result of a stronger tendency to electron-transfer (nucleophilicity) from the aromatic ring.⁴⁷

It is well-known that steric effects tend to reduce the proportion of the *ortho* products.^{30,40a,41} Therefore, if the bulky metal-complexed species **VII** or **VIII** are the reactive species during the cyclization step, it is reasonable that the preference for closing the ring takes place at the C-6 position rather than at the most hindered C-2 position (in between both substituents).^{30,48}

CONCLUSION

In summary, we have developed a highly efficient, versatile, and regioselective solvent-free methodology to prepare 1,2-DHQ **3** through a magnesium bromide-catalyzed condensation of the MCR, with 1 equiv of the aniline and 2 equiv of the ketone. The scope of the method includes a variety of symmetrical and nonsymmetrical alkyl ketones, cyclic ketones, acetophenones, and diverse substituted anilines. Interestingly, the latter substrates were reactive enough to activate the cyclization process, even when having either weak electron-donating or electron-withdrawing substituents or being attached in an inadequate position at the benzene ring. Among the evaluated brominated catalysts, magnesium bromide proved to be the most efficient, leading to the desired products in moderate to

high yields. The synthesis was designed under a solvent-free procedure⁴⁹ and with a nontoxic and inexpensive catalyst. All these advantageous properties of the current approach are attractive in comparison with many of the already known methods, the latter of which are modestly selective,^{16,17} use complex or scope-limited functionalized starting materials,^{23a,24,25,37} require expensive or highly toxic catalysts,^{16,19,20a,22a–c} and/or involve severe and mostly solvent-dependent reaction conditions.^{21,22d} However, our method suffers from low reaction conversion for strongly inactivated anilines and some acyclic hindered ketones. The mechanistic study provides evidence that the most likely pathway includes the formation of an imine derivative, **8**, as the key intermediate, which undergoes condensation by the second equivalent of the ketone, followed by the cyclization process to yield the heterocyclic ring. The reaction between anilines and two different carbonylic starting materials to obtain quinolines and more complex 1,2-DHQ is currently under study, and the results will be reported in due course.

EXPERIMENTAL SECTION⁵⁰

General Procedure for the MgBr₂-Catalyzed Synthesis of 2,2,4-Trialkyl-1,2-dihydroquinolines 3a–g/4a, 3l–o/4l, 3q–s, and 3x (Method A). In a 25-mL round-bottomed flask, a mixture of aniline **1b–g** or **1k** (1.0 molar equiv), ketone, **2a–g** or **2l** (5.0 molar equiv), Li₂CO₃ (0.5 molar equiv), and anhydrous MgBr₂ (1.0 molar equiv) was vigorously stirred at 60 °C for 1–7 h. A 10% aqueous solution of NH₄Cl (5 mL) was added, and the mixture was extracted with CH₂Cl₂ (3 × 10 mL). The organic layer was dried (Na₂SO₄) and the solvent removed under vacuum. The residue was purified by column chromatography over silica gel (15 g/g of crude, hexane) to give either the corresponding 1,2-dihydroquinoline **3a–g**, **3m–o**, **3q–s**, or **3x** or the isomeric mixture of 1,2-dihydroquinolines **3a/4a** or **3l/4l**.

General Procedure for the PyHBr₃-Catalyzed Synthesis of 2,2,4-Trialkyl-1,2-dihydroquinolines 3a–c/4a and 3g (Method B). In a 25-mL round-bottomed flask, a mixture of aniline **1b** (1.0 molar equiv), ketone, **2a–c** or **2g** (5.0 molar equiv), Li₂CO₃ (2.0 molar equiv), and Py·HBr₃ (0.5 molar equiv) was vigorously stirred at 60 °C for 12 h. A 10% aqueous solution of NH₄Cl (5 mL) was added, and the mixture was stirred for 10 min, and extracted with CH₂Cl₂ (3 × 20 mL). The organic layer was dried (Na₂SO₄) and the solvent removed under vacuum. The residue was purified by column chromatography over silica gel (15 g/g of crude, hexane) to give either the corresponding 1,2-dihydroquinoline **3b,c** or **3g** or the isomeric mixture of 1,2-dihydroquinolines **3a/4a**.

General Procedure for the MgBr₂-Catalyzed Synthesis of 2-Alkyl-2,4-diaryl-1,2-dihydroquinolines 3h–k, 3p, 3t–v, and 3w (Method C). In a 25-mL round-bottomed flask, a mixture of aniline **1b**, **1d**, **1h–i**, or **1k** (1.0 molar equiv), acetophenone, **2b**, **2h–k**, or **2l** (5.0 molar equiv), Li₂CO₃ (0.5 molar equiv), and MgBr₂ (1.0 molar equiv) was vigorously stirred at 90 °C for 5–19 h. A 10% aqueous solution of NH₄Cl (5 mL) was added, and the mixture was extracted with CH₂Cl₂ (3 × 10 mL). The organic layer was dried (Na₂SO₄) and the solvent removed under vacuum. The residue was purified by column chromatography over silica gel (15 g/g of crude, hexane/EtOAc, 99:1) to give the corresponding 1,2-dihydroquinoline **3h–k**, **3p**, **3t–v**, or **3w**.

7-Methoxy-2,2,4-trimethyl-1,2-dihydroquinoline (3a) and 5-Methoxy-2,2,4-trimethyl-1,2-dihydroquinoline (4a). Method A. A mixture of **1b** (0.100 g, 0.813 mmol), **2a** (0.235 g, 4.06 mmol), Li₂CO₃ (0.030 g, 0.41 mmol), and MgBr₂ (0.150 g, 0.815 mmol) was heated to 60 °C for 1 h, affording a mixture of **3a/4a** (98:2) which was purified over silica gel (15 g/g of crude, hexane) to give 0.143 g (87%) of **3a** as a white powder, and 0.003 g (2%) of **4a** as a pale yellow powder. **Method B.** A mixture of **1b** (0.150 g, 1.22 mmol), **2a** (0.348 g, 6.00 mmol), Li₂CO₃ (0.180 g, 2.43 mmol), and Py·HBr₃ (0.195 g, 0.61 mmol) afforded a mixture of **3a/4a** (98:2) which was purified over

silica gel (15 g/g of crude, hexane) to give 0.188 g (76%) of **3a** as a white powder and 0.003 g (1%) of **4a** as a pale yellow powder. **Method D.** A mixture of **1b** (0.100 g, 0.813 mmol), **7** (0.400 g, 4.07 mmol), Li_2CO_3 (0.030 g, 0.41 mmol), and MgBr_2 (0.150 g, 0.815 mmol) was vigorously stirred at 60 °C for 1 h. A 10% aqueous solution of NH_4Cl (5 mL) was added, and the mixture was extracted with CH_2Cl_2 (3 × 10 mL). The organic layer was dried (Na_2SO_4) and the solvent removed under vacuum. The residue was purified by column chromatography over silica gel (15 g/g of crude, hexane) to afford a mixture of **3a/4a** (97:3) which was purified over silica gel (15 g/g of crude, hexane) to give 0.079 g (48%) of **3a** as a white powder and 0.0017 g (1%) of **4a** as a pale yellow powder. Data for **3a**: R_f 0.72 (hexane/EtOAc, 7:3); mp 61–62 °C (lit.⁵¹ mp 67–69 °C); IR (KBr) 3386, 2966, 1651, 1612, 1577, 1511, 1483, 1452, 1381, 1336, 1276, 1259, 1208, 1162, 1026, 1000, 832, 811 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.26 (s, 6H, 2Me-C2), 1.95 (d, $J = 1.5$ Hz, 3H, Me-C4), 3.73 (br, 1H, NH), 3.74 (s, 3H, CH_3O), 5.18 (q, $J = 1.5$ Hz, 1H, H-3), 6.01 (d, $J = 2.7$ Hz, 1H, H-8), 6.19 (dd, $J = 8.4, 2.7$ Hz, 1H, H-6), 6.96 (d, $J = 8.4$ Hz, 1H, H-5); ^{13}C NMR (75.4 MHz, CDCl_3) δ 18.6 (C-3-C4), 31.0 (2 $\text{CH}_3\text{-C2}$), 51.9 (C-2), 55.0 (CH_3O), 98.5 (C-8), 102.2 (C-6), 115.3 (C-4a), 124.6 (C-5), 125.9 (C-3), 128.1 (C-4), 144.6 (C-8a), 160.1 (C-7); MS (70 eV) m/z 203 (M^+ , 8), 188 (100), 186 (2), 173 (6), 159 (2), 145 (24); HRMS (EI) m/z [M^+] calcd for $\text{C}_{13}\text{H}_{17}\text{NO}$ 203.1310, found 203.1313. Data of **4a**: R_f 0.75 (hexane/EtOAc, 7:3); mp 81–82 °C. IR (film) 2961, 1640, 1597, 1493, 1464, 1268, 1230, 1162, 1106, 777, 728 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.22 (s, 6H, 2Me-C2), 2.16 (d, $J = 1.5$ Hz, 3H, Me-C4), 3.70 (br, 1H, NH), 3.75 (s, 3H, OCH_3), 5.22 (q, $J = 1.5$ Hz, 1H, H-3), 6.15 (dd, $J = 8.1, 1.1$ Hz, 1H, H-6 or H-8), 6.23 (dd, $J = 8.1, 1.1$ Hz, 1H, H-8 or H-6), 6.92 (t, $J = 8.1$ Hz, 1H, H-7); ^{13}C NMR (75.4 MHz, CDCl_3) δ 23.0 ($\text{CH}_3\text{-C4}$), 29.6 (2 $\text{CH}_3\text{-C2}$), 50.7 (C-2), 55.2 (CH_3O), 101.3 (C-6 or C-8), 107.5 (C-8 or C-6), 111.1 (C-4a), 128.4 (C-7), 128.9 (C-3), 129.3 (C-4), 145.5 (C-8a), 157.6 (C-5); MS (70 eV) m/z 203 (M^+ , 5), 188 (100), 173 (50), 157 (2), 145 (21), 130 (5), 115 (4); HRMS (EI) m/z [M^+] calcd for $\text{C}_{13}\text{H}_{17}\text{NO}$ 203.1310, found 203.1309.

2,4-Diethyl-7-methoxy-2-methyl-1,2-dihydroquinoline (3b). **Method A.** A mixture of **1b** (0.100 g, 0.813 mmol), **2b** (0.292 g, 4.06 mmol), Li_2CO_3 (0.030 g, 0.41 mmol), and MgBr_2 (0.150 g, 0.815 mmol) was heated to 60 °C for 1 h to give 0.184 g (98%) of **3b** as a white powder. **Method B.** A mixture of **1b** (0.200 g, 1.62 mmol), **2b** (0.585 g, 8.12 mmol), Li_2CO_3 (0.060 g, 0.82 mmol), and Py-HBr_3 (0.259 g, 0.81 mmol) gave 0.32 g (85%) of **3b** as a white powder. R_f 0.81 (hexane/EtOAc, 7:3); mp 59–60 °C; IR (film) 3376, 2963, 1649, 1612, 1577, 1514, 1479, 1462, 1373, 1327, 1298, 1270, 1219, 1207, 1166, 1038, 826, 796 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.89 (t, $J = 7.5$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{-C2}$), 1.15 (t, $J = 7.5$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{-C4}$), 1.21 (s, 3H, $\text{CH}_3\text{-C2}$), 1.42–1.56 (m, 2H, $\text{CH}_3\text{CH}_2\text{-C2}$), 2.35 (qd, $J = 7.5, 1.2$ Hz, 2H, $\text{CH}_3\text{CH}_2\text{-C4}$), 3.62 (br s, 1H, NH), 3.74 (s, 3H, CH_3O), 5.07 (t, $J = 1.2$ Hz, 1H, H-3), 6.00 (d, $J = 2.4$ Hz, 1H, H-8), 6.15 (dd, $J = 8.4, 2.4$ Hz, 1H, H-6), 6.99 (d, $J = 8.4$ Hz, 1H, H-5); ^{13}C NMR (75.4 MHz, CDCl_3) δ 8.5 ($\text{CH}_3\text{CH}_2\text{-C2}$), 12.9 ($\text{CH}_3\text{CH}_2\text{-C4}$), 24.6 ($\text{CH}_3\text{CH}_2\text{-C4}$), 29.5 ($\text{CH}_3\text{-C2}$), 36.5 ($\text{CH}_3\text{CH}_2\text{-C2}$), 54.7 (C-2), 55.0 (CH_3O), 98.3 (C-8), 101.7 (C-6), 114.3 (C-4a), 122.6 (C-5), 124.2 (C-3), 134.4 (C-4), 145.3 (C-8a), 160.0 (C-7); MS (70 eV) m/z 231 (M^+ , 1), 216 (15), 202 (100), 187 (9), 158 (8); HRMS (EI) m/z [M^+] calcd for $\text{C}_{15}\text{H}_{21}\text{NO}$ 231.1623, found 231.1624.

7-Methoxy-2-methyl-2,4-di-n-propyl-1,2-dihydroquinoline (3c). **Method A.** A mixture of **1b** (0.051 g, 0.415 mmol), **2c** (0.190 g, 2.21 mmol), Li_2CO_3 (0.015 g, 0.20 mmol) and MgBr_2 (0.075 g, 0.41 mmol) was heated to 60 °C for 1 h to give 0.086 g (80%) of **3c** as a pale brown oil. **Method B.** A mixture of **1b** (0.05 g, 0.4 mmol), **2c** (0.172 g, 2.00 mmol), Li_2CO_3 (0.06 g, 0.8 mmol), and Py-HBr_3 (0.064 g, 0.2 mmol) gave 0.034 g (33%) of **3c** as a pale brown oil: R_f 0.78 (hexane/EtOAc, 7:3); IR (film) 3377, 2956, 2930, 2869, 1648, 1612, 1577, 1514, 1464, 1327, 1297, 1277, 1264, 1211, 1166, 1039, 820, 789 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.86–0.91 (m, 3H, $\text{CH}_3\text{CH}_2\text{CH}_2\text{-C2}$), 0.95 (t, $J = 7.2$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 1.21 (s, 3H, $\text{CH}_3\text{-C2}$), 1.35–1.49 (m, 4H, $\text{CH}_3\text{CH}_2\text{CH}_2\text{-C2}$), 1.56 (sext, $J = 7.2$ Hz, 2H, $\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 2.20–2.39 (m, 2H, $\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 3.61 (br s, 1H, NH), 3.73 (s, 3H, CH_3O), 5.07 (br s, 1H, H-3), 5.98

(d, $J = 2.6$ Hz, 1H, H-8), 6.14 (dd, $J = 8.3, 2.6$ Hz, 1H, H-6), 6.97 (d, $J = 8.3$ Hz, 1H, H-5); ^{13}C NMR (75.4 MHz, CDCl_3) δ 14.0 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 14.5 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C2}$), 17.5 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C2}$), 21.4 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 30.1 ($\text{CH}_3\text{-C2}$), 34.1 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 46.7 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C2}$), 54.6 (C-2), 55.0 (CH_3O), 98.2 (C-8), 101.6 (C-6), 114.1 (C-4a), 124.2 (C-3), 124.3 (C-5), 132.4 (C-4), 145.3 (C-8a), 159.9 (C-7); MS (70 eV) m/z 259 (M^+ , 4), 244 (11), 228 (23), 215 (100), 200 (7), 187 (29). HRMS (EI) m/z [M^+] calcd for $\text{C}_{17}\text{H}_{25}\text{NO}$: 259.1936; found: 259.1930.

2,4-Dibutyl-7-methoxy-2-methyl-1,2-dihydroquinoline (3d).^{18c} **Method A.** A mixture of **1b** (0.050 g, 0.41 mmol), **2d** (0.205 g, 2.05 mmol), Li_2CO_3 (0.015 g, 0.20 mmol) and MgBr_2 (0.075 g, 0.41 mmol) was heated to 60 °C for 1 h to give 0.073 g (62%) of **3d** as a pale yellow oil. R_f 0.76 (hexane/EtOAc, 7:3). IR (film) 3379, 2956, 2930, 2859, 1647, 1613, 1578, 1514, 1465, 1328, 1297, 1273, 1210, 1166, 1046, 825 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.88 (t, $J = 7.5$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{CH}_2\text{-C2}$), 0.93 (t, $J = 7.2$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 1.21 (s, 3H, $\text{CH}_3\text{-C2}$), 1.24–1.57 (m, 10H, 5CH_2), 2.27–2.39 (m, 2H, $\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 3.62 (br s, 1H, NH), 3.74 (s, 3H, CH_3O), 5.08 (s, 1H, H-3), 5.98 (d, $J = 2.4$ Hz, 1H, H-8), 6.15 (dd, $J = 8.4, 2.4$ Hz, 1H, H-6), 6.97 (d, $J = 8.4$ Hz, 1H, H-5); ^{13}C NMR (75.4 MHz, CDCl_3) δ 14.0 (CH_3), 14.1 (CH_3), 22.6 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 23.1 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C2}$), 26.4 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C2}$), 30.0 ($\text{CH}_3\text{-C2}$), 30.6 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 31.8 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C4}$), 43.9 ($\text{CH}_3\text{CH}_2\text{CH}_2\text{-C2}$), 54.5 (C-2), 55.0 (CH_3O), 98.3 (C-8), 101.7 (C-6), 114.2 (C-4a), 124.1 (C-3), 124.3 (C-5), 132.6 (C-4), 145.3 (C-8a), 159.9 (C-7); MS (70 eV) m/z 287 (M^+ , 5), 272 (8), 231 (17), 230 (100), 200 (5), 187 (21), 144 (6). HRMS (EI) m/z [M^+] calcd for $\text{C}_{19}\text{H}_{29}\text{NO}$: 287.2249; found: 287.2255.

2,2,4-Triethyl-7-methoxy-3-methyl-1,2-dihydroquinoline (3e). **Method A.** A mixture of **1b** (0.050 g, 0.41 mmol), **2e** (0.180 g, 2.09 mmol), Li_2CO_3 (0.015 g, 0.20 mmol) and MgBr_2 (0.075 g, 0.41 mmol) was heated to 60 °C for 1 h to give 0.007 g (6%) of **3e** as a pale yellow oil. R_f 0.80 (hexane/EtOAc, 7:3). IR (film) 2964, 2929, 1614, 1462, 1416, 1271, 1218, 1164, 1033, 852, 823 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.91 (t, $J = 7.2$ Hz, 6H, $2\text{CH}_3\text{CH}_2\text{-C2}$), 1.08 (t, $J = 7.5$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{-C4}$), 1.11–1.24 (m, 2H, $\text{CH}_3\text{CH}_2\text{-C2}$), 1.68 (br s, 3H, $\text{CH}_3\text{-C3}$), 1.72–1.85 (m, 2H, $\text{CH}_3\text{CH}_2\text{-C2}$), 2.42 (q, $J = 7.5$ Hz, 2H, $\text{CH}_3\text{CH}_2\text{-C4}$), 3.26 (br s, 1H, NH), 3.73 (s, 3H, CH_3O), 5.90 (d, $J = 2.5$ Hz, 1H, H-8), 6.06 (dd, $J = 8.5, 2.5$ Hz, 1H, H-6), 6.92 (d, $J = 8.5$ Hz, 1H, H-5); ^{13}C NMR (75.4 MHz, CDCl_3) δ 8.6 ($2\text{CH}_3\text{CH}_2\text{-C2}$), 13.1 ($\text{CH}_3\text{CH}_2\text{-C4}$), 13.2 ($\text{CH}_3\text{-C3}$), 20.8 ($\text{CH}_3\text{CH}_2\text{-C4}$), 37.7 ($2\text{CH}_3\text{CH}_2\text{-C2}$), 55.0 (CH_3O), 62.0 (C-2), 96.7 (C-8), 100.5 (C-6), 113.6 (C-4a), 123.7 (C-5), 124.2 (C-3), 132.4 (C-4), 145.8 (C-8a), 159.4 (C-7); MS (70 eV) m/z 259 (M^+ , 1), 230 (5), 229 (45), 228 (100), 212 (5), 200 (11), 184 (7). HRMS (EI) m/z [$\text{M}^+\text{-C}_2\text{H}_5$] calcd for $\text{C}_{15}\text{H}_{20}\text{NO}$: 230.1545; found: 230.1550.

7-Methoxy-1,2,3,5-tetrahydrospiro[cyclopenta[*c*]quinoline-4,1'-cyclopentane] (3f). **Method A.** A mixture of **1b** (0.050 g, 0.41 mmol), **2f** (0.172 g, 2.05 mmol), Li_2CO_3 (0.015 g, 0.20 mmol) and MgBr_2 (0.075 g, 0.41 mmol) was heated to 60 °C for 1 h, and after purification by column chromatography over pretreated silica gel gave 0.103 g (99%) of **3f** as a white solid. R_f 0.75 (hexane/EtOAc, 7:3); mp 107–108 °C [Lit.⁵² 113–116 °C]. IR (film) 3370, 2948, 2838, 1658, 1610, 1516, 1487, 1370, 1335, 1317, 1296, 1268, 1198, 1164, 1072, 1044, 842, 767 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.63–1.81 (m, 6H, H-2', 2H-3', 2H-4', H-5'), 1.82–1.92 (m, 2H, H-2', H-5'), 1.93–2.05 (m, 2H, H-2), 2.44–2.52 (m, 2H, H-1), 2.57–2.66 (m, 2H, H-3), 3.74 (s, 3H, CH_3O), 3.95 (br s, 1H, NH), 6.02 (d, $J = 2.3$ Hz, 1H, H-6), 6.16 (dd, $J = 8.3, 2.3$ Hz, 1H, H-8), 6.78 (d, $J = 8.3$ Hz, 1H, H-9); ^{13}C NMR (75.4 MHz, CDCl_3) δ 22.2 (C-2), 23.8 (C-3', C-4'), 31.1 (C-3), 31.9 (C-1), 39.7 (C-2', C-5'), 55.0 (CH_3O), 65.0 (C-4), 98.1 (C-6), 101.9 (C-8), 114.1 (C-9a), 123.9 (C-9), 131.6 (C-9b), 135.7 (C-3a), 143.8 (C-5a), 159.5 (C-7); MS (70 eV) m/z 255 (M^+ , 24), 254 (10), 227 (22), 212 (22), 213 (47), 183 (15), 135 (12). HRMS (EI) m/z [M^+] calcd for $\text{C}_{17}\text{H}_{21}\text{NO}$: 255.1623; found: 255.1622.

3'-Methoxy-7',8',9',10'-tetrahydro-5'-H-spiro[cyclohexane-1,6'-phenanthridine] (3g). **Method A.** A mixture of **1b** (0.050 g, 0.41 mmol), **2g** (0.201 g, 2.05 mmol), Li_2CO_3 (0.015 g, 0.20 mmol) and

MgBr₂ (0.075 g, 0.41 mmol) was heated to 60 °C for 1 h, and after purification by column chromatography over pretreated silica gel gave 0.115 g (99%) of **3g** as a white solid. **Method B:** A mixture of **1b** (0.050 g, 0.41 mmol), **2g** (0.201 g, 2.05 mmol), Li₂CO₃ (0.060 g, 0.82 mmol) and Py.HBr₃ (0.064 g, 0.2 mmol) gave 0.052 g (45%) of **3g** as a white powder. *R_f* 0.75 (hexane/EtOAc, 7:3); mp 97–98 °C [Lit.⁵³ 95–97 °C]. IR (KBr) 3383, 2925, 2848, 2832, 1640, 1612, 1582, 1518, 1479, 1461, 1320, 1302, 1256, 1207, 1164, 1045, 843 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.26–1.80 (m, 14H, H-2, H-3, H-4, H-5, H-6, H-8', H-9'), 1.98–2.06 (m, 2H, H-10'), 2.22–2.33 (m, 2H, H-7'), 3.69 (s, 3H, CH₃O), 4.32 (br s, 1H, NH), 6.07 (d, *J* = 2.5 Hz, 1H, H-8), 6.17 (dd, *J* = 8.5, 2.5 Hz, 1H, H-6), 6.90 (d, *J* = 8.5 Hz, 1H, H-5); ¹³C NMR (75.4 MHz, CDCl₃) δ 21.0 (C-3, C-5), 22.4 (C-8'), 23.1 (C-9'), 24.6 (C-10'), 25.2 (C-4), 25.3 (C-7'), 32.4 (C-2, C-6), 54.6 (C-6'), 54.9 (CH₃O), 99.3 (C-4'), 102.8 (C-2'), 117.9 (C-10b'), 123.0 (C-1'), 124.8 (C-10a'), 133.0 (C-6a'), 143.3 (C-4a'), 159.1 (C-3'); MS (70 eV) *m/z* 283 (M⁺, 25), 241 (20), 240 (100), 227 (30), 226 (24), 212 (18), 197 (8). HRMS (EI) *m/z* [M⁺] calcd for C₁₉H₂₅NO: 283.1936; found: 283.1936. Anal. Calcd for C₁₉H₂₅NO: C, 80.52; H, 8.89; N, 4.94. Found: C, 80.65; H, 8.97; N, 4.96.

7-Methoxy-2-methyl-2,4-diphenyl-1,2-dihydroquinoline (3h).^{18c} **Method C.** A mixture of **1b** (0.050 g, 0.41 mmol), **2h** (0.24 g, 2.0 mmol), Li₂CO₃ (0.015 g, 0.20 mmol) and MgBr₂ (0.075 g, 0.41 mmol) was heated to 90 °C for 5 h to give 0.13 g (98%) of **3h** as a white powder. **Method E:** In a 10 mL round-bottomed flask, a mixture of imine **8** (0.1210 g, 0.536 mmol), **2h** (0.257 g, 2.14 mmol), Li₂CO₃ (0.0190 g, 0.268 molar equiv) and MgBr₂ (0.098g, 0.536 mmol) was vigorously stirred at 90 °C under N₂ atmosphere for 5 h. A 10% aqueous solution of NH₄Cl (5 mL) was added and the mixture was extracted with CH₂Cl₂ (3 × 10 mL). The organic layer was dried (Na₂SO₄) and the solvent removed under vacuum. The residue was purified by column chromatography over silica gel (2.5 g/g of crude, hexane/EtOAc, 99:1) to give 0.173 (99%) of **3h** as a white powder. *R_f* 0.71 (hexane/EtOAc, 7:3); mp 114–115 °C [Lit.⁵⁴ 108–110 °C]. IR (KBr) 3370, 2961, 2930, 1607, 1575, 1512, 1463, 1443, 1350, 1276, 1223, 1204, 1174, 1125, 1030, 826, 771, 761, 699 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.75 (s, 3H, CH₃-C2), 3.74 (s, 3H, CH₃O), 4.22 (br s, 1H, NH), 5.50 (s, 1H, H-3), 6.09–6.15 (m, 2H, H-6, H-8), 6.82 (d, *J* = 9.0 Hz, 1H, H-5), 7.22 (tt, *J* = 7.4, 1.2 Hz, 1H, H-4'), 7.29–7.36 (m, 7H, H-3', H-2'', H-3'', H-4''), 7.52–7.56 (m, 2H, H-2'); ¹³C NMR (75.4 MHz, CDCl₃) δ 30.1 (CH₃-C2), 55.1 (CH₃O), 57.1 (C-2), 98.7 (C-8), 102.5 (C-6), 114.0 (C-4a), 125.3 (C-2'), 126.6 (C-4'), 126.7 (C-3), 127.2 (C-5), 127.3 (C-4''), 128.1 (PhH), 128.4 (PhH), 128.9 (PhH), 135.3 (C-4), 139.6 (C-1''), 144.5 (C-8a), 148.8 (C-1'), 160.5 (C-7); MS (70 eV) *m/z* 327 (M⁺, 4), 312 (100), 269 (17), 250 (25), 207 (11), 77 (4). HRMS (EI) *m/z* [M⁺] calcd for C₂₃H₂₁NO: 327.1623; found: 327.1624.

7-Methoxy-2-methyl-2,4-di-p-tolyl-1,2-dihydroquinoline (3i).^{18c} **Method C.** A mixture of **1b** (0.050 g, 0.41 mmol), **2i** (0.27 g, 2.0 mmol), Li₂CO₃ (0.015 g, 0.20 mmol) and MgBr₂ (0.075 g, 0.41 mmol) was heated to 90 °C for 5 h to give 0.105 g (73%) of **3i** as a pale yellow oil. *R_f* 0.70 (hexane/EtOAc, 7:3). IR (film) 3376, 2920, 1612, 1509, 1464, 1294, 1276, 1221, 1167, 1114, 1039, 814 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.65 (s, 3H, CH₃-C2), 2.23 (s, 3H, CH₃Ph), 2.29 (s, 3H, CH₃Ph), 3.66 (s, 3H, CH₃O), 4.11 (br s, 1H, NH), 5.40 (s, 1H, H-3), 6.01–6.06 (m, 2H, H-6, H-8), 6.73–6.78 (m, 1H, H-5), 7.02–7.08 (m, 2H, H-3'), 7.06–7.12 (m, 2H, H-3''), 7.12–7.20 (m, 2H, H-2''), 7.31–7.37 (m, 2H, H-2'); ¹³C NMR (75.4 MHz, CDCl₃) δ 20.9 (CH₃Ph-C2), 21.2 (CH₃Ph-C4), 30.2 (CH₃-C2), 55.1 (CH₃O), 56.8 (C-2), 98.6 (C-8), 102.4 (C-6), 114.1 (C-4a), 125.2 (C-2'), 126.5 (C-3), 127.2 (C-5), 128.8 (C-2'', C-3''), 129.0 (C-3'), 135.0 (C-4), 136.3 (C-4'), 136.7 (C-4''), 136.9 (C-1''), 144.6 (C-8a), 146.0 (C-1'), 160.4 (C-7). MS (70 eV) *m/z* 355 (M⁺, 4), 340 (100), 297 (16), 264 (24), 221 (15). HRMS (EI) *m/z* [M⁺] calcd for C₂₅H₂₅NO: 355.1936; found: 355.1943.

7-Methoxy-2-methyl-2,4-bis(4-methoxyphenyl)-1,2-dihydroquinoline (3j). **Method C.** A mixture of **1b** (0.050 g, 0.41 mmol), **2j** (0.300 g, 2.05 mmol), Li₂CO₃ (0.015 g, 0.20 mmol) and MgBr₂ (0.075 g, 0.41 mmol) was heated to 90 °C for 5 h to give 0.108 g (69%) of **3j** as a pale yellow oil. *R_f* 0.73 (hexane/EtOAc, 7:3). IR (film) 3372,

2959, 2931, 1610, 1509, 1463, 1291, 1246, 1176, 1034, 828 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.73 (s, 3H, CH₃-C2), 3.75 (s, 3H, CH₃O), 3.79 (s, 3H, CH₃O), 3.83 (s, 3H, CH₃O), 5.44 (s, 1H, H-3), 6.11 (d, *J* = 2.1 Hz, 1H, H-8), 6.12 (dd, *J* = 8.3, 2.1 Hz, 1H, H-6), 6.84 (d, *J* = 8.4 Hz, 1H, H-5), 6.82–6.88 (m, 2H, H-3'), 6.88–6.96 (m, 2H, H-3''), 7.22–7.31 (m, 2H, H-2'), 7.44–7.51 (m, 2H, H-2''); ¹³C NMR (75.4 MHz, CDCl₃) δ 30.0 (CH₃-C2), 55.08 (CH₃O), 55.22 (CH₃O), 55.25 (CH₃O), 56.6 (C-2), 98.6 (C-8), 102.4 (C-6), 113.5 (C-3' or C-3''), 113.6 (C-3'' or C-3'), 114.2 (C-4a), 126.4 (C-3), 126.5 (C-2'), 127.2 (C-5), 130.0 (C-2''), 132.0 (C-1''), 134.5 (C-4), 141.2 (C-1'), 144.5 (C-8a), 158.3 (C-4'), 158.9 (C-4''), 160.4 (C-7); MS (70 eV) *m/z* 387 (M⁺, 7), 330 (16), 263 (90), 262 (100), 236 (31), 219 (36), 204 (20), 177 (17). HRMS (EI) *m/z* [M⁺] calcd for C₂₅H₂₅NO₃: 387.1834; found: 387.1835.

7-Methoxy-2-methyl-2,4-bis(4-nitrophenyl)-1,2-dihydroquinoline (3k). **Method C.** A mixture of **1b** (0.050 g, 0.41 mmol), **2k** (0.33 g, 2.0 mmol), Li₂CO₃ (0.015 g, 0.20 mmol) and MgBr₂ (0.075 g, 0.41 mmol) was heated to 90 °C for 5 h to give 0.078 g (46%) of **3k** as a pale red oil. *R_f* 0.66 (hexane/EtOAc, 7:3). IR (film) 3381, 2925, 1613, 1596, 1516, 1465, 1346, 1289, 1275, 1223, 1169, 1108, 1037, 852, 698 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.82 (s, 3H, CH₃-C2), 3.79 (s, 3H, CH₃O), 4.38 (br s, 1H, NH), 5.57 (s, 1H, H-3), 6.19 (dd, *J* = 8.6, 2.4 Hz, 1H, H-6), 6.24 (d, *J* = 2.4 Hz, 1H, H-8), 6.70 (d, *J* = 8.6 Hz, 1H, H-5), 7.46–7.52 (m, 2H, H-2''), 7.67–7.73 (m, 2H, H-2'), 8.16–8.20 (m, 2H, H-3'), 8.20–8.26 (m, 2H, H-3''); ¹³C NMR (75.4 MHz, CDCl₃) δ 30.3 (CH₃-C2), 55.5 (CH₃O), 57.5 (C-2), 99.5 (C-8), 103.9 (C-6), 113.1 (C-4a), 123.8 (C-3'), 124.1 (C-3''), 126.3 (C-2'), 126.4 (C-3), 127.5 (C-5), 130.0 (C-2''), 135.53 (C-4a), 144.4 (C-8a), 146.3 (C-1''), 147.1 (C-4'), 147.6 (C-4''), 155.7 (C-1'), 161.5 (C-7); MS (70 eV) *m/z* 417 (M⁺, 2), 371 (100), 356 (18), 340 (28), 284 (46), 270 (32), 241 (66), 152 (25). HRMS (EI) *m/z* [M⁺-CH₃] calcd for C₂₂H₁₆N₃O₅: 402.1090; found: 402.1095.

Dimethyl 7-Methoxy-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate (3l). **Dimethyl 5-Methoxy-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate (4l).** **Method A.** A mixture of **1b** (0.050 g, 0.41 mmol), **2l** (0.23 g, 2.25 mmol), Li₂CO₃ (0.015 g, 0.20 mmol) and MgBr₂ (0.075 g, 0.41 mmol) was heated to 60 °C for 1 h to give 0.065 g (55%) of **3l** as a yellow solid and 0.017 g (15%) of **4l** as a white solid. Data of **3l**: *R_f* 0.55 (hexane/EtOAc, 7:3); mp 106–107 °C. IR (film) 3369, 2953, 1723, 1614, 1517, 1438, 1272, 1232, 1198, 1170, 1146, 1117, 1022, 838, 800 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.55 (s, 3H, CH₃-C2''), 3.74 (s, 3H, CO₂CH₃-1), 3.77 (s, 3H, CH₃O-7''), 3.85 (s, 3H, CO₂CH₃-1'), 4.52 (br s, 1H, NH), 6.18 (d, *J* = 2.6 Hz, 1H, H-8''), 6.30 (dd, *J* = 8.8, 2.6 Hz, 1H, H-6''), 6.54 (br s, 1H, H-3''), 7.73 (d, *J* = 8.8 Hz, 1H, H-5''); ¹³C NMR (75.4 MHz, CDCl₃) δ 27.5 (CH₃-C2''), 52.0 (CO₂CH₃-1'), 52.8 (CO₂CH₃-1), 55.1 (CH₃O-7''), 58.6 (C-2''), 99.4 (C-8''), 104.5 (C-6''), 109.9 (C-4a''), 118.0 (C-8a''), 127.9 (C-5''), 130.0 (C-3''), 144.1 (C-4''), 160.9 (C-7''), 166.3 (CO₂CH₃-1'), 174.6 (CO₂CH₃-1); MS (70 eV) *m/z* 291 (M⁺, 1), 276 (1), 246 (5), 232 (100), 189 (5), 173 (10), 130 (5). HRMS (EI) *m/z* [M⁺] calcd for C₁₅H₁₇NO₅: 291.1107; found: 291.1106. Data of **4l**: *R_f* 0.48 (hexane/EtOAc, 7:3); mp 144–145 °C. IR (KBr) 3368, 2947, 1725, 1636, 1602, 1502, 1469, 1435, 1358, 1324, 1274, 1246, 1198, 1173, 1127, 1089, 1026, 967, 758, 737 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.52 (s, 3H, CH₃-C2''), 3.73 (s, 3H, CO₂CH₃-1), 3.75 (s, 3H, CH₃O-5''), 3.80 (s, 3H, CO₂CH₃-1'), 4.57 (br s, 1H, NH), 6.04 (d, *J* = 1.8 Hz, 1H, H-3''), 6.27 (d, *J* = 8.1 Hz, 1H, H-6''), 6.33 (dd, *J* = 8.1, 0.9 Hz, 1H, H-8''), 7.05 (t, *J* = 8.1 Hz, 1H, H-7''); ¹³C NMR (75.4 MHz, CDCl₃) δ 26.7 (CH₃-C2''), 52.0 (CO₂CH₃-1), 52.7 (CO₂CH₃-1), 55.8 (CH₃O-5''), 57.8 (C-2''), 101.7 (C-6''), 104.7 (C-4a''), 107.8 (C-8''), 126.3 (C-3''), 129.1 (C-4''), 130.4 (C-7''), 144.1 (C-8a''), 155.7 (C-5''), 169.6 (CO₂CH₃-1'), 174.6 (CO₂CH₃-1); MS (70 eV) *m/z* 291 (M⁺, 1), 276 (2), 260 (3), 232 (100), 204 (2), 186 (3), 172 (7), 159 (12), 143 (4), 131 (4). HRMS (EI) *m/z* [M⁺-OMe] calcd for C₁₄H₁₄NO₄: 260.0923; found: 260.0916.

7-Chloro-2,4-diethyl-2-methyl-1,2-dihydroquinoline (3m).⁵⁵ **Method A.** A mixture of **1c** (0.050 g, 0.39 mmol), **2b** (0.140 g, 1.95 mmol), Li₂CO₃ (0.015 g, 0.20 mmol) and MgBr₂ (0.072 g, 0.39 mmol) was heated to 60 °C for 6 h to give 0.041 g (45%) of **3m** as a pale yellow oil. *R_f* 0.95 (hexane/EtOAc, 7:3). IR (film) 3391, 2965,

2924, 1650, 1597, 1569, 1493, 1478, 1454, 1376, 1305, 1161, 1091, 880, 837 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 0.89 (t, $J = 7.4$, 3H, $\text{CH}_3\text{CH}_2\text{-C}2$), 1.13 (t, $J = 7.2$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{-C}4$), 1.22 (s, 3H, $\text{CH}_3\text{-C}2$), 1.41–1.56 (m, 2H, $\text{CH}_3\text{CH}_2\text{-C}2$), 2.34 (qd, $J = 7.2$, 1.1 Hz, 2H, $\text{CH}_3\text{CH}_2\text{-C}4$), 3.63 (br s, 1H, NH), 5.17 (s, 1H, H-3), 6.39 (d, $J = 2.0$ Hz, 1H, H-8), 6.51 (dd, $J = 8.2$, 2.0 Hz, 1H, H-6), 6.95 (d, $J = 8.2$ Hz, 1H, H-5); ^{13}C NMR (75.4 MHz, CDCl_3) δ 8.5 ($\text{CH}_3\text{CH}_2\text{-C}2$), 12.7 ($\text{CH}_3\text{CH}_2\text{-C}4$), 24.5 ($\text{CH}_3\text{CH}_2\text{-C}4$), 29.7 ($\text{CH}_3\text{-C}2$), 36.7 ($\text{CH}_3\text{CH}_2\text{-C}2$), 54.9 (C-2), 112.1 (C-8), 116.3 (C-6), 119.0 (C-4a), 124.2 (C-5), 125.0 (C-3), 133.3 (C-4), 134.1 (C-7), 145.0 (C-8a); MS (70 eV) m/z 236 (M^+ , 1), 207 (38), 205 (100), 190 (25), 170 (13), 154 (22), 140 (12), 127 (17). HRMS (EI) m/z [M^+] calcd for $\text{C}_{14}\text{H}_{18}\text{ClN}$: 235.1128; found: 235.1139.

2,4-Diethyl-2,7-dimethyl-1,2-dihydroquinoline (3n). Method A. A mixture of **1d** (0.15 g, 1.4 mmol), **2b** (0.504 g, 7.00 mmol), Li_2CO_3 (0.052 g, 0.70 mmol) and MgBr_2 (0.258 g, 1.40 mmol) was heated to 60 °C for 7 h to give 0.295 g (98%) of **3n** as a pale yellow oil. R_f 0.80 (hexane/EtOAc, 7:3). IR (film) 3377, 2964, 2918, 1648, 1614, 1475, 1452, 1374, 1321, 841, 802 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 0.90 (t, $J = 7.5$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{-C}2$), 1.15 (t, $J = 7.5$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{-C}4$), 1.21 (s, 3H, $\text{CH}_3\text{-C}2$), 1.38–1.60 (m, 2H, $\text{CH}_3\text{CH}_2\text{-C}2$), 2.21 (s, 3H, $\text{CH}_3\text{-C}7$), 2.30–2.41 (m, 2H, $\text{CH}_3\text{CH}_2\text{-C}4$), 3.53 (br s, 1H, NH), 5.14 (s, 1H, H-3), 6.26 (d, $J = 0.9$ Hz, 1H, H-8), 6.41 (dd, $J = 7.8$, 0.9 Hz, 1H, H-6), 6.97 (d, $J = 7.8$ Hz, 1H, H-5); ^{13}C NMR (125 MHz, CDCl_3) δ 8.5 ($\text{CH}_3\text{CH}_2\text{-C}2$), 12.9 ($\text{CH}_3\text{CH}_2\text{-C}4$), 21.3 ($\text{CH}_3\text{-C}7$), 24.6 ($\text{CH}_3\text{CH}_2\text{-C}4$), 29.4 ($\text{CH}_3\text{-C}2$), 36.4 ($\text{CH}_3\text{CH}_2\text{-C}2$), 54.5 (C-2), 113.4 (C-8), 117.5 (C-6), 118.0 (C-4a), 123.1 (C-5), 124.1 (C-3), 134.7 (C-4), 138.1 (C-7), 143.8 (C-8a); MS (70 eV) m/z 215 (M^+ , 4), 200 (14), 186 (100), 171 (27), 154 (10), 144 (6), 128 (7). HRMS (EI) m/z [M^+] calcd for $\text{C}_{15}\text{H}_{21}\text{N}$: 215.1674; found: 215.1677.

7-Methyl-1,2,3,5-tetrahydrospiro[cyclopent[a]quinoline-4,1'-cyclopentane] (3o). Method A. A mixture of **1d** (0.100 g, 0.93 mmol), **2f** (0.391 g, 4.65 mmol), Li_2CO_3 (0.034 g, 0.46 mmol) and MgBr_2 (0.171 g, 0.93 mmol) was heated to 60 °C for 7 h to give 0.221 g (99%) of **3o** as a white solid. R_f 0.80 (hexane/EtOAc, 7:3); mp 102–103 °C. IR (film) 3391, 2951, 2864, 1658, 1610, 1595, 1509, 1458, 1366, 1168, 884, 808 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.60–1.80 (m, 6H, H-2', 2H-3', 2H-4', H-5'), 1.80–1.92 (m, 2H, H-2', H-5'), 1.94–2.06 (m, 2H, H-2), 2.20 (s, 3H, $\text{CH}_3\text{-C}7$), 2.44–2.54 (m, 2H, H-1), 2.59–2.68 (m, 2H, H-3), 3.85 (br, 1H, NH) 6.27 (d, $J = 0.6$ Hz, 1H, H-6), 6.42 (dd, $J = 7.2$, 0.6 Hz, 1H, H-8), 6.76 (d, $J = 7.2$ Hz, 1H, H-9); ^{13}C NMR (75.4 MHz, CDCl_3) δ 21.4 ($\text{CH}_3\text{-C}7$), 22.3 (C-2), 23.8 (C-3', C4'), 31.1 (C-3), 32.0 (C-1), 39.6 (C-2', C-5'), 65.0 (C-4), 112.8 (C-6), 117.9 (C-8), 123.1 (C-9), 132.0 (C-9b), 137.3 (C-7), 137.5 (C-3a), 142.5 (C-5a); MS (70 eV) m/z 239 (M^+ , 2), 224 (10), 210 (22), 181 (18), 167 (8). HRMS (EI) m/z [M^+] calcd for $\text{C}_{17}\text{H}_{21}\text{N}$: 239.1674; found: 239.1672.

2,7-Dimethyl-2,4-diphenyl-1,2-dihydroquinoline (3p). ^{18c,54} Method C. A mixture of **1d** (0.100 g, 0.93 mmol), **2h** (0.561 g, 4.67 mmol), Li_2CO_3 (0.034 g, 0.47 mmol) and MgBr_2 (0.171 g, 0.93 mmol) was heated to 90 °C for 19 h to give 0.285 g (98%) of **3p** as a yellow oil. R_f 0.78 (hexane/EtOAc, 7:3). IR (film) 3378, 3025, 2968, 2921, 1615, 1490, 1467, 1443, 1320, 1181, 1028, 805, 763, 699 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.75 (s, 3H, $\text{CH}_3\text{-C}2$), 2.22 (s, 3H, $\text{CH}_3\text{-C}7$), 4.12 (br s, 1H, NH), 5.50 (s, 1H, H-3), 6.36–6.37 (m, 2H, H-6, H-8), 6.78 (d, $J = 8.0$ Hz, 1H, H-5), 7.22 (tt, $J = 8.0$ Hz, 1H, H-4'), 7.28–7.38 (m, 7H, H-3', H-2'', H-3'', H-4''), 7.54–7.56 (m, 2H, H-2'); ^{13}C NMR (125 MHz, CDCl_3) δ 21.4 ($\text{CH}_3\text{-C}7$), 30.1 ($\text{CH}_3\text{-C}2$), 57.1 (C-2), 113.7 (C-8), 117.7 (C-4a), 118.1 (C-6), 125.4 (C-2'), 126.0 (C-5), 126.7 (PhH), 127.3 (PhH), 128.1 (C-3), 128.1 (PhH), 128.4 (PhH), 128.9 (PhH), 135.6 (C-4), 139.0 (C-7), 139.6 (C-1'), 143.1 (C-8a), 148.9 (C-1'); MS (70 eV) m/z 296 (M^+ -15, 54), 280 (8), 234 (100), 218 (23), 204 (16), 189 (8), 77 (44). HRMS (EI) m/z [M^+] calcd for $\text{C}_{23}\text{H}_{21}\text{N}$: 311.1674; found: 311.1675.

Dimethyl 3-Methyl-3,4-dihydrobenzo[f]quinoline-1,3-dicarboxylate (3q). Method A. A mixture of **1e** (0.100 g, 0.70 mmol), **2l** (0.356 g, 3.50 mmol), Li_2CO_3 (0.026 g, 0.35 mmol) and MgBr_2 (0.129 g, 0.70 mmol) was heated to 60 °C for 1 h to give 0.208 g (96%) of **3q** as a yellow solid. R_f 0.32 (hexane/EtOAc, 7:3); mp 179–180 °C. IR (film) 3365, 1720, 1625, 1519, 1446, 1386, 1243, 1195, 1173, 1120, 816, 748

cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 1.56 (s, 3H, $\text{CH}_3\text{-C}3$), 3.72 (s, 3H, $\text{CO}_2\text{CH}_3\text{-C}3$), 3.79 (s, 3H, $\text{CO}_2\text{CH}_3\text{-C}1$), 4.86 (br s, 1H, NH), 6.43 (s, 1H, H-2), 6.97 (d, $J = 8.3$ Hz, 1H, H-5), 7.23 (td, $J = 8.3$, 1.0 Hz, 1H, H-8), 7.34 (td, $J = 8.3$, 1.0 Hz, 1H, H-9), 7.38 (br d, $J = 8.3$ Hz, 1H, H-10), 7.63 (d, $J = 8.3$ Hz, 1H, H-6), 7.67 (d, $J = 8.3$ Hz, 1H, H-7); ^{13}C NMR (125 MHz, CDCl_3) δ 25.4 ($\text{CH}_3\text{-C}3$), 52.2 ($\text{CO}_2\text{CH}_3\text{-C}3$), 52.8 ($\text{CO}_2\text{CH}_3\text{-C}1$), 57.8 (C-3), 110.5 (C-10b), 117.0 (C-5), 122.5 (C-8), 123.2 (C-10), 126.2 (C-9), 128.6 (C-6a), 128.7 (C-7), 129.6 (C-2), 130.0 (C-10a), 130.4 (C-1), 131.0 (C-6), 142.1 (C-4a), 169.6 ($\text{CO}_2\text{CH}_3\text{-C}1$), 174.5 ($\text{CO}_2\text{CH}_3\text{-C}3$); MS (70 eV) m/z 311 (M^+ , 3), 269 (5), 253 (21), 252 (100), 236 (4), 192 (12), 91 (13), 86 (13). HRMS (EI) m/z [M^+] calcd for $\text{C}_{18}\text{H}_{17}\text{NO}_4$: 311.1158; found: 311.1151.

Dimethyl 6-Methoxy-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate (3r). ^{22b,d,23} Method A. A mixture of **1f** (0.100 g, 0.81 mmol), **2l** (0.414 g, 4.06 mmol), Li_2CO_3 (0.03 g, 0.4 mmol) and MgBr_2 (0.149 g, 0.81 mmol) was heated to 60 °C for 1 h to give 0.194 g (82%) of **3r** as a yellow oil. R_f 0.56 (hexane/EtOAc, 7:3). IR (film) 3368, 2916, 1723, 1626, 1498, 1436, 1220, 1154, 1115, 1035, 814, 737 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.55 (s, 3H, $\text{CH}_3\text{-C}2$), 3.72 (s, 3H, $\text{CO}_2\text{CH}_3\text{-C}2$), 3.75 (s, 3H, CH_3O), 3.86 (s, 3H, $\text{CO}_2\text{CH}_3\text{-C}4$), 4.41 (br s, 1H, NH), 6.69 (d, $J = 8.7$ Hz, 1H, H-8), 6.72 (dd, $J = 8.7$, 3.0 Hz, 1H, H-7), 6.75 (s, 1H, H-3), 7.50 (d, $J = 3.0$ Hz, 1H, H-5); ^{13}C NMR (75.4 MHz, CDCl_3) δ 26.8 ($\text{CH}_3\text{-C}2$), 52.0 ($\text{CO}_2\text{CH}_3\text{-C}2$), 55.6 (CH_3O), 58.5 (C-2), 111.4 (C-5), 115.1 (C-8), 116.1 (C-7), 117.3 (C-4a), 127.8 (C-4), 134.2 (C-3), 136.6 (C-8a), 152.5 (C-6), 165.3 ($\text{CO}_2\text{CH}_3\text{-C}4$), 174.6 ($\text{CO}_2\text{CH}_3\text{-C}2$); MS (70 eV) m/z 291 (M^+ , 1), 283 (1), 217 (16), 216 (100), 215 (52), 188 (16), 184 (22), 157 (32), 156 (65), 155 (23), 129 (17). HRMS (EI) m/z [M^+ -MeOH] calcd for $\text{C}_{14}\text{H}_{14}\text{NO}_4$: 260.0923; found: 260.0923.

Dimethyl 2,6-Dimethyl-1,2-dihydroquinoline-2,4-dicarboxylate (3s). ^{22b} Method A. A mixture of **1g** (0.100 g, 0.93 mmol), **2l** (0.476 g, 4.67 mmol), Li_2CO_3 (0.034 g, 0.46 mmol) and MgBr_2 (0.171 g, 0.93 mmol) was heated to 60 °C for 1 h to give 0.20 g (78%) of **3s** as a yellow oil. R_f 0.57 (hexane/EtOAc, 7:3). IR (film) 3370, 2952, 1724, 1627, 1499, 1437, 1272, 1221, 1154, 1116, 1035, 814, 780, 737 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.54 (s, 3H, $\text{CH}_3\text{-C}2$), 2.23 (s, 3H, $\text{CH}_3\text{-C}6$), 3.73 (s, 3H, $\text{CO}_2\text{CH}_3\text{-C}2$), 3.86 (s, 3H, $\text{CO}_2\text{CH}_3\text{-C}4$), 4.41 (br s, 1H, NH), 6.55 (d, $J = 8.0$ Hz, 1H, H-8), 6.65 (d, $J = 2.0$ Hz, 1H, H-3), 6.90 (dd, $J = 8.0$, 2.0 Hz, 1H, H-7), 7.60 (br s, 1H, H-5); ^{13}C NMR (125 MHz, CDCl_3) δ 20.8 ($\text{CH}_3\text{-C}6$), 27.2 ($\text{CH}_3\text{-C}2$), 52.0 ($\text{CO}_2\text{CH}_3\text{-C}4$), 52.7 ($\text{CO}_2\text{CH}_3\text{-C}2$), 58.5 (C-2), 114.2 (C-8), 116.5 (C-4a), 126.8 (C-5), 127.9 (C-6), 128.3 (C-4), 130.3 (C-7), 132.9 (C-3), 140.2 (C-8a), 166.2 ($\text{CO}_2\text{CH}_3\text{-C}4$), 174.6 ($\text{CO}_2\text{CH}_3\text{-C}2$); MS (70 eV) m/z 275 (M^+ , 2), 216 (60), 215 (80), 184 (37), 156 (100), 155 (36), 129 (26), 115 (14). HRMS (EI) m/z [M^+] calcd for $\text{C}_{15}\text{H}_{17}\text{NO}_4$: 275.1158; found: 275.1159.

Dimethyl 6-Cyano-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate (3t). ^{22d} Method C. A mixture of **1h** (0.300 g, 2.54 mmol), **2l** (1.301 g, 12.75 mmol), Li_2CO_3 (0.094 g, 1.27 mmol) and MgBr_2 (0.467 g, 2.54 mmol) was heated to 90 °C for 5 h to give 0.69 g (94%) of **3t** as a greenish solid. R_f 0.42 (hexane/EtOAc, 7:3); mp 139–140 °C. IR (film) 3357, 2954, 2217, 1720, 1629, 1602, 1498, 1438, 1282, 1217, 1149, 1116, 1032, 823, 779, 736 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.60 (s, 3H, $\text{CH}_3\text{-C}2$), 3.79 (s, 3H, $\text{CO}_2\text{CH}_3\text{-C}2$), 3.89 (s, 3H, $\text{CO}_2\text{CH}_3\text{-C}4$), 4.93 (br s, 1H, NH), 6.60 (d, $J = 8.3$ Hz, 1H, H-8), 6.77 (d, $J = 2.1$ Hz, 1H, H-3), 7.32 (dd, $J = 8.3$, 1.7 Hz, 1H, H-7), 8.20 (d, $J = 1.7$ Hz, 1H, H-5); ^{13}C NMR (75.4 MHz, CDCl_3) δ 28.3 ($\text{CH}_3\text{-C}2$), 52.4 ($\text{CO}_2\text{CH}_3\text{-C}4$), 53.2 ($\text{CO}_2\text{CH}_3\text{-C}2$), 58.9 (C-2), 100.7 (C-6), 114.1 (C-8), 115.6 (C-4a), 119.9 (CN), 126.4 (C-4), 131.1 (C-5), 133.5 (C-7), 133.6 (C-3), 145.8 (C-8a), 165.2 ($\text{CO}_2\text{CH}_3\text{-C}4$), 173.3 ($\text{CO}_2\text{CH}_3\text{-C}2$); MS (70 eV) m/z 286 (M^+ , 1), 228 (16), 227 (100), 213 (14), 199 (19), 185 (9), 168 (16). HRMS (EI) m/z [M^+] calcd for $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_4$: 286.0954, found 286.0966.

2,6-Diethyl-2-methyl-1,2-dihydroquinoline-6-carbonitrile (3u). Method C. A mixture of **1h** (0.300 g, 2.54 mmol), **2b** (0.918 g, 12.75 mmol), Li_2CO_3 (0.094 g, 1.27 mmol), and MgBr_2 (0.467 g, 2.54 mmol) was heated to 90 °C for 15 h to give 0.15 g (26%) of **3u** as a yellow solid. R_f 0.67 (hexane/EtOAc, 7:3); mp 105–106 °C. IR (film) 3342, 2965, 2210, 1652, 1599, 1503, 1450, 1333, 1309, 1185, 891, 816

cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.91 (t, *J* = 7.5 Hz, 3H, CH₃CH₂-C2), 1.16 (t, *J* = 7.5 Hz, 3H, CH₃CH₂-C4), 1.28 (s, 3H, CH₃-C2), 1.42–1.60 (m, 2H, CH₃CH₂-C2), 2.35 (qd, *J* = 7.5, 1.2 Hz, 2H, CH₃CH₂-C4), 4.05 (br s, 1H, NH), 5.21 (br s, 1H, H-3), 6.35 (d, *J* = 8.4 Hz, 1H, H-8), 7.19 (dd, *J* = 8.4, 2.1 Hz, 1H, H-7), 7.27 (d, *J* = 2.1 Hz, 1H, H-5); ¹³C NMR (75.4 MHz, CDCl₃) δ 8.5 (CH₃CH₂-C2), 12.5 (CH₃CH₂-C4), 24.2 (CH₃CH₂-C4), 30.8 (CH₃-C2), 37.5 (CH₃CH₂-C2), 55.7 (C-2), 97.7 (C-6), 112.1 (C-8), 119.8 (C-4a), 121.0 (CN), 125.5 (C-3), 127.2 (C-5), 132.6 (C-7), 133.5 (C-4), 147.5 (C-8a); MS (70 eV) *m/z* 226 (M⁺, 1), 211 (12), 198 (15), 197 (100), 196 (9), 183 (6), 182 (26), 181 (6). HRMS (EI) *m/z* [M⁺] calcd for C₁₃H₁₈N₂ 226.1470, found 226.1480.

2,6-Diethyl-2-methyl-6-nitro-1,2-dihydroquinoline (3v). Method C. A mixture of **1i** (0.300 g, 2.16 mmol), **2b** (0.778 g, 10.80 mmol), Li₂CO₃ (0.081 g, 1.08 mmol), and MgBr₂ (0.397 g, 2.16 mmol) was heated to 90 °C for 15 h to give 0.165 g (31%) of **3v** as reddish crystals: *R*_f 0.65 (hexane/EtOAc, 7:3); mp 150–151 °C. IR (film) 3324, 2966, 1656, 1602, 1582, 1533, 1505, 1460, 1368, 1283, 1251, 1154, 1108, 821, 748, 734, 651 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 0.91 (t, *J* = 7.5 Hz, 3H, CH₃CH₂-C2), 1.19 (t, *J* = 7.5 Hz, 3H, CH₃CH₂-C4), 1.32 (s, 3H, CH₃-C2), 1.48–1.61 (m, 2H, CH₃CH₂-C2), 2.42 (m, 2H, CH₃CH₂-C4), 4.53 (br s, 1H, NH), 5.25 (br s, 1H, H-3), 6.34 (d, *J* = 9.0 Hz, 1H, H-8), 7.89 (dd, *J* = 9.0, 2.5 Hz, 1H, H-7), 6.96 (d, *J* = 2.5 Hz, 1H, H-5); ¹³C NMR (125 MHz, CDCl₃) δ 8.5 (CH₃CH₂-C2), 12.4 (CH₃CH₂-C4), 24.3 (CH₃CH₂-C4), 31.1 (CH₃-C2), 37.6 (CH₃CH₂-C2), 56.3 (C-2), 111.1 (C-8), 118.6 (C-4a), 119.8 (C-5), 125.4 (C-3), 125.7 (C-7), 133.6 (C-4), 137.4 (C-6), 149.9 (C-8a); MS (70 eV) *m/z* 246 (M⁺, 1), 231 (9), 218 (14), 217 (100), 185 (24), 172 (14), 171 (78), 170 (15), 159 (8); HRMS (EI) *m/z* [M⁺] calcd for C₁₄H₁₈N₂O₂ 246.1368, found 246.1360.

2,6-Diethyl-8-methoxy-2-methyl-1,2-dihydroquinoline (3w). Method C. A mixture of **1k** (0.300 g, 2.44 mmol), **2b** (0.88 g, 12.2 mmol), Li₂CO₃ (0.090 g, 1.22 mmol), and MgBr₂ (0.449 g, 2.44 mmol) was heated to 90 °C for 15 h to give 0.24 g (42%) of **3w** as a greenish oil: *R*_f 0.81 (hexane/EtOAc, 7:3); IR (film) 3406, 2964, 2933, 1646, 1608, 1578, 1488, 1460, 1375, 1256, 1217, 1044, 993, 968, 915, 843, 777, 728, 635 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 0.90 (t, *J* = 7.5 Hz, 3H, CH₃CH₂-C2), 1.15 (t, *J* = 7.5 Hz, 3H, CH₃CH₂-C4), 1.23 (s, 3H, CH₃-C2), 1.45–1.57 (m, 2H, CH₃CH₂-C2), 2.35–2.45 (m, 2H, CH₃CH₂-C4), 3.82 (s, 3H, CH₃O), 4.16 (br s, 1H, NH), 5.19 (br s, 1H, H-3), 6.53 (t, *J* = 8.0 Hz, 1H, H-6), 6.64 (d, *J* = 8.0 Hz, 1H, H-7), 6.77 (d, *J* = 8.0 Hz, 1H, H-5); ¹³C NMR (125 MHz, CDCl₃) δ 8.5 (CH₃CH₂-C2), 13.0 (CH₃CH₂-C4), 24.9 (CH₃CH₂-C4), 29.5 (CH₃-C2), 36.4 (CH₃CH₂-C2), 54.1 (C-2), 55.6 (CH₃O), 109.4 (C-7), 115.0 (C-6), 115.9 (C-5), 120.5 (C-4a), 125.0 (C-3), 133.7 (C-8a), 134.9 (C-4), 145.4 (C-8); MS (70 eV) *m/z* 231 (M⁺, 8), 216 (10), 203 (15), 202 (100), 185 (8), 187 (44), 186 (9), 172 (6); HRMS (EI) *m/z* [M⁺] calcd for C₁₅H₂₁NO 231.1623, found 231.1615.

Dimethyl 8-Methoxy-2-methyl-1,2-dihydroquinoline-2,4-dicarboxylate (3x). Method A. A mixture of **1k** (0.300 g, 2.44 mmol), **2l** (1.24 g, 12.2 mmol), Li₂CO₃ (0.090 g, 1.22 mmol), and MgBr₂ (0.440 g, 2.39 mmol) was heated to 60 °C for 1 h to give 0.61 g (86%) of **3x** as a greenish solid: *R*_f 0.54 (hexane/EtOAc, 7:3); mp 99–100 °C; IR (KBr) 3372, 3080, 2953, 1726, 1630, 1580, 1452, 1378, 1238, 1121, 1082, 1030, 959, 841, 737 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.58 (s, 3H, CH₃-C2), 3.74 (s, 3H, CO₂CH₃-C2), 3.85 (s, 3H, CO₂CH₃-4), 3.87 (s, 3H, CH₃O), 5.12 (br s, 1H, NH), 6.67 (d, *J* = 2.1 Hz, 1H, H-3), 6.67 (t, *J* = 8.1 Hz, 1H, H-6), 6.74 (dd, *J* = 8.1, 1.5 Hz, 1H, H-7), 7.43 (br dd, *J* = 8.1, 0.9 Hz, 1H, H-5); ¹³C NMR (75.4 MHz, CDCl₃) δ 27.5 (CH₃-C2), 52.0 (CO₂CH₃-C4), 52.7 (CO₂CH₃-C2), 55.7 (CH₃O), 58.3 (C-2), 110.5 (C-7), 116.2 (C-4a), 117.2 (C-6), 118.6 (C-5), 128.2 (C-4), 132.6 (C-8a), 132.7 (C-3), 145.9 (C-8), 166.2 (CO₂CH₃-C4), 174.5 (CO₂CH₃-C2); MS (70 eV) *m/z* 291 (M⁺, 1), 276 (2), 246 (5), 232 (100), 217 (32), 159 (17), 131 (4), 103 (3); HRMS (EI) *m/z* [M⁺] calcd for C₁₅H₁₇NO₅ 291.1107, found 291.1121.

(E)-3-Methoxy-N-(1-phenylethylidene)aniline (8). In a 25-mL round-bottomed flask, a mixture of **1b** (0.388 g, 3.15 mmol), phenylacetylene (**9**) (0.082 g, 0.80 mmol), and mercury(I) chloride (0.035 g, 0.074 mmol) was vigorously stirred at 20 °C for 24 h. A 3 M

aqueous solution of KOH (10 mL) and NaBH₄ (0.0076 g, 0.20 mmol) was added, and the mixture was stirred at 20 °C for 10 min and then filtered followed by the extraction of the aqueous layer with EtOAc (3 × 20 mL). The combined organic layers were dried (Na₂SO₄), and the solvent was removed under vacuum. The excess of **1b** was removed by distillation with a Kugelrohr apparatus, giving a pure residue of **8** (0.179 g, 99%) as a pale yellow oil: *R*_f 0.75 (hexane/EtOAc, 7:3); IR (film) 1634, 1595, 1579, 1483, 1447, 1287, 1261, 1147, 1043, 912, 851, 764, 693 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 2.27 (s, 3H, CH₃C=N), 3.83 (s, 3H, CH₃O), 6.38–6.46 (m, 2H, H-2, H-4), 6.68 (dd, *J* = 8.1, 2.4 Hz, 1H, H-6), 7.29 (t, *J* = 8.1 Hz, 1H, H-5), 7.42–7.54 (m, 3H, H-3', H-4'), 7.96–8.05 (m, 2H, H-2'); ¹³C NMR (75.4 MHz, CDCl₃) δ 17.3 (CH₃C=N), 55.1 (CH₃O), 104.9 (C-2), 108.7 (C-6), 111.6 (C-4), 127.1 (C-2'), 128.3 (C-3'), 129.7 (C-4'), 130.4 (C-5), 139.2 (C-1'), 153.0 (C-1), 160.2 (C-3), 165.5 (C=N); MS (70 eV) *m/z* 225 (M⁺, 43), 210 (100), 195 (11), 180 (10), 167 (12), 148 (18), 103 (21), 92 (22), 77 (50); HRMS (EI) *m/z* [M⁺] calcd for C₁₅H₁₅NO 225.1154, found 225.1139.

■ ASSOCIATED CONTENT

■ Supporting Information

Copies of the ¹H NMR, ¹³C NMR, and MS spectra of all compounds. Crystallographic information for **3x**, including X-ray diffraction data, atomic coordinates, thermal parameters, torsion angles, and complete bond distances and angles. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Fax: +5255-5729-6300/46211. E-mail: jtamariz@woodward.encb.ipn.mx, jtamarizm@gmail.com.

Present Address

†Unidad Profesional Interdisciplinaria de Ingeniería, Campus Zacatecas (UPIIZ), Instituto Politécnico Nacional, Blvd. del Bote S/N, Cerro del Gato, Ejido La Escondida, Col. Ciudad Administrativa, C.P. 98160, Zacatecas, Zac., Mexico.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Bruce A. Larsen for reviewing the English of the manuscript. J.T. acknowledges SIP/IPN (Grant Nos. 20090519, 20100236, 20110172, 20120830, and 20130686) and CONACYT (Grant Nos. 43508Q, 83446, and 178319) for financial support. R.U.G., H.C.C., R.B., and A.V.J. thank CONACYT for awarding them graduate scholarships and also thank SIP/IPN (PIFI) and Ludwig K. Hellweg Foundation for scholarship complements. J.L.V. thanks SNI/CONACYT for a scholarship awarded. F.D. and J.T. are fellows of the EDI-IPN and COFAA-IPN programs.

■ REFERENCES

- (1) Johnson, J. V.; Rauckman, B. S.; Baccanari, D. P.; Roth, B. *J. Med. Chem.* **1989**, *32*, 1942–1949.
- (2) (a) Fotie, J.; Kaiser, M.; Delfin, D. A.; Manley, J.; Reid, C. S.; Paris, J.-M.; Wenzler, T.; Maes, L.; Mahasenan, K. V.; Li, C.; Werbovetz, K. A. *J. Med. Chem.* **2010**, *53*, 966–982. (b) Reid, C. S.; Patrick, D. A.; He, S.; Fotie, J.; Premalatha, K.; Tidwell, R. R.; Wang, M. Z.; Liu, Q.; Gershkovich, P.; Wasan, K. M.; Wenzler, T.; Brun, R.; Werbovetz, K. A. *Bioorg. Med. Chem.* **2011**, *19*, 513–523.
- (3) (a) Kumar, S.; Engman, L.; Valgimigli, L.; Amorati, R.; Fumo, M. G.; Pedulli, G. F. *J. Org. Chem.* **2007**, *72*, 6046–6055 and references cited therein. (b) Lockhart, B.; Bonhomme, N.; Roger, A.; Dorey, G.; Casara, P.; Lestage, P. *Eur. J. Pharmacol.* **2001**, *416*, 59–68.

- (4) Takahashi, H.; Bekkali, Y.; Capolino, A. J.; Gilmore, T.; Goldrick, S. E.; Kaplita, P. V.; Liu, L.; Nelson, R. M.; Terenzio, D.; Wang, J.; Zuvella-Jelaska, L.; Proudfoot, J.; Nabozny, G.; Thomson, D. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 5091–5095.
- (5) Tsushima, K.; Hatakoshi, M.; Matsuo, N.; Ohno, N.; Nakayama, I. *Agric. Biol. Chem.* **1985**, *49*, 2421–2423.
- (6) Balayer, A.; Sévenet, T.; Schaller, H.; Hadi, A. H. A.; Chiaroni, A.; Riche, C.; País, M. *Nat. Prod. Lett.* **1993**, *2*, 61–67.
- (7) (a) Ambrus, D.; Szabolcsi, T.; Hutás, I. (Chinoin Gyogyszer es Vegyeszeti Termekek Gyara R.T.) Hung. Teljes HU 26100A2, 1984; *Chem. Abstr.* **1984**, *100*, 66972h. (b) Spark, A. A. *J. Am. Oil Chem. Soc.* **1982**, *59*, 185–188. (c) He, P.; Ackman, R. G. *J. Sci. Food Agric.* **2000**, *80*, 10–16. (d) Shah, A. G.; Pierson, J. A.; Pavlostathis, S. G. *Water Res.* **2005**, *39*, 4251–4263.
- (8) (a) Yu, L.; Qinyu, G.; Lianxin, L.; Guangxia, S. *J. Korean Chem. Soc.* **2011**, *55*, 830–834. (b) Nikulin, S. S.; Verezhnikov, V. N.; Poyarkova, T. N.; Shmyreva, Zh. V. *Russ. J. Appl. Chem.* **2003**, *76*, 1319–1322.
- (9) (a) Arseniyadis, S.; Wagner, A.; Mioskowski, C. *Tetrahedron Lett.* **2004**, *45*, 2251–2253. (b) Sathe, M.; Derveni, M.; Allen, M.; Cullen, D. C. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 1792–1795. (c) Bandyopadhyay, P.; Sathe, M.; Sharma, P.; Kaushik, M. P. *Tetrahedron Lett.* **2012**, *53*, 4631–4635.
- (10) (a) Yao, M.; Inoue, H.; Yoshioka, N. *Chem. Phys. Lett.* **2005**, *402*, 11–16. (b) Yao, M.; Shibuya, H.; Kato, T.; Inoue, H.; Yoshioka, N. *Polyhedron* **2005**, *24*, 2828–2834.
- (11) (a) Skraup, Z. H. *Monatsh. Chem.* **1881**, *2*, 139–170. (b) Levin, J.; Riehm, P. *Chem. Ber* **1886**, *19*, 1394–1395. (c) Doebner, O.; Miller, W. v. *Chem. Ber.* **1881**, *14*, 2812–2817.
- (12) For a recent review on MCR, see: Dömling, A.; Wang, W.; Wang, K. *Chem. Rev.* **2012**, *112*, 3083–3135.
- (13) Reddelien, G.; Thurm, A. *Chem. Ber.* **1932**, *65*, 1511–1521.
- (14) Craig, D. J. *Am. Chem. Soc.* **1938**, *60*, 1458–1465.
- (15) Vaughan, W. R. *Organic Synthesis*; Wiley: New York, 1955; Collect. Vol. III, pp 329–330.
- (16) Theoclitou, M. E.; Robinson, L. A. *Tetrahedron Lett.* **2002**, *43*, 3907–3910.
- (17) Fotie, J.; Wangun, H. V. K.; Fronczek, F. R.; Massawe, N.; Bhattarai, B. T.; Rhodus, J. L.; Singleton, T. A.; Bohle, D. S. *J. Org. Chem.* **2012**, *77*, 2784–2790.
- (18) (a) Luo, Y.; Li, Z.; Li, C.-J. *Org. Lett.* **2005**, *7*, 2675–2678. (b) Yi, C. S.; Yun, S. Y. *J. Am. Chem. Soc.* **2005**, *127*, 17000–17006. (c) Liu, X.-Y.; Ding, P.; Huang, J.-S.; Che, C.-M. *Org. Lett.* **2007**, *9*, 2645–2648.
- (19) Kothandaraman, P.; Foo, S. J.; Chan, P. W. H. *J. Org. Chem.* **2009**, *74*, 5947–5952.
- (20) (a) Waldmann, H.; Karunakar, G. V.; Kumar, K. *Org. Lett.* **2008**, *10*, 2159–2162. (b) Gurunathan, S.; Perumal, P. T. *Tetrahedron Lett.* **2011**, *52*, 1783–1787. (c) Williamson, N. M.; Ward, A. D. *Tetrahedron* **2005**, *61*, 155–165. (d) Cossy, J.; Poitevin, C.; Gomez Pardo, D.; Peglion, J.-L.; Dessinges, A. J. *Org. Chem.* **1998**, *63*, 4554–4557.
- (21) Hegedüs, A.; Hell, Z.; Vargadi, T.; Potor, A.; Gresits, I. *Catal. Lett.* **2007**, *117*, 99–101.
- (22) (a) Li, Y.; Wu, C.; Huang, J.; Su, W. *Synth. Commun.* **2006**, *36*, 3065–3073. (b) Zhang, J.-C.; Ji, J.-X. *ACS Catal.* **2011**, *1*, 1360–1363. (c) Ranu, B. C.; Hajra, A.; Dey, S. S.; Jana, U. *Tetrahedron* **2003**, *59*, 813–819. (d) Gültekin, Z.; Frey, W. *ARKIVOC* **2012**, *viii*, 250–261.
- (23) (a) Hu, X.-Y.; Zhang, J.-C.; Wei, W.; Ji, J.-J. *Tetrahedron Lett.* **2011**, *52*, 2903–2905. (b) Zhu, Y.-W.; Qian, J.-L.; Yi, W.-B.; Cai, C. *Tetrahedron Lett.* **2013**, *54*, 638–641.
- (24) Makino, K.; Hara, O.; Takiguchi, Y.; Katano, T.; Asakawa, Y.; Hatano, K.; Hamada, Y. *Tetrahedron Lett.* **2003**, *44*, 8925–8929.
- (25) Ye, K.-Y.; He, H.; Liu, W.-B.; Dai, L.-X.; Helmchen, G.; You, S.-L. *J. Am. Chem. Soc.* **2011**, *133*, 19006–19014.
- (26) (a) Cruz, M. C.; Tamariz, J. *Tetrahedron Lett.* **2004**, *45*, 2377–2380. (b) Cruz, M. C.; Tamariz, J. *Tetrahedron* **2005**, *61*, 10061–10072. (c) Correa, C.; Cruz, M. C.; Jiménez, F.; Zepeda, L. G.; Tamariz, J. *Aust. J. Chem.* **2008**, *61*, 991–999. (d) Cruz, M. C.; Jiménez, F.; Delgado, F.; Tamariz, J. *Synlett* **2006**, 749–755.
- (e) Jerezano, A.; Jiménez, F.; Cruz, M. C.; Montiel, L. E.; Delgado, F.; Tamariz, J. *Helv. Chim. Acta* **2011**, *94*, 185–198. (f) González-Romero, C.; Bernal, P.; Jiménez, F.; Cruz, M. C.; Fuentes-Benites, A.; Benavides, A.; Bautista, R.; Tamariz, J. *Pure Appl. Chem.* **2007**, *79*, 181–191.
- (27) Yang, S.-H. *Synlett* **2009**, 1351–1352 and references cited therein.
- (28) (a) Fujisawa, H.; Sasaki, Y.; Mukaiyama, T. *Chem. Lett.* **2001**, 190–191. (b) Bouzide, A. *Org. Lett.* **2002**, *4*, 1347–1350. (c) Chakraborti, A. K.; Bhagat, S.; Rudrawar, S. *Tetrahedron Lett.* **2004**, *45*, 7641–7644. (d) Abaee, M. S.; Sharifi, R.; Mojtahedi, M. M. *Org. Lett.* **2005**, *7*, 5893–5895. (e) Karikomi, M.; Takayama, T.; Haga, K.; Hiratani, K. *Tetrahedron Lett.* **2005**, *46*, 6541–6544. (f) Salehi, H.; Guo, Q.-X. *Synth. Commun.* **2004**, *34*, 171–179. (g) Evans, D. A.; Downey, C. W.; Shaw, J. T.; Tedrow, J. S. *Org. Lett.* **2002**, *4*, 1127–1130. (h) Huang, J.-W.; Chen, C.-D.; Leung, M.-k. *Tetrahedron Lett.* **1999**, *40*, 8647–8650.
- (29) Although alkyl pyruvates have proven to be highly efficient and regioselective substrates for an analogous Brønsted-catalyzed multistep procedure to afford 1,2-DHQ^{22b,23} no meta-substituted anilines with strong electron-donating groups such as **1b** were evaluated.
- (30) (a) Kohnstam, G.; Williams, D. L. H. Directive and Activating Effects of Alkoxy Groups in Aromatic and Aliphatic Reactions. In *The Chemistry of the Ether Linkage*; Patai, S., Ed.; Wiley: New York, 1967; pp 132–150. (b) Kruse, L. I.; Cha, J. K. *J. Chem. Soc., Chem. Commun.* **1982**, 1333–1336.
- (31) Tapia, I.; Alcazar, V.; Grande, M.; Moran, J. R. *Tetrahedron* **1988**, *44*, 5113–5116.
- (32) (a) Denmark, S. E.; Venkatraman, S. *J. Org. Chem.* **2006**, *71*, 1668–1676. (b) Eisch, J. J.; Dluzniewski, T. *J. Org. Chem.* **1989**, *54*, 1269–1274. (c) Walter, H.; Sauter, H.; Winkler, T. *Helv. Chim. Acta* **1992**, *75*, 1274–1280. (d) Walter, H.; Schneider, J. *Heterocycles* **1995**, *41*, 1251–1269.
- (33) Chakraborti, A. K.; Sharma, L.; Gulhane, R.; Shivani. *Tetrahedron* **2003**, *59*, 7661–7668. Theoretical evidence has also shown that, independently of its structure, the Lewis acid plays an important role in activating the aromatic substrate, and as a result, promotes a lower activation energy of the interaction between the electrophile and the aromatic substrate: Tarakeshwar, P.; Lee, J. Y.; Kim, K. S. *J. Phys. Chem. A* **1998**, *102*, 2253–2255.
- (34) We found that this reaction was successful and efficiently catalyzed with mercury(I) chloride under solvent-free conditions, as opposed to the reported use of mercury(II) chloride in THF: Barluenga, J.; Aznar, F.; Liz, R.; Rodes, R. *J. Chem. Soc., Perkin Trans. 1* **1980**, 2732–2737.
- (35) Xiao, F.; Chen, W.; Liao, Y.; Deng, G.-J. *Org. Biomol. Chem.* **2012**, *10*, 8593–8596.
- (36) For the intracomplex reaction, as an alternative mechanism in electrophilic aromatic substitution, see: (a) Aschi, M.; Attinà, M.; Cacace, F. *J. Am. Chem. Soc.* **1995**, *117*, 12832–12839. (b) Tanaka, M.; Fujiwara, M.; Xu, Q.; Souma, Y.; Ando, H.; Laali, K. K. *J. Am. Chem. Soc.* **1997**, *119*, 5100–5105.
- (37) Edwards, J. P.; Ringgenberg, J. D.; Jones, T. K. *Tetrahedron Lett.* **1998**, *39*, 5139–5142.
- (38) (a) Forrest, T. P.; Dauphinee, G. A.; Miles, W. F. *Can. J. Chem.* **1974**, *52*, 884–887. (b) Forrest, T. P.; Dauphinee, G. A.; Deraniyagala, S. A. *Can. J. Chem.* **1985**, *63*, 412–417.
- (39) Kinetic experiments have shown that a conformational factor is involved in the preferential formation of the *para* versus *ortho* regioisomer, which explains the larger deprotonation rate in the *para* versus *ortho* σ -complex intermediate: Effenberger, F.; Maier, A. H. *J. Am. Chem. Soc.* **2001**, *123*, 3429–3433.
- (40) (a) Smith, M. B.; March, J. *March's Advanced Organic Chemistry, Reactions, Mechanism, and Structure*; Wiley-Interscience: Hoboken, 2007; pp 665–684. (b) Isaacs, N. S. *Physical Organic Chemistry*; Longman Scientific & Technical; Burnt Mill, 1987; pp 428–453.
- (41) (a) Carey, F. A.; Sundberg, R. J. *Advanced Organic Chemistry*; Kluwer Academic/Plenum Publishers: New York, 2000; Part A, pp 551–589. (b) Norman, R. O. C.; Taylor, R. *Electrophilic Substitution in*

- Benzenoid Compounds*; Elsevier: Amsterdam, 1965; pp 25–60.
- (c) Heaney, H. The Bimolecular Aromatic Friedel-Crafts Reaction. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Heathcock, C. H., Eds.; Pergamon Press: Oxford, 1991; pp 733–752.
- (42) For typical examples of *para* regioselectivity in a Friedel–Crafts reaction of substituted benzene rings with electron-donating groups, see: (a) Hashimoto, I.; Kawaji, T.; Badea, F. D.; Sawada, T.; Mataka, S.; Tashiro, M.; Fukata, G. *Res. Chem. Intermed.* **1996**, *22*, 855–869. (b) Ranu, B. C.; Ghosh, K.; Jana, U. *J. Org. Chem.* **1996**, *61*, 9546–9547.
- (43) Fleming, I. *Molecular Orbitals and Organic Chemical Reactions*, reference ed.; Wiley: Chichester, 2010; pp 167–177.
- (44) (a) Dewar, M. J. S. *J. Chem. Soc.* **1949**, 463–469. (b) Anderson, G. M., III; Kollman, P. A.; Domelsmith, L. N.; Houk, K. N. *J. Am. Chem. Soc.* **1979**, *101*, 2344–2352. (c) Kato, S. *Theor. Chem. Acc.* **2000**, *103*, 219–220. (d) Pankratov, A. N. *THEOCHEM* **2000**, 507, 239–244. (e) Hirao, H.; Ohwada, T. *J. Phys. Chem. A* **2003**, *107*, 2875–2881.
- (45) Galavov, B.; Koleva, G.; Schaefer, H. F., III; Schleyer, P. v. R. *J. Org. Chem.* **2010**, *75*, 2813–2819.
- (46) Geerlings, P.; De Proft, F.; Langenaeker, W. *Chem. Rev.* **2003**, *103*, 1793–1873 and references cited therein.
- (47) Pérez, P.; Domingo, L. R.; Duque-Noreña, M.; Chamorro, E. *THEOCHEM* **2009**, 895, 86–91.
- (48) Pearson, D. E.; Buehler, C. A. *Synthesis* **1971**, 455–477.
- (49) Tanaka, K.; Toda, F. *Chem. Rev.* **2000**, *100*, 1025–1074.
- (50) For general information, see the Supporting Information.
- (51) Rosowsky, A.; Modest, E. J. *J. Org. Chem.* **1965**, *30*, 1832–1837.
- (52) Walter, H.; Rigassi, T. Eur. Pat. Appl. EP 463996, 1992; *Chem. Abstr.* **1992**, *116*, 176101m.
- (53) Mammen, H.; Dietzel, S.; Martin, H.-D. *Z. Chem.* **1985**, *25*, 435–436. *Chem. Abstr.* **1987**, *106*, 49997b.
- (54) Arduini, A.; Bigi, F.; Casiraghi, G.; Casnati, G.; Sartori, G. *Synthesis* **1981**, 975–977.
- (55) Kamakshi, R.; Reddy, B. S. R. *Catal. Commun.* **2007**, *8*, 825–828.