

Microwave-Assisted or Cu–NHC-Catalyzed Cycloaddition of Azido-Disubstituted Alkynes: Bifurcation of Reaction Pathways

Yuyu Xia,[†] Ling-yan Chen,[†] Shang Lv,[†] Zhihua Sun,^{*,†} and Bing Wang^{*,†,§}


[†]College of Chemistry and Chemical Engineering, Shanghai University of Engineering Science, 333 Longteng Road, Shanghai 201620, China

[‡]Department of Chemistry, Fudan University, 220 Handan Road, Shanghai 200433, China

[§]Institutes of Biomedical Sciences, Fudan University, 138 Yixueyuan Road, Shanghai 200032, China

S Supporting Information

ABSTRACT: Microwave irradiation promoted the intramolecular cycloaddition of 2-azidoacetamides derived from α -chiral propargylic amines, affording 1,4,5-trisubstituted triazoles **4** bearing a chiral aminomethyl side chain at C5. In contrast, for the same substrates **3a–k**, Cu(I)–NHC complexes catalyzed the intermolecular cycloaddition in an unexpected desilylative fashion, leading to 1,4-disubstituted triazoles **5**. This demonstrates that 1-silyl alkynes can be employed as substrates for CuAAC with a suitable coupling partner.




1,2,3-Triazole is a well-known pharmacophore possessing a wide spectrum of interesting biological activities.¹ The thermal 1,3-dipolar cycloaddition of azides to alkynes to form 1,2,3-triazoles (Huisgen reaction) was first documented in 1893 and studied in greater detail in the late 1950s and the 1960s.² However, this reaction in general has low regioselectivities for even terminal alkynes. The copper-catalyzed variant of this transformation (CuAAC), first reported in 2002,³ marked the emergence of “click chemistry”⁴ and has since found widespread applications in medicinal chemistry, materials science, and chemical biology.⁵ A complementary ruthenium-catalyzed approach (RuAAC) exhibits the opposite regioselectivity.⁶ With the recent advances in N-heterocyclic carbene (NHC) chemistry,⁷ a few Cu–NHC complexes have been reported to catalyze azide cycloaddition to terminal alkynes.⁸ While the CuAAC of terminal alkynes enjoys excellent regioselectivity for the 1,4-disubstituted adducts, unsymmetric internal alkynes remain difficult substrates because of low reactivity and problematic regioselectivity control. In this regard, several groups have recently reported Huisgen reactions of well-designed internal alkynes with unambiguous regioselectivity by taking advantage of intramolecular reaction.⁹ Additionally, Schubert and co-workers reported that 1-trimethylsilyl-substituted alkynes are excellent substrates favoring the formal 1,5-regioselective intermolecular AAC under thermal conditions.¹⁰ However, the scope of CuAAC of unsymmetrically disubstituted alkynes is still limited to 1-halo- and 1-metallo (Al, Bi)-substituted alkynes.¹¹ As part our ongoing interest in novel Cu–NHC complexes for organic synthesis,¹² we designed a series of disubstituted alkynes with a pendant

azide function and explored their reactivity under both microwave dielectric heating and Cu–NHC catalysis. This enabled us to compare the scope and limitations of the two sets of conditions for unsymmetrically disubstituted alkynes and to determine whether the 1,5-regioselectivity of 1-silylated alkynes also holds for CuAAC.

The substrates were based on synthetically versatile α -chiral propargylic amines¹³ because of their ready availability, high enantiopurity, and structural diversity. The preparation was straightforward. After removal of the *N*-sulfinyl group, amidation with azidoacetic acid¹⁴ produced the desired azido alkynes in good yields (Scheme 1). The γ - and α -groups of the propargylic amines constitute the C4 and C5 substituents of the triazole.

Substrates **3** were first subjected to Cu-free microwave-assisted cyclization at 160 °C (power = 200 W, maximum internal pressure = 200 psi). The Huisgen cycloaddition proceeded smoothly to form 1,4,5-trisubstituted triazoles **4** in very good to excellent isolated yields (>85% in most cases;

Scheme 1. Synthesis of Azido Internal Alkynes




Received: June 11, 2014

Published: September 30, 2014

Table 1). Conventional heating at the same temperature proved ineffective, and the starting material was recovered unchanged.

Table 1. Intramolecular Cycloaddition of Azido Alkynes **3 under Microwave Irradiation**




entry	R'	R	product	yield (%) ^a
1	2-propyl	TMS	4a	95
2 ^b	2-propyl	TMS	4a	NR
3	cyclohexyl	TMS	4b	97
4	cyclopropyl	TMS	4c	96
5	1-adamantyl	TMS	4d	85
6	benzyl	TMS	4e	90
7	(<i>E</i>)-PhCH=CH	TMS	4f	87
8	4-Me-phenyl	TMS	4g	93
9	4-MeO-phenyl	TMS	4h	94
10	pyridin-3-yl	TMS	4i	75 ^c
11	furan-2-yl	TMS	4j	88
12 ^d	<i>N</i> -Boc-indol-3-yl	TMS	4k	85
13 ^e	4-MeO-phenyl	phenyl	4l	80
14 ^e	4-MeO-phenyl	cyclopropyl	4m	75

^aIsolated yields. ^bUpon oil bath heating to $160\text{ }^\circ\text{C}$ for 4 h, **3a** was recovered (>90%). ^cYield after desilylation. ^dThe indole *N*-Boc protection was simultaneously removed because of its lability. ^eThe reaction time was 3 h.

Very bulky 1-adamantyl or heterocyclic R' groups were all tolerated, except in the case of **3i**, where the silyl group was partially cleaved because of its decreased stability in the presence of the basic pyridine ring (entry 10). It is notable that substrates **3a–k** required only 1 h of microwave irradiation, while the reaction time for the azido alkynes used in Taddei's study was three times longer.^{9a} This suggests that placement of a substituent at the α -carbon of the propargylic amine rather than at the α -position of the carbonyl resulted in a more favorable conformation placing the azido group and the alkyne in proximity of each other to facilitate the cyclization. Replacing the silyl group with either an aromatic group (phenyl) or an alkyl group (cyclopropyl) still afforded satisfactory yields, although a longer reaction time was necessary (entries 13 and 14). Products **4** have the potential for further modifications such as hydrolytic ring opening, reduction of the lactam, and stereoselective α -alkylation. Moreover, the TMS group on the triazole ring can be easily converted to halides and used for subsequent cross-coupling.¹⁵

Then the reaction of azido alkynes **3** using Cu–NHC catalysis was examined at room temperature (Table 2). Different types of NHC precursors, including those widely represented in the literature (**6–8**; Figure 1) and the benzoimidazole-based compounds recently reported by our group (**9–11**), were screened using **3g** as a model substrate. It was found that the complex formed by the NHC derived from **9** exhibited the highest catalytic activity in 50% aqueous acetonitrile (Table 2, entry 5), while other Cu–NHC complexes were less effective. In anhydrous solvents such as MeCN, THF, *tert*-butanol, and DMF, no product was formed.

Table 2. Screening of Cu(I)–NHC Catalysts



entry	NHC precursor	yield (%) ^a
1	none	11
2	6	40
3	7	45
4	8	38
5	9	81
6	10	30
7	11	10
8 ^b	9	NR
9 ^c	9	66

^aIsolated yields. ^bIn MeCN. ^cIn 1:1 *t*-BuOH/ H_2O .





Figure 1. NHC precursors screened for the cycloaddition reaction.

Other mixed solvent systems such as 1:1 *t*-BuOH/ H_2O afforded lower yields (entry 9). Under the optimal conditions, a single isolable product was obtained in a satisfactory yield of 81% after 24 h at rt. NMR and MS analysis revealed that the isolated product **5g** was formed via an intermolecular reaction with formal 1,4-regioselectivity (as confirmed by HMBC experiments), opposite to that of the thermal AAC of silyl alkynes.¹⁰ Furthermore, it is interesting that the TMS group supposed to be on the heterocycle was absent, while the terminal TMS of the unreacted alkyne was intact. The spontaneous loss of TMS during an AAC reaction, without the assistance of desilylation reagents, has not been reported to date. In all cases with Cu–NHC catalysis, no intramolecular reaction product was detected by either TLC or LC–MS. It is also notable that further reaction of **5g**, which possesses both azide and alkyne functions, to form oligomers or polymers was minimal without resorting to high-dilution conditions. Unfortunately, the accelerating effect of carboxylic acids¹⁶ in CuAAC cannot be exploited in the present protocol, as HOAc could decompose the Cu–NHC catalyst. The use of a Lewis acid (ZnCl_2) alone or with an NHC was also unproductive.

This protocol was then extended to other azido alkynes **3**, and the results are summarized in Table 3. A series of 1,4-disubstituted 1,2,3-triazoles **5** carrying alkyne and azide functions on the side chains were obtained (Table 3, entries

Table 3. Intermolecular CuAAC of 1-Silylated Alkynes **3**

entry	R'	R	conversion (%) ^a	product	yield (%) ^b
1	2-propyl	TMS	73	5a	69
2	cyclohexyl	TMS	70	5b	65
3	benzyl	TMS	70	5e	68
4	(<i>E</i>)-PhCH=CH	TMS	75	5f	72
5	4-Me-phenyl	TMS	85	5g	81
6	4-MeO-phenyl	TMS	84	5h	79
7	furan-2-yl	TMS	78	5j	75
8	4-MeO-phenyl	phenyl	NR	–	–
9	4-MeO-phenyl	cyclopropyl	NR	–	–
10	4-Me-phenyl	H	100	polymer	–

^aBased on recovered **3**. ^bIsolated yields.

1–7). Although the reaction did not go to completion in 24 h (65–85% conversion), the yields based on reacted starting material were excellent (>90%). In all cases, the triazole ring did not bear a TMS group. Unfortunately, when the terminal silyl group was replaced by a carbogenic group, the cycloaddition did not proceed, and the starting material was recovered (entries 8 and 9). Interestingly, for an analogous terminal alkyne substrate (desilylated **3g**), polymer formation was observed immediately (entry 10).¹⁷ Therefore, compound **5** represents a class of triazole-based bifunctional molecules that is not readily accessible via the conventional AAC of terminal alkynes.

In order to gain some insight into the mechanism for the CuAAC of 1-silyl alkynes, several control experiments were carried out (Scheme 2). When 1 equiv of benzyl azide was reacted with **3g**, only 5% of the product was 1-benzyl triazole **12**, while 95% was the normal product **5g** (eq 1). In view of the fact that the intrinsic reactivities of azidoacetamide and benzyl azide toward a terminal alkyne under Cu–NHC catalysis are virtually the same (eq 2), this and the polymerization of terminal alkyne (Table 3, entry 10) essentially ruled out free terminal alkyne (formed by either Cu-catalyzed or simple base-induced hydrolysis) as the key intermediate as well as the presence of a desilylation step prior to CuAAC. Another possible mechanism is that the amide carbonyl group may participate in the reaction to facilitate removal of the TMS group for the formation of copper acetylide species without involving a free terminal alkyne. In order to test this possibility, reactions 3 and 4 were conducted. The results showed that the reaction between benzyl azide and 1-(trimethylsilyl)-phenylacetylene was very sluggish under our CuAAC conditions. However, the reaction of *N*-benzyl-2-azidoacetamide with the same alkyne afforded the triazole in a good yield (70%) in 24 h. Moreover, when the very bulky and hydrolytically stable 1-triisopropylsilyl group was employed to cap the alkyne, a moderate yield of 58% could still be achieved. This provides strong evidence that the adjacent functional group of the azide plays an important role in promoting the cleavage of the terminal silyl group regardless of its size. To further illustrate this possibility, eq 5 showed that when the azide and the carbonyl group were separated with more methylene units, the reaction was significantly less efficient than

using 2-azidoacetamide. This could occur because the corresponding transition states became slack and less favorable. Therefore, we speculate that a cyclic transition state was involved as a result of the dative interaction of the amide carbonyl with the silyl group (eq 6). Without this neighboring group, the formation of copper acetylide species could rely only on the assistance of water, and this process is apparently not efficient.

Compounds **4** were converted to unnatural amino acids **13** in very good yields upon microwave-assisted acid hydrolysis (Scheme 3). These products may find application in the study of conformationally constrained peptides and medicinal chemistry.

In summary, the [3 + 2] cycloaddition of a series of azidoacetamides derived from α -chiral propargylic amines was investigated. Microwave irradiation promoted intramolecular cycloaddition to afford 1,4,5-trisubstituted triazoles bearing a chiral aminomethyl side chain at C5, while Cu(I)–NHC complexes catalyzed intermolecular desilylative AAC of azido 1-silyl alkynes at rt, leading to 1,4-disubstituted triazoles **5** without the heterocyclic TMS group. Intramolecular reaction products were not detected under the present CuAAC protocol, while analogous terminal alkynes underwent facile polymerization. The controllable reaction pathways offer good opportunities for building molecular diversity from readily available chiral propargylic amines. The use of 1-silyl alkynes as substrates for CuAAC will also enhance the application of this reaction in organic synthesis.

EXPERIMENTAL SECTION

General Methods. All commercially available reagents were used without purification unless otherwise noted. Acetonitrile was refluxed with sodium under argon before use. Column chromatography was performed using silica gel (300–400 mesh). Visualization of the compounds was accomplished with UV light (254 nm) and iodine. ¹H and ¹³C NMR spectra were recorded in CDCl₃ operating at 400 and 100 MHz, respectively. Proton chemical shifts are reported relative to the residual proton signals of the deuterated solvent [CDCl₃ (7.26 ppm) or DMSO-*d*₆ (2.50 and 3.33 ppm)] or TMS. Carbon chemical shifts were internally referenced to the deuterated solvent signals in CDCl₃ (77.00 ppm) or DMSO-*d*₆ (40.0 ppm). Data are represented as follows: chemical shift (multiplicity, coupling constant in hertz, integration). Multiplicities are denoted as follows: bs = broad singlet, s

(*S*)-2-Azido-*N*-(1-cyclohexyl-3-(trimethylsilyl)prop-2-yn-1-yl)acetamide (**3b**). A colorless oil (800 mg, 71%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.54 (d, *J* = 8.4 Hz, 1H), 4.49–4.45 (m, 1H), 3.88–3.79 (m, 2H), 1.83–1.6 (m, 5H), 1.46–1.42 (m, 1H), 1.26–0.96 (m, 5H), 0.15 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.1, 105.3, 87.7, 50.9, 46.6, 42.2, 29.2, 28.8, 26.3, 25.8, 25.7, 0.4. HRMS (ESI-TOF) calcd for C₁₄H₂₄N₄NaOSi [M + Na]⁺ 315.1612, found 315.1607.

(*S*)-2-Azido-*N*-(1-cyclopropyl-3-(trimethylsilyl)prop-2-yn-1-yl)acetamide (**3c**). A colorless oil (660 mg, 69%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.70 (d, *J* = 8.4 Hz, 1H), 4.50–4.46 (m, 1H), 3.88–3.78 (m, 2H), 1.15–1.06 (m, 1H), 0.51–0.29 (m, 4H), 0.14 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.0, 104.4, 86.9, 50.9, 44.3, 15.5, 3.3, 2.6, 0.4. HRMS (ESI-TOF) calcd for C₁₁H₁₈N₄NaOSi [M + Na]⁺ 273.1142, found 273.1130.

N-(*S*)-1-(3*S*,5*S*,7*S*)-Adamantan-1-yl-3-(trimethylsilyl)prop-2-yn-1-yl-2-azidoacetamide (**3d**). A colorless oil (800 mg, 61%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.41 (d, *J* = 9.2 Hz, 1H), 4.35 (d, *J* = 9.6 Hz, 1H), 3.86 (s, 2H), 1.97 (s, 3H), 1.68–1.50 (m, 12H), 0.16 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.5, 104.4, 88.4, 51.1, 50.9, 38.3, 37.1, 36.8, 28.1, 0.1. HRMS (ESI-TOF) calcd for C₁₈H₂₈N₄NaOSi [M + Na]⁺ 367.1925, found 367.1929.

(*S*)-2-Azido-*N*-(1-phenyl-4-(trimethylsilyl)but-3-yn-2-yl)acetamide (**3e**). A yellow oil (800 mg, 69%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.67 (d, *J* = 8.4 Hz, 1H), 7.31–7.23 (m, 5H), 4.78–4.73 (m, 1H), 3.85–3.72 (m, 2H), 2.90–2.88 (m, 2H), 0.1 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.0, 137.4, 130.0, 128.5, 127.1, 105.9, 88.0, 51.0, 43.2, 41.4, 0.2. HRMS (ESI-TOF) calcd for C₁₅H₂₀N₄NaOSi [M + Na]⁺ 323.1299, found 323.1284.

(*S*,*E*)-2-Azido-*N*-(1-phenyl-5-(trimethylsilyl)pent-1-en-4-yn-3-yl)acetamide (**3f**). An orange-yellow solid (800 mg, 67%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.91 (d, *J* = 8.0 Hz, 1H), 7.47–7.27 (m, 5H), 6.73–6.69 (m, 1H), 6.27–6.22 (m, 1H), 5.45–5.41 (m, 1H), 3.94–3.84 (m, 2H), 0.19 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.0, 136.1, 131.5, 129.2, 128.6, 127.1, 126.6, 104.0, 88.9, 50.9, 43.0, 0.3. HRMS (ESI-TOF) calcd for C₁₆H₂₀N₄NaOSi [M + Na]⁺ 335.1299, found 335.1294.

(*S*)-2-Azido-*N*-(1-(*p*-tolyl)-3-(trimethylsilyl)prop-2-yn-1-yl)acetamide (**3g**). A light-yellow oil (850 mg, 74%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.09 (d, *J* = 8.4 Hz, 1H), 7.31 (d, *J* = 8.0 Hz, 2H), 7.20 (d, *J* = 8.0 Hz, 2H), 5.84 (d, *J* = 8.4 Hz, 1H), 3.92–3.80 (m, 2H), 2.30 (s, 3H), 0.17 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.0, 137.6, 136.4, 129.6, 127.2, 105.0, 88.6, 50.8, 44.5, 21.1, 0.3. HRMS (ESI-TOF) calcd for C₁₅H₂₀N₄NaOSi [M + Na]⁺ 323.1299, found 323.1292.

(*S*)-2-Azido-*N*-(1-(4-methoxyphenyl)-3-(trimethylsilyl)prop-2-yn-1-yl)acetamide (**3h**). A light-yellow oil (800 mg, 69%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.07 (d, *J* = 8.0 Hz, 1H), 7.35 (d, *J* = 8.8 Hz, 2H), 6.95 (d, *J* = 8.4 Hz, 2H), 5.82 (d, *J* = 8.4 Hz, 1H), 3.91–3.75 (m, 2H), 3.35 (s, 3H), 0.17 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 166.9, 159.4, 131.3, 128.6, 114.4, 105.1, 88.5, 55.6, 50.8, 44.2, 0.3. HRMS (ESI-TOF) calcd for C₁₅H₂₀N₄NaO₂Si [M + Na]⁺ 339.1248, found 339.1237.

(*S*)-2-Azido-*N*-(1-(pyridin-3-yl)-3-(trimethylsilyl)prop-2-yn-1-yl)acetamide (**3i**). A white solid (600 mg, 54%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.25 (d, *J* = 8.0 Hz, 1H), 8.65 (d, *J* = 0.2 Hz, 1H), 8.55–8.53 (m, 1H), 7.82 (d, *J* = 8.0 Hz, 1H), 7.46–7.43 (m, 1H), 5.97 (d, *J* = 8.4 Hz, 1H), 3.96–3.85 (m, 2H), 0.18 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.2, 149.5, 148.7, 135.1, 135.0, 124.1, 103.8, 50.9, 42.9, 31.1, 0.2. HRMS (ESI-TOF) calcd for C₁₀H₉N₅NaO [M + Na]⁺ 238.0699, found 238.0686.

(*R*)-2-Azido-*N*-(1-(furan-2-yl)-3-(trimethylsilyl)prop-2-yn-1-yl)acetamide (**3j**). A yellow oil (700 mg, 66%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.18 (d, *J* = 8.4 Hz, 1H), 7.67 (s, 1H), 6.45–6.40 (m, 2H), 5.94 (d, *J* = 8.0 Hz, 1H), 3.92–3.82 (m, 2H), 0.17 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.1, 150.9, 143.8, 111.0, 108.1, 102.2, 88.1, 50.7, 39.0, 0.2. HRMS (ESI-TOF) calcd for C₁₂H₁₆N₄NaO₂Si [M + Na]⁺ 299.0935, found 299.0928.

(*S*)-*tert*-Butyl 3-(1-(2-Azidoacetamido)-3-(trimethylsilyl)prop-2-yn-1-yl)-1*H*-indole-1-carboxylate (**3k**). A yellow oil (1.036 g, 60%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.11 (d, *J* = 8.4 Hz, 1H), 8.10 (d, *J* = 8.4 Hz, 1H), 7.69 (s, 1H), 7.60 (d, *J* = 8.0 Hz, 1H), 7.40–7.36 (m,

1H), 7.30–7.26 (m, 1H), 6.10 (d, *J* = 8.0 Hz, 1H), 3.94–3.81 (m, 2H), 1.63 (s, 9H), 0.20 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.2, 149.2, 135.7, 128.0, 125.4, 124.6, 123.3, 120.1, 119.1, 115.4, 103.9, 88.2, 84.6, 50.9, 37.6, 28.1, 0.2. HRMS (ESI-TOF) calcd for C₂₁H₂₇N₅NaO₃Si [M + Na]⁺ 448.1775, found 448.1766.

(*S*)-2-Azido-*N*-(1-(4-methoxyphenyl)-3-phenylprop-2-yn-1-yl)acetamide (**3l**). A white solid (860 mg, 70%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.16 (d, *J* = 8.4 Hz, 1H), 7.49–7.40 (m, 7H), 6.97 (d, *J* = 8.8 Hz, 2H), 6.03 (d, *J* = 8.4 Hz, 1H), 3.94–3.84 (m, 2H), 3.76 (s, 3H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 167.0, 159.4, 131.9, 131.6, 129.3, 129.2, 128.7, 122.4, 114.5, 88.8, 84.0, 55.7, 51.0, 44.2. HRMS (ESI-TOF) calcd for C₁₈H₁₆N₄NaO₂ [M + Na]⁺ 343.1165, found 343.1154.

(*S*)-2-Azido-*N*-(3-cyclopropyl-1-(4-methoxyphenyl)prop-2-yn-1-yl)acetamide (**3m**). A white solid (760 mg, 70%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.95 (d, *J* = 8.4 Hz, 1H), 7.32 (d, *J* = 8.8 Hz, 2H), 6.92 (d, *J* = 8.8 Hz, 2H), 5.72–5.69 (m, 1H), 3.88–3.77 (m, 2H), 3.74 (s, 3H), 1.40–1.34 (m, 1H), 0.82–0.77 (m, 2H), 0.63–0.59 (m, 2H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 166.7, 159.3, 132.2, 128.5, 114.3, 87.8, 74.5, 55.6, 50.9, 43.8, 8.3. HRMS (ESI-TOF) calcd for C₁₅H₁₆N₄NaO₂ [M + Na]⁺ 307.1165, found 307.1167.

General Procedure for Cu-Free Microwave-Assisted AAC: (*S*)-4-Isopropyl-3-(trimethylsilyl)-4,5-dihydro[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4a**). Compound **3a** (126 mg, 0.5 mmol) was dissolved in acetonitrile (20 mL) and water (5 mL). This solution was heated by microwave irradiation to 160 °C and maintained at this temperature for 1 h. The solvent was removed under vacuum. This gave sufficiently pure **4a** as a white solid (120 mg, 95%). Mp 176–177 °C. [α]_D²⁵ = +68.3 (c 0.23, MeOH). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.81 (d, *J* = 3.2 Hz, 1H), 5.15–5.00 (m, 2H), 4.59–4.58 (m, 1H), 1.99–1.95 (m, 1H), 1.01 (d, *J* = 7.2 Hz, 3H), 0.62 (d, *J* = 6.8 Hz, 3H), 0.30 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 165.3, 139.6, 137.9, 53.5, 48.7, 37.0, 19.8, 15.8, –0.4. MS (ESI) *m/z* 253.1 [M + H]⁺. HRMS (ESI-TOF) calcd for C₁₁H₂₁N₄O₂Si [M + H]⁺ 253.1479, found 253.1470.

(*S*)-4-Cyclohexyl-3-(trimethylsilyl)-4,5-dihydro[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4b**). A white solid (142 mg, 97%). Mp 213–214 °C. [α]_D²⁵ = +61.8 (c 0.17, MeOH). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.81 (d, *J* = 3.2 Hz, 1H), 5.12–4.99 (m, 2H), 4.54 (s, 1H), 1.77–1.59 (m, 5H), 1.32–0.92 (m, 6H), 0.31 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 165.2, 139.6, 137.6, 53.2, 48.8, 46.9, 29.9, 26.3, 26.1, 26.0, 25.8, –0.4. MS (ESI) *m/z* 293.2 [M + H]⁺. HRMS (ESI-TOF) calcd for C₁₄H₂₅N₄O₂Si [M + H]⁺ 293.1792, found 293.1785.

(*S*)-4-Cyclopropyl-3-(trimethylsilyl)-4,5-dihydro[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4c**). A white solid (120 mg, 96%). Mp 151–152 °C. [α]_D²⁵ = +50.2 (c 0.20, MeOH). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.79 (d, *J* = 3.6 Hz, 1H), 5.23–5.00 (m, 2H), 4.49–4.47 (m, 1H), 1.26–1.19 (m, 1H), 0.51–0.49 (m, 4H), 0.46 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 165.2, 139.7, 137.6, 50.0, 48.8, 19.0, 2.8, 2.3, –0.3. MS (ESI) *m/z* 251.1 [M + H]⁺. HRMS (ESI-TOF) calcd for C₁₁H₁₉N₄O₂Si [M + H]⁺ 251.1323, found 251.1318.

(4*S*)-4-(1*R*,3*R*,5*S*)-Adamantan-1-yl-3-(trimethylsilyl)-4,5-dihydro[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4d**). This was further purified by silica gel chromatography (eluent: 5:1 hexane/acetone) to give a white solid (152 mg, 85%). Mp 297–298 °C. [α]_D²⁵ = +78.4 (c 0.18, MeOH). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.85 (d, *J* = 4.0 Hz, 1H), 5.13–4.96 (m, 2H), 4.07 (d, *J* = 4.4 Hz, 1H), 1.96 (s, 3H), 1.65–1.44 (m, 12H), 0.33 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 165.8, 141.4, 135.2, 57.7, 49.9, 38.4, 36.5, 28.0, 0.1. MS (ESI) *m/z* 345.2 [M + H]⁺. HRMS (ESI-TOF) calcd for C₁₈H₂₉N₄O₂Si [M + H]⁺ 345.2105, found 345.2106.

(*S*)-4-Benzyl-3-(trimethylsilyl)-4,5-dihydro[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4e**). This was further purified by silica gel chromatography (eluent: 5:1 hexane/acetone) to give a light-yellow solid (135 mg, 90%). Mp 206–207 °C. [α]_D²⁵ = +169.1 (c 0.15, MeOH). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.75 (d, *J* = 3.6 Hz, 1H), 7.28–7.19 (m, 3H), 6.74–6.72 (m, 2H), 5.14–5.11 (m, 1H), 4.70 (d, *J* = 17.2 Hz, 1H), 3.29 (d, *J* = 17.6 Hz, 1H), 3.15–3.10 (m, 1H), 2.99–2.95 (m, 1H), 0.37 (s, 9H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ 164.8, 139.7, 137.1, 134.9, 130.5, 128.7, 127.6, 49.0, 48.0, 43.8, –0.4. MS

(ESI) m/z 301.1 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{15}H_{21}N_4OSi$ $[M + H]^+$ 301.1479, found 301.1483.

(*S,E*)-4-Styryl-3-(trimethylsilyl)-4,5-dihydro-[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4f**). This was further purified by silica gel chromatography (eluent: 5:1 hexane/acetone) to give an orange-yellow solid (136 mg, 87%). Mp 197–198 °C. $[\alpha]_D^{25} = +94.3$ (c 0.21, MeOH). 1H NMR (400 MHz, DMSO- d_6) δ 8.96 (d, $J = 3.2$ Hz, 1H), 7.46–7.29 (m, 5H), 6.65 (d, $J = 15.6$ Hz, 1H), 6.35–6.29 (m, 1H), 5.43–5.40 (m, 1H), 5.17–5.06 (m, 2H), 0.25 (s, 9H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 164.3, 140.5, 136.7, 136.0, 132.2, 129.3, 128.8, 128.1, 127.1, 51.0, 48.5, –0.5. MS (ESI) m/z 313.1 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{16}H_{21}N_4OSi$ $[M + H]^+$ 313.1479, found 301.1466.

(*S*)-4-(*p*-Tolyl)-3-(trimethylsilyl)-4,5-dihydro-[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4g**). A white solid (140 mg, 93%). Mp 199–200 °C. $[\alpha]_D^{25} = +124.1$ (c 0.19, MeOH). 1H NMR (400 MHz, DMSO- d_6) δ 9.07 (d, $J = 2.4$ Hz, 1H), 7.18 (d, $J = 8.0$ Hz, 2H), 7.08 (d, $J = 8.0$ Hz, 2H), 5.83 (s, 1H), 5.34–5.11 (m, 2H), 2.29 (s, 3H), 0.01 (s, 9H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 163.5, 140.4, 138.4, 138.4, 137.4, 129.8, 128.2, 52.5, 48.3, 21.2, –1.0. MS (ESI) m/z 301.1 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{15}H_{21}N_4OSi$ $[M + H]^+$ 301.1479, found 301.1480.

(*S*)-4-(4-Methoxyphenyl)-3-(trimethylsilyl)-4,5-dihydro-[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4h**). A white solid (149 mg, 94%). Mp 171–172 °C. $[\alpha]_D^{25} = +136.9$ (c 0.19, MeOH). 1H NMR (400 MHz, DMSO- d_6) δ 9.05 (s, 1H), 7.12 (d, $J = 8.8$ Hz, 2H), 6.92 (d, $J = 8.4$ Hz, 2H), 5.83 (s, 1H), 5.33–5.10 (m, 2H), 3.74 (s, 3H), –0.01 (s, 9H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 163.4, 159.8, 140.4, 137.6, 133.4, 129.6, 114.6, 55.7, 52.2, 48.3, 31.1, –1.0. MS (ESI) m/z 317.1 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{15}H_{21}N_4O_2Si$ $[M + H]^+$ 317.1428, found 317.1419.

(*S*)-4-(Pyridin-3-yl)-4,5-dihydro-[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4i**). This was further purified by silica gel chromatography (eluent: 2:1 hexane/acetone) to give a light-yellow oil (108 mg, 75%). $[\alpha]_D^{25} = +67.6$ (c 0.19, MeOH). 1H NMR (400 MHz, DMSO- d_6) δ 9.11 (s, 1H), 8.65 (s, 1H), 8.58 (s, 1H), 7.78 (d, $J = 8.0$ Hz, 1H), 7.54 (s, 1H), 7.45–7.42 (m, 1H), 6.09 (s, 1H), 5.31–5.11 (m, 2H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 164.4, 150.1, 148.9, 135.3, 132.1, 130.2, 49.7, 48.5. MS (ESI) m/z 216.1 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{10}H_{10}N_5O$ $[M + H]^+$ 216.0880, found 216.0877.

(*R*)-4-(Furan-2-yl)-3-(trimethylsilyl)-4,5-dihydro-[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4j**). This was further purified by silica gel chromatography (eluent: 5:1 hexane/acetone) to give a light-yellow solid (121 mg, 88%). Mp 160–161 °C. $[\alpha]_D^{25} = +85.5$ (c 0.24, MeOH). 1H NMR (400 MHz, DMSO- d_6) δ 9.17 (d, $J = 3.2$ Hz, 1H), 7.64 (d, $J = 0.8$ Hz, 1H), 6.47–6.42 (m, 2H), 6.03 (d, $J = 3.2$ Hz, 1H), 5.23–5.14 (m, 2H), 0.15 (s, 9H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 164.3, 152.0, 144.0, 140.6, 135.2, 111.3, 108.7, 48.5, 46.1, 31.1, –0.9. MS (ESI) m/z 277.1 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{12}H_{17}N_4O_2Si$ $[M + H]^+$ 277.1115, found 277.1103.

(*S*)-4-(1*H*-Indol-3-yl)-3-(trimethylsilyl)-4,5-dihydro-[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4k**). This was further purified by silica gel chromatography (eluent: 5:1 hexane/acetone) to give an orange-yellow solid (138 mg, 85%). Mp 256–257 °C. $[\alpha]_D^{25} = +116.2$ (c 0.10, MeOH). 1H NMR (400 MHz, DMSO- d_6) δ 11.19 (s, 1H), 8.95 (d, $J = 2.0$ Hz, 1H), 7.39–7.31 (m, 2H), 7.10–6.89 (m, 3H), 6.16 (d, $J = 1.6$ Hz, 1H), 5.35–5.14 (m, 2H), –0.16 (s, 9H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 163.6, 140.3, 138.1, 136.9, 125.5, 125.1, 122.0, 119.6, 118.7, 114.1, 112.3, 48.2, 46.0, –1.1. MS (ESI) m/z 326.1 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{16}H_{20}N_5OSi$ $[M + H]^+$ 326.1432, found 326.1426.

(*S*)-4-(4-Methoxyphenyl)-3-phenyl-4,5-dihydro-[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4l**). Reaction time: 3 h. This was further purified by silica gel chromatography (eluent: 5:1 hexane/acetone) to give a light-yellow oil (128 mg, 80%). $[\alpha]_D^{25} = +137.9$ (c 0.38, MeOH). 1H NMR (400 MHz, DMSO- d_6) δ 9.21 (d, $J = 2.4$ Hz, 1H), 7.62–7.60 (m, 2H), 7.37–7.28 (m, 3H), 7.17–7.15 (m, 2H), 6.85–6.82 (m, 2H), 6.22 (d, $J = 3.2$ Hz, 1H), 5.41–5.18 (m, 2H), 3.69 (s, 3H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 163.5, 159.6, 141.3, 132.3, 130.7, 129.1, 129.0, 128.4, 128.2, 126.9, 114.6, 55.6, 51.8, 48.9. MS (ESI) m/z 321.1

$[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{18}H_{17}N_4O_2$ $[M + H]^+$ 321.1346, found 321.1344.

(*S*)-3-Cyclopropyl-4-(4-methoxyphenyl)-4,5-dihydro-[1,2,3]triazolo[1,5-*a*]pyrazin-6(7*H*)-one (**4m**). Reaction time: 3 h. This was further purified by silica gel chromatography (eluent: 5:1 hexane/acetone) to give a white solid (107 mg, 75%). Mp 118–119 °C. $[\alpha]_D^{25} = +31.1$ (c 0.30, MeOH). 1H NMR (400 MHz, DMSO- d_6) δ 9.06 (d, $J = 2.4$ Hz, 1H), 7.27 (d, $J = 8.8$ Hz, 2H), 6.93 (d, $J = 8.4$ Hz, 2H), 5.90 (s, 1H), 5.26–5.02 (m, 2H), 3.75 (s, 3H), 0.77–0.73 (m, 1H), 0.63–0.56 (m, 2H), 0.53–0.48 (m, 2H). ^{13}C NMR (100 MHz, DMSO- d_6) δ 164.0, 159.6, 143.1, 132.5, 129.1, 128.8, 114.6, 55.7, 51.4, 48.6, 7.4, 7.0. MS (ESI) m/z 285.1 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{15}H_{17}N_4O_2$ $[M + H]^+$ 285.1346, found 285.1347.

General Procedure for Cu–NHC-Catalyzed AAC: 2-Azido-*N*-((*S*)-2-methyl-1-(1-(2-((*S*)-4-methyl-1-(trimethylsilyl)pent-1-yn-3-yl)-amino)-2-oxoethyl)-1*H*-1,2,3-triazol-4-yl)propyl)acetamide (**5a**). The NHC precursor **9** (15 mg, 0.05 mmol) was dissolved in acetonitrile (3 mL) under nitrogen. To this was added CuCl (5 mg, 0.05 mmol) and sodium *tert*-butoxide (4.8 mg, 0.05 mmol). The mixture was stirred for 3 h at rt. Then **3a** (150 mg, 0.5 mmol) and water (3 mL) were added successively to the reaction mixture, and the resulting solution was stirred at rt for 24 h. After removal of the solvent under vacuum, the crude product was purified by silica gel chromatography (eluent: 5:1 hexane/acetone) to give **5a** as a white solid (149 mg, 69%). Mp 108–109 °C. $[\alpha]_D^{25} = +170.8$ (c 0.20, MeOH). 1H NMR (400 MHz, CD₃OD) δ 7.91 (s, 1H), 5.18 (s, 2H), 4.97 (d, $J = 7.6$ Hz, 1H), 4.55 (d, $J = 6.0$ Hz, 1H), 3.93 (m, 2H), 2.30–2.23 (m, 1H), 1.95–1.86 (m, 1H), 1.04–0.99 (m, 9H), 0.92 (d, $J = 6.8$ Hz, 3H), 0.18 (s, 9H). ^{13}C NMR (100 MHz, CD₃OD) δ 168.3, 165.4, 124.1, 102.9, 87.9, 51.6, 51.5, 51.4, 32.8, 32.1, 18.3, 17.8, 17.6, 16.8, –1.5. MS (ESI) m/z 433.2 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{19}H_{33}N_8O_2Si$ $[M + H]^+$ 433.2490, found 433.2492.

2-Azido-*N*-((*S*)-cyclohexyl(1-(2-((*S*)-1-cyclohexyl-3-(trimethylsilyl)prop-2-yn-1-yl)amino)-2-oxoethyl)-1*H*-1,2,3-triazol-4-yl)methyl)acetamide (**5b**). A white solid (166 mg, 65%). Mp 158–159 °C. $[\alpha]_D^{25} = +82.1$ (c 0.17, MeOH). 1H NMR (400 MHz, CD₃OD) δ 7.90 (s, 1H), 5.16 (s, 2H), 4.98 (d, $J = 8.0$ Hz, 1H), 4.54 (d, $J = 6.8$ Hz, 1H), 3.91 (m, 2H), 1.92–1.56 (m, 12H), 1.31–0.99 (m, 10H), 0.18 (s, 9H). ^{13}C NMR (100 MHz, CD₃OD) δ 168.2, 165.4, 147.0, 124.2, 103.3, 87.9, 51.5, 51.5, 50.7, 42.4, 41.6, 29.6, 28.9, 28.8, 28.4, 25.9, 25.6, 25.6, 25.5, –1.5. MS (ESI) m/z 513.3 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{25}H_{41}N_8O_2Si$ $[M + H]^+$ 513.3116, found 513.3127.

2-Azido-*N*-((*S*)-1-(1-(2-oxo-2-((*S*)-1-phenyl-4-(trimethylsilyl)but-3-yn-2-yl)amino)ethyl)-1*H*-1,2,3-triazol-4-yl)-2-phenylethyl)acetamide (**5c**). A white syrup (179 mg, 68%). $[\alpha]_D^{25} = +119.7$ (c 0.15, MeOH). 1H NMR (400 MHz, CD₃OD) δ 7.77 (s, 1H), 7.31–7.18 (m, 10H), 5.43–5.39 (m, 1H), 5.13–5.03 (m, 2H), 4.88–4.86 (m, 1H), 3.81 (s, 2H), 3.33–3.29 (m, 1H), 3.20–3.14 (m, 1H), 2.97 (d, $J = 6.8$ Hz, 2H), 0.14 (s, 9H). ^{13}C NMR (100 MHz, CD₃OD) δ 168.0, 165.2, 137.4, 136.6, 129.4, 129.0, 128.0, 127.9, 126.5, 126.3, 123.8, 103.7, 88.2, 51.5, 51.5, 43.5, 41.1, 40.0, –1.6. MS (ESI) m/z 529.2 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{27}H_{33}N_8O_2Si$ $[M + H]^+$ 529.2492, found 529.2483.

2-Azido-*N*-((*S,E*)-1-(1-(2-oxo-2-((*S,E*)-1-phenyl-5-(trimethylsilyl)pent-1-en-4-yn-3-yl)amino)ethyl)-1*H*-1,2,3-triazol-4-yl)-3-phenylallyl)acetamide (**5d**). A white solid (198 mg, 72%). Mp 115–116 °C. $[\alpha]_D^{25} = -20.3$ (c 0.20, MeOH). 1H NMR (400 MHz, CD₃OD) δ 7.36 (s, 1H), 7.35–7.31 (m, 4H), 7.29–7.24 (m, 6H), 6.81 (d, $J = 15.6$ Hz, 1H), 6.69 (d, $J = 16.0$ Hz, 1H), 6.55–6.49 (m, 1H), 6.25–6.20 (m, 1H), 5.94 (d, $J = 6.4$ Hz, 1H), 5.49–5.47 (m, 1H), 5.23 (s, 2H), 3.98 (s, 2H), 0.22 (s, 9H). ^{13}C NMR (100 MHz, CD₃OD) δ 168.0, 165.2, 147.2, 136.4, 136.1, 132.2, 132.1, 128.3, 128.2, 127.8, 127.6, 126.3, 126.3, 126.2, 124.9, 124.2, 101.9, 89.0, 56.9, 51.6, 51.5, 43.3, –1.6. MS (ESI) m/z 553.2 $[M + H]^+$. HRMS (ESI-TOF) calcd for $C_{29}H_{32}N_8NaO_2Si$ $[M + Na]^+$ 575.2310, found 575.2296.

2-Azido-*N*-((*S*)-1-(2-oxo-2-((*S*)-1-(*p*-tolyl)-3-(trimethylsilyl)prop-2-yn-1-yl)amino)ethyl)-1*H*-1,2,3-triazol-4-yl)(*p*-tolyl)methyl)acetamide (**5e**). A white solid (214 mg, 81%). Mp 85–86 °C. $[\alpha]_D^{25} = -8.6$ (c 0.36, MeOH). 1H NMR (400 MHz, CD₃OD) δ 7.73 (s, 1H), 7.36 (d, $J = 8.0$ Hz, 2H), 7.27 (d, $J = 8.0$ Hz, 2H), 7.20–7.18 (m, 4H),

6.32 (s, 1H), 5.86 (s, 1H), 5.19–5.10 (m, 2H), 4.00–3.91 (m, 2H), 2.35 (m, 6H), 0.20 (s, 9H). ^{13}C NMR (100 MHz, CD_3OD) δ 167.9, 165.1, 137.8, 137.4, 136.8, 135.5, 129.0, 129.0, 126.9, 126.6, 124.6, 103.0, 88.7, 51.5, 51.4, 49.5, 45.0, 19.7, –1.6. MS (ESI) m/z 529.2 [$\text{M} + \text{H}$] $^+$. HRMS (ESI-TOF) calcd for $\text{C}_{27}\text{H}_{33}\text{N}_8\text{O}_2\text{Si}$ [$\text{M} + \text{H}$] $^+$ 529.2490, found 529.2498.

2-Azido-N-((S)-(4-methoxyphenyl)(1-(2-((S)-1-(4-methoxyphenyl)-3-(trimethylsilyl)prop-2-yn-1-yl)amino)-2-oxoethyl)-1H-1,2,3-triazol-4-yl)methyl)acetamide (5h). A white solid (221 mg, 79%). Mp 61–62 °C. $[\alpha]_{\text{D}}^{25} = -9.4$ (c 0.17, MeOH). ^1H NMR (400 MHz, CD_3OD) δ 7.73 (s, 1H), 7.40 (d, $J = 8.4$ Hz, 2H), 7.29 (d, $J = 8.8$ Hz, 2H), 6.93–6.90 (m, 4H), 6.30 (s, 1H), 5.85 (s, 1H), 5.18–5.05 (m, 2H), 3.99–3.89 (m, 2H), 3.80–3.79 (m, 6H), 0.20 (s, 9H). ^{13}C NMR (100 MHz, CD_3OD) δ 167.9, 165.0, 159.7, 159.5, 148.2, 131.8, 130.4, 128.3, 128.0, 124.5, 113.8, 113.7, 103.1, 88.7, 54.4, 54.4, 51.5, 51.4, 49.3, 44.7, –1.5. MS (ESI) m/z 561.2 [$\text{M} + \text{H}$] $^+$. HRMS (ESI-TOF) calcd for $\text{C}_{27}\text{H}_{32}\text{N}_8\text{NaO}_4\text{Si}$ [$\text{M} + \text{Na}$] $^+$ 583.2208, found 583.2203.

2-Azido-N-((R)-furan-2-yl(1-(2-((R)-1-(furan-2-yl)-3-(trimethylsilyl)prop-2-yn-1-yl)amino)-2-oxoethyl)-1H-1,2,3-triazol-4-yl)methyl)acetamide (5j). A light-yellow oil (180 mg, 75%). $[\alpha]_{\text{D}}^{25} = +13.9$ (c 0.33, MeOH). ^1H NMR (400 MHz, CD_3OD) δ 7.96 (s, 1H), 7.51–7.49 (m, 2H), 6.47–6.32 (m, 5H), 5.97 (m, 1H), 5.24–5.15 (m, 2H), 3.99–3.91 (m, 2H), 0.20 (s, 9H). ^{13}C NMR (100 MHz, CD_3OD) δ 168.0, 165.2, 151.7, 150.5, 145.6, 142.9, 142.9, 142.6, 124.7, 110.1, 107.6, 107.4, 100.3, 88.1, 51.4, 51.3, 43.9, 39.4, –1.7. MS (ESI) m/z 481.1 [$\text{M} + \text{H}$] $^+$. HRMS (ESI-TOF) calcd for $\text{C}_{21}\text{H}_{25}\text{N}_8\text{O}_4\text{Si}$ [$\text{M} + \text{H}$] $^+$ 481.1763, found 481.1766.

General Procedure for Microwave-Assisted Acid Hydrolysis: (S)-2-(5-(Amino(cyclohexyl)methyl)-1H-1,2,3-triazol-1-yl)acetic Acid (13b). Compound **4b** (55 mg, 0.22 mmol) was added to 6 M aqueous HCl (2.7 mL), and the mixture was heated under microwave conditions at 85 °C for three cycles of 1 h each. The solution was washed with CHCl_3 (4×3 mL), and the aqueous layer was evaporated under vacuum to give compound **13b** as a light-yellow solid (44.6 mg, 85% yield). Mp 96–97 °C. $[\alpha]_{\text{D}}^{25} = +11.1$ (c 0.75, MeOH). ^1H NMR (400 MHz, CD_3OD) δ 7.98 (s, 1H), 5.54–5.39 (AB, $J_{\text{AB}} = 18.0$ Hz, 2H), 4.45 (d, $J = 9.2$ Hz, 1H), 2.04–1.28 (m, 11H). ^{13}C NMR (100 MHz, CD_3OD) δ 168.4, 135.4, 132.1, 49.2 (2C), 41.0, 28.7, 28.6, 25.3, 25.1 (2C). HRMS (ESI-TOF) calcd for $\text{C}_{11}\text{H}_{18}\text{N}_4\text{NaO}_2$ [$\text{M} + \text{Na}$] $^+$ 261.1322, found 261.1319.

(S,E)-2-(5-(1-Amino-3-phenylallyl)-1H-1,2,3-triazol-1-yl)acetic Acid (13f). A yellow solid (45.4 mg, 80% yield). Mp 127–128 °C. $[\alpha]_{\text{D}}^{25} = +103.4$ (c 0.30, MeOH). ^1H NMR (400 MHz, D_2O) δ 7.91 (s, 1H), 7.46–7.44 (m, 2H), 7.37–7.32 (m, 3H), 6.82 (d, $J = 16.0$ Hz, 1H), 6.32 (dd, $J = 15.6, 8.0$ Hz, 1H), 5.33 (d, $J = 8.0$ Hz, 1H), 5.04 (AB, $J_{\text{AB}} = 17.6$ Hz, 2H). ^{13}C NMR (100 MHz, D_2O) δ 172.2, 136.9, 135.0, 134.7, 132.8, 129.3, 129.0, 127.1, 120.7, 52.1, 47.3. HRMS (ESI-TOF) calcd for $\text{C}_{13}\text{H}_{14}\text{N}_4\text{NaO}_2$ [$\text{M} + \text{Na}$] $^+$ 281.1009, found 281.1001.

ASSOCIATED CONTENT

Supporting Information

NMR spectra of all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Authors

*E-mail: sungaris@gmail.com.

*E-mail: wangbing@fudan.edu.cn.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors thank Professor Erkang Fan (University of Washington) for helpful discussions and are grateful for financial support from the Shanghai Municipal Education Commission (14ZZ159), the Shanghai Municipal Science and Technology Commission (12430501300), the National Natural Science Foundation of China (21072032 and 21272039), the

Special Scientific Foundation for Outstanding Young Teachers in Shanghai (ZZGJD13020), City Level Subjects of Shanghai University of Engineering and Technology (14XKCZ04), the Program for Special Appointment (Eastern Scholar) at Shanghai Institutions of Higher Education, and startup funding from Shanghai University of Engineering Science (E1-0501-14-0124 and NHRC-2014-05).

REFERENCES

- (1) For a review, see: Agalave, S. G.; Maujan, S. R.; Pore, V. S. *Chem.—Asian J.* **2011**, *6*, 2696–2718.
- (2) (a) Michael, A. J. *Prakt. Chem.* **1893**, *46*, 94. (b) Kirmse, W.; Horner, L. *Justus Liebigs Ann. Chem.* **1958**, *614*, 1–3. (c) Huisgen, R. *Proc. Chem. Soc.* **1961**, 357–396. (d) Huisgen, R. *Angew. Chem., Int. Ed. Engl.* **1963**, *2*, 565–598. (e) Huisgen, R. *Angew. Chem., Int. Ed. Engl.* **1963**, *2*, 633–645.
- (3) (a) Tormøe, C. W.; Christensen, C.; Meldal, M. *J. Org. Chem.* **2002**, *67*, 3057–3064. (b) Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2596–2599. For selected reviews, see: (c) Hein, J.; Fokin, V. V. *Chem. Soc. Rev.* **2010**, *39*, 1302–1315. (d) Kappe, C. O.; Van der Eycken, E. *Chem. Soc. Rev.* **2010**, *39*, 1280–1290. (e) Liang, L.; Astruc, D. *Coord. Chem. Rev.* **2011**, *255*, 2933–2945. (f) Jewett, J. C.; Bertozzi, C. R. *Chem. Soc. Rev.* **2010**, *39*, 1272–1279.
- (4) (a) Kolb, H. C.; Finn, M. G.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2001**, *40*, 2004–2021. (b) Evans, R. A. *Aust. J. Chem.* **2007**, *60*, 384–395.
- (5) (a) Kolb, H. C.; Sharpless, K. B. *Res. Focus* **2003**, *8*, 1128–1137. (b) Gil, M. V.; Arevalo, M. J.; Lopez, O. *Synthesis* **2007**, 1589–1620. (c) Aureggi, V.; Sedelmeier, G. *Angew. Chem., Int. Ed.* **2007**, *46*, 8440–8444. (d) Demko, Z. P.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2110–2113. (e) Lewis, W. G.; Green, L. G.; Grynszpan, F.; Radić, Z.; Carlier, P. R.; Taylor, P.; Finn, M. G.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 1053–1057. (f) Pachón, L. D.; Van Maarseveen, J. H.; Rothenberg, G. *Adv. Synth. Catal.* **2005**, *347*, 811–815. (g) Lipshutz, B. H.; Taft, B. R. *Angew. Chem., Int. Ed.* **2006**, *45*, 8235–8238.
- (6) (a) Zhang, L.; Chen, X.; Xue, P.; Sun, H. H. Y.; Williams, I. D.; Sharpless, K. B.; Fokin, V. V.; Jia, G. *J. Am. Chem. Soc.* **2005**, *127*, 15998–15999. (b) Boren, B. C.; Narayan, S.; Rasmussen, L. K.; Zhang, L.; Zhao, H.; Lin, Z.; Jia, G.; Fokin, V. V. *J. Am. Chem. Soc.* **2008**, *130*, 8923–8930. (c) Rasmussen, L. K.; Boren, B. C.; Fokin, V. V. *Org. Lett.* **2007**, *9*, 5337–5339. (d) Majireck, M. M.; Weinreb, S. M. *J. Org. Chem.* **2006**, *71*, 8680–8683.
- (7) (a) Díez-González, S.; Stevens, E. D.; Nolan, S. P. *Chem. Commun.* **2008**, 4747–4749. (b) Glorius, F. *Top. Organomet. Chem.* **2007**, *21*, 159–192. (c) Dröge, T.; Glorius, F. *Angew. Chem., Int. Ed.* **2010**, *49*, 6940–6952. (d) Mahatthananchai, J.; Bode, J. W. *Acc. Chem. Res.* **2014**, *47*, 696–707. (e) *N-Heterocyclic Carbenes in Transition Metal Catalysis and Organocatalysis*; Cazin, C. S. J., Ed.; Springer: Dordrecht, The Netherlands, 2010.
- (8) (a) Díez-González, S.; Correa, A.; Cavallo, L.; Nolan, S. P. *Chem.—Eur. J.* **2006**, *12*, 7558–7564. Two examples using a simple symmetric alkyne were reported in this paper. (b) Nakamura, T.; Terashima, T.; Ogata, K.; Fukuzawa, S. *Org. Lett.* **2011**, *13*, 620–623. (c) Díez-González, S.; Collinson, J. M.; Wilton-Ely, J. D. E. *T. Chem. Commun.* **2013**, *49*, 11358–11360.
- (9) For cycloaddition of azides to internal alkynes, see: (a) Balducci, E.; Bellucci, L.; Petricci, E.; Taddei, M.; Tafi, A. *J. Org. Chem.* **2009**, *74*, 1314–1321. (b) Mont, N.; Mehta, V. P.; Appukkuttan, P.; Beryozkina, T.; Toppet, S.; Van Hecke, K.; Van Meervelt, L.; Voet, A.; DeMaeyer, M.; Van der Eycken, E. *J. Org. Chem.* **2008**, *73*, 7509–7516. (c) Brawn, R. A.; Welzel, M.; Lowe, T.; Panek, J. S. *Org. Lett.* **2010**, *12*, 336–339. (d) De Moliner, F.; Crosignani, S.; Galatini, A.; Riva, R.; Basso, A. *ACS Comb. Sci.* **2011**, *13*, 453–457. (e) Fiandanese, V.; Maurantonio, S.; Punzi, A.; Rafaschieri, G. *Org. Biomol. Chem.* **2012**, *10*, 1186–1195. For an interesting stepwise formal cycloaddition, see: (f) Zhang, H.;

Tanimoto, H.; Morimoto, T.; Nishiyama, Y.; Kakiuchi, K. *Org. Lett.* **2013**, *15*, 5222–5225.

(10) Kloss, F.; Köhn, U.; Jahn, B. O.; Hager, M. D.; Görls, H.; Schubert, U. S. *Chem.—Asian J.* **2011**, *6*, 2816–2824.

(11) (a) Hein, J. E.; Tripp, J. C.; Krasnova, L. B.; Sharpless, K. B.; Fokin, V. V. *Angew. Chem., Int. Ed.* **2009**, *48*, 8018–8021. (b) Panteleev, J.; Geyer, K.; Aguilar-Aguilar, A.; Wang, L.; Lautens, M. *Org. Lett.* **2010**, *12*, 5092–5095. (c) Schulman, J. M.; Friedman, A. A.; Panteleev, J.; Lautens, M. *Chem. Commun.* **2012**, *48*, 55–57. For an overview, see: (d) Spiteri, C.; Moses, J. E. *Angew. Chem., Int. Ed.* **2010**, *49*, 31–33. For RuAAC of 1-iodoalkynes, see: (e) Oakdale, J. S.; Sit, R. K.; Fokin, V. V. *Chem.—Eur. J.* **2014**, *20*, 11101–11110. For reactions of 1-metalated alkyne intermediates, see: (f) Zhou, Y.; Lecourt, T.; Micouin, L. *Angew. Chem., Int. Ed.* **2010**, *49*, 2607–2610. (g) Worrell, B. T.; Ellery, S. P.; Fokin, V. V. *Angew. Chem., Int. Ed.* **2013**, *52*, 13037–13041. For an Ir-catalyzed cycloaddition to 1-thioalkynes, see: (h) Ding, S.; Jia, G.; Sun, J. *Angew. Chem., Int. Ed.* **2014**, *53*, 1877–1880.

(12) (a) Wen, K.; Wang, H.; Cheng, J.; Zhang, H.; Cui, X.; Wei, C.; Fan, E.; Sun, Z. *J. Org. Chem.* **2013**, *78*, 3405–3409. (b) Wang, H.; Xia, Y.; Lv, S.; Xu, J.; Sun, Z. *Tetrahedron Lett.* **2013**, *54*, 2124–2127. (c) Cheng, J.; Chen, Y.-L.; Zheng, Y.; Sun, Z. *RSC Adv.* **2014**, *4*, 21131–21133.

(13) Chen, B.-L.; Wang, B.; Lin, G.-Q. *J. Org. Chem.* **2010**, *75*, 941–944.

(14) Boyer, J. H.; Hamer, J. *J. Am. Chem. Soc.* **1955**, *77*, 951–954.

(15) (a) Mehta, S.; Larock, R. C. *J. Org. Chem.* **2010**, *75*, 1652–1658. (b) Huang, J.; Macdonald, S. J. F.; Harrity, J. P. A. *Chem. Commun.* **2009**, 436–438. (c) Reference 9e.

(16) (a) Shao, C.; Wang, X.; Xu, J.; Zhao, J.; Zhang, Q.; Hu, Y. *J. Org. Chem.* **2010**, *75*, 7002–7005. (b) Shao, C.; Zhu, R.; Luo, S.; Zhang, Q.; Wang, X.; Hu, Y. *Tetrahedron Lett.* **2011**, *52*, 3782–3785.

(17) Slow addition of a dilute solution of the same substrate to the Cu–NHC catalyst produced the “dimer” and “trimer” in a 65:35 ratio (as determined by LC–MS). We thank one of the reviewers for the constructive suggestion of the possibility of reaction kinetics involving a quasi-stationary low concentration of the desilylated alkyne intermediate.