

# Synthesis of Vinyl Trifluoromethyl Thioethers via Copper-Mediated Trifluoromethylthiolation of Vinyl Bromides


Yangjie Huang,<sup>†</sup> Jianping Ding,<sup>†</sup> Chuyi Wu,<sup>†</sup> Huidong Zheng,<sup>\*,†</sup> and Zhiqiang Weng<sup>\*,†,‡</sup>

<sup>†</sup>Department of Chemistry, Fuzhou University, Fuzhou, Fujian 350108, P. R. China

<sup>‡</sup>State Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Fuzhou, Fujian 350002, P. R. China

**S** Supporting Information

**ABSTRACT:** A copper-mediated trifluoromethylthiolation of vinyl bromides has been developed. This method provides ready access to vinyl trifluoromethyl thioethers in good to high yields from simple, inexpensive starting materials. A broad substrate scope is achieved, and the reaction is compatible with various functional groups, including cyano, nitro, trifluoromethyl, alkoxy, amino, halide, and heterocyclic groups.




Fluorine containing compounds are ubiquitous in pharmaceuticals, agrochemicals, and advanced materials.<sup>1,2</sup> Their fluorine moieties offer unusual chemical and physical properties, which can profoundly enhance the biological activity and metabolic stability.<sup>3,4</sup> The trifluoromethylthio group (CF<sub>3</sub>S–) is of special interest, as it has one of the highest Hansch hydrophobicity parameter values ( $\pi_R = 1.44$ ).<sup>5</sup> As a consequence, great synthetic efforts have been exerted to develop efficient methods for the incorporation of the CF<sub>3</sub>S– group into organic molecules.<sup>6–11</sup> In particular, transition-metal-catalyzed or -mediated trifluoromethylthiolation have experienced impressive progress and growth in recent years. For instance, Cu-catalyzed and -mediated trifluoromethylthiolation of aryl<sup>12–14</sup> and alkyl boronic acids,<sup>15</sup> aryl halides,<sup>16</sup> and arenediazonium salts,<sup>17</sup> related Cu<sup>18–20</sup> and Pd-promoted<sup>21</sup> trifluoromethylthiolation through the direct functionalization of C–H bonds, Ag-catalyzed decarboxylative trifluoromethylthiolation of aliphatic carboxylic acids,<sup>22</sup> and Pd-<sup>23</sup> and Ni-catalyzed trifluoromethylthiolation of aryl halides<sup>24</sup> have provided attractive access to compounds bearing C<sub>sp2</sub>–SCF<sub>3</sub> and C<sub>sp3</sub>–SCF<sub>3</sub> as well as C<sub>sp</sub>–SCF<sub>3</sub> bonds.

However, synthetic processes for the direct construction of C<sub>vinyl</sub>–SCF<sub>3</sub> bonds via transition-metal-catalyzed or -mediated trifluoromethylthiolation have not been well developed (Scheme 1). The Vivic,<sup>13</sup> Shen,<sup>25</sup> and Rueping<sup>14</sup> groups independently reported Cu-mediated and -catalyzed trifluoromethylthiolation of vinyl boronic acids to form vinyl trifluoromethyl thioethers. Most recently, Rueping and co-workers also developed a Cu-catalyzed trifluoromethylthiolation of vinyl iodides.<sup>26</sup> However, many of these procedures often require vinylboron compounds or vinyl iodides as substrates and/or the use of an electrophilic type trifluoromethylthiolation reagent.

We have therefore sought to develop a highly facile and cost-effective route to vinyl trifluoromethyl thioethers via copper-mediated trifluoromethylthiolation of vinyl bromides.

Copper-promoted coupling of vinyl bromides or iodides with S-nucleophiles has proven to be a powerful method for the

## Scheme 1. Methods for Preparation of Vinyl Trifluoromethyl Thioethers




synthesis of vinyl sulfides.<sup>27–32</sup> However, the coupling of vinyl bromides with SCF<sub>3</sub><sup>–</sup> to give vinyl trifluoromethyl thioether products is a challenging task because of the reduced nucleophilicity of the SCF<sub>3</sub><sup>–</sup> anion.<sup>26</sup>

As part of our continuing studies on copper-mediated trifluoromethylthiolation,<sup>33–38</sup> we recently developed a novel C<sub>aryl</sub>–SCF<sub>3</sub> bond formation mediated by a copper(I) trifluoromethylthiolate complex (bpy)Cu(SCF<sub>3</sub>) (**2**; bpy = 2,2′-bipyridine).<sup>33</sup> This complex reacted with various aryl iodides or activated aryl bromides to afford aryl trifluoromethyl thioethers in modest to excellent yields. These results encourage us to probe the feasibility of performing the trifluoromethylthiolation of more challenging substrates, such as vinyl bromides, using the copper reagent **2**.

Initially, trifluoromethylthiolation of phenylvinyl bromide (**1a**) with **2** to form styrenyl trifluoromethyl thioethers (**3a**) was investigated. The reaction first was conducted in CH<sub>3</sub>CN at 100 °C for 16 h providing the desired product **3a** in 23% yield (Table 1, entry 1). Next, we examined the possible influence of

Received: January 20, 2015

Published: February 12, 2015

**Table 1. Optimization of the Trifluoromethylthiolation of Phenylvinyl Bromide<sup>a</sup>**



entry	additive (equiv)	solvent	temp (°C)	time (h)	yield (%) <sup>b</sup>
1		CH <sub>3</sub> CN	100	16	23
2		DMF	100	16	28
3		DMSO	100	16	41
4		toluene	100	16	50
5		THF	100	16	59
6		dioxane	100	16	35
7		diglyme	100	16	64
8	LiF (2)	diglyme	100	16	63
9	NaF (2)	diglyme	100	16	57
10	KF (2)	diglyme	100	16	74
11	CsF (2)	diglyme	100	16	69
12	AgF (2)	diglyme	100	16	52
13	<i>n</i> -Bu <sub>4</sub> NF (2)	diglyme	100	16	13
14	KF (4)	diglyme	100	16	80
15	KF (4)	diglyme	100	24	92
16	KF (4)	diglyme	80	24	71

<sup>a</sup>Reaction conditions: **1a** (0.050 mmol), **2** (0.060 mmol), solvent (1.0 mL), N<sub>2</sub>. <sup>b</sup>The yield was determined by <sup>19</sup>F NMR spectroscopy with PhOCF<sub>3</sub> as internal standard.

some parameters such as the solvent and additive on the outcome of the reaction. Other solvents, such as DMF, DMSO, and dioxane, were also ineffective (Table 1, entries 2, 3, and 6). When the solvent was changed to toluene and THF, the desired product **3a** was obtained in yields of 50% and 59%, respectively (Table 1, entries 4, and 5). It was found that diglyme was an optimal solvent, affording **3a** in 64% yield (Table 1, entry 7). Thus, diglyme was chosen as a solvent of choice for further studies. Since the addition of alkali metal fluorides could significantly enhance the reaction efficiency probably by facilitating the formation of an active intermediate,<sup>36,37</sup> various fluorides were screened in the reaction (Table 1, entries 8–13). It was found that KF was superior to all other screened additives for the reaction (Table 1, entry 10). Increasing the quantity of KF from 2 to 4 equiv resulted in a higher yield of 80% (Table 1, entry 14). The reaction time was also examined. The results indicated that the yield of **3a** increased from 80% to 92% when the reaction time was prolonged from 16 to 24 h (Table 1, entry 15). Furthermore, lowering the reaction temperature to 80 °C resulted in a decreased yield of 71% (Table 1, entry 16).

Subsequently, a comparison with other trifluoromethylthiolation reagents was performed (Table 2). Reaction with (phen)Cu(SCF<sub>3</sub>) (phen = phenanthroline) generated product **3a** in moderate yield (55%; Table 2, entry 2). Similar behaviors were also observed in the previously reported trifluoromethylthiolation of aryl iodides.<sup>35</sup> Moreover in the case of the Cu(I) catalyzed trifluoromethylthiolation using the [Bu<sub>4</sub>N][SCF<sub>3</sub>] reagent, the reaction produced a lower yield of **3a** (30%; Table 2, entry 3). This result shows a higher reactivity of (bpy)Cu(SCF<sub>3</sub>) (**2**) compared to other reagents (Table 2, entry 1).

The generality of the reaction conditions was next tested on a variety of alkenyl bromides as illustrated in Table 3. A structurally diverse series of substituted  $\beta$ -bromostyrene underwent trifluoromethylthiolation with **2** by this procedure to produce the corresponding styrenyl trifluoromethyl thioethers in good to excellent yields. Moreover, no aryl acetylenes or homocoupled

**Table 2. Comparison of Copper-Mediated/-Catalyzed Trifluoromethylthiolation of Phenylvinyl Bromide**


entry	conditions	yield (%) <sup>a</sup>
1	(bpy)Cu(SCF <sub>3</sub> ) ( <b>2</b> ) (1.2 equiv)	92
2	(phen)Cu(SCF <sub>3</sub> ) (1.2 equiv)	55
3	[Bu <sub>4</sub> N][SCF <sub>3</sub> ] (2 equiv), CuI (20 mol %), phen (20 mol %)	30

<sup>a</sup>The yield was determined by <sup>19</sup>F NMR spectroscopy with PhOCF<sub>3</sub> as internal standard.

side products were observed. The products were characterized by IR, <sup>1</sup>H NMR, <sup>13</sup>C NMR, <sup>19</sup>F NMR, MS, and HRMS spectra. The stereochemistry, easily assignable by the <sup>1</sup>H NMR coupling constant value of the olefinic protons (*trans* configuration: *J* = 15.0–15.4 Hz; *cis* configuration: *J* = 10.6–10.7 Hz), is in accordance with the results obtained in closely related molecules.<sup>13,25,26</sup> It also is worth noting that the *E/Z* stereochemistry of the styrenyl trifluoromethyl thioether bond corresponds to the starting vinyl bromide (Table 3, entries 1–16).


For example, phenylvinyl bromides **1a** and **1b** afforded the corresponding products **3a** and **3b** in 90% and 92% yields, respectively (Table 3, entries 1 and 2).  $\beta$ -Bromostyrene having electron-donating groups on the phenyl ring were successfully converted to the desired products **3c–3f** in good to high yields (75%–93%; Table 3, entries 3–6). Furthermore, the introduction of electron-withdrawing substituents, including cyano, trifluoromethyl, and nitro groups, were well tolerated to give **3g–3j** in good yields (76–81%; Table 3, entries 7–10). This observation is consistent with our previous studies on trifluoromethylthiolation of  $\alpha$ -halo- $\alpha,\beta$ -unsaturated carbonyl substrates.<sup>36</sup> Interestingly, the fluoro and chloro groups at the *ortho*-, *meta*-, or *para*-position of aromatic rings were also tolerated and the respective products **3k–3o** were afforded in 62–85% yields (Table 3, entries 11–15). This method can be further extended to the heteroarylvinyl bromide systems. The reactions of (*Z*)-2-(2-bromovinyl)pyridine (**1p**) with **2** gave **3p** in 78% yield (Table 3, entry 16). Likewise,  $\alpha$ -bromostyrene (**1q**) could also be reactive under the standard reaction conditions, leading to the desired product **3q** in 73% yield (Table 3, entry 17). Moreover, even 2-bromo-1,1,2-triphenylethylene (**1r**), a sterically highly demanding arylvinyl bromide, was a suitable substrate providing good yield of product **3r** (72%; Table 3, entry 18). Unfortunately, trifluoromethylthiolation of a cyclic vinyl bromide **1s** under the standard conditions was less efficient, producing a low yield of product **3s** (23% of <sup>19</sup>F NMR yield; Table 3, entry 19), along with dehalogenated and other unidentified side products. In addition to vinyl bromides, nonconjugated vinyl iodides are also suitable for the reaction. (1-Iodovinyl)cyclohexane (**1t**) and 2-iodooct-1-ene (**1u**) reacted with **2** to deliver good yields of the desired products **3t** and **3u** (70% and 61% of <sup>19</sup>F NMR yields, respectively; Table 3, entries 20 and 21).

To demonstrate the scalability and practicability of our synthesis, the trifluoromethylthiolation was conducted on a gram scale. Using 1.0 g of **1a** as the substrate, the trifluoromethylthiolated product **3a** was obtained in 86% yield (Scheme 2).

Table 3. Substrate Scope of Trifluoromethylthiolation of Vinyl Bromides<sup>a</sup>

entry	vinyl halides <b>1</b> ( <i>E/Z</i> )	products <b>3</b> ( <i>E/Z</i> ) <sup>b</sup>	yield (%)	entry	vinyl halides <b>1</b> ( <i>E/Z</i> )	products <b>3</b> ( <i>E/Z</i> ) <sup>b</sup>	yield (%)
1	<b>1a</b> (94:6)	<b>3a</b> (94:6)	90	12	<b>1l</b> (93:7)	<b>3l</b> (93:7)	85
2	<b>1b</b> (95:5)	<b>3b</b> (95:5)	92	13	<b>1m</b> (70:30)	<b>3m</b> (70:30)	79
3	<b>(E)-1c</b>	<b>(E)-3c</b>	89	14	<b>1n</b> (85:15)	<b>3n</b> (85:15)	83
4	<b>(E)-1d</b>	<b>(E)-3d</b>	91	15	<b>(E)-1o</b>	<b>(E)-3o</b>	62
5	<b>(E)-1e</b>	<b>(E)-3e</b>	75	16	<b>(Z)-1p</b>	<b>(Z)-3p</b>	78
6	<b>1f</b> (95:5)	<b>3f</b> (95:5)	93	17	<b>1q</b>	<b>3q</b>	73
7	<b>1g</b> (55:45)	<b>3g</b> (55:45)	80	18	<b>1r</b>	<b>3r</b>	72
8	<b>1h</b> (70:30)	<b>3h</b> (70:30)	79	19	<b>1s</b>	<b>3s</b>	23 <sup>c</sup>
9	<b>1i</b> (40:60)	<b>3i</b> (40:60)	81	20	<b>1t</b>	<b>3t</b>	70 <sup>c</sup>
10	<b>1j</b> (80:20)	<b>3j</b> (80:20)	76	21	<b>1u</b>	<b>3u</b>	61 <sup>c</sup>
11	<b>1k</b> (80:20)	<b>3k</b> (80:20)	77				

<sup>a</sup>Reaction conditions: **1** (0.50 mmol), **2** (0.60 mmol), KF (2.0 mmol), diglyme (5.0 mL), N<sub>2</sub>. <sup>b</sup>The *E/Z* isomer ratio was determined from the <sup>19</sup>F NMR analysis. <sup>c</sup>The yield was determined by <sup>19</sup>F NMR spectroscopy with PhOCF<sub>3</sub> as internal standard.

Scheme 2. Scalability of the Trifluoromethylthiolation of **1a**

A series of experiments were conducted to probe whether any radical intermediates are involved in this transformation (Table 4). The reaction proceeds cleanly in the presence of the radical inhibitor 2,6-bis(1,1-dimethylethyl)-4-methylphenol (BHT, 1.0 equiv) to afford the desired product **3a** in 93% yield. Similarly, carrying out the reaction in the presence of 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO), a radical scavenger, gave **3a** in 88% yield. Taken together, these two results and the complete retention of the double-bond stereochemistry after reaction

Table 4. Inhibition Experiment on the Trifluoromethylthiolation of **1a**<sup>a</sup>

entry	additive (equiv)	yield (%) <sup>b</sup>
1	BHT (1.0)	93
2	TEMPO (1.0)	88

<sup>a</sup>Reaction conditions: **1a** (0.050 mmol), **2** (0.060 mmol), KF (0.20 mmol), diglyme (1.0 mL), N<sub>2</sub>. <sup>b</sup>The yield was determined by <sup>19</sup>F NMR spectroscopy with PhOCF<sub>3</sub> as internal standard.

appear inconsistent with a radical chain mechanism.<sup>39</sup> Thus, it is likely that the reaction proceeds via an oxidative addition–reductive elimination pathway.<sup>26,33</sup>

In conclusion, we have demonstrated a copper-mediated trifluoromethylthiolation of vinyl bromides that gives vinyl trifluoromethyl thioethers in good to excellent yields. The reaction demonstrates a broad substrate scope and a variety of functional groups, including cyano, nitro, trifluoromethyl, alkoxy, amino, halide, and heterocyclic groups that are tolerated. The complete retention of the olefin geometry, functional group tolerance, and scalability of this reaction make it an attractive protocol for the practical synthesis of these potentially important molecules.

## EXPERIMENTAL SECTION

**General Methods.** All manipulations were carried out under an inert atmosphere using a nitrogen-filled glovebox or standard Schlenk techniques. All glassware was oven or flame-dried immediately prior to use. Solvents were freshly dried and degassed according to the procedures in Purification of Laboratory Chemicals prior to use. Deuterated solvents were purchased commercially and were degassed and stored over activated 4 Å molecular sieves.  $\beta$ -Bromostyrene **1b-o**,<sup>40,41</sup> **1p**,<sup>42</sup> alkenyl iodides **1t-u**,<sup>43</sup> (bpy)Cu(SCF<sub>3</sub>), (phen)Cu(SCF<sub>3</sub>),<sup>35</sup> and [Bu<sub>4</sub>N][SCF<sub>3</sub>]<sup>44</sup> were prepared according to the published procedures. All other reagents were obtained from commercial sources and used without further purification. The <sup>1</sup>H, <sup>19</sup>F, and <sup>13</sup>C{<sup>1</sup>H} NMR spectra were recorded at 400, 376, and 101 MHz, respectively. <sup>19</sup>F NMR chemical shifts were determined relative to CFCl<sub>3</sub> as the outside standard and low field is positive. Mass spectrometry was performed on a GC/MS spectrometer with the electron impact (EI) ionization technique. HRMS data were recorded on a GC-TOF instrument using the EI technique.

**General Procedure for Trifluoromethylthiolation of Phenylvinyl Bromide with (bpy)Cu(SCF<sub>3</sub>).** Phenylvinyl bromide **1** (0.50 mmol), [(bpy)Cu(SCF<sub>3</sub>)]**2** (192 mg, 0.60 mmol, 1.2 equiv), KF (116 mg, 2.0 mmol), and diglyme (5.0 mL) were added to a reaction tube with a Teflon screw cap equipped with a stir bar. The mixture was stirred at 100 °C for 24 h. The reaction mixture was filtered through a pad of Celite. The filtrate was added to water (10 mL) at 0 °C. The resulting mixture was extracted with Et<sub>2</sub>O (3 × 15 mL), and the combined organic layers was washed with water and then dried over MgSO<sub>4</sub>. The solvent was removed by rotary evaporation in an ice bath, and the resulting product was purified by column chromatography on silica gel with pentane/Et<sub>2</sub>O.

**Styryl(trifluoromethyl)sulfane<sup>14</sup> (3a).** Obtained as a yellow oil in 90% yield (91 mg) with an *E/Z* ratio of 94:6. *R<sub>f</sub>* (*n*-pentane/diethyl ether 10:1) = 0.87. NMR of the major isomer: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.61–7.29 (m, 5H), 7.02 (d, *J* = 15.3 Hz, 1H), 6.75 (d, *J* = 15.3 Hz, 1H). <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) δ –42.7 (s, 3F). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>) δ 141.2 (q, *J* = 1.2 Hz), 135.1 (s), 129.6 (q, *J* = 307.9 Hz), 129.2 (s), 128.9 (s), 126.8 (s), 111.7 (q, *J* = 3.3 Hz). IR(KBr): ν 1716, 1652, 1541, 1497, 1447, 1109, 953, 739, 757, 698, 590, 467 cm<sup>-1</sup>. GC-MS *m/z* 204 (M<sup>+</sup>), 135 (M<sup>+</sup>–CF<sub>3</sub>). HRMS (EI): calcd for C<sub>9</sub>H<sub>7</sub>F<sub>3</sub>S, 204.0221; found, 204.0217.

**(4-Methylstyryl)(trifluoromethyl)sulfane<sup>26</sup> (3b).** Obtained as a yellow oil in 92% yield (100 mg) with an *E/Z* ratio of 95:5. *R<sub>f</sub>* (*n*-pentane/diethyl ether 10:1) = 0.87. NMR of the major isomer: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.28 (d, *J* = 8.1 Hz, 2H), 7.16 (d, *J* = 8.1 Hz, 2H), 6.98 (d, *J* = 15.3 Hz, 1H), 6.66 (d, *J* = 15.3 Hz, 1H), 2.35 (s, 3H). <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) δ –42.9 (s, 3F). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>) δ 141.8 (q, *J* = 1.3 Hz), 139.4 (s), 132.4 (s), 129.7 (q, *J* = 307.9 Hz), 129.6 (s), 126.8 (s), 110.3 (q, *J* = 3.2 Hz), 21.3 (s). IR(KBr): ν 2926, 2359, 1512, 1157, 1110, 958, 824, 787, 755, 505 cm<sup>-1</sup>. GC-MS *m/z* 218 (M<sup>+</sup>), 149 (M<sup>+</sup>–CF<sub>3</sub>). HRMS (EI): calcd for C<sub>10</sub>H<sub>9</sub>F<sub>3</sub>S, 218.0377; found, 218.0379.

**(E)-(4-Methoxystyryl)(trifluoromethyl)sulfane<sup>25</sup> (3c).** Obtained as a yellow oil in 89% yield (104 mg). *R<sub>f</sub>* (*n*-pentane/diethyl ether 10:1) = 0.74. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.35 (d, *J* = 8.8 Hz, 2H), 6.99 (d, *J* = 15.2 Hz, 1H), 6.89 (d, *J* = 8.8 Hz, 2H), 6.57 (d, *J* = 15.2 Hz, 1H), 3.83 (s, 3H). <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) δ –43.1 (s, 3F). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>) δ 160.5 (s), 142.2 (q, *J* = 1.1 Hz),

129.7 (q, *J* = 307.9 Hz), 128.4 (s), 127.9 (s), 114.3 (s), 108.4 (q, *J* = 3.0 Hz), 55.3 (s). IR(KBr): ν 2961, 2839, 2359, 1607, 1511, 1305, 1258, 1240, 1107, 1033, 959, 837, 796, 754 cm<sup>-1</sup>. GC-MS *m/z* 234 (M<sup>+</sup>), 165 (M<sup>+</sup>–CF<sub>3</sub>). HRMS (EI): calcd for C<sub>10</sub>H<sub>9</sub>F<sub>3</sub>OS, 234.0326; found, 234.0323.

**(E)-(2-Methoxystyryl)(trifluoromethyl)sulfane (3d).** Obtained as a yellow oil in 91% yield (106 mg). *R<sub>f</sub>* (*n*-pentane/diethyl ether 10:1) = 0.65. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.42 (dd, *J* = 7.6, 1.6 Hz, 1H), 7.37–7.26 (m, 2H), 7.01–6.92 (m, 2H), 6.87 (d, *J* = 15.4 Hz, 1H), 3.91 (s, 3H). <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) δ –42.9 (s, 3F). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>) δ 157.1 (s), 136.8 (q, *J* = 1.2 Hz), 130.3 (s), 129.8 (q, *J* = 307.7 Hz), 128.0 (s), 124.0 (s), 120.7 (s), 112.3 (q, *J* = 3.1 Hz), 111.0 (s), 55.5 (s). IR(KBr): ν 2965, 2840, 2360, 2341, 1487, 1250, 1106, 1027, 959, 749 cm<sup>-1</sup>. GC-MS *m/z* 234 (M<sup>+</sup>), 165 (M<sup>+</sup>–CF<sub>3</sub>). HRMS (EI): calcd for C<sub>10</sub>H<sub>9</sub>F<sub>3</sub>OS, 234.0326; found, 234.0327.

**(E)-(Trifluoromethyl)(3,4,5-trimethoxystyryl)sulfane (3e).** Obtained as a yellow oil in 75% yield (110 mg). *R<sub>f</sub>* (*n*-pentane/diethyl ether 10:1) = 0.34. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 6.94 (d, *J* = 15.3 Hz, 1H), 6.63 (d, *J* = 15.3 Hz, 1H), 6.61 (s, 2H), 3.88 (s, 6H), 3.86 (s, 3H). <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) δ –42.8 (s, 3F). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>) δ 153.5 (s), 141.7 (q, *J* = 1.1 Hz), 139.2 (s), 130.7 (s), 129.6 (q, *J* = 308.1 Hz), 110.7 (q, *J* = 3.2 Hz), 104.0 (s), 61.0 (s), 56.2 (s). IR(KBr): ν 2941, 1576, 1541, 1507, 1457, 1417, 1334, 1242, 1125, 1005, 953, 799, 755 cm<sup>-1</sup>. GC-MS *m/z* 294 (M<sup>+</sup>), 225 (M<sup>+</sup>–CF<sub>3</sub>). HRMS (EI): calcd for C<sub>12</sub>H<sub>13</sub>F<sub>3</sub>O<sub>3</sub>S, 294.0537; found, 294.0532.

***N,N*-Dimethyl-4-(2-(trifluoromethylthio)vinyl)aniline (3f).** Obtained as a yellow oil in 93% yield (114 mg) with an *E/Z* ratio of 95:5. *R<sub>f</sub>* (*n*-pentane/diethyl ether 10:1) = 0.69. NMR of the major isomer: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.29 (d, *J* = 8.4 Hz, 2H), 6.97 (d, *J* = 15.0 Hz, 1H), 6.65 (d, *J* = 8.4 Hz, 2H), 6.43 (d, *J* = 15.0 Hz, 1H), 2.99 (s, 6H). <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) δ –43.6 (s, 3F). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>) δ 151.1 (s), 144.3 (s), 129.8 (q, *J* = 308.0 Hz), 128.3 (s), 127.2 (s), 123.2 (d, *J* = 0.9 Hz), 112.0 (s), 40.3 (s). IR(KBr): ν 2925, 2854, 1608, 1522, 1446, 1362, 1185, 1153, 1110, 792 cm<sup>-1</sup>. GC-MS *m/z* 247 (M<sup>+</sup>), 178 (M<sup>+</sup>–CF<sub>3</sub>). HRMS (EI): calcd for C<sub>11</sub>H<sub>12</sub>F<sub>3</sub>NS, 247.0643; found, 247.0642.

**4-(2-(Trifluoromethylthio)vinyl)benzonitrile (3g).** Obtained as a yellow oil in 80% yield (91 mg) with an *E/Z* ratio of 55:45. *R<sub>f</sub>* (*n*-pentane/diethyl ether 10:1) = 0.58. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) *E* + *Z*: δ 7.67 (d, *J* = 7.2 Hz, 2H), 7.64 (d, *J* = 7.2 Hz, 2H), 7.47 (d, *J* = 7.2 Hz, 2H), 7.42 (d, *J* = 7.2 Hz, 2H), 6.92 (s, 2H), 6.83 (d, *J* = 10.7 Hz, 1H), 6.59 (d, *J* = 10.7 Hz, 1H). <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) *E* + *Z*: δ –42.2 (s, 3F) –42.9 (s, 3F). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>) *E* + *Z*: δ 139.3/139.2 (s), 136.4 (q, *J* = 1.2 Hz), 132.7 (s), 132.4 (s), 129.12/129.11 (q, *J* = 308.0 Hz), 129.12 (s), 127.1 (s), 117.9/117.0 (q, *J* = 3.8 Hz). IR(KBr): ν 2977, 2916, 2848, 2230, 1463, 1114, 734 cm<sup>-1</sup>. GC-MS *m/z* 229 (M<sup>+</sup>), 160 (M<sup>+</sup>–CF<sub>3</sub>). HRMS (EI): calcd for C<sub>10</sub>H<sub>6</sub>F<sub>3</sub>NS, 229.0173; found, 229.0169.

**(Trifluoromethyl)(4-(trifluoromethyl)styryl)sulfane<sup>26</sup> (3h).** Obtained as a yellow oil in 79% yield (107 mg) with an *E/Z* ratio of 70:30. *R<sub>f</sub>* (*n*-pentane/diethyl ether 10:1) = 0.83. NMR of the major isomer: <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.62 (d, *J* = 8.0 Hz, 2H), 7.49 (d, *J* = 8.0 Hz, 2H), 6.98 (d, *J* = 15.0 Hz, 1H), 6.86 (d, *J* = 15.0 Hz, 1H). <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) δ –42.4 (s, 3F), –62.8 (s, 3F). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>) δ 138.3 (q, *J* = 1.3 Hz), 137.8 (s), 130.6 (s), 129.4 (q, *J* = 308.1 Hz), 128.8 (s), 126.9 (s), 125.9 (q, *J* = 3.8 Hz), 115.4 (q, *J* = 3.1 Hz). IR(KBr): ν 2955, 2924, 2854, 1457, 1326, 1168, 1135, 1110, 1068 cm<sup>-1</sup>. GC-MS *m/z* 272 (M<sup>+</sup>), 203 (M<sup>+</sup>–CF<sub>3</sub>). HRMS (EI): calcd for C<sub>10</sub>H<sub>6</sub>F<sub>6</sub>S, 272.0094; found, 272.0095.

**(3-Nitrostyryl)(trifluoromethyl)sulfane (3i).** Obtained as a yellow oil in 81% yield (100 mg) with an *E/Z* ratio of 40:60. *R<sub>f</sub>* (*n*-pentane/diethyl ether 10:1) = 0.64. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) *E* + *Z*: δ 8.26–8.14 (m, 4H), 7.73–7.64 (m, 2H), 7.63–7.53 (m, 2H), 7.04–6.87 (m, 2H), 6.61 (d, *J* = 10.6 Hz, 2H). <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>) δ –42.2 (s) (*E/Z*), –42.8 (s) (*E/Z*). <sup>13</sup>C{<sup>1</sup>H} NMR (101 MHz, CDCl<sub>3</sub>) *E* + *Z*: δ 134.2 (s), 132.4 (s), 129.9 (s), 129.6 (s), 129.3/129.1 (q, *J* = 308.2 Hz), 123.4/123.3 (s), 122.8 (s), 121.2 (s), 117.5/116.3 (q, *J* = 3.8 Hz). IR(KBr): ν 2926, 2855, 1609, 1532, 1458, 1351, 1111, 940, 806, 750, 737, 669 cm<sup>-1</sup>. GC-MS *m/z* 249 (M<sup>+</sup>), 180 (M<sup>+</sup>–CF<sub>3</sub>). HRMS (EI): calcd for C<sub>9</sub>H<sub>6</sub>F<sub>3</sub>NO<sub>2</sub>S, 249.0071; found, 249.0066.

**(4-Nitrostyryl)(trifluoromethyl)sulfane (3j).** Obtained as a yellow solid in 76% yield (94 mg) with an *E/Z* ratio 80:20.  $R_f$  (*n*-pentane/diethyl ether 10:1) = 0.67. NMR of the major isomer:  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.21 (d,  $J$  = 8.8 Hz, 2H), 7.53 (d,  $J$  = 8.8 Hz, 2H), 6.97 (s, 2H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -42.0 (s, 3F).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  135.6 (q,  $J$  = 1.3 Hz), 129.3 (s), 129.2 (q,  $J$  = 308.2 Hz), 127.2 (s), 124.3 (s), 123.9 (s), 118.1 (q,  $J$  = 3.4 Hz). IR(KBr):  $\nu$  2955, 2924, 2854, 1457, 1326, 1168, 1135, 1110, 1068  $\text{cm}^{-1}$ . GC-MS  $m/z$  249 ( $\text{M}^+$ ), 180 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_9\text{H}_6\text{F}_3\text{NO}_2\text{S}$ , 249.0071; found, 249.0073.

**(3-Fluorostyryl)(trifluoromethyl)sulfane (3k).** Obtained as a yellow oil in 77% yield (85 mg) with *E/Z* ratio 80:20.  $R_f$  (*n*-pentane/diethyl ether 10:1) = 0.82. NMR of the major isomer:  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.45–7.28 (m, 1H), 7.20–6.99 (m, 3H), 6.94 (d,  $J$  = 15.4 Hz, 1H), 6.77 (d,  $J$  = 15.4 Hz, 1H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -42.6 (s, 3F), -112.5 to -112.6 (m, 1F).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  164.3 (s), 161.8 (s), 138.9 (q,  $J$  = 1.2 Hz), 130.5 (s), 130.4 (s), 129.5 (q,  $J$  = 308.0 Hz), 122.7 (s), 115.9 (s), 113.4 (s). IR(KBr):  $\nu$  1715, 1683, 1652, 1614, 1580, 1557, 1557, 1540, 1506, 1488, 1446, 1223, 1111, 949, 775, 757, 680  $\text{cm}^{-1}$ . GC-MS  $m/z$  222 ( $\text{M}^+$ ), 153 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_9\text{H}_6\text{F}_4\text{S}$ , 222.0126; found, 222.0123.

**(4-Fluorostyryl)(trifluoromethyl)sulfane (3l).** Obtained as a yellow oil in 85% yield (94 mg) with *E/Z* ratio 93:7.  $R_f$  (*n*-pentane/diethyl ether 10:1) = 0.85. NMR of the major isomer:  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.58–7.28 (m, 2H), 7.05 (t,  $J$  = 8.6 Hz, 2H), 6.96 (d,  $J$  = 15.3 Hz, 1H), 6.65 (d,  $J$  = 15.3 Hz, 1H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -42.8 (s), -111.4 to -111.5 (m, 1F).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  164.4 (s), 162.0 (s), 140.1–140.0 (m), 129.6 (q,  $J$  = 307.8 Hz), 128.6 (s), 116.1 (s), 111.6–111.2 (m). IR(KBr):  $\nu$  2926, 2856, 1602, 1508, 1457, 1234, 1159, 1111, 956, 933, 839, 798, 756, 527, 515  $\text{cm}^{-1}$ . GC-MS  $m/z$  222 ( $\text{M}^+$ ), 153 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_9\text{H}_6\text{F}_4\text{S}$ , 222.0126; found, 222.0128.

**(3-Chlorostyryl)(trifluoromethyl)sulfane (3m).** Obtained as a yellow oil in 79% yield (94 mg) with an *E/Z* ratio of 70:30.  $R_f$  (*n*-pentane/diethyl ether 10:1) = 0.83. NMR of the major isomer:  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.40 (s, 1H), 7.37–7.21 (m, 3H), 6.93 (d,  $J$  = 15.2 Hz, 1H), 6.80 (d,  $J$  = 15.2 Hz, 1H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -42.5 (s, 3F).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  138.6 (q,  $J$  = 1.2 Hz), 136.8 (s), 134.9 (s), 130.1 (s), 129.4 (q,  $J$  = 307.9 Hz), 129.0 (s), 126.6 (s), 124.9 (d,  $J$  = 18.3 Hz), 113.8 (q,  $J$  = 3.2 Hz). IR(KBr):  $\nu$  1716, 1683, 1647, 1594, 1560, 1541, 1507, 1473, 1418, 1237, 1111, 950, 881, 792, 773, 756, 692, 679  $\text{cm}^{-1}$ . GC-MS  $m/z$  237 ( $\text{M}^+$ ), 168 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_9\text{H}_6\text{ClF}_3\text{S}$ , 237.9831; found, 237.9834.

**(4-Chlorostyryl)(trifluoromethyl)sulfane<sup>13</sup> (3n).** Obtained as a yellow oil in 83% yield (98 mg) with an *E/Z* ratio of 85:15.  $R_f$  (*n*-pentane/diethyl ether 10:1) = 0.82. NMR of the major isomer:  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.30–7.27 (m, 4H), 6.89 (d,  $J$  = 15.4 Hz, 1H), 6.70 (d,  $J$  = 15.4 Hz, 1H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -42.6 (s, 3F).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  139.3 (s), 135.0 (s), 133.6 (s), 129.6 (q,  $J$  = 307.6 Hz), 129.1 (s), 128.0 (s), 112.6 (q,  $J$  = 3.2 Hz). IR(KBr):  $\nu$  2956, 2925, 2854, 1541, 1508, 1490, 1457, 1260, 1163, 1111, 1094, 1013, 792, 794  $\text{cm}^{-1}$ . GC-MS  $m/z$  237 ( $\text{M}^+$ ), 168 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_9\text{H}_6\text{ClF}_3\text{S}$ , 237.9831; found, 237.9829.

**(E)-(2,6-Dichlorostyryl)(trifluoromethyl)sulfane (3o).** Obtained as a yellow oil in 62% yield (84 mg).  $R_f$  (*n*-pentane/diethyl ether 10:1) = 0.83.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.34 (d,  $J$  = 8.1 Hz, 2H), 7.22–7.09 (m, 1H), 7.05–6.88 (m, 2H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -42.1 (s, 3F).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  134.4 (s), 132.3 (q,  $J$  = 1.6 Hz), 129.4 (d,  $J$  = 308.1 Hz), 129.3 (s), 128.7 (s), 121.8 (q,  $J$  = 3.4 Hz). IR(KBr):  $\nu$  1647, 1557, 1541, 1508, 1487, 1428, 1163, 1109, 946, 774, 757, 708  $\text{cm}^{-1}$ . GC-MS  $m/z$  271 ( $\text{M}^+$ ), 202 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_9\text{H}_5\text{Cl}_2\text{F}_3\text{S}$ , 271.9441; found, 271.9445.

**(Z)-2-(2-(Trifluoromethylthio)vinyl)pyridine (3p).** Obtained as a yellow solid in 78% yield (79 mg).  $R_f$  (*n*-pentane/diethyl ether 10:1) = 0.68.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.65 (d,  $J$  = 4.5 Hz, 1H), 7.67 (td,  $J$  = 7.7, 1.7 Hz, 1H), 7.23 (d,  $J$  = 7.8 Hz, 1H), 7.17–7.10 (m, 1H), 6.73 (s, 2H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -46.6 (s, 3F).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  154.3 (s), 148.0 (s), 136.5 (s), 130.1 (q,  $J$  = 310.9 Hz), 124.6 (q,  $J$  = 1.2 Hz), 123.6 (s), 121.5 (s), 121.4 (s). IR(KBr):  $\nu$  1716, 1605, 1586, 1557, 1541, 1507, 1473, 1457, 1430, 1119, 997, 868, 795,

743, 688, 625, 541  $\text{cm}^{-1}$ . GC-MS  $m/z$  205 ( $\text{M}^+$ ), 136 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_8\text{H}_6\text{F}_3\text{NS}$ , 205.0173; found, 205.0175.

**(1-Phenylvinyl)(trifluoromethyl)sulfane (3q).** Obtained as a yellow oil in 73% yield (74 mg).  $R_f$  (*n*-pentane/diethyl ether 10:1) = 0.83.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.65–7.61 (m, 2H), 7.41–7.36 (m, 3H), 6.08 (s, 1H), 5.99 (s, 1H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -41.2 (s, 3F).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  135.5 (q,  $J$  = 1.9 Hz), 129.4 (q,  $J$  = 309.1 Hz), 129.1 (s), 128.5 (s), 127.4 (q,  $J$  = 1.8 Hz), 127.2 (s), 117.7 (s). IR(KBr):  $\nu$  1868, 1748, 1734, 1716, 1697, 1683, 1541, 1507, 1121, 771, 701  $\text{cm}^{-1}$ . GC-MS  $m/z$  204 ( $\text{M}^+$ ), 135 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_9\text{H}_7\text{F}_3\text{S}$ , 204.0221; found, 204.0222.

**(Trifluoromethyl)(1,2,2-triphenylvinyl)sulfane (3r).** Obtained as a white solid in 72% yield (128 mg).  $R_f$  (*n*-pentane/diethyl ether 10:1) = 0.79.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.45–7.29 (m, 7H), 7.22–7.15 (m, 3H), 7.11–7.01 (m, 3H), 6.98–6.88 (m, 2H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -39.5 (s, 3F).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  152.4 (q,  $J$  = 1.5 Hz), 142.6 (s), 141.5 (s), 139.4 (s), 130.5 (s), 130.4 (s), 129.6 (s), 129.4 (q,  $J$  = 311.8 Hz), 128.3 (s), 128.0 (s), 127.9 (s), 127.8 (s), 127.7 (s), 127.4 (s), 126.1 (q,  $J$  = 1.5 Hz). IR(KBr):  $\nu$  1868, 1791, 1772, 1748, 1716, 1692, 1683, 1558, 1541, 1507, 1489, 1444, 1126, 1108, 1075, 755, 745, 696  $\text{cm}^{-1}$ . GC-MS  $m/z$  356 ( $\text{M}^+$ ), 287 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_{21}\text{H}_{15}\text{F}_3\text{S}$ , 356.0847; found, 356.0843.

**(1H-Inden-2-yl)(trifluoromethyl)sulfane (3s).**  $^{19}\text{F}$  NMR analysis of the filtrate indicated that 3s was produced in 23% yield.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{CN}$ )  $\delta$  7.55–7.39 (m, 2H), 7.37–7.25 (m, 3H), 3.70 (s, 2H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CD}_3\text{CN}$ )  $\delta$  -42.6 (s).  $^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_3\text{CN}$ )  $\delta$  144.8 (s), 142.8 (s), 142.2 (q,  $J$  = 1.5 Hz), 129.6 (q,  $J$  = 307.9 Hz), 128.2 (q,  $J$  = 2.0 Hz), 126.9 (s), 123.8 (s), 121.8 (s), 117.3 (s), 43.8 (q,  $J$  = 1.0 Hz). HRMS (EI): calcd for  $\text{C}_{10}\text{H}_7\text{F}_3\text{S}$ , 216.0221; found, 216.0225.

**(1-Cyclohexylvinyl)(trifluoromethyl)sulfane (3t).** Obtained as a yellow oil in 70% yield (73 mg).  $R_f$  (*n*-pentane) = 0.85.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  5.66 (s, 1H), 5.64 (s, 1H), 2.18 (t,  $J$  = 11.4 Hz, 1H), 1.91 (dd,  $J$  = 58.1, 12.5 Hz, 4H), 1.43–1.09 (m, 6H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -41.0 (s).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  141.8 (q,  $J$  = 1.0 Hz), 129.9 (q,  $J$  = 307.7 Hz), 124.0 (q,  $J$  = 1.2 Hz), 46.2 (d,  $J$  = 0.9 Hz), 32.0 (s), 26.2 (s), 26.0 (s). IR(KBr):  $\nu$  2932, 2856, 2360, 1615, 1451, 1261, 1125, 1101, 929, 892, 806, 757, 606  $\text{cm}^{-1}$ . GC-MS  $m/z$  210 ( $\text{M}^+$ ), 141 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_9\text{H}_{13}\text{F}_3\text{S}$ , 210.0690; found, 210.0692.

**Oct-1-en-2-yl(trifluoromethyl)sulfane (3u).**  $^{19}\text{F}$  NMR analysis of the filtrate indicated that 3u was produced in 61% yield. This volatile compound was inevitably contaminated with a small amount of unidentified products.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  5.61 (s, 1H), 5.59 (s, 1H), 2.36 (t,  $J$  = 7.2 Hz, 2H), 1.37–1.22 (m, 8H), 0.89 (s, 3H).  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ )  $\delta$  -41.4 (s).  $^{13}\text{C}\{^1\text{H}\}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  142.9 (q,  $J$  = 1.2 Hz), 129.8 (q,  $J$  = 308.1 Hz), 125.7 (q,  $J$  = 1.3 Hz), 38.4 (s), 31.5 (s), 28.4 (s), 27.6 (s), 22.6 (s), 14.0 (s). GC-MS  $m/z$  212 ( $\text{M}^+$ ), 143 ( $\text{M}^+ - \text{CF}_3$ ). HRMS (EI): calcd for  $\text{C}_9\text{H}_{15}\text{F}_3\text{S}$ , 212.0847; found, 212.0852.

**Gram Scale Reactions for Synthesis of Styrenyl Trifluoromethyl Thioethers (3a).** Phenylvinyl bromide **1** (1.0 g, 5.5 mmol), [(bpy)Cu(SCF<sub>3</sub>)] **2** (2.1 g, 6.6 mmol, 1.2 equiv), KF (1.28 g, 22.0 mmol), and diglyme (20.0 mL) were added to a reaction tube with a Teflon screw cap equipped with a stir bar. The mixture was stirred at 100 °C for 24 h. The reaction mixture was filtered through a pad of Celite. The filtrate was added to water (60 mL) at 0 °C. The resulting mixture was extracted with Et<sub>2</sub>O (3 × 30 mL), and the combined organic layers were washed with water and then dried over MgSO<sub>4</sub>. The solvent was removed by rotary evaporation in an ice bath, and the resulting product was purified by column chromatography on silica gel with pentane/Et<sub>2</sub>O. **3a** was obtained as a yellow oil in 86% yield (0.96 g).

**Procedure for the Reaction of Phenylvinyl Bromide 1 with 2 in the Presence of 1.0 equiv of BHT.** Phenylvinyl bromide **1** (0.050 mmol), [(bpy)Cu(SCF<sub>3</sub>)] **2** (19.2 mg, 0.060 mmol, 1.2 equiv), KF (11.6 mg, 0.20 mmol), BHT (11.0 mg, 0.050 mmol, 1.0 equiv), and diglyme (1.0 mL) were added to a oven-dried 5 mL test tube with a Teflon screw cap. The tube was sealed, and the reaction solution was placed into a preheated 100 °C oil bath for 24 h. The tube was removed from the oil bath and cooled to room temperature, and then 10  $\mu\text{L}$  (trifluoro-

methoxy)benzene were added as an internal standard. The resulting mixture was filtered through a layer of Celite. The filtrate was analyzed by  $^{19}\text{F}$  NMR and GC-MS. The yield of the styrenyl trifluoromethyl thioethers (**3a**) was calculated to be 93%.

**Procedure for the Reaction of Phenylvinyl Bromide 1 with 2 in the Presence of 1.0 equiv of TEMPO.** Phenylvinyl bromide **1** (0.050 mmol), [(bpy)Cu(SCF<sub>3</sub>)] **2** (19.2 mg, 0.060 mmol, 1.2 equiv), KF (11.6 mg, 0.20 mmol), TEMPO (7.80 mg, 0.050 mmol, 1.0 equiv), and diglyme (1.0 mL) were added to a oven-dried 5 mL test tube with a Teflon screw cap. The tube was sealed, and the reaction solution was placed into a preheated 100 °C oil bath for 24 h. The tube was removed from the oil bath and cooled to room temperature, and then 10  $\mu\text{L}$  (trifluoromethoxy)benzene were added as an internal standard. The resulting mixture was filtered through a layer of Celite. The filtrate was analyzed by  $^{19}\text{F}$  NMR and GC-MS. The yield of the styrenyl trifluoromethyl thioethers (**3a**) was calculated to be 88%.

## ■ ASSOCIATED CONTENT

### ■ Supporting Information

Full NMR spectra of new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Authors

\*E-mail: [youngman@fzu.edu.cn](mailto:youngman@fzu.edu.cn) (H.Z.).

\*E-mail: [zweng@fzu.edu.cn](mailto:zweng@fzu.edu.cn) (Z.W.).

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

Financial support from National Natural Science Foundation of China (21372044), Research Fund for the Doctoral Program of Higher Education of China (No. 20123514110003), the SRF for ROCS, SEM, China (2012-1707), the Science Foundation of the Fujian Province, China (2013J01040), and Fuzhou University (022318, 022494) is gratefully acknowledged.

## ■ REFERENCES

- (1) Kirsch, P. *Modern fluoroorganic chemistry: synthesis, reactivity, applications*; Wiley-VCH: Weinheim, 2004.
- (2) Ojima, I., Ed. *Fluorine in Medicinal Chemistry and Chemical Biology*; John Wiley & Sons Ltd.: 2009.
- (3) Wang, J.; Sánchez-Roselló, M.; Aceña, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. *Chem. Rev.* **2014**, *114*, 2432.
- (4) Zhu, W.; Wang, J.; Wang, S.; Gu, Z.; Aceña, J. L.; Izawa, K.; Liu, H.; Soloshonok, V. A. *J. Fluorine Chem.* **2014**, *167*, 37.
- (5) Hansch, C.; Leo, A.; Taft, R. W. *Chem. Rev.* **1991**, *91*, 165.
- (6) Tlili, A.; Billard, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 6818.
- (7) Toulgoat, F.; Alazet, S.; Billard, T. *Eur. J. Org. Chem.* **2014**, 2415.
- (8) Chu, L.; Qing, F.-L. *Acc. Chem. Res.* **2014**, *47*, 1513.
- (9) Wang, H.; Vivic, D. A. *Synlett* **2013**, *24*, 1887.
- (10) He, W.; Weng, Z. *Prog. Chem.* **2013**, *25*, 1071.
- (11) Xu, X.-H.; Matsuzaki, K.; Shibata, N. *Chem. Rev.* **2015**, *115*, 731.
- (12) Chen, C.; Xie, Y.; Chu, L.; Wang, R.-W.; Zhang, X.; Qing, F.-L. *Angew. Chem., Int. Ed.* **2012**, *51*, 2492.
- (13) Zhang, C.-P.; Vivic, D. A. *Chem.—Asian J.* **2012**, *7*, 1756.
- (14) Pluta, R.; Nikolaienko, P.; Rueping, M. *Angew. Chem., Int. Ed.* **2014**, *53*, 1650.
- (15) Shao, X.; Liu, T.; Lu, L.; Shen, Q. *Org. Lett.* **2014**, *16*, 4738.
- (16) Xu, J.; Mu, X.; Chen, P.; Ye, J.; Liu, G. *Org. Lett.* **2014**, *16*, 3942.
- (17) Danoun, G.; Bayarmagnai, B.; Gruenberg, M. F.; Goossen, L. J. *Chem. Sci.* **2014**, *5*, 1312.
- (18) Tran, L. D.; Popov, I.; Daugulis, O. *J. Am. Chem. Soc.* **2012**, *134*, 18237.
- (19) Xu, C.; Ma, B.; Shen, Q. *Angew. Chem., Int. Ed.* **2014**, *53*, 9316.

- (20) Chen, C.; Xu, X.-H.; Yang, B.; Qing, F.-L. *Org. Lett.* **2014**, *16*, 3372.
- (21) Xu, C.; Shen, Q. *Org. Lett.* **2014**, *16*, 2046.
- (22) Hu, F.; Shao, X.; Zhu, D.; Lu, L.; Shen, Q. *Angew. Chem., Int. Ed.* **2014**, *53*, 6105.
- (23) Teverovskiy, G.; Surry, D. S.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2011**, *50*, 7312.
- (24) Zhang, C.-P.; Vivic, D. A. *J. Am. Chem. Soc.* **2012**, *134*, 183.
- (25) Shao, X.; Wang, X.; Yang, T.; Lu, L.; Shen, Q. *Angew. Chem., Int. Ed.* **2013**, *52*, 3457.
- (26) Rueping, M.; Tolstoluzhsky, N.; Nikolaienko, P. *Chem.—Eur. J.* **2013**, *19*, 14043.
- (27) Bates, C. G.; Saejueng, P.; Doherty, M. Q.; Venkataraman, D. *Org. Lett.* **2004**, *6*, 5005.
- (28) Zheng, Y.; Du, X.; Bao, W. *Tetrahedron Lett.* **2006**, *47*, 1217.
- (29) Kabir, M. S.; Van Linn, M. L.; Monte, A.; Cook, J. M. *Org. Lett.* **2008**, *10*, 3363.
- (30) Kabir, M. S.; Lorenz, M.; Van Linn, M. L.; Namjoshi, O. A.; Ara, S.; Cook, J. M. *J. Org. Chem.* **2010**, *75*, 3626.
- (31) Kao, H.-L.; Lee, C.-F. *Org. Lett.* **2011**, *13*, 5204.
- (32) Xu, R.-S.; Yue, L.; Pan, Y.-J. *Tetrahedron* **2012**, *68*, 5046.
- (33) Weng, Z.; He, W.; Chen, C.; Lee, R.; Tan, D.; Lai, Z.; Kong, D.; Yuan, Y.; Huang, K.-W. *Angew. Chem., Int. Ed.* **2013**, *52*, 1548.
- (34) Kong, D.; Jiang, Z.; Xin, S.; Bai, Z.; Yuan, Y.; Weng, Z. *Tetrahedron* **2013**, *69*, 6046.
- (35) Tan, J.; Zhang, G.; Ou, Y.; Yuan, Y.; Weng, Z. *Chin. J. Chem.* **2013**, *31*, 921.
- (36) Zhu, P.; He, X.; Chen, X.; You, Y.; Yuan, Y.; Weng, Z. *Tetrahedron* **2014**, *70*, 672.
- (37) Lin, Q.; Chen, L.; Huang, Y.; Rong, M.; Yuan, Y.; Weng, Z. *Org. Biomol. Chem.* **2014**, *12*, 5500.
- (38) Hou, C.; Lin, X.; Huang, Y.; Chen, Z.; Weng, Z. *Synthesis* **2015**, *47*, doi: 10.1055/s-0034-1379972.
- (39) Cohen, T.; Cristea, I. *J. Am. Chem. Soc.* **1976**, *98*, 748.
- (40) Telvekar, V. N.; Arote, N. D.; Herlekar, O. P. *Synlett* **2005**, 2495.
- (41) Fursule, R. A.; Patil, P. O.; Shewale, B. D.; Kosalge, S. B.; Deshmukh, P. K.; Patil, D. A. *Chem. Pharm. Bull.* **2009**, *57*, 1243.
- (42) Kaloko, J.; Hayford, A. *Org. Lett.* **2005**, *7*, 4305.
- (43) Cheung, L. L. W.; Yudin, A. K. *Org. Lett.* **2009**, *11*, 1281.
- (44) Adams, D. J.; Clark, J. H. *J. Org. Chem.* **2000**, *65*, 1456.