


Metal-Free Iodine-Catalyzed Direct Arylthiation of Substituted Anilines with Thiols

Daoshan Yang, Kelu Yan, Wei Wei, Jing Zhao, Mengqi Zhang, Xuguang Sheng, Guoqing Li, Shenglei Lu, and Hua Wang*

The Key Laboratory of Life-Organic Analysis and Key Laboratory of Pharmaceutical Intermediates and Analysis of Natural Medicine, School of Chemistry and Chemical Engineering, Qufu Normal University, Qufu 273165, Shandong, P. R. China

S Supporting Information

ABSTRACT: Iodine-catalyzed direct arylation of substituted anilines for the synthesis of various diaryl sulfides has been developed under metal- and solvent-free conditions. The present method uses readily available thiols as the arylation reagents, and environmentally friendly and inexpensive I_2 as the catalyst. Importantly, no base or ligand was necessary. Such a simple, efficient, and economical transformation provides an



attractive approach to various diaryl sulfides in good to excellent yields.


INTRODUCTION

Diaryl sulfides are ubiquitous in natural products, pharmacologically active compounds, and organic materials.¹ Therefore, the development of novel, practical, and highly efficient methods for the construction of diaryl sulfide motifs continues to stimulate a large number of research groups. The classical methods for the formation of diaryl sulfides traditionally involve two approaches. One is the transition-metal-catalyzed cross-couplings of disulfides or thiols with aryl halides, pseudo halides or arylboronic acids,² where transition-metal catalysts mainly focus on palladium,³ copper,⁴ iron,⁵ nickel,⁶ cobalt,⁷ and rhodium⁸ (Scheme 1, a). Another method typically relies on the coupling of sulfonyl chlorides with organozinc⁹ or Grignard reagents¹⁰ (Scheme 1, b). In recent years, seeking mild and selective methods for the direct functionalization of inert C–H bonds has received much more attention in organic

chemistry.¹¹ Undoubtedly, direct arylation of C–H bonds is more economical and practical.¹² However, research surveys of this synthetic strategy for the preparation of diaryl sulfides are rather limited,¹³ and in this respect, several examples using diaryl disulfides or 1-(substituted phenylthio)pyrrolidine-2,5-dione as the arylation reagents under Pd,¹⁴ Cu,¹⁵ and Fe¹⁶ catalytic conditions have been reported (Scheme 1, c). Very recently, Fu and co-workers demonstrated an elegant work for the synthesis of diaryl sulfides via iron- or boron-catalyzed C–H arylation of phenols at room temperature.¹⁷ Despite some great advantages, these reactions could encounter certain limitations, including readily unavailable precursors, harsh reaction conditions, and toxic metal salt catalysts. As a consequence, it remains a challenging, but very attractive, task to develop more efficient, economical, and practical synthesis methods for constructing the diaryl sulfide derivatives.

Recently, there are increasing demands in metal-free transformations owing to that trace-metal impurities might be avoided in the target molecules.¹⁸ As a consequence, the development of a metal-free protocol for the C–S bond formation via direct aryl C–H bond functionalization appears desirable and synthetically attractive. On the other hand, molecular iodine as an inexpensive, green, and efficient reagent has been extensively used in organic transformations.¹⁹ In 2012, Bolm and co-workers reported a facile procedure for the synthesis of thioethers via transition-metal-free thiolation of 1,3,4-oxadiazoles.²⁰ In 2013, Deng and co-workers described an iodine-promoted method for the synthesis of 2-arylsulfonylphenols under metal-free conditions.²¹ In 2014, Huang' group reported the direct use of arylsulfonyl hydrazide as the arylation reagents for iodine-mediated thiolation of substituted naphthols and naphthylamines.²² Herein, we report a


Scheme 1. Strategies for the Synthesis of Diaryl Sulfides



Received: March 10, 2015

Published: June 1, 2015

Table 1. I₂-Catalyzed Coupling Reaction of 4-Methoxybenzenethiol (1a) with Aniline (2a) Leading to 4-(4-Methoxyphenylthio)aniline (3a): Optimization of Conditions^a



entry	catalyst	oxidant	temp. [°C]	yield (3a) [%] ^b	yield (3a') [%] ^b
1	I ₂	DTBP	90	51	10
2	ⁿ Bu ₄ NI	DTBP	90	trace	trace
3	KI	DTBP	90	trace	trace
4	NIS	DTBP	90	42	7
5	I ₂	H ₂ O ₂	90	47	9
6	I ₂	TBHP	90	16	trace
7	I ₂	DTBP	100	59	12
8	I ₂	DTBP	110	66	14
9	I ₂	DTBP	120	75	11
10	I ₂	DTBP	120	68 ^c	14
11	I ₂	DTBP	120	56 ^d	15
12	I ₂	DTBP	130	75	10
13	I ₂	DTBP	110	65 ^e	11
14	none	DTBP	120	0	0


^aReaction conditions: 4-methoxybenzenethiol (1a) (0.5 mmol), aniline (2a) (1.0 mmol), catalyst (0.075 mmol), oxidant (2.5 mmol), reaction time (24 h) under nitrogen atmosphere. ^bIsolated yield. ^cIn the presence of DTBP (3 equiv). ^dIn the presence of DTBP (2 equiv). ^eIn the presence of DTBP (8 equiv).

metal-free molecular iodine-catalyzed approach to diaryl sulfides containing amino groups from readily available thiols and substituted anilines under solvent free-conditions (Scheme 1, d).

RESULTS AND DISCUSSION

First, 4-methoxybenzenethiol (1a) and aniline (2a) were selected as the model substrates to optimize the reaction conditions, including the catalysts, oxidants, reaction temperatures, and the amounts of oxidant under a nitrogen atmosphere. As shown in Table 1, four catalysts, such as I₂, ⁿBu₄NI, KI, and NIS, were investigated at 90 °C by using 5.0 equiv of DTBP (di-*tert*-butyl peroxide) as the oxidant, and I₂ provided 4-(4-methoxyphenylthio)aniline (3a) in 51% yield with the corresponding *ortho*-substituted product (3a') appearing in 10% yield (entries 1–4). We have compared different oxidants (compare entries 1, 5, and 6), and DTBP was superior to the other oxidants (entry 1). In addition, various temperatures were investigated (entries 7–9), and 120 °C was discovered to be more suitable for this transformation (entry 9). Elevating the reaction temperature did not enhance the yield (entry 12, Table 1). Besides that, different amounts of DTBP were used in the reactions (entries 9–11), and the 5 equiv of DTBP provided the highest yield (entry 9). Furthermore, 8 equiv of DTBP at 110 °C gave a 65% yield, indicating that the temperature is crucial for this reaction (entry 13, Table 1). Notably, no conversion was observed in the absence of iodine (entry 14). After the optimization process for catalysts, temperature, and oxidants, the various diaryl sulfide derivatives were synthesized under our standard conditions: 15 mol % I₂ as the catalyst, 5 equiv of DTBP as the oxidant under solvent-free conditions at 120 °C.

With the optimized conditions in hand, a variety of substituted thiols were employed for coupling with aniline derivatives (Tables 2–4). It was found that the moderate to excellent yields were obtained under the standard conditions. It should be pointed out that the arythiation of aniline derivatives could depend on the position of the substituents of anilines. As shown in Table 2, the arythiation mainly occurred at the *para* site of the amino group in anilines, due to the *ortho* steric hindrance effects (Table 2, 3a–3d'). When the amino group was substituted by other groups, such as methyl and amino, the arythiation took place only at the *para* site of substituted amino groups in anilines (Table 2, 3e–3p). Heteroaromatic thiol (i.e., thiophene-2-thiol) could be tolerated in this transformation, affording the desired products in 76% yield (Table 2, 3p). Although aromatic thiols exhibited high reactivity, unfortunately, aliphatic ones were poor substrates (Table 2, 3n and 3o). Importantly, phenylhydrazine was an excellent substrate in the transformation and afforded the only *para*-substituted products (Table 2, 3k–3m). Besides, the reaction could be performed only at the *para* site of amino when the amino *ortho* site was occupied by a substituent, as shown in Table 3. On the other hand, the reaction could occur at the amino *ortho* site when the amino *para* site was occupied (Table 4). For substituted thiols, the substrates containing electron-donating functional groups showed higher reactivity than the others (Table 2, 3a, 3b, 3k, and 3m; Table 3, 5a, 5b, 5f, and 5g; Table 4, 7c, 7d, 7e, and 7f). The catalytic efficiency was not affected by steric hindrance in thiols (Table 2, 3j and 3l; Table 3, 5d; Table 4, 7b, 7h, and 7k). Further, for substituted anilines, the substrates containing electron-donating groups showed higher reactivity than those with electron-withdrawing groups (Table 3, 5a, 5e, 5c, and 5h; Table 4, 7a, 7g, 7e, and 7m). Notably, when naphthalen-2-amine was used as the substrate, the arythiation reaction occurred on the *ortho*

Table 2. I₂-Catalyzed Synthesis of Diaryl Sulfides from Phenylthiols (1) with Anilines (2)^{a,b,c}

^aReaction conditions: substituted thiols (0.5 mmol), anilines (1.0 mmol), I₂ (0.075 mmol), DTBP (2.5 mmol). ^bIsolated yield. ^cReaction time (24 h).

α -carbon of NH₂ due to the highest electron density of α -carbon (Table 4, 7o–7r). In addition, the I₂-catalyzed reactions could tolerate some functional groups such as amino groups (Tables 2–4), hydrazine (products 3k–3m), methyl ether (products 3m, 5f, and 7k), C–Cl bond (products 3g, 5g, and 7g), and C–Br bond (products 3h, 5h, and 7f).


The arythiation reactions were investigated using different amounts of anilines. The results are shown in Scheme 2. Under the standard conditions, the control experiments with 4-chloroaniline **6b** alone gave the messy yields detected by TLC, indicating that 4-chloroaniline **6b** could be oxidized into other substances in this transformation [eq 1, Scheme 2]. Furthermore, different amounts of **6b** ranging from 0.5 to 3.0 mmol were applied in the reactions. The obtained yields increased as the **6b** amounts increased up to 1.5 mmol to reach the maximum [eq 2, Scheme 2]. 1.0 mmol of **6b** gave a slightly lower yield than 1.5 mmol of **6b**; however, it is economical and of practical application in the arythiation. Notably, the yields could decrease dramatically when the amount of **6b** was changed from 1.0 to 0.5 mmol. It is thereby thought that the aniline could be oxidized before it could react with the thiols under the given conditions. As a result, 2 equiv of aniline

should be sufficient for this reaction, whereas the excess of the aniline might make up for the loss.

Further, we explored the synthetic applicability of the present method. The gram-scale reaction was performed between **1f** and **6b**, and the reaction afforded **7c** in 89% yield (Scheme 3). Therefore, this simple protocol could be served as an efficient and practical method for the synthesis of various diaryl sulfide derivatives containing amino groups.

Inspired by these excellent results above, we applied this I₂-catalyzed method to construct some potential medicinal agents. As shown in Scheme 4, coupling of 2,4-dichlorobenzenethiol (**1r**) with 3-chloroaniline (**8b**) provided **9a** in 92% yield under the standard conditions, which was used as inhibitors of LFA-1/CAM-1,²³ suggesting that the developed method would be applied in the medical industry.

In order to explore the possible mechanism of I₂-catalyzed direct C–H arythiation of substituted anilines, several control experiments were performed, as shown in Scheme 5. When 4-methylbenzenethiol (**1f**) was added independently under the standard conditions, 1,2-di-*p*-tolylidysulfane (**10**) was obtained in 97% yield. Furthermore, treatment of 1,2-di-*p*-tolylidysulfane (**10**) with 4-chloroaniline (**6b**) under the standard conditions

Table 3. I₂-Catalyzed Synthesis of Diaryl Sulfides from Coupling Phenylthiols (1) with *Ortho*-Substituted Anilines (4)^{a,b,c}

^aReaction conditions: substituted thiols (0.5 mmol), *ortho*-substituted anilines (1.0 mmol), I₂ (0.075 mmol), DTBP (2.5 mmol). ^bIsolated yield. ^cReaction time (24 h).

led to 7c in 90% yield, indicating that 1,2-diphenyldisulfane might be intermediates for this transformation. Additionally, the reaction of 1,2-di-*p*-tolylidysulfane (10) with 4-chloroaniline (6b) was tested without iodine. As expected, no conversion was observed. To investigate the mechanism further, the reaction of 1,2-di-*p*-tolylidysulfane (10) with 4-chloroaniline (6b) was tested in the presence of 1 equiv of 2,2,6,6-tetramethylpiperidinyl-1-oxy (TEMPO), a classical radical inhibitor, the C–H arylation reaction was completely suppressed, indicating that the present reaction might involve a radical process.

On the basis of these preliminary results above and together with previous reports in the literature,²⁴ a possible mechanism for I₂-catalyzed synthesis of diaryl sulfide derivatives was proposed in Scheme 6. Initially, 4-methylbenzenethiol (1f) was oxidized to 1,2-di-*p*-tolylidysulfane (10) under the standard conditions. The reaction of 10 with I₂ formed 2 equiv of an electrophilic species *p*-MePhSI (I),²¹ which attacked 4-chloroaniline (6b) to yield II. Then, the subsequent loss of HI resulted in the desirable product (7c). Finally, treatment of *t*-BuO• radical with HI led to the catalyst I₂. Further investigations on the more detailed mechanism are ongoing in our laboratory.

In conclusion, a novel and efficient method has been developed for the synthesis of diaryl sulfide derivatives containing amino groups via the I₂-catalyzed direct arylation of substituted anilines under solvent-free conditions. A series of diaryl sulfide derivatives could be efficiently obtained in moderate to excellent yields. This method can enjoy the following advantages: (a) commercially available and nontoxic molecular iodine as the catalyst; (b) solvent-free conditions; (c)


no addition of any ligand, base, or additive; and (d) easy workup procedure. The developed synthesis approach would extend the scope of synthetic methods for diverse diaryl sulfides in the academic and industrial fields. Further investigation on the practical application of this method is in progress.

EXPERIMENTAL SECTION

General. All commercially available reagent grade chemicals were purchased from chemical suppliers and used as received without further purification. ¹H NMR and ¹³C NMR spectra were recorded in CDCl₃ on a 400 MHz spectrometer with TMS as internal standard (400 MHz ¹H, 100 MHz ¹³C) at room temperature, the chemical shifts (δ) were expressed in ppm, and *J* values were given in Hz. The following abbreviations are used to indicate the multiplicity: singlet (s), doublet (d), triplet (t), quartet (q) and multiplet (m). Column chromatography was performed on silica gel (200–300 mesh). Mass analyses and HRMS were obtained by ESI on a TOF mass analyzer.


General Experimental Procedures. A 25 mL Schlenk tube equipped with a magnetic stirring bar was charged with I₂ (19 mg, 0.075 mmol), substituted various thiols (1) (0.5 mmol), and anilines or phenylhydrazine (1.0 mmol). The tube was evacuated twice and backfilled with nitrogen, and DTBP (2.5 mmol) was added into the tube. The tube was sealed, and then the mixture was allowed to stir under a nitrogen atmosphere at 120 °C for 24 h. After completion of the reaction, the resulting solution was cooled down to room temperature, and the solvent was removed with the aid of a rotary evaporator. The desired product was obtained by column chromatography on silica gel using petroleum ether/ethyl acetate as eluent.

4-(4-Methoxyphenylthio)aniline (3a).²⁵ Compound 3a was obtained in 75% yield (87 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (50:1). ¹H NMR (CDCl₃, 400 MHz, ppm) δ 7.25 (d, 2H, *J* = 2.4 Hz), 7.23 (d, 2H, *J* = 2.4 Hz), 6.83


Table 4. I₂-Catalyzed Synthesis of Diaryl Sulfides from Coupling Phenylthiols (1) with *Para*-Substituted Anilines (6)^{a,b,c}

^aReaction conditions: substituted thiols (0.5 mmol), *para*-substituted anilines (1.0 mmol), I₂ (0.075 mmol), DTBP (2.5 mmol). ^bIsolated yield. ^cReaction time (24 h).


Scheme 2. Investigation of the Amount of Anilines in the Present Method



Scheme 3. Synthesis of 7c on Gram Scale




Scheme 4. Synthesis of 3-Chloro-4-(2,4-dichlorophenylthio)aniline (9a) under the Standard Conditions




(d, 2H, *J* = 8.0 Hz), 6.65 (d, 2H, *J* = 8.0 Hz), 3.80 (s, 3H), 3.75 (s, br, 2H). ¹³C NMR (CDCl₃, 100 MHz, ppm) δ 158.5, 146.2, 133.9, 131.5, 128.9, 123.6, 115.8, 114.7, 55.4; HRMS *m/z* calcd. for C₁₃H₁₃NOS [M + H]⁺: 232.0796, found: 232.0785.

2-(4-Methoxyphenylthio)aniline (3a').²⁶ Compound 3a' was obtained in 11% yield (13 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (50:1). ¹H NMR (CDCl₃, 400 MHz, ppm) δ 7.44 (d, 1H, *J* = 8.0 Hz), 7.21 (t, 1H, *J* = 8.0 Hz), 7.16

Scheme 5. Control Experiments



Scheme 6. A Proposed Mechanism for the Direct Transformation



(d, 2H, $J = 8.0$ Hz), 6.83 (d, 2H, $J = 8.0$ Hz), 6.79–6.74 (m, 2H), 4.29 (s, br, 2H), 3.79 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 158.4, 148.1, 136.3, 130.4, 129.7, 126.9, 118.7, 116.8, 115.4, 114.8, 55.4; HRMS m/z calcd. for C₁₃H₁₃NOS [M + H]⁺: 232.0796, found: 232.0785.

4-(4-Chlorophenylthio)aniline (3b).²⁷ Compound 3b was obtained in 65% yield (77 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (50:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.32 (d, 2H, $J = 8.0$ Hz), 7.20 (d, 2H, $J = 8.0$ Hz), 7.06 (d, 2H, $J = 8.0$ Hz), 6.70 (d, 2H, $J = 8.0$ Hz), 3.87 (s, br, 2H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 158.5, 146.2, 133.9, 131.5, 128.9, 123.6, 115.8, 114.7; HRMS m/z calcd. for C₁₂H₁₀ClNS [M + H]⁺: 236.0301, found: 236.0312.

2-(4-Chlorophenylthio)aniline (3b').²⁶ Compound 3b' was obtained in 9% yield (11 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (50:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.47 (d, 2H, $J = 8.0$ Hz), 7.28 (t, 1H, $J = 8.0$ Hz), 7.21 (d, 2H, $J = 8.0$ Hz), 7.02 (d, 2H, $J = 8.0$ Hz), 6.83–6.77 (m, 2H), 4.32 (s, br, 2H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 148.8, 137.5, 135.5, 131.5, 131.3, 129.1, 127.7, 118.9, 115.5, 113.8; HRMS m/z calcd. for C₁₂H₁₀ClNS [M + H]⁺: 236.0301, found: 236.0312.

4-(3-Methoxyphenylthio)aniline (3c).²⁸ Compound 3c was obtained in 71% yield (82 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (50:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.35 (d, 2H, $J = 8.0$ Hz), 7.15 (t, 1H, $J = 8.0$ Hz), 6.74–6.66 (m, 5H), 3.88 (s, br, 2H), 3.75 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 159.9, 147.2, 141.3, 136.4, 129.6, 119.9, 119.4, 115.9, 112.5, 110.8, 55.2; HRMS m/z calcd. for C₁₃H₁₃NOS [M + H]⁺: 232.0796, found: 232.0790.

2-(3-Methoxyphenylthio)aniline (3c').²⁹ Compound 3c' was obtained in 7% yield (8 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (50:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.43 (d, 1H, $J = 8.0$ Hz), 7.21 (t, 1H, $J = 8.0$ Hz), 7.16 (d, 2H, $J = 8.0$ Hz), 6.83 (d, 2H, $J = 8.0$ Hz), 6.79–6.74 (m, 2H), 4.29 (s, br, 2H), 3.79 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 158.4, 148.1, 136.3, 130.4, 129.7, 126.9, 118.7, 116.8, 115.4, 114.8, 55.4; HRMS m/z calcd. for C₁₃H₁₃NOS [M + H]⁺: 232.0796, found: 232.0790.

4-(Naphthalen-2-ylthio)aniline (3d).³⁰ Compound 3d was obtained in 68% yield (86 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (50:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 8.45 (d, 1H, $J = 8.0$ Hz), 7.90 (d, 1H, $J = 8.0$ Hz), 7.43 (d, 1H, $J = 8.0$ Hz), 7.62–7.54 (m, 2H), 7.39–7.34 (m, 3H), 7.27 (d, 1H, $J = 8.0$ Hz), 6.69 (d, 2H, $J = 8.0$ Hz), 3.76 (s, br, 2H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 146.8, 136.3, 135.3, 133.9, 131.7, 128.5, 126.7, 126.6, 126.4, 126.3, 125.8, 124.7, 120.8, 116.1; HRMS m/z calcd. for C₁₆H₁₃NS [M + H]⁺: 252.0847, found: 252.0846.

2-(Naphthalen-2-ylthio)aniline (3d').²⁶ Compound 3d' was obtained in 6% yield (7 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (50:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 8.38 (d, 1H, $J = 8.0$ Hz), 7.89 (d, 1H, $J = 8.0$ Hz), 7.68 (d, 1H, $J = 8.0$ Hz), 7.63–7.49 (m, 2H), 7.50 (d, 1H, $J = 8.0$ Hz), 7.33–7.28 (m, 2H), 6.99 (d, 2H, $J = 8.0$ Hz), 6.86–6.80 (m, 2H), 4.30 (s, br, 2H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 148.7, 137.3, 133.9, 133.6, 131.1, 131.0, 128.6, 126.3, 126.2, 126.1, 125.9, 124.0, 123.7, 119.0, 115.5, 114.1; HRMS m/z calcd. for C₁₆H₁₃NS [M + H]⁺: 252.0847, found: 252.0846.

4-(4-Methoxyphenylthio)-N-methylaniline (3e). Compound 3e was obtained in 73% yield (90 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.32 (d, 2H, $J = 8.0$ Hz), 7.24 (d, 2H, $J = 8.0$ Hz), 6.84 (d, 2H, $J = 8.0$ Hz), 6.59 (d, 2H, $J = 8.0$ Hz), 3.87 (s, br, 1H), 3.80 (s, 3H), 2.85 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 148.7, 137.3, 133.9, 133.6, 131.1, 131.0, 128.6, 126.3, 126.2, 126.0, 125.9, 124.0, 123.7, 119.0, 115.5, 114.1; HRMS m/z calcd. for C₁₄H₁₅NOS [M + H]⁺: 246.0953, found: 246.0961.

N-Methyl-4-(p-tolylthio)aniline (3f). Compound 3f was obtained in 69% yield (79 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.36 (d, 2H, $J = 8.0$ Hz), 7.08 (d, 2H, $J = 8.0$ Hz), 7.06 (d, 2H, $J = 8.0$ Hz), 6.60 (d, 2H, $J = 8.0$ Hz), 3.91 (s, br, 1H), 2.88 (s, 3H), 2.31 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 149.5, 136.2, 135.7, 135.1, 129.6, 127.8, 119.5, 113.1, 30.6, 20.9; HRMS m/z calcd. for C₁₄H₁₅NS [M + H]⁺: 230.1003, found: 230.1011.

4-(4-Chlorophenylthio)-N-methylaniline (3g). Compound 3g was obtained in 61% yield (76 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.37 (d, 2H, $J = 8.0$ Hz), 7.19 (d, 2H, $J = 8.0$ Hz), 7.05 (d, 2H, $J = 8.0$ Hz), 6.63 (d, 2H, $J = 8.0$ Hz), 3.96 (s, br, 1H), 2.89 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 150.0, 139.1, 136.5, 130.8, 128.8, 128.0, 117.3, 113.2, 30.5; HRMS m/z calcd. for C₁₃H₁₂ClNS [M + H]⁺: 250.0457, found: 250.0449.

4-(4-Bromophenylthio)-N-methylaniline (3h). Compound 3h was obtained in 64% yield (94 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.37 (d, 2H, $J = 8.0$ Hz), 7.34 (d, 2H, $J = 8.0$ Hz), 6.99 (d, 2H, $J = 8.0$ Hz), 6.63 (d, 2H, $J = 8.0$ Hz), 3.98 (s, br, 1H), 2.88 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 150.0, 139.9, 136.6, 131.7, 128.2, 118.5, 117.4, 113.2, 30.5; HRMS m/z calcd. for C₁₃H₁₂BrNS [M + H]⁺: 293.9952, 295.9932, found: 293.9943, 295.9921.

N-Methyl-4-(naphthalen-1-ylthio)aniline (**3i**). Compound **3i** was obtained in 78% yield (104 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 8.43 (d, 1H, $J = 8.0$ Hz), 7.88 (d, 1H, $J = 8.0$ Hz), 7.69 (d, 1H, $J = 8.0$ Hz), 7.61–7.55 (d, 2H, $J = 8.0$ Hz), 7.40 (d, 2H, $J = 8.0$ Hz), 7.34 (t, 1H, $J = 8.0$ Hz), 7.18 (d, 1H, $J = 8.0$ Hz), 6.64 (d, 1H, $J = 8.0$ Hz), 3.90 (s, br, 1H), 2.87 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 149.6, 137.0, 135.8, 133.8, 131.4, 128.5, 126.3, 126.2, 126.1, 125.8, 125.7, 124.5, 118.5, 113.3, 30.6; HRMS m/z calcd. for $\text{C}_{17}\text{H}_{15}\text{NS}$ [$\text{M} + \text{H}$] $^+$: 266.1003, found: 266.1012.

N,N-Dimethyl-4-(*o*-tolylthio)aniline (**3j**). Compound **3j** was obtained in 47% yield (57 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.38 (d, 2H, $J = 8.0$ Hz), 7.17 (d, 1H, $J = 8.0$ Hz), 7.06–7.04 (m, 2H), 6.88–6.85 (m, 1H), 6.75 (d, 2H, $J = 8.0$ Hz), 3.08 (s, 6H), 2.42 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 150.5, 139.2, 136.0, 135.4, 129.9, 126.9, 126.3, 125.0, 117.4, 113.1, 40.4, 20.1; HRMS m/z calcd. for $\text{C}_{15}\text{H}_{17}\text{NS}$ [$\text{M} + \text{H}$] $^+$: 244.1160, found: 244.1151.

4-(4-Chlorophenylthio)phenylhydrazine (**3k**). Compound **3k** was obtained in 52% yield (65 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.32 (d, 2H, $J = 8.0$ Hz), 7.20 (d, 2H, $J = 8.0$ Hz), 7.06 (d, 2H, $J = 8.0$ Hz), 6.70 (d, 2H, $J = 8.0$ Hz), 3.08 (s, br, 2H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 147.4, 138.5, 136.3, 131.0, 128.9, 128.4, 119.8, 115.9. HRMS m/z calcd. for $\text{C}_{12}\text{H}_{11}\text{ClN}_2\text{S}$ [$\text{M} + \text{H}$] $^+$: 251.0410, found: 251.0411.

4-(*m*-Tolylthio)phenylhydrazine (**3l**). Compound **3l** was obtained in 47% yield (54 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.36 (d, 2H, $J = 8.0$ Hz), 7.18 (t, 1H, $J = 8.0$ Hz), 7.02 (s, 1H), 6.97 (d, 2H, $J = 8.0$ Hz), 6.70 (d, 2H, $J = 8.0$ Hz), 3.85 (s, br, 2H), 2.31 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 147.0, 139.4, 138.7, 136.1, 128.8, 128.0, 126.3, 124.5, 120.6, 115.9; HRMS m/z calcd. for $\text{C}_{13}\text{H}_{14}\text{N}_2\text{S}$ [$\text{M} + \text{H}$] $^+$: 231.0956, found: 231.0967.

4-(4-Methoxyphenylthio)phenylhydrazine (**3m**). Compound **3m** was obtained in 62% yield (77 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.25–7.23 (m, 4H), 6.84 (d, 2H, $J = 8.0$ Hz), 6.65 (d, 2H, $J = 8.0$ Hz), 3.80 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 158.5, 146.2, 134.0, 131.5, 128.8, 123.5, 115.8, 114.6, 55.4; HRMS m/z calcd. for $\text{C}_{13}\text{H}_{14}\text{N}_2\text{OS}$ [$\text{M} + \text{H}$] $^+$: 247.0905, found: 247.0911.

4-(Butylthio)aniline (**3n**).²⁵ Compound **3n** was obtained in 13% yield (12 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.26 (d, 2H, $J = 8.0$ Hz), 6.64 (d, 2H, $J = 8.0$ Hz), 3.74 (s, br, 2H), 2.79 (t, 2H, $J = 8.0$ Hz), 1.57 (dt, 2H, $J = 8.0$ Hz), 1.42 (dt, 2H, $J = 8.0$ Hz), 0.91 (t, 3H, $J = 8.0$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 145.7, 133.7, 123.9, 115.6, 36.1, 31.5, 21.8, 13.7; HRMS m/z calcd. for $\text{C}_{10}\text{H}_{13}\text{NS}$ [$\text{M} + \text{H}$] $^+$: 182.1003, found: 182.1012.

4-(Butylthio)-*N*-methylaniline (**3o**). Compound **3o** was obtained in 22% yield (22 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.30 (d, 2H, $J = 8.0$ Hz), 6.58 (d, 2H, $J = 8.0$ Hz), 3.78 (s, br, 1H), 2.85 (s, 3H), 2.79 (t, 2H, $J = 8.0$ Hz), 1.57 (dt, 2H, $J = 8.0$ Hz), 1.42 (dt, 2H, $J = 8.0$ Hz), 0.93 (t, 3H, $J = 8.0$ Hz). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 148.7, 134.1, 122.0, 112.4, 36.5, 31.6, 30.7, 21.9, 13.7; HRMS m/z calcd. for $\text{C}_{11}\text{H}_{17}\text{NS}$ [$\text{M} + \text{H}$] $^+$: 196.1160, found: 196.1147.

4-(Thiophen-2-ylthio)aniline (**3p**). Compound **3p** was obtained in 79% yield (13 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.34 (d, 2H, $J = 4.0$ Hz), 7.24 (d, 2H, $J = 8.0$ Hz), 7.19 (d, 1H, $J = 4.0$ Hz), 7.00 (dd, 1H, $J = 4.0$ Hz), 6.63 (d, 2H, $J = 8.0$ Hz), 3.74 (s, br, 2H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 148.7, 134.1, 122.0, 112.4, 36.5, 31.6, 30.7, 21.9, 13.7; HRMS m/z calcd. for $\text{C}_{10}\text{H}_9\text{NS}_2$ [$\text{M} + \text{H}$] $^+$: 208.0255, found: 208.0261.

2-Methyl-4-(*p*-tolylthio)aniline (**5a**). Compound **5a** was obtained in 99% yield (114 mg) according to the general procedure. Eluent

petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.25 (s, 1H), 7.22 (d, 1H, $J = 8.0$ Hz), 7.12 (d, 2H, $J = 8.0$ Hz), 7.08 (d, 2H, $J = 8.0$ Hz), 6.68 (d, 1H, $J = 8.0$ Hz), 3.75 (s, br, 2H), 2.35 (s, 3H), 2.17 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 145.1, 136.4, 135.9, 135.3, 133.2, 129.7, 128.1, 123.3, 121.2, 115.6, 21.0, 17.3; HRMS m/z calcd. for $\text{C}_{14}\text{H}_{15}\text{NS}$ [$\text{M} + \text{H}$] $^+$: 230.1003, found: 230.1017.

4-(4-Chlorophenylthio)-2-methylaniline (**5b**). Compound **5b** was obtained in 88% yield (110 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.25–7.19 (s, 1H), 7.07 (d, 2H, $J = 8.0$ Hz), 6.69 (d, 1H, $J = 8.0$ Hz), 3.80 (s, br, 2H), 2.18 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 145.7, 138.8, 137.1, 134.0, 130.9, 128.9, 128.3, 123.4, 119.4, 115.7, 17.3. HRMS m/z calcd. for $\text{C}_{13}\text{H}_{12}\text{ClNS}$ [$\text{M} + \text{H}$] $^+$: 250.0457, found: 250.0462.

4-(4-Bromophenylthio)-2-methylaniline (**5c**). Compound **5c** was obtained in 91% yield (134 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.34 (d, 2H, $J = 8.0$ Hz), 7.25 (s, 1H), 7.23 (d, 1H, $J = 8.0$ Hz), 7.00 (d, 2H, $J = 8.0$ Hz), 6.69 (d, 2H, $J = 8.0$ Hz), 3.81 (s, br, 2H), 2.18 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 145.8, 139.6, 137.2, 134.1, 131.8, 128.5, 123.5, 119.2, 118.7, 115.7, 17.3; HRMS m/z calcd. for $\text{C}_{13}\text{H}_{12}\text{BrNS}$ [$\text{M} + \text{H}$] $^+$: 293.9952, 295.9932, found: 293.9941, 295.9926.

2-Methyl-4-(*o*-tolylthio)aniline (**5d**). Compound **5d** was obtained in 86% yield (99 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.26 (s, 1H), 7.24–7.20 (m, 2H), 7.12–7.09 (m, 2H), 6.72 (d, 1H, $J = 8.0$ Hz), 3.76 (s, br, 2H), 2.47 (s, 3H), 2.20 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 145.3, 138.8, 136.8, 135.7, 133.7, 130.1, 127.3, 126.4, 125.3, 123.5, 119.8, 115.8, 20.3, 17.4. HRMS m/z calcd. for $\text{C}_{14}\text{H}_{15}\text{NS}$ [$\text{M} + \text{H}$] $^+$: 230.1003, found: 230.1015.

2-Chloro-4-(*p*-tolylthio)aniline (**5e**). Compound **5e** was obtained in 95% yield (119 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.42 (d, 1H, $J = 4.0$ Hz), 7.21 (dd, 1H, $J = 8.0$ Hz), 7.17 (d, 2H, $J = 8.0$ Hz), 7.11 (d, 2H, $J = 8.0$ Hz), 6.74 (d, 1H, $J = 8.0$ Hz), 4.16 (s, br, 2H), 2.35 (s, 3H), 2.20 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 143.0, 136.2, 134.4, 134.3, 133.2, 129.9, 129.1, 123.0, 119.5, 116.3, 21.1; HRMS m/z calcd. for $\text{C}_{13}\text{H}_{12}\text{ClNS}$ [$\text{M} + \text{H}$] $^+$: 250.0457, found: 250.0462.

2-Chloro-4-(4-methoxyphenylthio)aniline (**5f**). Compound **5f** was obtained in 86% yield (114 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.33 (s, 1H), 7.32 (d, 2H, $J = 8.0$ Hz), 7.13 (d, 1H, $J = 8.0$ Hz), 6.86 (d, 2H, $J = 8.0$ Hz), 6.70 (d, 1H, $J = 8.0$ Hz), 4.13 (s, br, 2H), 3.81 (s, 3H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 158.9, 142.5, 132.8, 132.4, 131.7, 127.6, 124.9, 119.5, 116.3, 114.8, 55.4; HRMS m/z calcd. for $\text{C}_{13}\text{H}_{12}\text{ClNOS}$ [$\text{M} + \text{H}$] $^+$: 266.0406, found: 266.0411.

2-Chloro-4-(4-chlorophenylthio)aniline (**5g**). Compound **5g** was obtained in 75% yield (101 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.44 (s, 1H), 7.23–7.20 (m, 3H), 7.10 (d, 2H, $J = 8.0$ Hz), 6.76 (d, 1H, $J = 8.0$ Hz), 4.26 (s, br, 2H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 143.7, 137.4, 135.2, 134.1, 131.6, 129.1, 129.0, 121.0, 119.5, 116.4; HRMS m/z calcd. for $\text{C}_{12}\text{H}_9\text{Cl}_2\text{NS}$ [$\text{M} + \text{H}$] $^+$: 269.9911, found: 269.9923.

4-(4-Bromophenylthio)-2-chloroaniline (**5h**). Compound **5h** was obtained in 78% yield (122 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.44 (s, 1H), 7.36 (d, 2H, $J = 8.0$ Hz), 7.22 (d, 2H, $J = 8.0$ Hz), 7.02 (d, 2H, $J = 8.0$ Hz), 6.77 (d, 2H, $J = 8.0$ Hz), 4.26 (s, br, 2H). ^{13}C NMR (CDCl_3 , 100 MHz, ppm) δ 143.8, 138.2, 135.4, 134.3, 132.0, 129.2, 120.7, 119.5, 119.4, 116.4; HRMS m/z calcd. for $\text{C}_{12}\text{H}_9\text{BrClNS}$ [$\text{M} + \text{H}$] $^+$: 313.9406, 315.9385, found: 313.9417, 315.9392.

2-Chloro-4-(*o*-tolylthio)aniline (**5i**). Compound **5i** was obtained in 79% yield (99 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl_3 , 400 MHz, ppm) δ 7.43 (s, 1H), 7.25–4.13 (m, 4H), 7.05 (d, 2H, $J = 8.0$ Hz),

6.76 (d, 2H, $J = 8.0$ Hz), 4.22 (s, br, 2H), 2.48 (s, br, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 143.2, 137.3, 136.7, 134.6, 133.5, 130.4, 128.6, 126.7, 126.2, 121.7, 119.7, 116.5, 20.4; HRMS m/z calcd. for C₁₃H₁₂CIN₃ [M + H]⁺: 250.0457, found: 250.0462.

2-Ethyl-4-(*p*-tolylthio)aniline (5j). Compound 5j was obtained in 99% yield (121 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.31 (s, 1H), 7.24 (d, 1H, $J = 8.0$ Hz), 7.15 (d, 2H, $J = 8.0$ Hz), 7.10 (d, 2H, $J = 8.0$ Hz), 6.70 (d, 1H, $J = 8.0$ Hz), 3.76 (s, br, 2H), 2.52 (dd, 2H, $J = 8.0$ Hz), 2.35 (s, 3H), 1.30 (t, 3H, $J = 8.0$ Hz). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 144.6, 136.8, 136.1, 135.3, 134.4, 133.1, 129.7, 129.0, 128.0, 116.1, 23.9, 21.0, 12.9; HRMS m/z calcd. for C₁₅H₁₇NS [M + H]⁺: 244.1160, found: 244.1154.

2-Ethyl-4-(4-methoxyphenylthio)aniline (5k). Compound 5k was obtained in 99% yield (128 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.26 (d, 2H, $J = 8.0$ Hz), 7.22 (s, 1H), 7.15 (d, 2H, $J = 8.0$ Hz), 6.85 (d, 2H, $J = 8.0$ Hz), 6.65 (d, 1H, $J = 8.0$ Hz), 3.81 (s, 3H), 3.74 (s, br, 2H), 2.50 (dd, 2H, $J = 8.0$ Hz), 1.26 (t, 3H, $J = 8.0$ Hz). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 158.4, 144.1, 133.1, 131.7, 131.1, 129.2, 128.9, 123.1, 116.1, 114.7, 55.4, 23.9, 12.9; HRMS m/z calcd. for C₁₅H₁₇NOS [M + H]⁺: 259.1031, found: 259.1022.

4-(4-Chlorophenylthio)-2-ethylaniline (5l). Compound 5l was obtained in 97% yield (128 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.30 (s, 1H), 7.26–7.21 (m, 3H), 7.09 (d, 2H, $J = 8.0$ Hz), 6.71 (d, 2H, $J = 8.0$ Hz), 3.84 (s, br, 2H), 2.53 (dd, 2H, $J = 8.0$ Hz), 1.29 (t, 3H, $J = 8.0$ Hz). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 145.2, 138.9, 136.8, 135.5, 135.2, 133.9, 128.9, 128.2, 117.4, 116.2, 23.9, 12.8; HRMS m/z calcd. for C₁₄H₁₄CIN₃ [M + H]⁺: 264.0614, found: 264.0603.

4-(4-Bromophenylthio)-2-ethylaniline (5m). Compound 5m was obtained in 97% yield (150 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.35 (d, 2H, $J = 8.0$ Hz), 7.29 (s, 1H), 7.24 (d, 1H, $J = 8.0$ Hz), 7.02 (d, 2H, $J = 8.0$ Hz), 6.71 (d, 1H, $J = 8.0$ Hz), 3.85 (s, br, 2H), 2.53 (dd, 2H, $J = 8.0$ Hz), 1.29 (t, 3H, $J = 8.0$ Hz). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 145.3, 139.7, 136.8, 135.2, 134.0, 131.8, 129.1, 128.4, 23.8, 12.8; HRMS m/z calcd. for C₁₄H₁₄BrNS [M + H]⁺: 308.0109, 310.0088, found: 309.0121, 310.0095.

4-(4-Chlorophenylthio)-2,6-dimethylaniline (5n). Compound 5n was obtained in 99% yield (130 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.22–7.19 (m, 4H), 7.09 (d, 2H, $J = 8.0$ Hz), 3.72 (s, br, 2H), 2.21 (s, 6H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 144.0, 139.1, 136.5, 135.1, 128.9, 128.2, 122.8, 118.5, 17.6; HRMS m/z calcd. for C₁₄H₁₄CIN₃ [M + H]⁺: 263.0535, found: 263.0528.

2,6-Dimethyl-4-(*p*-tolylthio)aniline (5o). Compound 5o was obtained in 99% yield (121 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.19 (s, 2H), 7.16–7.09 (m, 4H), 3.76 (s, br, 2H), 2.36 (s, 3H), 2.22 (s, 6H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 143.4, 136.2, 135.2, 134.4, 129.7, 127.9, 122.6, 120.2, 21.0, 17.6. HRMS m/z calcd. for C₁₅H₁₇NS [M + H]⁺: 244.1160, found: 244.1154.

4-Methyl-2-(*m*-tolylthio)aniline (7a). Compound 7a was obtained in 84% yield (97 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.35 (s, 1H), 7.16 (t, 1H, $J = 8.0$ Hz), 7.10 (d, 1H, $J = 8.0$ Hz), 7.01–6.79 (m, 2H), 6.93 (d, 1H, $J = 8.0$ Hz), 6.76 (d, 1H, $J = 8.0$ Hz), 4.17 (s, br, 2H), 2.33 (s, 3H), 2.31 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 146.5, 138.9, 137.5, 136.8, 131.9, 128.9, 128.1, 127.1, 126.4, 123.6, 115.5, 114.5, 21.5, 20.3; HRMS m/z calcd. for C₁₄H₁₅NS [M + H]⁺: 230.1003, found: 230.1017.

4-Methyl-2-(*o*-tolylthio)aniline (7b). Compound 7b was obtained in 81% yield (93 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.34 (s, 1H), 7.25 (d, 1H, $J = 8.0$ Hz), 7.16–7.12 (m, 3H), 6.81 (t, 2H, $J = 8.0$ Hz), 4.18 (s, br, 2H), 2.53 (s, br, 3H), 2.34 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 146.6, 137.5, 136.0, 135.2,

131.9, 130.3, 128.3, 126.7, 125.5, 125.2, 115.6, 114.0, 20.4, 20.2; HRMS m/z calcd. for C₁₄H₁₅NS [M + H]⁺: 230.1003, found: 230.1017.

4-Chloro-2-(*p*-tolylthio)aniline (7c). Compound 7c was obtained in 86% yield (108 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.42 (d, 1H, $J = 4.0$ Hz), 7.17 (d, 1H, $J = 8.0$ Hz), 7.11–7.06 (m, 4H), 6.72 (d, 1H, $J = 8.0$ Hz), 4.28 (s, br, 2H), 2.32 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 146.9, 136.2, 135.6, 131.8, 130.5, 130.0, 127.7, 122.5, 117.2, 116.3, 21.0; HRMS m/z calcd. for C₁₃H₁₂CIN₃ [M + H]⁺: 250.0457, found: 250.0462.

4-Chloro-2-(4-methoxyphenylthio)aniline (7d). Compound 7d was obtained in 72% yield (96 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.38 (d, 1H, $J = 4.0$ Hz), 7.21 (d, 2H, $J = 8.0$ Hz), 7.14 (d, 1H, $J = 8.0$ Hz), 6.86 (d, 2H, $J = 8.0$ Hz), 6.68 (d, 1H, $J = 8.0$ Hz), 4.29 (s, br, 2H), 3.80 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 155.8, 146.3, 134.6, 130.7, 130.0, 125.5, 122.6, 118.9, 116.3, 115.0, 55.4; HRMS m/z calcd. for C₁₃H₁₂CINOS [M + H]⁺: 266.0406, found: 266.0411.

4-Chloro-2-(4-chlorophenylthio)aniline (7e).²³ Compound 7e was obtained in 88% yield (118 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.45 (s, 1H), 7.24–7.20 (m, 3H), 7.05 (d, 2H, $J = 8.0$ Hz), 6.74 (d, 1H, $J = 8.0$ Hz), 4.31 (s, br, 2H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 147.3, 136.2, 134.4, 131.8, 131.3, 129.3, 128.2, 122.6, 116.4, 115.5; HRMS m/z calcd. for C₁₂H₉Cl₂NS [M + H]⁺: 269.9911, found: 269.9923.

2-(4-Bromophenylthio)-4-chloroaniline (7f). Compound 7f was obtained in 84% yield (132 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.46 (s, 1H), 7.37 (d, 2H, $J = 8.0$ Hz), 7.22 (d, 1H, $J = 8.0$ Hz), 6.98 (d, 2H, $J = 8.0$ Hz), 6.73 (d, 1H, $J = 8.0$ Hz), 4.31 (s, br, 2H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 147.3, 136.2, 135.1, 132.2, 131.3, 129.3, 128.4, 122.6, 119.7, 116.5. HRMS m/z calcd. for C₁₂H₉BrClNS [M + H]⁺: 313.9406, 315.9385, found: 313.9417, 315.9392.

4-Chloro-2-(*m*-tolylthio)aniline (7g). Compound 7g was obtained in 69% yield (86 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.48 (s, 1H), 7.22–7.16 (m, 2H), 7.01 (d, 2H, $J = 8.0$ Hz), 6.94 (d, 1H, $J = 8.0$ Hz), 6.74 (d, 1H, $J = 8.0$ Hz), 4.31 (s, br, 2H), 2.33 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 147.3, 139.1, 136.1, 135.4, 130.8, 129.1, 127.7, 127.0, 124.2, 122.5, 116.4, 116.3, 21.5; HRMS m/z calcd. for C₁₃H₁₂CIN₃ [M + H]⁺: 250.0457, found: 250.0462.

4-Chloro-2-(*o*-tolylthio)aniline (7h). Compound 7h was obtained in 62% yield (78 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.39 (s, 1H), 7.21 (d, 2H, $J = 8.0$ Hz), 7.14–7.07 (m, 2H), 6.82 (d, 1H, $J = 8.0$ Hz), 6.75 (d, 1H, $J = 8.0$ Hz), 4.27 (s, br, 2H), 2.45 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 147.1, 135.9, 135.7, 134.5, 130.6, 130.4, 126.7, 126.3, 125.9, 122.7, 116.3, 116.0, 20.1; HRMS m/z calcd. for C₁₃H₁₂CIN₃ [M + H]⁺: 250.0457, found: 250.0462.

4-Methoxy-2-(*p*-tolylthio)aniline (7i). Compound 7i was obtained in 64% yield (79 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.10–7.04 (m, 4H), 7.04 (s, 1H), 6.88 (d, 1H, $J = 8.0$ Hz), 6.77 (d, 1H, $J = 8.0$ Hz), 4.01 (s, br, 2H), 3.77 (s, 3H), 2.32 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 152.4, 142.4, 135.7, 132.7, 129.9, 127.4, 120.6, 117.7, 116.7, 116.4, 55.9, 21.0; HRMS m/z calcd. for C₁₄H₁₅NOS [M + H]⁺: 246.0953, found: 246.0961.

4-Methoxy-2-(*m*-tolylthio)aniline (7j). Compound 7j was obtained in 64% yield (79 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). ^1H NMR (CDCl₃, 400 MHz, ppm) δ 7.16 (t, 1H, $J = 8.0$ Hz), 7.06 (s, 1H), 6.98 (d, 2H, $J = 8.0$ Hz), 6.94–6.88 (m, 2H), 6.79 (d, 1H, $J = 8.0$ Hz), 3.99 (s, br, 2H), 3.78 (s, 3H), 2.31 (s, 3H). ^{13}C NMR (CDCl₃, 100 MHz, ppm) δ 152.3, 142.7, 138.9, 136.3, 129.0, 127.4, 126.6, 123.8, 120.9, 118.0, 116.7, 115.6,

55.9, 21.5. HRMS m/z calcd. for $C_{14}H_{15}NOS$ $[M + H]^+$: 246.0953, found: 246.0961.

4-Methoxy-2-(*o*-tolylthio)aniline (7k). Compound **7k** was obtained in 61% yield (75 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). 1H NMR ($CDCl_3$, 400 MHz, ppm) δ 7.23 (d, 1H, $J = 8.0$ Hz), 7.13–7.09 (m, 2H), 7.03 (s, 1H), 6.93 (d, 2H, $J = 8.0$ Hz), 6.85–6.80 (m, 2H), 3.97 (s, br, 2H), 3.70 (s, 3H), 2.49 (s, 3H). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm) δ 152.6, 142.8, 135.6, 135.4, 130.3, 126.7, 125.9, 125.6, 120.6, 117.9, 116.8, 115.2, 55.9, 20.2. HRMS m/z calcd. for $C_{14}H_{15}NOS$ $[M + H]^+$: 246.0953, found: 246.0961.

4-Methoxy-2-(4-methoxyphenylthio)aniline (7l). Compound **7l** was obtained in 57% yield (75 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). 1H NMR ($CDCl_3$, 400 MHz, ppm) δ 7.19 (d, 2H, $J = 8.0$ Hz), 6.98 (d, 1H, $J = 4.0$ Hz), 6.84 (d, 2H, $J = 8.0$ Hz), 6.82 (d, 1H, $J = 4.0$ Hz), 6.74 (d, 1H, $J = 8.0$ Hz), 6.85–6.80 (m, 2H), 3.95 (s, br, 2H), 3.79 (s, 3H), 3.75 (s, 3H). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm) δ 158.5, 152.4, 141.8, 130.2, 126.4, 119.9, 118.2, 116.9, 116.7, 114.9, 55.8, 55.4; HRMS m/z calcd. for $C_{14}H_{15}NO_2S$ $[M + H]^+$: 262.0902, found: 262.0911.

2-(4-Chlorophenylthio)-4-methoxyaniline (7m). Compound **7m** was obtained in 87% yield (116 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). 1H NMR ($CDCl_3$, 400 MHz, ppm) δ 7.21 (d, 2H, $J = 8.0$ Hz), 7.05–7.03 (m, 3H), 6.88 (d, 1H, $J = 4.0$ Hz), 6.78 (d, 1H, $J = 8.0$ Hz), 4.01 (s, br, 2H), 3.77 (s, 3H). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm) δ 152.4, 142.7, 135.2, 131.4, 129.2, 127.9, 120.9, 118.5, 116.8, 114.8, 55.9. HRMS m/z calcd. for $C_{13}H_{12}ClNOS$ $[M + H]^+$: 266.0406, found: 266.0411.

2-(4-Bromophenylthio)-4-methoxyaniline (7n). Compound **7n** was obtained in 85% yield (132 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). 1H NMR ($CDCl_3$, 400 MHz, ppm) δ 7.36 (d, 2H, $J = 8.0$ Hz), 7.05 (s, 1H), 6.98 (d, 2H, $J = 8.0$ Hz), 6.91 (d, 1H, $J = 8.0$ Hz), 6.78 (d, 1H, $J = 8.0$ Hz), 4.01 (s, br, 2H), 3.77 (s, 3H). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm) δ 152.4, 142.8, 136.0, 132.1, 128.1, 120.9, 119.2, 118.6, 116.9, 114.5, 55.9. HRMS m/z calcd. for $C_{13}H_{12}BrNOS$ $[M + H]^+$: 309.9901, 31.9881, found: 309.9912, 31.9891.

1-(4-Chlorophenylthio)naphthalen-2-amine (7o).²² Compound **7o** was obtained in 97% yield (139 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). 1H NMR ($CDCl_3$, 400 MHz, ppm) δ 8.27 (d, 1H, $J = 8.0$ Hz), 7.82–7.76 (m, 2H), 7.49 (t, 1H, $J = 8.0$ Hz), 7.32 (t, 1H, $J = 8.0$ Hz), 7.16 (d, 2H, $J = 8.0$ Hz), 7.08 (d, 1H, $J = 8.0$ Hz), 6.98 (d, 2H, $J = 8.0$ Hz), 4.76 (s, br, 2H). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm) δ 148.5, 136.5, 135.5, 132.1, 130.9, 129.1, 128.5, 128.4, 128.0, 127.1, 124.0, 122.8, 117.7, 104.0; HRMS m/z calcd. for $C_{16}H_{12}ClNS$ $[M + H]^+$: 286.0457, found: 286.0451.

1-(4-Bromophenylthio)naphthalen-2-amine (7p).²² Compound **7p** was obtained in 94% yield (155 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). 1H NMR ($CDCl_3$, 400 MHz, ppm) δ 8.27 (d, 1H, $J = 8.0$ Hz), 7.82–7.76 (m, 2H), 7.49 (t, 1H, $J = 8.0$ Hz), 7.32 (t, 1H, $J = 8.0$ Hz), 7.16 (d, 2H, $J = 8.0$ Hz), 7.08 (d, 1H, $J = 8.0$ Hz), 6.98 (d, 2H, $J = 8.0$ Hz), 4.76 (s, br, 2H). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm) δ 148.5, 136.5, 135.5, 132.1, 130.9, 129.1, 128.5, 128.4, 128.0, 127.1, 124.0, 122.8, 117.7, 104.0; HRMS m/z calcd. for $C_{16}H_{12}BrNS$ $[M + H]^+$: 329.9952, 331.9932, found: 329.9937, 331.9923.

1-(*p*-Tolylthio)naphthalen-2-amine (7q).²² Compound **7q** was obtained in 90% yield (119 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). 1H NMR ($CDCl_3$, 400 MHz, ppm) δ 8.40 (d, 1H, $J = 8.0$ Hz), 7.82–7.79 (m, 2H), 7.50 (t, 1H, $J = 8.0$ Hz), 7.34 (t, 1H, $J = 8.0$ Hz), 7.09–7.01 (m, 5H), 4.75 (s, br, 2H), 2.31 (s, 3H). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm) δ 148.5, 136.7, 135.0, 133.3, 131.8, 129.9, 128.5, 127.9, 126.1, 124.4, 123.1, 122.6, 117.8, 105.2; HRMS m/z calcd. for $C_{17}H_{15}NS$ $[M + H]^+$: 266.1003, found: 266.1012.

1-(4-Methoxyphenylthio)naphthalen-2-amine (7r).²² Compound **7r** was obtained in 89% yield (125 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). 1H NMR

($CDCl_3$, 400 MHz, ppm) δ 8.39 (d, 1H, $J = 8.0$ Hz), 7.78–7.75 (m, 2H), 7.50 (t, 1H, $J = 8.0$ Hz), 7.31 (t, 1H, $J = 8.0$ Hz), 7.06 (d, 3H, $J = 8.0$ Hz), 6.77 (d, 1H, $J = 8.0$ Hz), 4.73 (s, br, 2H), 3.74 (s, 3H). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm) δ 157.8, 148.3, 136.7, 131.6, 128.5, 128.4, 127.9, 127.8, 127.5, 124.4, 122.6, 117.7, 114.8, 106.2, 55.3. HRMS m/z calcd. for $C_{17}H_{15}NOS$ $[M + H]^+$: 282.0953, found: 282.0947.

3-Chloro-4-(2,4-dichlorophenylthio)aniline (9a).²³ Compound **9a** was obtained in 92% yield (140 mg) according to the general procedure. Eluent petroleum ether/ethyl acetate (20:1). 1H NMR ($CDCl_3$, 400 MHz, ppm) δ 7.39–7.37 (m, 2H), 7.05 (d, 1H, $J = 8.0$ Hz), 6.88 (s, 1H), 6.59 (t, 2H, $J = 8.0$ Hz), 4.03 (s, br, 2H). ^{13}C NMR ($CDCl_3$, 100 MHz, ppm) δ 149.4, 140.7, 138.8, 136.4, 131.3, 131.0, 129.2, 127.7, 127.4, 116.3, 116.1, 114.4. HRMS m/z calcd. for $C_{12}H_8Cl_3NS$ $[M + H]^+$: 303.9521, found: 303.9532.

■ ASSOCIATED CONTENT

📄 Supporting Information

1H and ^{13}C NMR spectra for all compounds. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.5b00540.

■ AUTHOR INFORMATION

✉ Corresponding Author

*E-mail: huawang_qfnu@126.com (H.W.).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The authors gratefully acknowledge the financial support from the National Natural Science Foundation of China (Nos. 21302110, 21302109, and 21375075), the Taishan Scholar Foundation of Shandong Province, the Project of Shandong Province Higher Educational Science and Technology Program (J13LD14), and the Natural Science Foundation of Shandong Province (ZR2013BQ017).

■ REFERENCES

- (a) Nakazawa, T.; Xu, J.; Nishikawa, T.; Oda, T.; Fujita, A.; Ukai, K.; Mangindaan, R. E. P.; Rotinsulu, H.; Kobayashi, H.; Namikoshi, M. *J. Nat. Prod.* **2007**, *70*, 439–442. (b) Nielsen, S. F.; Nielsen, E. O.; Olsen, G. M.; Liljefors, T.; Peters, D. *J. Med. Chem.* **2000**, *43*, 2217–2226. (c) Gangjee, A.; Zeng, Y. B.; Talreja, T.; McGuire, J. J.; Kisliuk, R. L.; Queener, S. F. *J. Med. Chem.* **2007**, *50*, 3046–3053. (d) Zhou, Y.; Liu, W.-J.; Ma, Y.; Wang, H.; Qi, L.; Cao, Y.; Wang, J.; Pei, J. *J. Am. Chem. Soc.* **2007**, *129*, 12386–12387. (e) Takimiya, K.; Shinamura, S.; Osaka, I.; Miyazaki, E. *Adv. Mater.* **2011**, *23*, 4347–4370. (f) Mori, T.; Nishimura, T.; Yamamoto, T.; Doi, I.; Miyazaki, E.; Osaka, I.; Takimiya, K. *J. Am. Chem. Soc.* **2013**, *135*, 13900–13913.
- (a) Kondo, T.; Mitsudo, T.-a. *Chem. Rev.* **2000**, *100*, 3205–3220. (b) Li, G. Y. *Angew. Chem., Int. Ed.* **2001**, *40*, 1513–1516.
- (a) Migita, T.; Shimizu, T.; Asami, Y.; Shiobara, J.; Kato, Y.; Kosugi, M. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1385–1398. (b) Mann, G.; Baranano, D.; Hartwig, J. F.; Rheingold, A. L.; Guzei, I. A. *J. Am. Chem. Soc.* **1998**, *120*, 9205–9219. (c) Baranano, D.; Hartwig, J. F. *J. Am. Chem. Soc.* **1995**, *117*, 2937–2938. (d) Jiang, Z.; She, J.; Lin, X. *Adv. Synth. Catal.* **2009**, *351*, 2558–2562.
- (a) Ma, D.; Cai, Q. *Acc. Chem. Res.* **2008**, *41*, 1450–1460. (b) Kwong, F. Y.; Buchwald, S. L. *Org. Lett.* **2002**, *4*, 3517–3520. (c) Chen, C. F.; Chen, Y. W.; Lin, C. H.; Lin, H. P.; Lee, C. F. *Chem. Commun.* **2010**, *46*, 282–284. (d) Bhadra, S.; Sreedhar, B.; Ranu, B. C. *Adv. Synth. Catal.* **2009**, *351*, 2369–2378.
- (a) Wu, W.-Y.; Wang, J.-C.; Tsai, F.-Y. *Green Chem.* **2009**, *11*, 326–329. (b) Correa, A.; Carril, M.; Bolm, C. *Angew. Chem., Int. Ed.* **2008**, *47*, 2880–2883.

- (6) (a) Zhang, Y.; Ngeow, K. C.; Ying, J. Y. *Org. Lett.* **2007**, *9*, 3495–3499. (b) Joyce, L. L.; Evindar, G.; Batey, R. A. *Chem. Commun.* **2004**, 446–447. (c) Prasad, D. J. C.; Naidu, A. B.; Sekar, G. *Tetrahedron Lett.* **2009**, *50*, 1411–1415. (d) Bates, C. G.; Saejueng, P.; Doherty, M. Q.; Venkataraman, D. *Org. Lett.* **2004**, *6*, 5005–5008. (e) Rout, L.; Sen, T. K.; Punniyamurthy, T. *Angew. Chem., Int. Ed.* **2007**, *46*, 5583–5586. (f) Ramana, T.; Saha, P.; Das, M.; Punniyamurthy, T. *Org. Lett.* **2010**, *12*, 84–87.
- (7) Wong, Y. C.; Jayanth, T. T.; Cheng, C. H. *Org. Lett.* **2006**, *8*, 5613–5616.
- (8) Arisawa, M.; Suzuki, T.; Ishikawa, T.; Yamaguchi, M. *J. Am. Chem. Soc.* **2008**, *130*, 12214–12215.
- (9) Yonova, I. M.; Osborne, C. A.; Morrisette, N. S.; Jarvo, E. R. *J. Org. Chem.* **2014**, *79*, 1947–1953.
- (10) (a) Stoll, H.; Krasovskiy, A.; Knochel, P. *Angew. Chem., Int. Ed.* **2006**, *45*, 606–609. (b) Cheng, J.-H.; Ramesh, C.; Kao, H.-L.; Wang, Y.-J.; Chan, C.-C.; Lee, C.-F. *J. Org. Chem.* **2012**, *77*, 10369–10374.
- (11) For recent reviews, see: (a) Sun, C.-L.; Li, B.-J.; Shi, Z.-J. *Chem. Rev.* **2011**, *111*, 1293–1314. (b) Lewis, J. C.; Bergman, R. G.; Ellman, J. A. *Acc. Chem. Res.* **2008**, *41*, 1013–1025. (c) Engle, K. M.; Mei, T.-S.; Wasa, M.; Yu, J.-Q. *Acc. Chem. Res.* **2012**, *45*, 788–802. (d) Mkhaliid, I. A. I.; Barnard, J. H.; Marder, T. B.; Murphy, J. M.; Hartwig, J. F. *Chem. Rev.* **2010**, *110*, 890–931. (e) Colby, D. A.; Tsai, A. S.; Bergman, R. G.; Ellman, J. A. *Acc. Chem. Res.* **2012**, *45*, 814–825. (f) Neufeldt, S. R.; Sanford, M. S. *Acc. Chem. Res.* **2012**, *45*, 936–946.
- (12) Anbarasan, P.; Neumann, H.; Beller, M. *Chem. Commun.* **2011**, *47*, 3233–3235.
- (13) (a) Prasad, C. D.; Balkrishna, S. J.; Kumar, A.; Bhakuni, B. S.; Shrimali, K.; Biswas, S.; Kumar, S. *J. Org. Chem.* **2013**, *78*, 1434–1443. (b) Tudge, M.; Tamiya, M.; Savarin, C.; Humphrey, G. R. *Org. Lett.* **2006**, *8*, 565–568. (c) Silveira, C. S.; Medes, S. R.; Wolf, L.; Marins, G. M. *Tetrahedron Lett.* **2010**, *51*, 2014–2016.
- (14) (a) Saravanan, P.; Anbarasan, P. *Org. Lett.* **2014**, *16*, 848–851. (b) Anbarasan, P.; Neumann, H.; Beller, M. *Chem. Commun.* **2011**, *47*, 3233–3235.
- (15) (a) Chen, X.; Hao, X.-S.; Goodhue, C. E.; Yu, J.-Q. *J. Am. Chem. Soc.* **2006**, *128*, 6790–6791. (b) Zhang, S.; Qian, P.; Zhang, M.; Hu, M.; Cheng, J. *J. Org. Chem.* **2010**, *75*, 6732–6735. (c) Chu, L.; Yue, X.; Qing, F.-L. *Org. Lett.* **2010**, *12*, 1644–1666. (d) Ranjit, S.; Lee, R.; Heryadi, D.; Shen, C.; Wu, J. E.; Zhang, P.; Huang, K.-W.; Liu, X. *J. Org. Chem.* **2011**, *76*, 8999–9007.
- (16) (a) Zhang, M.; Zhang, S.; Pan, C.; Chen, F. *Synth. Commun.* **2012**, *42*, 2844–2853. (b) Fang, X.-L.; Tang, R.-Y.; Zhang, X.-G.; Li, J.-H. *Synthesis* **2011**, 1099–1105.
- (17) Tian, H.; Zhu, C.; Yang, H.; Fu, H. *Chem. Commun.* **2014**, *50*, 8875–8877.
- (18) (a) Thome, I.; Besson, C.; Kleine, T.; Bolm, C. *Angew. Chem., Int. Ed.* **2013**, *52*, 7509–7513. (b) Kano, T.; Shirozu, F.; Maruoka, K. *J. Am. Chem. Soc.* **2013**, *135*, 18036–18039. (c) Sun, C.-L.; Shi, Z.-J. *Chem. Rev.* **2014**, *114*, 9219–9280.
- (19) (a) Zmitek, K.; Zupan, M.; Stavber, S.; Iskra, J. *Org. Lett.* **2006**, *8*, 2491–2494. (b) Nobuta, T.; Tada, N.; Fujiya, A.; Kariya, A.; Miura, T.; Itoh, A. *Org. Lett.* **2013**, *15*, 574–577. (c) Pan, X.; Boussonnière, A.; Curran, D. P. *J. Am. Chem. Soc.* **2013**, *135*, 14433–14437. (d) Humne, V.; Dangat, Y.; Vank, K.; Lokhande, P. *Org. Biomol. Chem.* **2014**, *12*, 4832–4836.
- (20) Zou, L.-H.; Reball, J.; Mottweiler, J.; Bolm, C. *Chem. Commun.* **2012**, *48*, 11307–11309.
- (21) (a) Liao, Y.; Jiang, P.; Chen, S.; Qia, H.; Deng, G.-J. *Green Chem.* **2013**, *15*, 3302–3306. (b) Du, H.-A.; Tang, R.-Y.; Deng, C.-L.; Liu, Y.; Li, J.-H.; Zhang, X.-G. *Adv. Synth. Catal.* **2011**, *353*, 2739–2748. (c) Hiebel, M.-A.; Berteina-Raboin, S. *Green Chem.* **2015**, *17*, 937–944.
- (22) Kang, X.; Yan, R.; Yu, G.; Pang, X.; Liu, X.; Li, X.; Xiang, L.; Huang, G. *J. Org. Chem.* **2014**, *79*, 10605–10610.
- (23) Kerry, F.; Mark, O.; Donald E. S.; Janet, A. (Icos Corp., USA). Patent WO0059878, 2000.
- (24) Du, H. A.; Tang, R. Y.; Deng, C. L.; Liu, Y.; Li, J. H.; Zhang, X. G. *Adv. Synth. Catal.* **2011**, *353*, 2739–2748.
- (25) Duan, Z.; Ranjit, S.; Liu, X. *Org. Lett.* **2010**, *12*, 2430–2433.
- (26) Feng, Y.-S.; Qi, H.-X.; Wang, W.-C.; Liang, Y.-F.; Xu, H.-J. *Tetrahedron Lett.* **2012**, *53*, 2914–2917.
- (27) Cabrero-Antonino, J. R.; García, T.; Rubio-Marqués, P.; Vidal-Moya, J. A.; Leyva-Pérez, A.; Al-Deyab, S. S.; Al-Resayes, S. I.; Díaz, U.; Corma, A. *ACS Catal.* **2011**, *1*, 147–158.
- (28) Kabir, M. S.; Lorenz, M.; Van Linn, M. L.; Namjoshi, O. A.; Ara, S.; Cook, J. M. *J. Org. Chem.* **2010**, *75*, 3626–3643.
- (29) Wang, C.; Ma, Z.; Sun, X.-L.; Gao, Y.; Guo, Y.-H.; Tang, Y.; Shi, L.-P. *Organometallics* **2006**, *25*, 3259–3266.
- (30) Sivan, V.; Reena, V.; Amarendar, C.; Ravindra, P. P.; Ajayan, V. *Synlett* **2010**, *18*, 2733–2736.