

# Iron(III)-Mediated Radical Nitration of Bisarylfonyl Hydrazones: Synthesis of BisarylNitromethyl Sulfones

Dinabandhu Sar,<sup>†</sup> Raghunath Bag,<sup>†</sup> Debajyoti Bhattacharjee,<sup>‡</sup> Ramesh Chandra Deka,<sup>‡</sup> and Tharmalingam Punniyamurthy\*,<sup>†</sup>

<sup>†</sup>Department of Chemistry, Indian Institute of Technology Guwahati, Guwahati-781039, India

<sup>‡</sup>Department of Chemistry, Tezpur University, Tezpur-784028, India

## Supporting Information




**ABSTRACT:** Iron(III)-mediated radical nitration of bisarylfonyl hydrazones is described. In this protocol, the nontoxic and inexpensive  $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$  plays a dual role as catalyst as well as nitro source. The mild conditions, broad substrate scope, and the functional group compatibility are the significant features. The reaction pathway has been demonstrated using DFT calculations, and the products can be subsequently converted into oximes using  $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$  in high yields.

## INTRODUCTION


Iron compounds have recently attracted significant interest in organic synthesis as nontoxic and inexpensive green elements.<sup>1</sup> Several studies have employed iron complexes as replacements for the expensive transition-metal catalysts such as Pd and Rh for the formation of carbon–heteroatom bonds.<sup>2</sup> In addition,  $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$  is well-known to produce  $\text{NO}_2$  radical on heating,<sup>3,4</sup> however, its application in synthetic chemistry is limited (Scheme 1a).<sup>5</sup> In continuation of our studies on

### Scheme 1. Iron(III)-Mediated Radical Nitrations

#### a) Previous study



#### b) This study



bisarylhydrazones,<sup>6</sup> we report here an  $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ -mediated radical nitration of bisarylfonyl hydrazones at moderate temperature (Scheme 1b). The broad substrate scope, use of nontoxic iron salt as the nitro source, and the readily accessible aryl aldehydes and arylsulfonyl hydrazides as the substrate precursors are the significant practical features. The reaction pathway is analyzed using density functional theory (DFT).

The chemistry of nitroalkanes<sup>7</sup> and sulfones<sup>8</sup> continues to attract the attention of chemists as they comprise a synthetically important class of compounds for the generation of molecules

of pharmaceutical relevance.<sup>9</sup> They are widely used in different efficient stereoselective carbon–carbon bond forming reactions and subsequent numerous functional group transformations. For examples, nitroalkanes are utilized in stereoselective Henry reaction<sup>10</sup> and Michael addition,<sup>11</sup> and for the generation of oximes, amines, and carbonyl compounds,<sup>12</sup> while aryl sulfones are substrate precursors for Julia olefination,<sup>13</sup> aldol reaction,<sup>14</sup> and Michael addition,<sup>15</sup> and for the generation of ketones and alkanes by desulfonylation.<sup>16</sup> Aryl nitroalkanes are demonstrated as key intermediates in the total synthesis of manzamine A<sup>17</sup> and (−)-nutlin-3.<sup>18</sup> Development of effective methods for the construction of nitromethane<sup>19,20</sup> with aryl and sulfonyl groups<sup>21</sup> would thus be valuable as they may serve as a precursor to an array of useful structural scaffolds (Scheme 2).<sup>22</sup>


## RESULTS AND DISCUSSION

We commenced the optimization studies with aldehyde **1a** and *p*-toluenesulfonyl hydrazide **2a** as test substrates using a series of nitro sources in different solvents at varied temperature (Table 1). To our delight, the reaction occurred to give bisarylNitromethyl sulfone **3a** in 63% yield when the substrates **1a** and **2a** were stirred at 60 °C for 0.5 h in 1,2-dichloroethane (DCE) followed by at 80 °C for 1.5 h with  $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ . In a set of nitro sources screened,  $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$  gave **3a** in 51% yield, while  $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$  afforded the target product in 38% yield (entries 1–2). In contrast,  $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ ,  $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$ ,  $^t\text{BuONO}$ , and  $\text{AgNO}_3$  failed to react, and the formation of **3a** was not observed. DCE was found to be a solvent of choice, while  $\text{CH}_3\text{CN}$ , toluene and 1,4-dioxane gave

Received: April 30, 2015

Published: June 3, 2015

Scheme 2. Some of the Possible Utilities of the Products

Table 1. Optimization of the Reaction Conditions<sup>a</sup>

entry	nitro source (mol %)	solvents	yield (%)
1	Cu(NO <sub>3</sub> ) <sub>2</sub> ·3H <sub>2</sub> O (50)	DCE	38
2	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O (35)	DCE	51
3	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O (35)	toluene	47
4	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O (35)	CH <sub>3</sub> CN	38
5	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O (35)	dioxane	36
6	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O (35)	THF	trace
7	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O (35)	DMF	trace
8	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O (35)	DCE	36 <sup>b</sup>
9	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O (35)	DCE	32 <sup>c</sup>
10	Fe(NO <sub>3</sub> ) <sub>3</sub> ·9H <sub>2</sub> O (35)	DCE	63 <sup>d</sup>

<sup>a</sup>Reaction conditions: aldehyde **1a** (1.0 mmol), sulfonhydrazide **2a** (1.2 mmol), DCE (3 mL), 60 °C, 0.5 h; Fe(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O (35 mol %), 80 °C, air, 1.5 h. <sup>b</sup>Temp. 60 °C. <sup>c</sup>Temp. 100 °C. <sup>d</sup>MS 4 Å (50 mg) used.

**3a** in 36–47% yields (entries 2–5). In contrast, THF and DMF yielded inferior results (entries 6–7). The optimum temperature was found to be 80 °C, and the addition of the molecular sieves of 4 Å led to an increase in the yield to 63% (entries 8–10).

With the optimal conditions, the scope of the protocol was explored for the reactions of a series of substituted aldehydes **1b–r** with *p*-toluenesulfonyl hydrazide **2a** as a standard substrate (Scheme 3). The reaction of benzaldehyde **1b** afforded bisarylnitromethyl sulfone **3b** in 66% yield. *o*-Tolualdehyde **1c** underwent the reaction to provide the target product **3c** in 57% yield. The reactions of the aldehydes **1d–g** with 3-chloro, 3-methyl, 3-nitro, and 3-trifluoromethyl substituents afforded the corresponding substituted bisarylnitromethyl sulfones **3d–g** in 51–61% yields. The aldehydes **1h–m** bearing 4-bromo, 4-chloro, 4-cyano, 4-fluoro, 4-methoxy, and 4-ester substituents underwent the reaction to produce the target products **3h–m** in 52–69% yields. In addition, the aldehyde **1n** with 2-methoxy and 3-bromo substituents produced the desired **3n** in 65% yield, while the reactions of the aldehydes **1o** and **1p** with 2,4,5-trimethoxy and 3,4,5-trimethoxy groups furnished **3o** and **3p** in 72% and 70% yields, respectively. Furthermore, 2-naphthaldehyde **1q** under-

went the reaction to give **3q** in 73% yield. However, the reaction of aliphatic aldehyde **1r** was not successful due to the decomposition of the hydrazone generated from the aldehyde and sulfonyl hydrazide. Compound **3o** yielded single crystals in CH<sub>3</sub>CN whose structure was determined by single crystal X-ray analysis (see the Supporting Information).


Next, the protocol was extended for the reaction of various arylsulfonyl hydrazides with the aldehydes **1i** and **1l** having 4-chloro and 4-methoxy substituents as the representative examples (Scheme 4). Arylsulfonyl hydrazide **2b** underwent the reaction to furnish the target bisarylnitromethyl sulfones **3s** and **3t** in 66% and 68% yields, respectively. Similarly, the reactions of the arylsulfonyl hydrazides **2c–e** bearing 4-chloro, 4-methoxy, and 4-nitro substituents occurred with 54–64% yields. Furthermore, arylsulfonyl hydrazide **2f** with 2,4,6-trimethyl substituents underwent the reaction with 66% and 64% yields, respectively. These results suggest that the protocol has broad substrate scope to afford an effective route for the construction of functionalized bisarylnitromethyl sulfones.

Finally, the reaction of the aldehyde **1a** and sulfonyl hydrazide **2a** was examined in gram-scale as representative examples (Scheme 5). The reaction readily took place to afford **3a** in good yields, which suggests that the protocol is scalable. Furthermore, the products can be readily converted into oximes in high yields (Scheme 6). For example, the Sn-mediated reaction of **3b**, **3h**, and **3l** afforded the corresponding oximes **4a–c** in high yields. The compound **4a** afforded single crystals in chloroform whose structure was determined by single crystal X-ray analysis (see the Supporting Information).

To gain insight into the mechanism, the reaction of bisarylsulfonyl hydrazone **5** was independently investigated (Scheme 7). As above, the nitration took place efficiently to afford **3a** in 64% yield, which suggests that the condensation of the aldehyde with sulfonyl hydrazide may first give bisarylsulfonyl hydrazone that may undergo nitration to afford the target products. Next, the reaction of the aldehyde **1b** and arylsulfonyl hydrazide **2a** was examined with Fe(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O as the representative examples in the presence of radical scavengers, 1,1-diphenylethylene and *N*-*tert*-butyl-*α*-phenylnitrone (Scheme 8).<sup>23</sup> However, no reaction was observed, which suggests that the reaction may involve the radical intermediates. Furthermore, the substrates showed no nitration in the presence of base, which suggests that the reaction may not proceed via the carbene intermediate (Scheme 9).<sup>24</sup> In addition, *N*-benzyl bisarylsulfonyl hydrazone failed to react, and the substrate was isolated intact, which suggests that N-H is essential for the reaction (Scheme 10). Thus, the reaction of the aldehyde with arylsulfonyl hydrazide can give sulfonyl hydrazone **a** that can react with Fe(NO<sub>3</sub>)<sub>3</sub> via single electron transfer to produce Fe(NO<sub>3</sub>)<sub>2</sub>, NO<sub>3</sub><sup>−</sup>, and intermediate **b** (Scheme 11).<sup>25</sup> Isomerization of **b** can give **c** that can react with the NO<sub>2</sub> radical<sup>3,4</sup> to give **d** and the target products **3** via path **a**. Alternatively, the intermediate **c** may give **e** via path **b**, which with the NO<sub>2</sub> radical can give the target product.<sup>3,4</sup> Oxidation of the Fe(II) species with HNO<sub>3</sub> can regenerate Fe(III) species to complete the catalytic cycle.

The optimized structures of different species involved in the reactions are shown in Figure 1.

The vibrational frequencies of all of the optimized structures are found to be positive, confirming them to be at energy minima (see the Supporting Information).<sup>26,27</sup> To understand the feasibility of the reaction paths, we have calculated the standard reaction enthalpy ( $\Delta H_r$ ) and Gibbs free energy ( $\Delta G_r$ )

Scheme 3. Reaction of Various Aldehydes with *p*-Toluenesulfonyl Hydrazide<sup>a</sup>

<sup>a</sup>Reaction conditions: aldehyde (1.0 mmol), sulfonyl hydrazide **2a** (1.2 mmol), DCE (3 mL), 60 °C, 0.5 h; Fe(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O (35 mol %), MS 4 Å (50 mg), air, 80 °C, 1.5 h. n.o. = not observed.

for both the paths including zero point correction. The  $\Delta G_r$  value for path **a** is found to be -1.76 eV, which indicates that this path is feasible. The  $\Delta H_r$  value is found also to be negative (-13.95 eV) indicating that the path is exothermic in nature. For path **b**, both the thermochemical parameters  $\Delta G_r$  and  $\Delta H_r$  are found to be highly positive, which reveals that this pathway is not feasible. Hence, path **a** may be predominant in nature.

## CONCLUSIONS


In summary, iron(III)-mediated radical nitration of bisarylsulfonyl hydrazones is reported at moderate temperature. The reaction is free from the use of additives and effective under neutral conditions. The mild reaction conditions, broad substrate scope, and the use of nontoxic iron salt as the nitro source are the important practical features. The products can be

converted into oximes using SnCl<sub>2</sub> in high yields. The reaction pathway has been analyzed using DFT calculations.

## EXPERIMENTAL SECTION

**General Information.** Arylsulfonyl chlorides, aryl aldehydes, hydrazine hydrate (99–100%), and SnCl<sub>2</sub>·2H<sub>2</sub>O (97%) were purchased from a commercial source and used as received. Fe(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O (98%) was purchased from a chemical supplier and used as received. Sulfonyl hydrazides,<sup>28</sup> aryl aldehydes,<sup>29</sup> and sulfonyl hydrazone<sup>30</sup> were prepared according to a reported procedure. TLC analysis was performed on silica gel G/GF 254 plates. Silica gel (60–120 mesh) was used for column chromatography. NMR spectra were recorded on 400 and 600 MHz spectrometers using Me<sub>4</sub>Si as an internal standard. Melting points were determined with a melting point apparatus and are uncorrected. FT-IR spectra were collected using an IR spectrometer. A Q-ToF ESI-MS instrument was used for recording high-resolution mass spectra (HRMS). Single crystal X-ray

**Scheme 4. Reaction of Different Sulfonyl Hydrazides with Aldehydes **1i** and **1l**<sup>a</sup>**




<sup>a</sup>Reaction conditions: aldehyde (1.0 mmol), sulfonyl hydrazide (1.2 mmol), DCE (3 mL), 60 °C, 0.5 h; Fe(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O (35 mol %), MS 4 Å (50 mg), air, 80 °C, 1.5 h.

data were collected using a CCD diffractometer, which is equipped with 1.75 kW sealed-tube Mo-K $\alpha$  irradiation ( $\lambda = 0.71073 \text{ \AA}$ ) at 298(2) K. The crystal structure was solved by a direct method using SHEXL-97 (Göttingen, Germany).

**General Procedure for the Iron(III)-Mediated Nitration of Bisarylsulfonyl hydrazones.** Aryl aldehyde **1** (1 mmol) and arylsulfonyl hydrazide **2** (1.2 mmol) were stirred in 1,2-dichloroethane (3 mL) for 0.5 h at 60 °C under air. The reaction mixture was cooled to room temperature and then treated with Fe(NO<sub>3</sub>)<sub>3</sub>·9H<sub>2</sub>O (35 mol %, 0.35 mmol, 141 mg) and molecular sieves (4 Å, 50 mg). The resultant mixture was stirred for 1.5 h at 80 °C under air. The progress of the reaction was monitored by TLC using ethyl acetate and hexane as eluent. After completion, the reaction mixture was cooled to room temperature, and the solvent was evaporated under reduced pressure. The residue was treated with saturated NaHCO<sub>3</sub> (3 mL) and extracted with ethyl acetate (3 × 15 mL). The combined organic solution was successively washed with brine (1 × 10 mL) and water (1 × 10 mL). Drying (Na<sub>2</sub>SO<sub>4</sub>) and evaporation of the solvent gave a residue that was purified on silica gel column chromatography using hexane and ethyl acetate as eluent.

**1-Bromo-3-(nitro(tosyl)methyl)benzene **3a**.** Analytical TLC on silica gel, 1:3 ethyl acetate/hexane  $R_f = 0.41$ ; white solid (231 mg,

**Scheme 6. Synthesis of Oximes**



63%); mp 79–80 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.65 (d,  $J = 7.6$  Hz, 1H), 7.62 (d,  $J = 8.0$  Hz, 2H), 7.56 (s, 1H), 7.45 (d,  $J = 7.6$  Hz, 1H), 7.35 (d,  $J = 8.0$  Hz, 2H), 7.29 (t,  $J = 8.4$  Hz, 1H), 6.40 (s, 1H), 2.48 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  147.4, 135.0, 132.6, 130.7, 130.5, 130.1, 128.5, 127.1, 123.0, 103.0, 22.0; FT-IR (KBr) 2982, 1594, 1561, 1473, 1337, 1257, 1188, 1153, 1081, 1039, 1014, 661 cm<sup>-1</sup>; HRMS (APCI) *m/z* [M – H]<sup>+</sup> calcd for C<sub>14</sub>H<sub>12</sub>BrNO<sub>4</sub>S, 367.9598; found, 367.9591.

**1-Methyl-4-(nitro(phenyl)methylsulfonyl)benzene **3b**.** Analytical TLC on silica gel, 1:9 ethyl acetate/hexane  $R_f = 0.52$ ; white solid (192 mg, 66%); mp 163–164 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.59 (d,  $J = 7.8$  Hz, 2H), 7.51 (t,  $J = 7.8$  Hz, 1H), 7.47 (d,  $J = 7.2$  Hz, 2H), 7.39 (t,  $J = 7.8$  Hz, 2H), 7.31 (d,  $J = 8.4$  Hz, 2H), 6.46 (s, 1H), 2.46 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>)  $\delta$  147.0, 131.8, 131.2, 130.7, 130.0, 129.7, 129.1, 125.3, 103.9, 22.0; FT-IR (KBr) 2984, 2359, 1592, 1559, 1488, 1454, 1333, 1303, 1184, 1155, 1083, 1030, 1016 cm<sup>-1</sup>; HRMS (APCI) *m/z* [M – H]<sup>+</sup> calcd for C<sub>14</sub>H<sub>13</sub>NO<sub>4</sub>S, 290.0493; found, 290.0486.





**1-Methyl-2-(nitro(tosyl)methyl)benzene **3c**.** Analytical TLC on silica gel, 1:3 ethyl acetate/hexane  $R_f = 0.38$ ; white solid (173 mg, 57%); mp 198–199 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.64 (d,  $J = 7.8$  Hz, 2H), 7.54 (d,  $J = 7.8$  Hz, 1H), 7.37 (t,  $J = 7.8$  Hz, 1H), 7.33 (d,  $J = 7.8$  Hz, 2H), 7.26 (d,  $J = 8.4$  Hz, 1H), 7.18 (t,  $J = 7.8$  Hz, 1H), 6.81 (s, 1H), 2.47 (s, 3H), 2.39 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>)  $\delta$  147.0, 138.9, 131.8, 131.7, 131.2, 130.8, 130.0, 128.4, 126.8, 124.2, 99.2, 22.0, 19.8; FT-IR (KBr) 2999, 1689, 1595, 1557, 1491, 1456, 1406, 1355, 1331, 1292, 1264, 1153, 1084, 1037, 1018 cm<sup>-1</sup>; HRMS (APCI) *m/z* [M – H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>15</sub>NO<sub>4</sub>S, 304.0649; found, 304.0658.

**1-Chloro-3-(nitro(tosyl)methyl)benzene **3d**.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f = 0.39$ ; white solid (198 mg, 61%); mp 82–84 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  7.62 (d,  $J = 7.8$  Hz, 2H), 7.50 (d,  $J = 7.8$  Hz, 1H), 7.45 (s, 1H), 7.40 (d,  $J = 7.8$  Hz, 1H), 7.36–7.34 (m, 3H), 6.41 (s, 1H), 2.48 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>)  $\delta$  147.4, 135.2, 132.0, 130.9, 130.7, 130.3, 130.1, 129.8, 128.1, 127.0, 103.1, 22.0; FT-IR (KBr) 2990, 1597, 1562, 1477, 1408, 1384, 1340, 1261, 1187, 1157, 1083, 1017, 663 cm<sup>-1</sup>; HRMS (APCI) *m/z* [M – H]<sup>+</sup> calcd for C<sub>14</sub>H<sub>12</sub>ClNO<sub>4</sub>S, 324.0103; found, 324.0110.

**1-Methyl-3-(nitro(tosyl)methyl)benzene **3e**.** Analytical TLC on silica gel, 1:6 ethyl acetate/hexane  $R_f = 0.49$ ; white solid (177 mg, 58%); mp 116–117 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.61 (d,  $J = 7.6$  Hz, 2H), 7.32 (d,  $J = 7.6$  Hz, 2H), 7.26–7.23 (m, 4H), 6.42 (s, 1H), 2.47 (s, 3H), 2.33 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$

**Scheme 5. Gram-Scale Synthesis**



**Scheme 7. Nitration of Bisarylsulfonyl Hydrazone****Scheme 8. Effect of Radical Scavengers****Scheme 9. Effect of Base****Scheme 10. Effect of N-Protecting Group****Scheme 11. Proposed Reaction Pathway**

146.9, 139.1, 132.6, 131.2, 130.8, 130.2, 129.9, 128.9, 126.8, 125.1, 103.9, 22.0, 21.4; FT-IR (KBr) 2983, 1633, 1592, 1554, 1484, 1333, 1305, 1155, 1083, 1039, 1017  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M - H]<sup>-</sup> calcd for C<sub>15</sub>H<sub>15</sub>NO<sub>4</sub>S, 304.0649; found, 304.0641.

**1-Nitro-3-(nitro(tosyl)methyl)benzene 3f.** Analytical TLC on silica gel, 1:3 ethyl acetate/hexane  $R_f$  = 0.28; yellow solid (184 mg, 55%); mp 132–133 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  8.41 (d, *J* = 8.4 Hz, 1H), 8.33 (s, 1H), 8.00 (d, *J* = 7.2 Hz, 1H), 7.70–7.66 (m, 3H), 7.39 (d, *J* = 7.6 Hz, 2H), 6.59 (s, 1H), 2.49 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (100

**Figure 1.** Optimized structures of organic species (white, hydrogen; gray, carbon; yellow, sulfur; red, oxygen; and blue, nitrogen).

MHz, CDCl<sub>3</sub>)  $\delta$  148.3, 147.9, 135.9, 130.5, 130.4, 130.3, 127.0, 126.6, 125.2, 102.3, 22.0; FT-IR (KBr) 2925, 1707, 1618, 1533, 1481, 1446, 1352, 1291, 1158, 1124, 1035, 1010, 815, 720, 668  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M - H]<sup>-</sup> calcd for C<sub>14</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>S, 335.0343; found, 335.0349.

**1-(Nitro(tosyl)methyl)-3-(trifluoromethyl)benzene **3g**.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f = 0.29$ ; yellow liquid (183 mg, 51%);  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.78 (d,  $J = 7.8$  Hz, 2H), 7.60–7.58 (m, 4H), 7.34 (d,  $J = 8.4$  Hz, 2H), 6.51 (s, 1H), 2.47 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  147.6, 133.5, 133.3, 130.7, 130.2, 129.8, 129.4, 128.6, 127.3, 126.6, 126.3, 103.1, 22.0; FT-IR (neat) 2926, 1596, 1566, 1452, 1384, 1332, 1261, 1159, 1133, 1101, 1081, 814, 703, 659  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{15}\text{H}_{12}\text{F}_3\text{NO}_4\text{S}$ , 358.0361; found, 358.0367.

**1-Bromo-4-(nitro(tosyl)methyl)benzene **3h**.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f = 0.40$ ; white solid (222 mg, 60%); mp 129–130  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.62 (d,  $J = 8.4$  Hz, 2H), 7.55 (d,  $J = 8.4$  Hz, 2H), 7.37–7.33 (m, 4H), 6.41 (s, 1H), 2.47 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  147.3, 132.5, 131.3, 131.0, 130.7, 130.2, 126.8, 124.2, 103.2, 22.0; FT-IR (KBr) 2923, 1626, 1562, 1488, 1408, 1384, 1340, 1155, 1075, 1012, 707, 658  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{14}\text{H}_{12}\text{BrNO}_4\text{S}$ , 367.9598; found, 367.9591.

**1-Chloro-4-(nitro(tosyl)methyl)benzene **3i**.** Analytical TLC on silica gel, 1:3 ethyl acetate/hexane  $R_f = 0.50$ ; white solid (201 mg, 62%); mp 128–129  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.62 (d,  $J = 7.8$  Hz, 2H), 7.44 (d,  $J = 8.4$  Hz, 2H), 7.39 (d,  $J = 8.4$  Hz, 2H), 7.34 (d,  $J = 8.4$  Hz, 2H), 6.43 (s, 1H), 2.47 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  147.3, 138.5, 131.2, 131.0, 130.6, 130.2, 129.5, 123.7, 103.1, 22.0; FT-IR (KBr) 2979, 1665, 1594, 1568, 1494, 1413, 1340, 1310, 1291, 1177, 1152, 1082, 1018, 818, 662  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{14}\text{H}_{12}\text{ClNO}_4\text{S}$ , 324.0103; found, 324.0108.

**4-(Nitro(tosyl)methyl)benzonitrile **3j**.** Analytical TLC on silica gel, 1:2 ethyl acetate/hexane  $R_f = 0.40$ ; white solid (164 mg, 52%); mp 125–126  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.72 (d,  $J = 8.4$  Hz, 2H), 7.66 (d,  $J = 8.4$  Hz, 2H), 7.63 (d,  $J = 7.8$  Hz, 2H), 7.36 (d,  $J = 7.8$  Hz, 2H), 6.49 (s, 1H), 2.48 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  147.8, 132.8, 130.8, 130.7, 130.4, 129.7, 117.7, 115.9, 102.9, 22.1; FT-IR (KBr) 2924, 2853, 2231, 1741, 1628, 1563, 1462, 1338, 1152, 1081, 1017, 812  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_4\text{S}$ , 315.0445; found, 315.0451.

**1-Fluoro-4-(nitro(tosyl)methyl)benzene **3k**.** Analytical TLC on silica gel, 1:2 ethyl acetate/hexane  $R_f = 0.56$ ; yellow solid (182 mg, 59%); mp 107–108  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.62 (d,  $J = 7.8$  Hz, 2H), 7.52–7.49 (m, 2H), 7.34 (d,  $J = 8.4$  Hz, 2H), 7.10 (t,  $J = 8.4$  Hz, 2H), 6.44 (s, 1H), 2.47 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  166.1 ( $J_{\text{C}-\text{F}} = 251.7$  Hz), 147.2, 132.2 ( $J_{\text{C}-\text{F}} = 9.2$  Hz), 131.0, 130.6, 130.1, 121.2 ( $J_{\text{C}-\text{F}} = 3.8$  Hz), 116.5 ( $J_{\text{C}-\text{F}} = 22.1$  Hz), 103.0, 22.0; FT-IR (KBr) 2979, 1605, 1566, 1512, 1406, 1384, 1339, 1233, 1154, 1083, 1016, 820, 665  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{14}\text{H}_{12}\text{FNO}_4\text{S}$ , 308.0398; found, 308.0396.

**1-Methoxy-4-(nitro(tosyl)methyl)benzene **3l**.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f = 0.51$ ; yellow solid (221 mg, 69%); mp 126–127  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.61 (d,  $J = 8.4$  Hz, 2H), 7.41 (d,  $J = 8.4$  Hz, 2H), 7.32 (d,  $J = 7.8$  Hz, 2H), 6.89 (d,  $J = 9.0$  Hz, 2H), 6.40 (s, 1H), 3.83 (s, 3H), 2.46 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  162.4, 146.9, 131.5, 131.4, 130.7, 130.0, 117.0, 114.5, 103.6, 55.6, 22.0; FT-IR (KBr) 2995, 1609, 1596, 1552, 1515, 1460, 1341, 1309, 1278, 1263, 1182, 1157, 1085, 1019  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{15}\text{H}_{15}\text{NO}_5\text{S}$ , 320.0598; found, 320.0599.

**Methyl-4-(nitro(tosyl)methyl)benzoate **3m**.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f = 0.22$ ; white solid (202 mg, 58%); mp 110–111  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  8.05 (d,  $J = 8.4$  Hz, 2H), 7.59–7.54 (m, 4H), 7.32 (d,  $J = 8.4$  Hz, 2H), 6.52 (s, 1H), 3.94 (s, 3H), 2.46 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  166.1, 147.4, 133.2, 130.9, 130.7, 130.2, 130.1, 129.8, 129.5, 103.4, 52.7, 22.0; FT-IR (KBr) 2995, 1714, 1593, 1555, 1432, 1341, 1291, 1187, 1154, 1114, 1083, 1018  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{16}\text{H}_{15}\text{NO}_6\text{S}$ , 348.0547; found, 348.0541.

**1-Bromo-2-methoxy-3-(nitro(tosyl)methyl)benzene **3n**.** Analytical TLC on silica gel, 1:3 ethyl acetate/hexane  $R_f = 0.35$ ; white solid (260 mg, 65%); mp 153–154  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.76 (d,  $J = 2.4$  Hz, 1H), 7.69 (d,  $J = 7.8$  Hz, 2H), 7.56 (dd,  $J = 9.0, 2.4$  Hz, 1H), 7.37 (d,  $J = 8.4$  Hz, 2H), 7.12 (s, 1H), 6.80 (d,  $J = 9.0$  Hz, 1H), 3.76

(s, 3H), 2.48 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  157.0, 147.0, 136.0, 132.9, 131.8, 130.5, 130.0, 115.7, 113.3, 112.9, 95.1, 56.5, 22.0; FT-IR (KBr) 2981, 1595, 1556, 1487, 1462, 1441, 1403, 1352, 1329, 1305, 1282, 1260, 1213, 1186, 1155, 1116, 1085, 1019, 820, 647  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{15}\text{H}_{14}\text{BrNO}_5\text{S}$ , 397.9703; found, 397.9710.

**1,2,4-Trimethoxy-5-(nitro(tosyl)methyl)benzene **3o**.** Analytical TLC on silica gel, 1:2 ethyl acetate/hexane  $R_f = 0.64$ ; yellow solid (274 mg, 72%); mp 148–149  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.69 (d,  $J = 8.4$  Hz, 2H), 7.34 (d,  $J = 8.4$  Hz, 2H), 7.15 (s, 1H), 7.11 (s, 1H), 6.46 (s, 1H), 3.91 (s, 3H), 3.74 (s, 3H), 3.73 (s, 3H), 2.46 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  153.5, 153.0, 146.5, 143.4, 132.4, 130.5, 129.9, 112.1, 104.4, 96.6, 95.9, 56.9, 56.4, 56.2, 21.9; FT-IR (KBr) 2977, 1599, 1525, 1406, 1384, 1332, 1305, 1232, 1214, 1153, 1112, 1083, 1039, 1027  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{17}\text{H}_{19}\text{NO}_5\text{S}$ , 380.0809; found, 380.0807.

**1,2,3-Trimethoxy-5-(nitro(tosyl)methyl)benzene **3p**.** Analytical TLC on silica gel, 1:2 ethyl acetate/hexane  $R_f = 0.59$ ; yellow solid (267 mg, 70%); mp 150–151  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.64 (d,  $J = 8.4$  Hz, 2H), 7.34 (d,  $J = 7.8$  Hz, 2H), 6.63 (s, 2H), 6.37 (s, 1H), 3.86 (s, 3H), 3.75 (s, 6H), 2.46 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  153.4, 147.0, 140.9, 131.1, 130.8, 129.9, 120.1, 107.0, 103.8, 61.1, 56.3, 21.9; FT-IR (KBr) 2945, 1593, 1557, 1509, 1465, 1424, 1336, 1277, 1244, 1230, 1155, 1129, 1079, 1018  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{17}\text{H}_{19}\text{NO}_7\text{S}$ , 380.0809; found, 380.0815.

**2-(Nitro(tosyl)methyl)naphthalene **3q**.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f = 0.71$ ; white solid (249 mg, 73%); mp 175–176  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.95 (s, 1H), 7.86 (t,  $J = 9.0$  Hz, 2H), 7.82 (d,  $J = 7.8$  Hz, 1H), 7.61–7.52 (m, 5H), 7.29 (d,  $J = 8.4$  Hz, 2H), 6.63 (s, 1H), 2.45 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  147.0, 134.6, 132.7, 131.3, 130.9, 130.7, 130.0, 129.0, 128.8, 128.4, 127.9, 127.3, 125.3, 122.5, 104.2, 22.0; FT-IR (KBr) 2987, 1593, 1559, 1507, 1338, 1306, 1211, 1156, 1125, 1083, 1018, 818, 665  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{18}\text{H}_{15}\text{NO}_4\text{S}$ , 340.0649; found, 340.0653.

**1-Chloro-4-(nitro(phenylsulfonyl)methyl)benzene **3s**.** Analytical TLC on silica gel, 1:6 ethyl acetate/hexane  $R_f = 0.29$ ; white solid (205 mg, 66%); mp 117–118  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.76–7.74 (m, 3H), 7.56 (t,  $J = 7.6$  Hz, 2H), 7.44–7.38 (m, 4H), 6.46 (s, 1H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  138.6, 135.7, 134.0, 131.1, 130.6, 129.6, 129.5, 123.5, 103.0; FT-IR (KBr) 2979, 1565, 1488, 1448, 1411, 1355, 1337, 1289, 1180, 1154, 1080, 1012, 770, 728  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{13}\text{H}_{10}\text{ClNO}_4\text{S}$ , 309.9943; found, 309.9943.

**1-Methoxy-4-(nitro(phenylsulfonyl)methyl)benzene **3t**.** Analytical TLC on silica gel, 1:6 ethyl acetate/hexane  $R_f = 0.31$ ; white solid (209 mg, 68%); mp 143–144  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.73 (d,  $J = 8.4$  Hz, 2H), 7.70 (d,  $J = 8.4$  Hz, 1H), 7.52 (t,  $J = 7.8$  Hz, 2H), 7.39 (d,  $J = 8.4$  Hz, 2H), 6.89 (d,  $J = 9.0$  Hz, 2H), 6.43 (s, 1H), 3.82 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  162.4, 135.4, 134.4, 131.4, 130.6, 129.3, 116.8, 114.5, 103.5, 55.6; FT-IR (KBr) 2990, 1607, 1578, 1554, 1513, 1460, 1445, 1338, 1309, 1276, 1262, 1181, 1160, 1113, 1084, 1018  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M – H] $^-$  calcd for  $\text{C}_{14}\text{H}_{13}\text{NO}_5\text{S}$ , 306.0442; found, 306.0448.

**1-Chloro-4-((4-chlorophenyl)(nitro)methyl)sulfonyl)benzene **3u**.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f = 0.45$ ; white solid (197 mg, 57%); mp 117–118  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  7.67 (d,  $J = 9.0$  Hz, 2H), 7.53 (d,  $J = 8.4$  Hz, 2H), 7.41 (s, 4H), 6.44 (s, 1H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (150 MHz,  $\text{CDCl}_3$ )  $\delta$  143.0, 138.8, 132.4, 132.1, 131.0, 129.9, 129.7, 123.3, 103.1; FT-IR (KBr) 2901, 2072, 1639, 1430, 1372, 1339, 1319, 1282, 1205, 1163, 1059, 1032, 896, 707  $\text{cm}^{-1}$ ; HRMS (APCI)  $jkm/z$  [M – H] $^-$  calcd for  $\text{C}_{13}\text{H}_9\text{Cl}_2\text{NO}_4\text{S}$ , 343.9557; found, 343.9550.

**1-Chloro-4-((4-methoxyphenyl)(nitro)methyl)sulfonyl)benzene **3v**.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f = 0.40$ ; yellow solid (208 mg, 61%); mp 113–114  $^\circ\text{C}$ ;  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.63 (d,  $J = 8.4$  Hz, 2H), 7.49 (d,  $J = 8.4$  Hz, 2H), 7.35 (d,  $J = 8.4$  Hz, 2H), 6.89 (d,  $J = 8.4$  Hz, 2H), 6.39 (s, 1H), 3.82 (s, 3H);  $^{13}\text{C}$  { $^1\text{H}$ }NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  162.5, 142.5, 132.7, 132.1, 131.3,

129.6, 116.6, 114.7, 103.6, 55.6; FT-IR (KBr) 2958, 1608, 1579, 1556, 1513, 1469, 1447, 1427, 1395, 1356, 1339, 1310, 1182, 1152, 1142, 1116, 1090, 1077, 1028, 657  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M - H]<sup>-</sup> calcd for C<sub>14</sub>H<sub>12</sub>ClNO<sub>3</sub>S, 340.0052; found, 340.0057.

**1-Chloro-4-((4-methoxyphenyl)sulfonyl)(nitro)methylbenzene 3w.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f$  = 0.34; yellow solid (218 mg, 64%); mp 112–113 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 7.65 (d,  $J$  = 9.0 Hz, 2H), 7.43 (d,  $J$  = 8.4 Hz, 2H), 7.39 (d,  $J$  = 8.4 Hz, 2H), 6.99 (d,  $J$  = 9.0 Hz, 2H), 6.42 (s, 1H), 3.91 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>) δ 165.5, 138.4, 133.1, 131.1, 129.5, 124.9, 123.9, 114.7, 103.2, 56.1; FT-IR (KBr) 2992, 1594, 1562, 1496, 1461, 1409, 1332, 1316, 1263, 1150, 1083, 1019, 830, 663  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M - H]<sup>-</sup> calcd for C<sub>14</sub>H<sub>12</sub>ClNO<sub>3</sub>S, 340.0052; found, 340.0057.

**1-Methoxy-4-((4-methoxyphenyl)(nitro)methylsulfonyl)benzene 3x.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f$  = 0.50; yellow solid (212 mg, 63%); mp 118–119 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.63 (d,  $J$  = 6.8 Hz, 2H), 7.39 (d,  $J$  = 7.2 Hz, 2H), 6.96 (d,  $J$  = 6.8 Hz, 2H), 6.88 (d,  $J$  = 6.8 Hz, 2H), 6.41 (s, 1H), 3.88 (s, 3H), 3.81 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (100 MHz, CDCl<sub>3</sub>) δ 165.2, 162.3, 133.0, 131.3, 125.3, 117.2, 114.5, 114.4, 103.7, 55.9, 55.6; FT-IR (KBr) 2993, 2389, 1609, 1592, 1561, 1513, 1499, 1461, 1441, 1330, 1310, 1267, 1179, 1148, 1084, 1023  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M - H]<sup>-</sup> calcd for C<sub>15</sub>H<sub>15</sub>NO<sub>6</sub>S, 336.0547; found, 336.0542.

**1-Chloro-4-(nitro((4-nitrophenyl)sulfonyl)methyl)benzene 3y.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f$  = 0.29; yellow solid (192 mg, 54%); mp 198–199 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.39 (d,  $J$  = 9.0 Hz, 2H), 7.95 (d,  $J$  = 9.0 Hz, 2H), 7.44–7.40 (m, 4H), 6.51 (s, 1H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>) δ 152.0, 139.7, 139.2, 132.3, 130.9, 129.9, 124.4, 122.7, 103.1; FT-IR (KBr) 2989, 1639, 1593, 1554, 1533, 1491, 1412, 1341, 1315, 1290, 1157, 1108, 1093, 1080, 1030, 740  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M - H]<sup>-</sup> calcd for C<sub>13</sub>H<sub>9</sub>ClN<sub>2</sub>O<sub>6</sub>S, 354.9792; found, 354.9799.

**1-Methoxy-4-(nitro((4-nitrophenyl)sulfonyl)methyl)benzene 3z.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f$  = 0.28; white solid (204 mg, 58%); mp 172–173 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 8.35 (d,  $J$  = 9.0 Hz, 2H), 7.92 (d,  $J$  = 9.0 Hz, 2H), 7.35 (d,  $J$  = 8.4 Hz, 2H), 6.91 (d,  $J$  = 9.0 Hz, 2H), 6.48 (s, 1H), 3.84 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>) δ 162.8, 151.8, 140.0, 132.4, 131.2, 124.2, 116.0, 114.9, 103.6, 55.7; FT-IR (KBr) 2959, 1610, 1582, 1556, 1526, 1513, 1402, 1344, 1307, 1252, 1179, 1144, 1111, 1079, 1026  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M - H]<sup>-</sup> calcd for C<sub>14</sub>H<sub>12</sub>N<sub>2</sub>O<sub>7</sub>S, 351.0292; found, 351.0289.

**2-((4-Chlorophenyl)(nitro)methylsulfonyl)-1,3,5-trimethylbenzene 3aa.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f$  = 0.36; white solid (233 mg, 66%); mp 164–165 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 7.59 (d,  $J$  = 9.0 Hz, 2H), 7.44 (d,  $J$  = 8.4 Hz, 2H), 7.00 (s, 2H), 6.43 (s, 1H), 2.53 (s, 6H), 2.33 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>) δ 145.8, 142.1, 138.5, 132.9, 132.1, 129.4, 129.2, 123.3, 102.5, 23.2, 21.4; FT-IR (KBr) 2997, 1598, 1567, 1489, 1443, 1411, 1350, 1322, 1288, 1144, 1089, 714, 651  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M - H]<sup>-</sup> calcd for C<sub>16</sub>H<sub>16</sub>ClNO<sub>4</sub>S, 352.0416; found, 352.0420.

**2-((4-Methoxyphenyl)(nitro)methylsulfonyl)-1,3,5-trimethylbenzene 3ab.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f$  = 0.54; yellow solid (223 mg, 64%); mp 134–135 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 7.56 (d,  $J$  = 9.0 Hz, 2H), 6.98 (s, 2H), 6.95 (d,  $J$  = 9.0 Hz, 2H), 6.40 (s, 1H), 3.84 (s, 3H), 2.52 (s, 6H), 2.32 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>) δ 162.5, 145.4, 142.0, 132.8, 132.3, 129.6, 116.7, 114.5, 103.1, 55.6, 23.2, 21.3; FT-IR (KBr) 2979, 1608, 1556, 1515, 1443, 1410, 1384, 1357, 1329, 1267, 1176, 1153, 1025  $\text{cm}^{-1}$ ; HRMS (APCI)  $m/z$  [M - H]<sup>-</sup> calcd for C<sub>17</sub>H<sub>19</sub>NO<sub>5</sub>S, 348.0911; found, 348.0905.

**General Procedure for the Synthesis of Oximes.** To a solution of bisarylnitromethyl sulfone (0.25 mmol) in ethanol (1 mL) was added SnCl<sub>2</sub>·2H<sub>2</sub>O (0.75 mmol, 169 mg). The reaction mixture was stirred for 3 h at 50 °C under air. The progress of the reaction was monitored by TLC using ethyl acetate and hexane as eluent. After completion, the reaction mixture was cooled to room temperature, and the solvent was evaporated under reduced pressure. The residue was treated with saturated NaHCO<sub>3</sub> (3 mL) and extracted with ethyl

acetate (3 × 15 mL). The combined organic solution was successively washed with brine (1 × 10 mL) and water (1 × 10 mL). Drying (Na<sub>2</sub>SO<sub>4</sub>) and evaporation of the solvent gave a residue that was purified by column chromatography using hexane and ethyl acetate as eluent.

**(E)-Phenyl(tosyl)methanone Oxime 4a.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f$  = 0.49; white solid (61 mg, 90%); mp 145–146 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 9.52 (s, 1H), 7.67 (d,  $J$  = 7.8 Hz, 2H), 7.46–7.44 (m, 1H), 7.41–7.40 (m, 4H), 7.27 (d,  $J$  = 7.8 Hz, 2H), 2.41 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>) δ 158.6, 145.6, 134.7, 130.9, 130.0, 129.6, 129.3, 128.6, 125.7, 21.9; FT-IR (KBr) 3373, 2922, 2854, 1593, 1443, 1385, 1289, 1145, 1086, 1014, 970  $\text{cm}^{-1}$ ; HRMS (ESI)  $m/z$  [M + H]<sup>+</sup> calcd for C<sub>14</sub>H<sub>13</sub>NO<sub>3</sub>S, 276.0694; found, 276.0698.

**(E)-(4-Bromophenyl)(tosyl)methanone Oxime 4b.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f$  = 0.42; white solid (76 mg, 86%); mp 160–161 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 9.13 (s, 1H), 7.68 (d,  $J$  = 8.4 Hz, 2H), 7.56 (d,  $J$  = 8.4 Hz, 2H), 7.32 (d,  $J$  = 8.4 Hz, 2H), 7.31 (d,  $J$  = 7.8 Hz, 2H), 2.43 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>) δ 158.5, 145.8, 134.5, 132.0, 130.1, 129.4, 125.8, 124.4, 21.9; FT-IR (KBr) 3325, 2921, 2851, 1594, 1586, 1484, 1315, 1215, 1147, 1073, 1019, 998, 810, 674  $\text{cm}^{-1}$ ; HRMS (ESI)  $m/z$  [M + H]<sup>+</sup> calcd for C<sub>14</sub>H<sub>12</sub>BrNO<sub>3</sub>S, 353.9800; found, 353.9796.

**(E)-(4-Methoxyphenyl)(tosyl)methanone Oxime 4c.** Analytical TLC on silica gel, 1:4 ethyl acetate/hexane  $R_f$  = 0.48; white solid (63 mg, 83%); mp 143–144 °C; <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>) δ 7.69 (d,  $J$  = 8.4 Hz, 2H), 7.46 (d,  $J$  = 9.0 Hz, 2H), 7.28 (d,  $J$  = 7.8 Hz, 2H), 6.92 (d,  $J$  = 9.0 Hz, 2H), 3.83 (s, 3H), 2.41 (s, 3H); <sup>13</sup>C {<sup>1</sup>H}NMR (150 MHz, CDCl<sub>3</sub>) δ 161.5, 158.5, 145.4, 135.0, 131.4, 129.9, 129.2, 117.5, 114.1, 55.5, 21.9; FT-IR (KBr) 3448, 2961, 2924, 2854, 1604, 1507, 1455, 1314, 1293, 1258, 1173, 1149, 1088, 1019, 969, 835  $\text{cm}^{-1}$ ; HRMS (ESI)  $m/z$  [M + H]<sup>+</sup> calcd for C<sub>15</sub>H<sub>15</sub>NO<sub>4</sub>S, 306.0800; found, 306.0800.

## ASSOCIATED CONTENT

### Supporting Information

Crystal data and structures of 3o and 4a, computational details, and NMR spectra (<sup>1</sup>H and <sup>13</sup>C) of the products 3a–q, 3s–ab, and 4a–c. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.5b00820.

## AUTHOR INFORMATION

### Corresponding Author

\*E-mail: tpunni@iitg.ernet.in.

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

We gratefully acknowledge Department of Science and Technology (SR/S1/OC-55/2011), Council of Scientific and Industrial Research (02(0088)/12/EMR-II), and Ministry of Human Resource and Development (COE) for financial support. D.S. thanks CSIR for SRF Fellowship. We also thank Central Instrumental Facility, Indian Institute of Technology Guwahati for NMR facilities.

## REFERENCES

- (1) For reviews on iron-catalyzed reactions, see: (a) *Iron-Catalysis in Organic Chemistry*; Plietker, B., Ed.; Wiley-VCH: Weinheim, Germany, 2008. (b) Bolm, C.; Legros, J.; LePaih, J.; Zani, L. *Chem. Rev.* **2004**, 104, 6217. (c) Fürstner, A. *Angew. Chem., Int. Ed.* **2009**, 48, 1364. (d) Sahan, A. A.; Bolm, C. *Chem. Soc. Rev.* **2009**, 38, 2730.
- (2) For examples, see: (a) Sylvester, K. T.; Chirik, P. J. *J. Am. Chem. Soc.* **2009**, 131, 8772. (b) Wu, J. Y.; Moreau, B.; Ritter, T. *J. Am. Chem. Soc.* **2009**, 131, 12915. (c) Yoshikai, N.; Mieczkowski, A.; Matsumoto,

- A.; Ilies, L.; Nakamura, E. *J. Am. Chem. Soc.* **2010**, *132*, 5568.  
 (d) Junge, K.; Wundt, B.; Shaikh, N.; Beller, M. *Chem. Commun.* **2010**, *1769*.
- (3) (a) Hill, W. D., Jr. *Inorg. Chem. Acta* **1986**, *121*, L33.  
 (b) Wieczorek-Ciurowa, K.; Kozak, A. *J. J. Therm. Anal. Calorim.* **1999**, *58*, 647.
- (4) Varma, R. S.; Naicker, K. P.; Liesen, P. *J. Tetrahedron Lett.* **1998**, *39*, 3977.
- (5) For nitration of alkenes, see: Taniguchi, T.; Fujii, T.; Ishibashi, H. *J. Org. Chem.* **2010**, *75*, 8126.
- (6) Guru, M. M.; Punniyamurthy, T. *J. Org. Chem.* **2012**, *77*, 5063.
- (7) For examples, see: (a) Feuer, H.; Nielsen, A. T. *Nitro Compounds: Recent Advances in Synthesis and Chemistry*; VCH: New York, 1990.  
 (b) Ono, N. *The Nitro group in Organic Synthesis*; Wiley-VCH: New York, 2001.  
 (c) Kudyba, I.; Raczkowski, J.; Jurczak, J. *J. Org. Chem.* **2004**, *69*, 2844.  
 (d) Ballini, R.; Bosica, G.; Fiorini, D.; Palmieri, A.; Petrini, M. *Chem. Rev.* **2005**, *105*, 933.  
 (e) Fioravanti, S.; Pellacani, L.; Tardella, P. A.; Vergari, M. C. *Org. Lett.* **2008**, *10*, 1449.  
 (f) Bui, T.; Syed, S.; Barbas, C. F., III. *J. Am. Chem. Soc.* **2009**, *131*, 8758.  
 (g) Marques-Lopez, E.; Merino, P.; Tejero, T.; Herrera, R. P. *Eur. J. Org. Chem.* **2009**, 2401.  
 (h) Ando, K.; Shimazu, Y.; Seki, N.; Yamataka, H. *J. Org. Chem.* **2011**, *76*, 3937.
- (8) For examples, see: (a) Simpkins, N. S. *Sulfones in Organic Synthesis*; Pergamon: Oxford, U.K., 1993.  
 (b) Toru, T.; Bolm, C. *Organosulfur Chemistry in Asymmetric Synthesis*; Wiley-VCH: Weinheim, Germany, 2008.  
 (c) El-Awa, A.; Noshi, M. N.; Mollat du Jourdin, X.; Fuchs, P. L. *Chem. Rev.* **2009**, *109*, 2315.  
 (d) Liu, C.-R.; Li, M.-B.; Cheng, D.-J.; Yang, C.-F.; Tian, S.-K. *Org. Lett.* **2009**, *11*, 2543.  
 (e) Prakash, G. K. S.; Chacko, S.; Vaghoo, H.; Shao, N.; Gurung, L.; Mathew, T.; Olah, G. A. *Org. Lett.* **2009**, *11*, 1127.
- (9) For examples, see: (a) Mulzer, J.; Zuhse, R.; Schmiechen, R. *Angew. Chem., Int. Ed.* **1992**, *31*, 870.  
 (b) Lovejoy, B.; Welch, A. R.; Carr, S.; Luong, C.; Broka, C.; Hendricks, R. T.; Campbell, J. A.; Walker, K. A. M.; Martin, R.; Van Wart, H.; Browner, M. F. *Nat. Struct. Biol.* **1999**, *6*, 217.  
 (c) Lakshmaiah, G.; Kawabata, T.; Shang, M.; Fuji, K. *J. Org. Chem.* **1999**, *64*, 1699.  
 (d) Mulzer, J.; Riether, D. *Org. Lett.* **2000**, *2*, 3139.  
 (e) Kudyba, I.; Raczkowski, J.; Jurczak, J. *J. Org. Chem.* **2004**, *69*, 2844.
- (10) For some selected examples, see: (a) Bulut, A.; Aslan, A.; Dogan, O. *J. Org. Chem.* **2008**, *73*, 7373.  
 (b) Kowalczyk, R.; Kwiatkowski, P.; Skarzewski, J.; Jurczak, J. *J. Org. Chem.* **2009**, *74*, 753.  
 (c) Fabris, M.; Noë, M.; Perosa, A.; Selva, M.; Ballini, R. *J. Org. Chem.* **2012**, *77*, 1805.  
 (d) Hashimoto, K.; Kumagai, N.; Shibasaki, M. *Org. Lett.* **2014**, *16*, 3496.
- (11) For some selected examples, see: (a) Ballini, R.; Bosica, G.; Fiorini, D.; Palmieri, A.; Petrini, M. *Chem. Rev.* **2005**, *105*, 933.  
 (b) Zhu, Q.; Lu, Y. *Org. Lett.* **2009**, *11*, 1721.  
 (c) García, J. M.; Maestro, M. A.; Oiarbide, M.; Odriozola, J. M.; Razkin, J.; Palomo, C. *Org. Lett.* **2009**, *11*, 3826.  
 (d) Sahu, B.; Gururaja, G. N.; Mobin, S. M.; Namboothiri, I. N. N. *J. Org. Chem.* **2009**, *74*, 2601.  
 (e) Ogawa, T.; Mouri, S.; Yazaki, R.; Kumagai, N.; Shibasaki, M. *Org. Lett.* **2012**, *14*, 110.  
 (f) Qian, H.; Yu, X.; Zhang, J.; Sun, J. *J. Am. Chem. Soc.* **2013**, *135*, 18020.  
 (g) Quintavalla, A.; Lanza, F.; Montroni, E.; Lombardo, M.; Trombini, C. *J. Org. Chem.* **2013**, *78*, 12049.
- (12) (a) Kurti, L.; Czako, B. *Strategic Applications of Named Reactions in Organic Synthesis*; Elsevier Academic Press: Amsterdam, The Netherlands, 2005.  
 (b) Ballini, R.; Palmieri, A.; Righi, P. *Tetrahedron* **2007**, *63*, 12099.
- (13) For recent examples, see: (a) Julia, M.; Paris, J.-M. *Tetrahedron Lett.* **1973**, *14*, 4833.  
 (b) Kelly, S. E. *Comp. Org. Syn.* **1991**, *1*, 792.  
 (c) Aïssa, C. *J. Org. Chem.* **2006**, *71*, 360.  
 (d) Aleman, J.; Reyes, E.; Richter, B.; Overgaard, J.; Jorgensen, K. A. *Chem. Commun.* **2007**, *3921*.
- (14) For examples, see: (a) Ni, C.; Wang, F.; Hu, J. *Beilstein J. Org. Chem.* **2008**, *4*, 21.  
 (b) Cassani, C.; Bernardi, L.; Fini, F.; Ricci, A. *Angew. Chem., Int. Ed.* **2009**, *48*, 5694.
- (15) Nielsen, M.; Jacobsen, C. B.; Holub, N.; Paixao, M. W.; Jorgensen, K. A. *Angew. Chem., Int. Ed.* **2010**, *49*, 2668.
- (16) For examples, see: (a) Najera, C.; Yus, M. *Tetrahedron* **1999**, *55*, 10547.  
 (b) Mosse, S.; Alexakis, A. *Org. Lett.* **2005**, *7*, 4361.  
 (c) Zhu, Q.; Lu, Y. *Org. Lett.* **2008**, *10*, 4803.
- (17) Jakubec, P.; Hawkins, A.; Felzmann, W.; Dixon, D. J. *J. Am. Chem. Soc.* **2012**, *134*, 17482.
- (18) Davis, T. A.; Johnston, J. N. *Chem. Sci.* **2011**, *2*, 1076.
- (19) For the synthesis of aryl nitromethane via nucleophilic substitution, see: Greger, J. G.; Yoon-Miller, S. J. P.; Bechtold, N. R.; Flewelling, S. A.; MacDonald, J. P.; Downey, C. R.; Cohen, E. A.; Pelkey, E. T. *J. Org. Chem.* **2011**, *76*, 8203.
- (20) For the synthesis aryl nitromethane via cross-coupling, see: (a) Fox, J. N.; Huang, X.; Chieffi, A.; Buchwald, S. L. *J. Am. Chem. Soc.* **2000**, *122*, 1360.  
 (b) Walvoord, R.; Berrett, S.; Kozlowski, M. C. *Org. Lett.* **2012**, *14*, 4086.  
 (c) Metz, A. E.; Kozlowski, M. C. *J. Org. Chem.* **2013**, *78*, 717.  
 (d) Walvoord, R. R.; Kozlowski, M. C. *J. Org. Chem.* **2013**, *78*, 8859.
- (21) For examples, see: (a) Pulkkinen, J.; Aburel, P. S.; Halland, N.; Jorgensen, K. A. *Adv. Synth. Catal.* **2004**, *346*, 1077.  
 (b) Cid, M. B.; Lopez-Cantarero, J.; Duce, S.; Garcia Ruano, J. L. *J. Org. Chem.* **2009**, *74*, 431.
- (22) For base mediated synthesis of bisarylnitromethyl sulfone, see: Truce, W. E.; Klingler, T. C.; Paar, J. E.; Feuer, H.; Wu, D. K. *J. Org. Chem.* **1969**, *34*, 3104.
- (23) (a) Antonchick, A. P.; Samanta, R.; Kulikov, K.; Lategahn, J. *Angew. Chem., Int. Ed.* **2011**, *50*, 8605.  
 (b) Huang, J.; He, Y.; Wang, Y.; Zhu, Q. *Chem.—Eur. J.* **2012**, *18*, 13964.
- (24) Barluenga, J.; Tomas-Gamasa, M.; Aznar, F.; Valdes, C. *Angew. Chem., Int. Ed.* **2010**, *49*, 4993.
- (25) Taniguchi, T.; Sugiura, Y.; Zaimoku, H.; Ishibashi, H. *Angew. Chem., Int. Ed.* **2010**, *49*, 10154.
- (26) (a) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785.  
 (b) Becke, A. D. *J. Chem. Phys.* **1992**, *96*, 2155.
- (27) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. *Gaussian 09*; Revision B.01; Gaussian, Inc.: Wallingford, CT, 2009.
- (28) Yang, F.-L.; Ma, X.-T.; Tian, S.-K. *Chem.—Eur. J.* **2012**, *18*, 1582.
- (29) Doshi, J. M.; Tian, D.; Xing, C. *J. Med. Chem.* **2006**, *49*, 7731.
- (30) Guru, M. M.; Ali, M. A.; Punniyamurthy, T. *J. Org. Chem.* **2011**, *76*, 5295.