


# One-Pot Synthesis of $\beta,\beta$ -Disubstituted $\alpha,\beta$ -Unsaturated Carbonyl Compounds

Masaharu Sugiura,\* Yasuhiko Ashikari, and Makoto Nakajima

Graduate School of Pharmaceutical Sciences, Kumamoto University, 5-1 Oe-honmachi, Chuo-ku, Kumamoto 862-0973, Japan

**S** Supporting Information

**ABSTRACT:**  $\text{TiCl}_4$ -promoted aldol reaction of ketones as aldol acceptors followed by elimination of the titanoxo group from the Ti-aldolates affords  $\beta,\beta$ -disubstituted  $\alpha,\beta$ -unsaturated carbonyl compounds in a one-pot procedure. The use of additives, such as DMF,  $N,N,N',N'$ -tetramethylethylenediamine, and pyridine, in the elimination step was found to be important.



$\beta,\beta$ -Disubstituted  $\alpha,\beta$ -unsaturated carbonyl compounds are useful compounds themselves and as starting materials for the construction of tetra- or trisubstituted carbon stereogenic centers at the carbonyl  $\beta$ -position. The total syntheses of various natural products have been accomplished starting from these functional compounds.<sup>1</sup> We also reported the Lewis base-catalyzed enantioselective conjugate reduction of a  $\beta,\beta$ -disubstituted  $\alpha,\beta$ -unsaturated ketone (enone) with trichlorosilane.<sup>2</sup>


The combination of the Horner–Wadsworth–Emmons reaction of ketones and subsequent transformation to enones via Weinreb amides is widely used to prepare acyclic  $\beta,\beta$ -disubstituted enones.<sup>3</sup> Sulfenylation/desulfenylation of ketones,<sup>4</sup> addition of organometallic reagents to  $\beta$ -amino-,  $\beta$ -alkylthio- or  $\beta$ -alkoxyenones,<sup>5</sup> conjugate addition of organometallic reagents to ynones,<sup>6</sup> rearrangement of tertiary propargylic alcohols or their acetates,<sup>7</sup> and oxidative rearrangement of tertiary allylic alcohols<sup>8</sup> have also been utilized. Recently, Donohoe and co-workers reported an efficient stereoselective method based on the Mizoroki–Heck reaction of vinyl Weinreb amides.<sup>9</sup> However, these precedents all require several reaction steps.

The aldol addition to simple ketones followed by dehydration (the aldol condensation) may offer a straightforward strategy for the preparation of  $\beta,\beta$ -disubstituted  $\alpha,\beta$ -unsaturated carbonyl compounds from readily available starting materials; however, this has not yet been systematically studied because of the relatively low reactivity of ketones as electrophiles (aldol acceptors), its propensity for the retroaldol reaction, and generation of undesired self- and/or cross-aldol products. In one of the few examples, Tanabe and co-workers reported highly useful diastereoselective  $\text{TiCl}_4/\text{Bu}_3\text{N}$ -promoted aldol reactions between two ketones or between phenyl esters/*S*-phenyl thioesters and ketones.<sup>10–13</sup> They also reported the synthesis of a muscone precursor<sup>10c</sup> and dihydrojasmonone<sup>11b</sup> by dehydration of the corresponding isolated aldol products. Inspired by their work, we envisaged that elimination of the titanoxo group directly from the Ti-aldolate intermediate (without isolation of the aldol product) might be feasible by

optimizing the reaction conditions. Here, we report the one-pot synthesis of  $\beta,\beta$ -disubstituted  $\alpha,\beta$ -unsaturated carbonyl compounds by using additives in the elimination step.

Initially, we investigated the reaction of methyl isopropyl ketone as the aldol donor and acetophenone as the aldol acceptor (Table 1). According to Tanabe's procedure with

**Table 1. Screening of Additives<sup>a</sup>**



entry	additive (equiv)	time (h)	yield (%) <sup>b</sup>	<i>E/Z</i> <sup>c</sup>
1	none	20	18	97:3
2	DMF (2.0)	20	58	96:4
3	DMF (5.0)	20	80	97:3
4	DMF (10)	20	74	98:2
5	NMP <sup>d</sup> (5.0)	21	19	97:3
6	DMPU <sup>e</sup> (5.0)	23	48	97:3
7	$\text{Bu}_3\text{N}$ (1.0)	20	65	99:1
8	$\text{Et}_3\text{N}$ (1.0)	20	48	97:3
9	$i\text{-Pr}_2\text{NEt}$ (1.0)	20	39	97:3
10	TMEDA <sup>f</sup> (1.0)	20	72	97:3
11	pyridine (3.0)	1.5	63	87:13
12	pyridine (5.0)	5	87	86:14
13	$\alpha$ -picoline (5.0)	2	82	90:10
14	2,6-lutidine (5.0)	21	71	88:12

<sup>a</sup>The reaction was carried out by using methyl isopropyl ketone (1.0 mmol),  $\text{TiCl}_4$  (1.1 mmol),  $\text{Bu}_3\text{N}$  (1.2 mmol), acetophenone (1.0 mmol), and additive in dichloromethane (2 mL). <sup>b</sup>Isolated yield. <sup>c</sup>Determined by <sup>1</sup>H NMR spectroscopic analysis of the crude product. <sup>d</sup>*N*-Methyl-2-pyrrolidone. <sup>e</sup>*N,N'*-Dimethylpropyleneurea. <sup>f</sup>*N,N,N',N'*-Tetramethylethylenediamine.


Received: May 30, 2015

Published: August 18, 2015

slight modification of the amounts of substrate and reagent,<sup>10</sup> methyl isopropyl ketone (1.0 equiv) in dichloromethane was treated successively with  $\text{TiCl}_4$  (1.1 equiv) and  $\text{Bu}_3\text{N}$  (1.2 equiv) at  $-78^\circ\text{C}$ , and acetophenone (1.0 equiv) was added to the mixture. After 30 min, formation of the aldol product was confirmed by TLC analysis. If the reaction was quenched at this stage, the aldol product was obtained in 96% yield. However, when the temperature was increased to  $\text{rt}$ <sup>14</sup> without workup to facilitate the elimination, the desired enone was obtained only in 18% yield (Table 1, entry 1). A large amount of acetophenone was generated, which indicated progress of the retroaldol reaction. We reasoned that the Ti-enolate was regenerated because Ti strongly coordinated to the carbonyl oxygen atom in the Ti-aldolate intermediate. To cleave the coordinate bond, DMF was added as a ligand of titanium in the elimination step. As expected, the yield of the enone increased (Table 1, entries 2–4); DMF (5 equiv) provided the best result. *N*-Methyl-2-pyrrolidone (NMP) and *N,N'*-dimethylpropyleneurea (DMPU) were inferior alternatives to DMF (Table 1, entries 5 and 6).<sup>15</sup> In addition to coordination to titanium, deprotonation of a carbonyl  $\alpha$ -proton would facilitate the elimination step. Thus, tertiary amines were examined as additives. Monoamines were moderately effective (Table 1, entries 7–9),<sup>16</sup> whereas *N,N,N',N'*-tetramethylethylenediamine (TMEDA), which can bidentately coordinate to titanium, afforded a good yield (Table 1, entry 10). The use of larger amounts of TMEDA (2.0 equiv) decreased the yield.<sup>17</sup> Next, pyridine and its derivatives were examined (Table 1, entries 11–14).<sup>18</sup> Unlike aliphatic amines, larger amounts (5.0 equiv) of pyridine gave higher yield than lower amounts (3.0 equiv) (entries 11 and 12). Although the *E* selectivity was lowered, significant rate enhancement was observed with pyridine and  $\alpha$ -picoline. Thus, we identified DMF, TMEDA, and pyridine as effective additives for the elimination step. It should also be noted that the titanium residue, which precipitates naturally or on addition of diethyl ether and hexane, can be removed by simple filtration through a Celite pad without aqueous workup. After evaporation, the filtrate can be directly purified by column chromatography on silica gel.

One-pot reactions of other substrates were investigated with these additives (Table 2). The reaction tolerated a variety of aromatic and aliphatic ketones. In most cases, pyridine resulted in a shorter reaction time and higher yield but lower *E* selectivity than DMF or TMEDA.<sup>19,20</sup> In most cases, high *E/Z* selectivity (>95:5) was observed with DMF and TMEDA additives. When propiophenone was used as aldol acceptor for the reaction with methyl isopropyl ketone, low *E/Z* selectivities were observed (Scheme 1).


On the basis of the above observation for the *E/Z* selectivity, we speculated that the stereochemical pathway would be different for the reaction with pyridine versus that with DMF or TMEDA. To verify this hypothesis, several control experiments were performed (Table 3). First, TMEDA or pyridine, and then (*E*)- or (*Z*)-enone (Ar = *p*- $\text{ClC}_6\text{H}_4$ , isolated by chromatography after one-pot reaction) were added to a mixture of  $\text{TiCl}_4$ ,  $\text{Bu}_3\text{N}$  and water in dichloromethane.<sup>21</sup> With TMEDA (Table 3, entries 1 and 2), the *E/Z* ratio became 97:3 irrespective of the geometry of the starting enone. However, with pyridine (Table 3, entries 3 and 4), the *E/Z* ratio of the starting enone was almost retained. Thus, the *E/Z* selectivity is controlled thermodynamically in the former case and kinetically in the latter case. Second, hydrogen chloride, a possible side product of the one-pot reaction, also caused isomerization (Table 3,

Table 2. Substrate Scope<sup>a</sup>

entry	R <sup>1</sup>	R <sup>2</sup>	additive <sup>c</sup>	time (h)	yield (%) <sup>b</sup>	<i>E/Z</i> <sup>d</sup>
1	Ph	Ph	D	20	45	98:2
2	Ph	Ph	T	2	64	97:3
3	Ph	Ph	P	2	67	87:13
4	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	Ph	P	2	54	90:10
5	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	Ph	P	2	68	92:8
6 <sup>e</sup>	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	Ph	P	2	87	92:8
7	Ph	<i>i</i> -Pr	D	24	45	86:14
8	Ph	<i>i</i> -Pr	T	1	86	87:13
9	Ph	<i>i</i> -Pr	P	2	73	88:12
10	<i>c</i> -Pr	Ph	D	7	19	95:5
11	<i>c</i> -Pr	Ph	T	20	38	95:5
12	<i>c</i> -Pr	Ph	P	1	77	89:11
13	<i>c</i> -Hex	Ph	D	20	73	97:3
14	<i>c</i> -Hex	Ph	T	20	72	97:3
15	<i>c</i> -Hex	Ph	P	1	86	85:15
16	<i>t</i> -Bu	Ph	D	20	38	98:2
17	<i>t</i> -Bu	Ph	T	20	78	98:2
18	<i>t</i> -Bu	Ph	P	1	87	86:14
19	<i>i</i> -Pr	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	D	4	70	>99:1
20	<i>i</i> -Pr	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	T	18	62	98:2
21	<i>i</i> -Pr	<i>p</i> -MeOC <sub>6</sub> H <sub>4</sub>	P	1	67	82:18
22	<i>i</i> -Pr	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	D	20	38	96:4
23	<i>i</i> -Pr	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	T	20	71	96:4
24	<i>i</i> -Pr	<i>p</i> -ClC <sub>6</sub> H <sub>4</sub>	P	1	67	73:27
25	<i>i</i> -Pr	<i>t</i> -Bu	P	2	99	73:27


<sup>a</sup>The reaction was carried out by using  $\text{R}^1\text{COCH}_3$  (0.5 mmol),  $\text{TiCl}_4$  (1.1 mmol),  $\text{Bu}_3\text{N}$  (1.2 mmol),  $\text{R}^2\text{COCH}_3$  (0.5 mmol), and additive [DMF (2.5 mmol), TMEDA (0.5 mmol), or pyridine (2.5 mmol)] in dichloromethane (1 mL). <sup>b</sup>Isolated yield. <sup>c</sup>D: DMF (5.0 equiv); T: TMEDA (1.0 equiv); P: pyridine (5.0 equiv). <sup>d</sup>Determined by <sup>1</sup>H NMR spectroscopic analysis of the crude product. <sup>e</sup>The aldol step was carried out at  $-45^\circ\text{C}$  for 1.0 h.

Scheme 1. Propiophenone as Aldol Acceptor



entry 5), which suggested that the excess of pyridine (5.0 equiv) trapped HCl and prevented the isomerization (equilibration).


The one-pot reaction was also applicable to the preparation of  $\beta,\beta$ -disubstituted  $\alpha,\beta$ -unsaturated carboxylic acid derivatives (Scheme 2). The reaction of *S*-phenyl thioacetate<sup>22</sup> with acetophenone using TMEDA or pyridine in the elimination step afforded the desired unsaturated thioester in good yield with good *E* selectivity. For the reaction with methyl isopropyl ketone, pyridine provided a much higher yield than TMEDA. In addition, the unsaturated thioester product (100% *E*, isolated by column chromatography) could be transformed to the corresponding *N*-methyl amide, aldehyde, or ethyl ketone by

Table 3. Control Experiments<sup>a</sup>


entry	E/Z (start)	additive (equiv)	time (h)	E/Z (end) <sup>b</sup>
1	98:2	TMEDA (1.0)	12	97:3
2	0:100	TMEDA (1.0)	12	97:3
3	98:2	pyridine (5.0)	1	98:2
4	0:100	pyridine (5.0)	1	3:97
5	0:100	HCl (1.0) <sup>c</sup>	1	97:3

<sup>a</sup>Additive and enone were successively added to a mixture of TiCl<sub>4</sub> (1.1 equiv), Bu<sub>3</sub>N (1.2 equiv) and water (1.0 equiv) in dichloromethane. <sup>b</sup>Determined by <sup>1</sup>H NMR spectroscopic analysis of the crude product. <sup>c</sup>Hydrogen chloride in Et<sub>2</sub>O was used instead of a mixture of TiCl<sub>4</sub>, Bu<sub>3</sub>N, and water.

Scheme 2. Transformation of S-Phenyl Thioacetate



treatment with MeNH<sub>2</sub>,<sup>23</sup> diisobutylaluminum hydride (DIBAL-H),<sup>24</sup> and EtZnI/palladium catalyst,<sup>25</sup> respectively.<sup>26</sup> It should be noted that these products are difficult to obtain from the one-pot reaction of the corresponding carbonyl compounds (i.e., *N*-methylacetamide, acetaldehyde, and 2-butanone).


The one-pot method was applied to the one-pot synthesis of simple natural products, *ar*-atlantone<sup>27</sup> and  $\alpha$ -atlantone<sup>28</sup> (Scheme 3). The Ti-aldol reaction of mesityl oxide with the required ketones followed by elimination with pyridine provided *ar*-atlantone and  $\alpha$ -atlantone in good yield and *E* selectivity.

In summary, we have demonstrated a one-pot method to synthesize  $\beta,\beta$ -disubstituted  $\alpha,\beta$ -unsaturated carbonyl compounds by sequential TiCl<sub>4</sub>-promoted aldol reaction to simple ketones and base-promoted elimination.

## EXPERIMENTAL SECTION

**General Methods.** Dry dichloromethane (dehydrated) was stored over 4-Å MS prior to use. Titanium(IV) chloride was distilled under reduced pressure. Tributylamine, *N,N,N',N'*-Tetramethylethylenediamine (TMEDA), and pyridine were distilled from CaH<sub>2</sub>. *N,N*-Dimethylformamide (DMF) was stored over 4 Å MS prior to use. All

Scheme 3. Synthesis of Atlantones



other solvents and chemicals were purified based on standard procedures. Melting points (mp) were uncorrected. <sup>1</sup>H and <sup>13</sup>C{<sup>1</sup>H} NMR spectra were measured in CDCl<sub>3</sub> with 400 or 600 MHz spectrometer. Tetramethylsilane (TMS) ( $\delta$  = 0 ppm) and CDCl<sub>3</sub> ( $\delta$  = 77.0 ppm) served as internal standards for <sup>1</sup>H and <sup>13</sup>C NMR, respectively. Infrared spectra were recorded on an FT-IR spectrometer. High-resolution mass spectra were recorded on a double-focusing magnetic-sector mass analyzer operating in a FAB mode. Thin-layer chromatography (TLC) was visualized with UV light, phosphomolybdic acid and/or anisaldehyde. Column chromatography was performed using silica gel (spherical, neutral, 63–210 nm). The reactions under anhydrous conditions were carried out using oven- and heating gun-dried glassware with a rubber septum and a magnetic stirring bar under argon atmosphere.

**Typical Procedure: One-Pot Synthesis of  $\beta,\beta$ -Disubstituted Enone (Table 1, Entry 12).** TiCl<sub>4</sub> (2.82 M in CH<sub>2</sub>Cl<sub>2</sub>; 0.39 mL) and Bu<sub>3</sub>N (0.29 mL, 1.2 mmol) were successively added to a stirred solution of methyl isopropyl ketone (87.0 mg, 1.0 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.0 mL) at –78 °C. After 30 min, acetophenone (119.7 mg, 1.0 mmol) was added to the mixture at –78 °C. Formation of the aldol product was detected by TLC analysis after 1 h. Then pyridine (0.40 mL, 5.0 mmol) was added at –78 °C and the reaction mixture was warmed to room temperature. After being stirred for 5 h, the reaction mixture was diluted with Et<sub>2</sub>O (5 mL) and hexane (5 mL). The mixture was filtered through a Celite pad and the filtrate was concentrated. The residue was purified by column chromatography on silica gel (hexane/acetone = 30:1) to give 2-methyl-5-phenylhex-4-en-3-one (163.8 mg, 87% yield, E/Z = 86:14).

**(E)-2-Methyl-5-phenylhex-4-en-3-one.**<sup>29</sup> TLC *R*<sub>f</sub> 0.38 (hexane/acetone = 10:1, stained dark green with anisaldehyde). <sup>1</sup>H NMR (400 MHz)  $\delta$  = 1.17 (d, *J* = 7.0 Hz, 6H), 2.55 (s, 3H), 2.73 (sept, *J* = 7.0 Hz, 1H), 6.55 (s, 1H), 7.33–7.43 (m, 3H), 7.44–7.58 (m, 2H).

**(Z)-2-Methyl-5-phenylhex-4-en-3-one.**<sup>29</sup> TLC *R*<sub>f</sub> 0.35 (hexane/acetone = 10:1, stained dark green with anisaldehyde). <sup>1</sup>H NMR (400 MHz)  $\delta$  = 0.99 (d, *J* = 6.9 Hz, 6H), 2.18 (d, *J* = 1.4 Hz, 3H), 2.45 (sept, *J* = 6.9 Hz, 1H), 6.20 (q, *J* = 1.4 Hz, 1H), 7.13–7.20 (m, 2H), 7.29–7.36 (m, 3H).

**(E)-1,3-Diphenylbut-2-en-1-one.**<sup>3f</sup> According to the typical procedure, the reaction of acetophenone (57.7 mg) and acetophenone (60.1 mg) at rt for 2 h gave 1,3-diphenylbut-2-en-1-one (75.1 mg, 67%, E/Z = 87:13). TLC *R*<sub>f</sub> 0.30 (hexane/AcOEt = 10:1, stained green with anisaldehyde). <sup>1</sup>H NMR (400 MHz)  $\delta$  = 2.60 (d, *J* = 0.9 Hz, 3H), 7.17 (q, *J* = 0.9 Hz, 1H), 7.37–7.45 (m, 3H), 7.47 (t, *J* = 7.8 Hz, 2H), 7.53–7.59 (m, 3H), 7.98–8.02 (m, 2H).

**(Z)-1,3-Diphenylbut-2-en-1-one.**<sup>3f</sup> TLC *R*<sub>f</sub> 0.24 (hexane/AcOEt = 10:1, stained green with anisaldehyde). <sup>1</sup>H NMR (400 MHz)  $\delta$  = 2.32 (s, 3H), 6.69 (brs, 1H), 7.14–7.27 (m, 5H), 7.34 (t, *J* = 7.8 Hz, 2H), 7.44 (t, *J* = 7.8 Hz, 1H), 7.83 (d, *J* = 7.8 Hz, 2H).

**(E)-1-(4-Methoxyphenyl)-3-phenylbut-2-en-1-one.**<sup>3f</sup> According to the typical procedure, the reaction of 4-methoxyacetophenone (150.2 mg) and acetophenone (123.2 mg) at rt for 2 h gave 1-(4-methoxyphenyl)-3-phenylbut-2-en-1-one (136.2 mg, 54%, E/Z = 90:10). TLC *R*<sub>f</sub> 0.36 (hexane/AcOEt = 7:1, stained yellow green with anisaldehyde). <sup>1</sup>H NMR (400 MHz)  $\delta$  = 2.57 (d, *J* = 1.4 Hz, 3H), 3.88 (s, 3H), 6.95 (d, *J* = 8.7 Hz, 2H), 7.12 (q, *J* = 1.4 Hz, 1H), 7.38–7.45 (m, 3H), 7.55–7.59 (m, 2H), 8.00 (d, *J* = 8.7 Hz, 2H).

**(E)-1-(4-Chlorophenyl)-3-phenylbut-2-en-1-one.**  $\text{TiCl}_4$  (3.0 M  $\text{CH}_2\text{Cl}_2$ ; 0.18 mL) and  $\text{Bu}_3\text{N}$  (0.14 mL, 0.6 mmol) were successively added to a stirred solution of 4-chloroacetophenone (71.2 mg, 0.46 mmol) in  $\text{CH}_2\text{Cl}_2$  (1.0 mL) at  $-78^\circ\text{C}$ . After 30 min, acetophenone (60.9 mg, 0.51 mmol) was added to the mixture at  $-78^\circ\text{C}$ . After being stirred at  $-45^\circ\text{C}$  for 1 h, pyridine (0.20 mL, 2.5 mmol) was added at  $-45^\circ\text{C}$  and the reaction mixture was warmed to room temperature. After being stirred for 2 h, the reaction mixture was diluted with  $\text{Et}_2\text{O}$  (5 mL) and hexane (5 mL). The mixture was filtered through a Celite pad and the filtrate was concentrated. The residue was purified by column chromatography on silica gel (hexane/AcOEt = 30/1) to give 1-(4-chlorophenyl)-3-phenylbut-2-en-1-one (102.7 mg, 87%,  $E/Z$  = 92:8). TLC  $R_f$  0.42 (hexane/AcOEt = 7:1, stained orange with anisaldehyde). IR (film) 3061, 1657, 1593, 1211, 1090, 762  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 2.60 (d,  $J$  = 1.4 Hz, 3H), 7.12 (q,  $J$  = 1.4 Hz, 1H), 7.40–7.47 (m, 5H), 7.55–7.59 (m, 2H), 7.93 (d,  $J$  = 8.2 Hz, 2H).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  = 19.0, 121.4, 126.5, 128.6, 128.8, 129.3, 129.7, 137.6, 138.9, 142.6, 156.0, 190.4. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  257 ( $\text{M} + \text{H}^+$ , 100), 220 ( $\text{M}^+ - \text{Cl}$ , 16), 139 ( $\text{C}_6\text{H}_4\text{CO}^+$ , 48). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{16}\text{H}_{14}\text{OCl}$  ( $\text{M} + \text{H}^+$ ) 257.0733, found 257.0732.

**(E)-3,4-Dimethyl-1-phenylpent-2-en-1-one.**<sup>6</sup> According to the typical procedure, the reaction of acetophenone (115.4 mg) and methyl isopropyl ketone (88.9 mg) at  $-78^\circ\text{C}$  for 2 h gave 3,4-dimethyl-1-phenyl-2-penten-1-one (131.9 mg, 73%,  $E/Z$  = 88:12). TLC  $R_f$  0.18 (hexane/ $\text{CH}_2\text{Cl}_2$  = 2:1, stained orange with anisaldehyde).  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 1.15 (d,  $J$  = 7.2 Hz, 6H), 2.17 (s, 3H), 2.49 (sept,  $J$  = 7.2 Hz, 1H), 6.74 (s, 1H), 7.43–7.47 (m, 2H), 7.47–7.56 (m, 1H), 7.92 (d,  $J$  = 7.2 Hz, 2H).

**(Z)-3,4-Dimethyl-1-phenyl-2-penten-1-one.**<sup>6</sup> TLC  $R_f$  0.24 (hexane/ $\text{CH}_2\text{Cl}_2$  = 2:1, stained orange with anisaldehyde).  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 1.08 (d,  $J$  = 7.0 Hz, 6H), 1.92 (s, 3H), 3.78 (sept, 1H), 6.61 (s, 1H), 7.40–7.50 (m, 2H), 7.50–7.56 (m, 1H), 7.93 (d,  $J$  = 7.3 Hz, 2H).

**(E)-1-Cyclopropyl-3-phenylbut-2-en-1-one.** According to the typical procedure, the reaction of cyclopropyl methyl ketone (46.2 mg) and acetophenone (60.7 mg) at rt for 1 h gave 1-cyclopropyl-3-phenylbut-2-en-1-one (71.7 mg, 77%,  $E/Z$  = 89:11). TLC  $R_f$  0.49 (hexane/AcOEt = 10:1, stained dark green with anisaldehyde). IR (ATR) 3008, 1668, 1597, 1574, 1381, 1102, 1094, 974, 895, 760, 693  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (600 MHz)  $\delta$  = 0.89–0.94 (m, 2H), 1.09–1.13 (m, 2H), 2.03–2.08 (m, 1H), 2.54 (d,  $J$  = 1.3 Hz, 3H), 6.66 (q,  $J$  = 1.3 Hz, 3H), 7.34–7.41 (m, 3H), 7.49–7.52 (m, 2H).  $^{13}\text{C}$  NMR (150 MHz)  $\delta$  = 11.2, 18.5, 22.8, 124.9, 126.5, 128.5, 128.9, 142.7, 152.9, 201.0. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  187 ( $\text{M} + \text{H}^+$ , 100), 145 ( $\text{PhMeC} = \text{CHCO}^+$ , 33). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{13}\text{H}_{15}\text{O}$  ( $\text{M} + \text{H}^+$ ) 187.1123, found 187.1119.

**(E)-1-Cyclohexyl-3-phenylbut-2-en-1-one.** According to the typical procedure, the reaction of cyclohexyl methyl ketone (60.5 mg) and methyl acetophenone (61.0 mg) at rt for 1 h gave 1-cyclohexyl-3-phenylbut-2-en-1-one (94.1 mg, 86%,  $E/Z$  = 85:15). TLC  $R_f$  0.59 (hexane/acetone = 10:1, stained dark green with anisaldehyde). IR (ATR) 2926, 2852, 1678, 1598, 1573, 1446, 1145, 966, 757, 694  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 1.15–1.46 (m, 5H), 1.60–1.72 (m, 1H), 1.77–1.85 (m, 2H), 1.86–1.94 (m, 2H), 2.40–2.49 (m, 1H), 2.53 (d,  $J$  = 1.4 Hz, 3H), 6.54 (q,  $J$  = 1.4 Hz, 1H), 7.34–7.42 (m, 3H), 7.46–7.51 (m, 2H).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  = 18.4, 25.8, 25.9, 28.6, 52.1, 123.6, 126.4, 128.5, 128.9, 142.8, 154.0, 204.6. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  229 ( $\text{M} + \text{H}^+$ , 96), 145 ( $\text{PhMeC} = \text{CHCO}^+$ , 100). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{16}\text{H}_{21}\text{O}$  ( $\text{M} + \text{H}^+$ ) 229.1592, found 229.1591.

**(E)-2,2-Dimethyl-5-phenylhex-4-en-3-one.** According to the typical procedure, the reaction of *tert*-butyl methyl ketone (49.4 mg) and acetophenone (60.0 mg) at rt for 1 h gave 2,2-dimethyl-5-phenylhex-4-en-3-one (86.3 mg, 87%,  $E/Z$  = 86:14). TLC  $R_f$  0.47 (hexane/acetone = 10:1, stained black with anisaldehyde). IR (ATR) 2966, 1674, 1597, 1573, 1446, 1088, 958, 764, 694  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 1.21 (s, 9H), 2.50 (d,  $J$  = 1.4 Hz, 3H), 6.74 (q,  $J$  = 1.4 Hz, 1H), 7.34–7.42 (m, 3H), 7.46–7.51 (m, 2H).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  = 18.6, 26.6, 44.2, 120.8, 126.4, 128.5, 128.8, 143.2, 154.1,

206.7. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  203 ( $\text{M} + \text{H}^+$ , 61), 145 ( $\text{PhMeC} = \text{CHCO}^+$ , 100). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{14}\text{H}_{19}\text{O}$  ( $\text{M} + \text{H}^+$ ) 203.1436, found 203.1433.

**(E)-5-(4-Methoxyphenyl)-2-methylhex-4-en-3-one.** According to the typical procedure, the reaction of methyl isopropyl ketone (40.7 mg) and *p*-methoxyacetophenone (74.1 mg) at rt for 1 h gave 5-(4-methoxyphenyl)-2-methylhex-4-en-3-one (68.7 mg, 67%,  $E/Z$  = 82:18). TLC  $R_f$  0.47 (hexane/AcOEt = 7:1, stained dark green with anisaldehyde). IR (film) 2966, 1680, 1593, 1512, 1252, 1180, 829  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 1.16 (d,  $J$  = 6.9 Hz, 6H), 2.54 (d,  $J$  = 1.4 Hz, 3H), 2.71 (sept,  $J$  = 6.9 Hz, 1H), 3.83 (s, 3H), 6.53 (q,  $J$  = 1.4 Hz, 1H), 6.90 (d,  $J$  = 8.7 Hz, 2H), 7.47 (d,  $J$  = 8.7 Hz, 2H).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  = 18.2, 18.5, 42.0, 55.3, 113.8, 121.6, 127.8, 134.8, 153.8, 160.4, 205.0. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  219 ( $\text{M} + \text{H}^+$ , 75), 175 ( $\text{ArMeC} = \text{CHCO}^+$ , 100). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{14}\text{H}_{19}\text{O}_2$  ( $\text{M} + \text{H}^+$ ) 219.1385, found 219.1384.

**(E)-5-(4-Chlorophenyl)-2-methylhex-4-en-3-one.** According to the typical procedure, the reaction of methyl isopropyl ketone (41.3 mg) and *p*-chloroacetophenone (72.8 mg) at rt for 1 h gave 5-(4-chlorophenyl)-2-methylhex-4-en-3-one (70.1 mg, 67%,  $E/Z$  = 73:27). TLC  $R_f$  0.57 (hexane/AcOEt = 10:1, stained dark green with anisaldehyde). IR (ATR) 2968, 1682, 1600, 1562, 1489, 1071, 1063, 1011, 967, 821, 770  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 1.15 (d,  $J$  = 6.9 Hz, 6H), 2.51 (d,  $J$  = 1.4 Hz, 3H), 2.71 (sept,  $J$  = 6.9 Hz, 1H), 6.52 (q,  $J$  = 1.4 Hz, 1H), 7.35 (d,  $J$  = 8.7 Hz, 2H), 7.42 (d,  $J$  = 8.7 Hz, 2H).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  = 18.1, 18.2, 41.9, 123.3, 127.6, 128.5, 134.8, 141.0, 152.5, 204.7. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  223 ( $\text{M} + \text{H}^+$ , 97), 179 ( $\text{ArMeC} = \text{CHCO}^+$ , 100). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{13}\text{H}_{16}\text{OCl}$  ( $\text{M} + \text{H}^+$ ) 223.0890, found 223.0895.

**(Z)-5-(4-Chlorophenyl)-2-methylhex-4-en-3-one.** TLC  $R_f$  0.49 (hexane/AcOEt = 10:1, stained dark green with anisaldehyde). IR (ATR) 2970, 1689, 1616, 1594, 1490, 1092, 1013, 843, 823, 765  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (600 MHz)  $\delta$  = 1.02 (d,  $J$  = 6.9 Hz, 6H), 2.15 (d,  $J$  = 1.4 Hz, 3H), 2.51 (sept,  $J$  = 6.9 Hz, 1H), 6.24 (q,  $J$  = 1.4 Hz, 1H), 7.11 (d,  $J$  = 8.3 Hz, 2H), 7.30 (d,  $J$  = 8.3 Hz, 2H).  $^{13}\text{C}$  NMR (150 MHz)  $\delta$  = 18.2, 27.0, 40.8, 125.4, 128.3, 128.4, 133.8, 139.4, 151.2, 204.6. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  223 ( $\text{M} + \text{H}^+$ , 100), 179 ( $\text{ArMeC} = \text{CHCO}^+$ , 95). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{13}\text{H}_{16}\text{OCl}$  ( $\text{M} + \text{H}^+$ ) 223.0890, found 223.0892.

**(E)-2,5,6,6-Tetramethylhept-4-en-3-one.**<sup>5b</sup> According to the typical procedure, the reaction of methyl isopropyl ketone (40.5 mg) and *tert*-butyl methyl ketone (48.3 mg) at rt for 2 h gave 2,5,6,6-tetramethylhept-4-en-3-one (79.1 mg, 99%,  $E/Z$  = 73:27). TLC  $R_f$  0.50 (hexane/AcOEt = 10:1, stained black with anisaldehyde).  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 1.09 (s, 9H), 1.12 (d,  $J$  = 6.9 Hz, 6H), 2.12 (s, 3H), 2.63 (sept,  $J$  = 6.9 Hz, 1H), 6.17 (s, 1H).

**(E)-2-Methyl-5-phenylhept-4-en-3-one.** According to the typical procedure, the reaction of methyl isopropyl ketone (44.1 mg) and propiophenone (69.0 mg) with TMEDA (150  $\mu\text{L}$ ) at rt for 24 h gave 2-methyl-5-phenylhept-4-en-3-one (88.8 mg, 86%,  $E/Z$  = 56:44). TLC  $R_f$  0.52 (hexane/AcOEt = 10:1, stained dark green with anisaldehyde). IR (ATR) 2968, 1682, 1595, 1574, 1463, 1445, 1068, 1056, 764, 695.  $^1\text{H}$  NMR (600 MHz)  $\delta$  = 1.06 (t,  $J$  = 7.2 Hz, 3H), 1.15 (d,  $J$  = 7.2 Hz, 6H), 2.70 (sept,  $J$  = 7.2 Hz, 1H), 3.04 (q,  $J$  = 7.2 Hz, 2H), 6.42 (s, 1H), 7.37–7.39 (m, 3H), 7.44–7.46 (m, 2H).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  = 13.6, 18.4, 24.7, 42.0, 123.0, 126.8, 128.5, 128.8, 141.7, 160.7, 204.8. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  203 ( $\text{M} + \text{H}^+$ , 69), 159 ( $\text{PhEtC} = \text{CHCO}^+$ , 100), 77 ( $\text{Ph}^+$ , 87). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{14}\text{H}_{19}\text{O}$  ( $\text{M} + \text{H}^+$ ) 203.1436, found 203.1427.

**(Z)-2-Methyl-5-phenylhept-4-en-3-one.**  $R_f$  0.45 (hexane/AcOEt = 10:1, stained dark green with anisaldehyde). IR (ATR) 2968, 1693, 1672, 1618, 1598, 1464, 1442, 1381, 1092, 1013, 832, 765, 697.  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 0.97 (d,  $J$  = 7.2 Hz, 6H), 1.05 (d,  $J$  = 7.8 Hz, 3H), 2.38–2.52 (m, 3H), 6.15 (s, 1H), 7.10–7.16 (m, 2H), 7.26–7.38 (m, 3H).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  = 12.3, 18.3, 33.2, 40.4, 124.0, 127.2, 127.7, 128.1, 140.5, 157.6, 206.2. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  203 ( $\text{M} + \text{H}^+$ , 100), 159 ( $\text{PhEtC} = \text{CHCO}^+$ , 100), 77 ( $\text{Ph}^+$ , 90), 71 ( $^i\text{PrCO}^+$ , 85). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{14}\text{H}_{19}\text{O}$  ( $\text{M} + \text{H}^+$ ) 203.1436, found 203.1442.

**S-Phenyl (E)-3-phenylbut-2-enethioate.** According to the typical procedure, the reaction of S-phenyl thioacetate (76.3 mg) and acetophenone (67.2 mg) at rt for 5 h gave S-phenyl 3-phenylbut-2-enethioate (108.3 mg, 85%, *E/Z* = 91:9). TLC  $R_f$  0.57 (hexane/AcOEt = 10:1, stained pale red with anisaldehyde). mp 63–64 °C. IR (ATR) 3056, 1682, 1597, 1572, 1438, 1047, 936, 833, 735, 690, 566  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 2.55 (d,  $J$  = 1.4 Hz, 3H), 6.49 (q,  $J$  = 1.4 Hz, 1H), 7.37–7.54 (m, 10H).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  = 18.9, 122.6, 126.6, 128.3, 128.6, 129.1, 129.3, 129.5, 134.6, 141.5, 154.1, 187.7. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  255 ( $\text{M} + \text{H}^+$ , 19), 145 ( $\text{PhMeC}=\text{CHCO}^+$ , 100). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{16}\text{H}_{15}\text{OS}$  ( $\text{M} + \text{H}^+$ ) 255.0844, found 255.0847.

**S-Phenyl (E)-3,4-dimethylpent-2-enethioate.** According to the typical procedure, the reaction of S-phenyl thioacetate (79.0 mg) and methyl isopropyl ketone (41.0 mg) at rt for 20 h gave S-phenyl 3,4-dimethylpent-2-enethioate (102.5 mg, 97%, *E/Z* = 87:13). TLC  $R_f$  0.62 (hexane/ $\text{CH}_2\text{Cl}_2$  = 1:1, stained pale red with anisaldehyde). IR (ATR) 2964, 1683, 1616, 1440, 1031, 1023, 883, 830, 729, 688  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 1.09 (d,  $J$  = 6.9 Hz, 6H), 2.11 (s, 3H), 2.38 (sept,  $J$  = 6.9 Hz, 1H), 6.07 (s, 1H), 7.38–7.46 (m, 5H).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  = 17.6, 20.8, 38.2, 119.7, 128.5, 129.0, 129.1, 134.7, 164.8, 187.7. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  221 ( $\text{M} + \text{H}^+$ , 23), 111 ( $^i\text{PrMeC}=\text{CHCO}^+$ , 100). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{13}\text{H}_{17}\text{OS}$  ( $\text{M} + \text{H}^+$ ) 221.1000, found 221.1003.

**Control Experiments for *E/Z* Isomerization.** Table 3, entry 1 (from *E* isomer):  $\text{TiCl}_4$  (3.00 M in  $\text{CH}_2\text{Cl}_2$ ; 0.12 mL) was added to a stirred solution of  $\text{H}_2\text{O}$  (6.1  $\mu\text{L}$ , 0.34 mmol) and  $\text{Bu}_3\text{N}$  (0.1 mL, 0.41 mmol) in  $\text{CH}_2\text{Cl}_2$  (0.68 mL) at  $-78$  °C. After 0.5 h, TMEDA (0.05 mL, 0.34 mmol) was added at  $-78$  °C and the reaction mixture was warmed to room temperature. After 0.5 h, (*E*)-5-(4-chlorophenyl)-2-methylhex-4-en-3-one (76.7 mg, 0.34 mmol, *E/Z* = 97:3) was added to the mixture. After being stirred for 12 h, the reaction mixture was diluted with  $\text{Et}_2\text{O}$  (5 mL) and hexane (5 mL). The mixture was filtered through a Celite pad. The filtrate was concentrated to give the sufficiently pure crude product, which  $^1\text{H}$  NMR spectroscopic analysis indicated the *E/Z* ratio to be 97:3.

Table 3, entry 2 (from *Z* isomer): Similarly to the above procedure, (*Z*)-5-(4-chlorophenyl)-2-methylhex-4-en-3-one (6.8 mg, *E/Z* = 0:100) was treated with the mixture of  $\text{TiCl}_4$  (3.00 M  $\text{CH}_2\text{Cl}_2$ ; 0.01 mL),  $\text{H}_2\text{O}$  (0.5  $\mu\text{L}$ ),  $\text{Bu}_3\text{N}$  (8.5  $\mu\text{L}$ ) and TMEDA (4.5  $\mu\text{L}$ ) in  $\text{CH}_2\text{Cl}_2$  (0.06 mL) at room temperature for 12 h. The  $^1\text{H}$  NMR spectroscopic analysis of the crude product indicated the *E/Z* ratio to be 97:3.

**Transformation of  $\alpha,\beta$ -Unsaturated S-Phenyl Thioester to (*E*)-*N*-Methyl-3-phenylbut-2-enamide.** According to the reported procedure,<sup>23</sup> 40%  $\text{MeNH}_2$  in  $\text{H}_2\text{O}$  (12  $\mu\text{L}$ , 0.24 mmol) was added over 30 min to a stirred solution of S-phenyl (*E*)-3,4-dimethylpent-2-enethioate (50.9 mg, *E/Z* = 100:0) in MeOH (2.6 mL) at 0 °C. After 2 h, the reaction mixture was concentrated. The residue was purified by column chromatography on silica gel (hexane/AcOEt = 10:1) to give (*E*)-*N*-methyl-3-phenylbut-2-enamide (30.5 mg, 87%, *E/Z* = 100:0). TLC  $R_f$  0.14 (hexane/AcOEt = 10:1, stained white with anisaldehyde). mp 119–120 °C. IR (ATR) 3251, 3077, 2919, 1651, 1619, 1563, 1447, 1365, 1287, 1234, 875, 764, 689  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (600 MHz)  $\delta$  = 2.55 (d,  $J$  = 1.1 Hz, 3H), 2.89 (d,  $J$  = 8.7 Hz, 3H), 5.70 (brs, 1H), 6.00 (q,  $J$  = 1.1 Hz, 1H), 7.30–7.37 (m, 3H), 7.40–7.43 (m, 2H).  $^{13}\text{C}$  NMR (150 MHz)  $\delta$  = 17.6, 26.1, 119.8, 126.1, 128.4, 142.7, 150.5, 167.6. LRMS (FAB+,  $\text{CHCl}_3$ +NBA)  $m/z$  176 ( $\text{M} + \text{H}^+$ , 100), 145 ( $\text{PhMeC}=\text{CHCO}^+$ , 24). HRMS (FAB+,  $\text{CHCl}_3$ +NBA) Calcd for  $\text{C}_{11}\text{H}_{14}\text{NO}$  ( $\text{M} + \text{H}^+$ ) 176.1075, found 176.1077.

**Transformation of  $\alpha,\beta$ -Unsaturated S-Phenyl Thioester to (*E*)-3-Phenylbut-2-enal.** DIBAL-H (1.0 M in hexane, 0.30 mL) was added to a stirred solution of S-phenyl (*E*)-3,4-dimethylpent-2-enethioate (50.4 mg, *E/Z* = 100:0) in methylcyclohexane (2.0 mL) at  $-78$  °C. The reaction mixture was stirred at  $-78$  °C for 0.5 h and quenched with  $\text{NaOH}/\text{H}_2\text{O}/\text{EtOH}$  (0.3 g/1.25 mL/1.75 mL). The mixture was extracted with AcOEt (3  $\times$  15 mL). The combined organic layers were washed with brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated. The residue was purified by column chromatography on silica gel (hexane/AcOEt = 20:1) to give (*E*)-3-phenylbut-2-enal<sup>30</sup> (23.9 mg, 82%, *E/Z* = 93:7). TLC  $R_f$  0.47 (hexane/AcOEt = 10:1,

stained black with anisaldehyde).  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 2.58 (d,  $J$  = 1.4 Hz, 3H), 6.40 (dd,  $J$  = 7.8, 1.4 Hz, 1H), 7.36–7.48 (m, 3H), 7.50–7.60 (m, 2H), 10.19 (d,  $J$  = 7.8 Hz, 1H).

**Transformation of  $\alpha,\beta$ -Unsaturated S-Phenyl Thioester to (*E*)-5-Phenylhex-4-en-3-one.** According to the reported procedure with a modification,<sup>25</sup>  $\text{EtZnI}$  (0.50 M in THF/ $\text{Et}_2\text{O}$ /hexane; 1.2 mL, prepared from diethylzinc and iodine according to the reported procedure<sup>31</sup>) was added to a stirred solution of S-phenyl (*E*)-3,4-dimethylpent-2-enethioate (50.9 mg, *E/Z* = 100:0) and  $\text{PdCl}_2(\text{PPh}_3)_2$  (14.0 mg, 10 mol %) in toluene (0.67 mL) at room temperature. After being stirred for 2 h, the reaction mixture was diluted with AcOEt and concentrated. The residue was purified by column chromatography on silica gel (hexane/AcOEt = 30:1) to give (*E*)-5-phenylhex-4-en-3-one<sup>3f</sup> (31.3 mg, 90%, *E/Z* = 100:0). TLC  $R_f$  0.52 (hexane/AcOEt = 10:1, stained black with anisaldehyde).  $^1\text{H}$  NMR (400 MHz)  $\delta$  = 1.13 (t,  $J$  = 7.3 Hz, 3H), 2.55 (d,  $J$  = 0.9 Hz, 3H), 2.57 (q,  $J$  = 7.3 Hz, 2H), 6.50 (q,  $J$  = 0.9 Hz, 1H), 7.33–7.42 (m, 3H), 7.45–7.52 (m, 2H).

**Synthesis of *ar*-Atlantone.** According to the typical procedure, the reaction of mesityl oxide (98.4 mg) and *p*-methylacetophenone (134.2 mg) at rt for 1 h gave *ar*-atlantone<sup>27b</sup> (167.2 mg, 78%, *E/Z* = 91:9). TLC  $R_f$  0.57 (hexane/AcOEt = 7:1, stained black with anisaldehyde).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 1.92 (s, 3H), 2.22 (s, 3H), 2.37 (s, 3H), 2.56 (s, 3H), 6.19 (brs, 1H), 6.49 (s, 1H), 7.18 (d,  $J$  = 8.0 Hz, 2H), 7.40 (d,  $J$  = 8.0 Hz, 2H).

**Synthesis of  $\alpha$ -Atlantone.** According to the typical procedure, the reaction of mesityl oxide (51.2 mg) and 1-(4-methyl-3-cyclohexen-1-yl)ethanone<sup>28b</sup> (66.0 mg) at rt for 2 h gave  $\alpha$ -atlantone<sup>28c</sup> (74.4 mg, 71%, *E/Z* = 90:10). TLC  $R_f$  0.56 (hexane/AcOEt = 10:1, stained black with anisaldehyde).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  = 1.62 (s, 3H), 1.65–2.20 (m, 7H), 1.85 (s, 3H), 2.13 (s, 6H), 5.41 (brs, 1H), 6.06 (brs, 1H), 6.08 (brs, 1H).

## ■ ASSOCIATED CONTENT

### 📄 Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.5b01217.

$^1\text{H}$  NMR spectra of the known compounds (the products in Table 1, the products in entries 1–4, 7–9, and 25 of Table 2, (*E*)-3-phenylbut-2-enal and (*E*)-5-phenylhex-4-en-3-one in Scheme 2, *ar*-atlantone and  $\alpha$ -atlantone in Scheme 3).  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of the new compounds (the products in entries 5, 6 and 10–24 of Table 2, the products in Scheme 1, and the thioester products in Scheme 2). (PDF)

## ■ AUTHOR INFORMATION

### ✉ Corresponding Author

\*E-mail: msugiura@kumamoto-u.ac.jp.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work was partially supported by JSPS KAKENHI Grant Number 23590009 and 26460010.

## ■ REFERENCES

- (1) For selected examples, see: (a) Salomon, R. G.; Lal, K.; Mazza, S. M.; Zarate, E. A.; Youngs, W. J. *J. Am. Chem. Soc.* **1988**, *110*, 5213.
- (b) Salomon, R. G.; Mazza, S. M.; Lal, K. *J. Org. Chem.* **1989**, *54*, 1562.
- (c) Li, S.; Chiu, P. *Tetrahedron Lett.* **2008**, *49*, 1741.
- (d) Esumi, T.; Mori, T.; Zhao, M.; Toyota, M.; Fukuyama, Y. *Org. Lett.* **2010**, *12*, 888.
- (e) Esumi, T.; Yamamoto, C.; Fukuyama, Y. *Synlett* **2013**, *24*, 1845.
- (f) Ciesielski, J.; Gadon, V.; Frontier, A. J. *J. Org. Chem.* **2013**, *78*, 9541.
- (g) McGrath, K. P.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2014**,

53, 1910. (h) Radomkit, S.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2014**, *53*, 3387.

(2) Sugiura, M.; Sato, N.; Kotani, S.; Nakajima, M. *Chem. Commun.* **2008**, 4309.

(3) (a) Wadsworth, W. S., Jr.; Emmons, W. D. *J. Am. Chem. Soc.* **1961**, *83*, 1733. (b) Appella, D. H.; Moritani, Y.; Shintani, R.; Ferreira, E. M.; Buchwald, S. L. *J. Am. Chem. Soc.* **1999**, *121*, 9473. (c) Tsuchiya, Y.; Hamashima, Y.; Sodeoka, M. *Org. Lett.* **2006**, *8*, 4851. (d) Kanazawa, Y.; Tsuchiya, Y.; Kobayashi, K.; Shiomi, T.; Itoh, J.; Kikuchi, M.; Yamamoto, Y.; Nishiyama, H. *Chem. - Eur. J.* **2006**, *12*, 63. (e) Lu, S.-M.; Bolm, C. *Chem. - Eur. J.* **2008**, *14*, 7513. (f) Nishikawa, Y.; Yamamoto, H. *J. Am. Chem. Soc.* **2011**, *133*, 8432.

(4) (a) Trost, B. M.; Salzmann, T. N. *J. Am. Chem. Soc.* **1973**, *95*, 6840. (b) Trost, B. M.; Salzmann, T. N.; Hiroi, K. *J. Am. Chem. Soc.* **1976**, *98*, 4887.

(5) (a) Kashima, C.; Yamamoto, Y. *Heterocycles* **1982**, *19*, 1211. (b) Dieter, R. K.; Silks, L. A. *J. Org. Chem.* **1986**, *51*, 4687. (c) Bartoli, G.; Marcantoni, E.; Petrini, M.; Sambri, L. *Chem. - Eur. J.* **1996**, *2*, 913. (d) Lipshutz, B. H.; Amorelli, B.; Unger, J. B. *J. Am. Chem. Soc.* **2008**, *130*, 14378.

(6) Tanaka, Y.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2010**, *132*, 8862.

(7) (a) Sugawara, Y.; Yamada, W.; Yoshida, S.; Ikeno, T.; Yamada, T. *J. Am. Chem. Soc.* **2007**, *129*, 12902. (b) Yu, M.; Li, G.; Wang, S.; Zhang, L. *Adv. Synth. Catal.* **2007**, *349*, 871. (c) Engel, D. A.; Dudley, G. B. *Org. Lett.* **2006**, *8*, 4027. (d) Egi, M.; Yamaguchi, Y.; Fujiwara, N.; Akai, S. *Org. Lett.* **2008**, *10*, 1867. (e) Pennell, M. N.; Turner, P. G.; Sheppard, T. D. *Chem. - Eur. J.* **2012**, *18*, 4748.

(8) (a) Shibuya, M.; Ito, S.; Takahashi, M.; Iwabuchi, Y. *Org. Lett.* **2004**, *6*, 4303. (b) Li, J.; Tan, C.; Gong, J.; Yang, Z. *Org. Lett.* **2014**, *16*, 5370.

(9) Baker, D. B.; Gallagher, P. T.; Donohoe, T. J. *Tetrahedron* **2013**, *69*, 3690.

(10) For titanium-promoted aldol reactions between simple ketones, see: (a) Yoshida, Y.; Hayashi, R.; Sumihara, H.; Tanabe, Y. *Tetrahedron Lett.* **1997**, *38*, 8727. (b) Yoshida, Y.; Matsumoto, N.; Hamasaki, R.; Tanabe, Y. *Tetrahedron Lett.* **1999**, *40*, 4227. (c) Tanabe, Y.; Matsumoto, N.; Higashi, T.; Misaki, T.; Itoh, T.; Yamamoto, M.; Mitarai, K.; Nishii, Y. *Tetrahedron* **2002**, *58*, 8269.

(11) For titanium-promoted aldol reactions of phenyl esters or S-phenyl thioesters and simple ketones, see: (a) Tanabe, Y.; Matsumoto, N.; Funakoshi, S.; Manta, N. *Synlett* **2001**, 1959. (b) Nagase, R.; Matsumoto, N.; Hosomi, K.; Higashi, T.; Funakoshi, S.; Misaki, T.; Tanabe, Y. *Org. Biomol. Chem.* **2007**, *5*, 151. Tanabe and his co-workers also reported synthesis of tri- and tetrasubstituted  $\alpha,\beta$ -unsaturated esters by cross-coupling reactions of enol tosylates, see: (c) Nakatsuji, H.; Ueno, K.; Misaki, T.; Tanabe, Y. *Org. Lett.* **2008**, *10*, 2131. (d) Nakatsuji, H.; Nishikado, H.; Ueno, K.; Tanabe, Y. *Org. Lett.* **2009**, *11*, 4258. (e) Nishikado, H.; Nakatsuji, H.; Ueno, K.; Nagase, R.; Tanabe, Y. *Synlett* **2010**, 2087. (f) Ashida, Y.; Sato, Y.; Suzuki, T.; Ueno, K.; Kai, K.; Nakatsuji, H.; Tanabe, Y. *Chem. - Eur. J.* **2015**, *21*, 5934.

(12) For other aldol additions to simple ketones, see: (a) Nozaki, H.; Oshima, K.; Takai, K.; Ozawa, S. *Chem. Lett.* **1979**, 379. (b) Tsuji, J.; Yamamda, T.; Kaito, M.; Mandai, T. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1417. (c) Allan, J. F.; Henderson, K. W.; Kennedy, A. R. *Chem. Commun.* **1999**, 1325. (d) Cergol, K. M.; Turner, P.; Coster, M. J. *Tetrahedron Lett.* **2005**, *46*, 1505. (e) Cergol, K. M.; Jensen, P.; Turner, P.; Coster, M. J. *Chem. Commun.* **2007**, 1363. (f) Alcobarro, S.; Gómez-Palomino, A.; Solà, R.; Romea, P.; Urpí, F.; Font-Bardia, M. *Org. Lett.* **2014**, *16*, 584. For Mukaiyama aldol reactions to ketones, see: (g) Ishihara, K.; Hiraiwa, Y.; Yamamoto, H. *Synlett* **2001**, 1851. (h) Yanai, H.; Yoshino, Y.; Takahashi, A.; Taguchi, T. *J. Org. Chem.* **2010**, *75*, 5375.

(13) We have reported the first enantioselective cross-aldol reaction between simple ketones, see: (a) Aoki, S.; Kotani, S.; Sugiura, M.; Nakajima, M. *Chem. Commun.* **2012**, 48, 5524. For the first enantioselective aldol addition of silyl ketene acetals to simple ketones, see: (b) Denmark, S. E.; Fan, Y. *J. Am. Chem. Soc.* **2002**,

*124*, 4233. (c) Denmark, S. E.; Fan, Y.; Eastgate, M. D. *J. Org. Chem.* **2005**, *70*, 5235. For a review of catalytic enantioselective aldol additions to ketones, see: (d) Adachi, S.; Harada, T. *Eur. J. Org. Chem.* **2009**, *22*, 3661.

(14) Lower temperatures than rt (even 0 °C) for the elimination step significantly decreased the reaction rate and did not improve the *E/Z* selectivity.

(15) NMP has been used as additive for TiCl<sub>4</sub>-promoted asymmetric aldol reactions, see: (a) Crimmins, M. T.; She, J. *Synlett* **2004**, 1371. (b) Sreenithya, A.; Sunoj, R. B. *Org. Lett.* **2012**, *14*, 5752.

(16) When the reaction was initiated with Bu<sub>3</sub>N (2.2 equiv) at beginning, the yield of the desired enone was decreased to 52% yield.

(17) In this case, the desired enone, the aldol product (5-hydroxy-2-methyl-5-phenylhexan-3-one), and a self-aldol product (3-hydroxy-1,3-diphenylbutan-1-one) were obtained in 59, 15, and 21% yields, respectively, according to the <sup>1</sup>H NMR analysis of the crude product. When TMEDA (1.0 equiv) was used (Table 1, entry 10), no self-aldol product but the aldol product was obtained in 3% yield along with the desired enone. These results indicate that the excess amount of TMEDA suppressed the elimination and promoted the retro aldol reaction.

(18) Pyridine and DMAP have been used as effective additives for acetylation of aluminium hemiacetals (intermediates of DIBAL-H reduction of esters), see: (a) Dahanukar, V. H.; Rychnovsky, S. D. *J. Org. Chem.* **1996**, *61*, 8317. (b) Kopecky, D. J.; Rychnovsky, S. D. *J. Org. Chem.* **2000**, *65*, 191.

(19) In Table 2, entry 7,  $\beta,\gamma$ -unsaturated ketone 3,4-dimethyl-1-phenylpent-3-en-1-one was obtained in 38% yield, which indicated that DMF did not deprotonate the carbonyl  $\alpha$ -proton. In Table 2, entry 10, ring-opened product (*E*)-7-chloro-2-phenylhept-2-en-4-one was obtained in 57% yield.

(20) The configurations of the known enone products were assigned on comparison with the literature data, and those of the unknown enones were determined by analogy.

(21) Water was added to reproduce the reaction conditions after the aldol reaction followed by the elimination of water.

(22) The reaction of phenyl acetate resulted in self-Claisen condensation. Tanabe et al. did not employ phenyl acetate and S-phenyl thioacetate for their TiCl<sub>4</sub>-promoted aldol reaction, see ref 11.

(23) Bae, H. Y.; Sim, J. H.; Lee, J.-W.; List, B.; Song, C. E. *Angew. Chem., Int. Ed.* **2013**, *52*, 12143.

(24) Corey, E. J.; Huang, H.-C. *Tetrahedron Lett.* **1989**, *30*, 5235.

(25) (a) Fukuyama, T.; Tokuyama, H. *Aldrichimica Acta* **2004**, *37*, 87. (b) Tokuyama, H.; Yokoshima, S.; Yamashita, T.; Fukuyama, T. *Tetrahedron Lett.* **1998**, *39*, 3189. (c) Zhou, G.; Lim, D.; Coltart, D. M. *Org. Lett.* **2008**, *10*, 3809.

(26) To the best of our knowledge, these functional group transformations have not been applied to  $\beta,\beta$ -disubstituted  $\alpha,\beta$ -unsaturated S-phenyl thioesters before. Especially, our modification for the DIBAL-H reduction (use of methylcyclohexane as the solvent and workup under basic conditions) was found to be important for retaining the geometry of the starting material.

(27) Synthesis of *ar*-atlantone: (a) Sakai, T.; Miyata, K.; Ishikawa, M.; Takeda, A. *Tetrahedron Lett.* **1985**, *26*, 4727. (b) Strunz, G. M.; Ya, L. *Can. J. Chem.* **1992**, *70*, 1317.

(28) Synthesis of  $\alpha$ -atlantone: (a) Crawford, R. J.; Erman, W. F.; Broaddus, C. D. *J. Am. Chem. Soc.* **1972**, *94*, 4298. (b) Babler, J. H.; Olsen, D. O.; Arnold, W. H. *J. Org. Chem.* **1974**, *39*, 1656. (c) Andrianome, M.; Delmond, B. *J. Org. Chem.* **1988**, *53*, 542.

(29) Barbee, T. R.; Albizati, K. F. *J. Org. Chem.* **1991**, *56*, 6764.

(30) Takaya, J.; Sasano, K.; Iwasawa, N. *Org. Lett.* **2011**, *13*, 1698.

(31) Charette, A. B.; Gagnon, A.; Fournier, J.-F. *J. Am. Chem. Soc.* **2002**, *124*, 386.