

An Optimized Glutamate Receptor Photoswitch with Sensitized Azobenzene Isomerization

Marta Gascón-Moya,[†] Arnau Pejoan,[†] Mercè Izquierdo-Serra,[‡] Silvia Pittolo,[‡] Gisela Cabré,[†] Jordi Hernando,[†] Ramon Alibés,[†] Pau Gorostiza,^{*,‡,§,||} and Félix Busqué^{*,†}


[†]Departament de Química, Universitat Autònoma de Barcelona, 08193 Bellaterra, Spain

[‡]Institut de Bioenginyeria de Catalunya (IBEC), Barcelona, Spain

[§]Centro de Investigación Biomédica en Red en Bioingeniería, Biomateriales y Nanomedicina (CIBER-BBN), Zaragoza, Spain

^{||}Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

Supporting Information



ABSTRACT: A new azobenzene-based photoswitch, **2**, has been designed to enable optical control of ionotropic glutamate receptors in neurons via sensitized two-photon excitation with NIR light. In order to develop an efficient and versatile synthetic route for this molecule, a modular strategy is described which relies on the use of a new linear fully protected glutamate derivative stable in basic media. The resulting compound undergoes one-photon *trans*–*cis* photoisomerization via two different mechanisms: direct excitation of its azoaromatic unit and irradiation of the pyrene sensitizer, a well-known two-photon sensitive chromophore. Moreover, **2** presents large thermal stability of its *cis* isomer, in contrast to other two-photon responsive switches relying on the intrinsic nonlinear optical properties of push–pull substituted azobenzenes. As a result, the molecular system developed herein is a very promising candidate for evoking large photoinduced biological responses during the multiphoton operation of neuronal glutamate receptors with NIR light, which require accumulation of the protein-bound *cis* state of the switch upon repeated illumination.

INTRODUCTION

Neuroscience has been revolutionized in recent years by the emergence of optogenetic¹ and optochemical tools,² a series of techniques that allow the optical control of neural networks by means of light-sensitive proteins and photoresponsive small molecules. An important strategy in this field is the use of photoswitchable tethered ligands (PTLs), in which a pharmacologically active moiety is covalently attached to a neuronal receptor through a photoisomerizable tether.^{2,3} This

enables the activity of the resulting protein–PTL construct to be modulated upon photoisomerization of the switch, which brings the ligand closer to or away from the binding site and, as such, triggers a reversible light-induced recognition event and the subsequent biological response. Owing to their synthetic accessibility, excellent photochemical properties, and

Received: June 19, 2015

Published: September 28, 2015




Figure 1. (a) Structure of a MAG molecular switch composed of maleimide (green), azobenzene (red), and glutamate (blue) units.⁷ (b) Operation mode of a cysteine-tagged ionotropic glutamate receptor functionalized with a photoswitchable tethered ligand. Reversible isomerization of the azoaromatic core of the switch modulates the separation distance between the glutamate ligand and the binding site of the receptor to induce channel opening (*trans-to-cis*) and closing (*cis-to-trans*). In this work, an additional building block is introduced into the structure of the PTL, an antenna enabling two-photon sensitization of azobenzene isomerization (MAGA-type switch). (c) Structures of the MAGA photoswitched tethered ligands 1 and 2, whose additional sensitizer unit is shown in magenta.

large geometrical change between their photointerconvertible *trans* and *cis* configurations, azobenzene derivatives have become the most common light-responsive building blocks for the preparation of PTLs,⁴ and they have been exploited to optically control a number of different neuronal receptors.^{2,3,5}

Since glutamate is the major excitatory neurotransmitter in the central nervous system, glutamate receptors play a central role in the regulation of neural activity⁶ and are an important target of optochemical techniques. Much attention has therefore been focused in recent years on the development of PTLs for the light-induced operation of these proteins, especially in the case of the ionotropic glutamate receptors (iGluR) controlling the channel-regulated flow of ions through the cell membrane.^{2,3} The most successful approach toward this goal is based on maleimide–azobenzene–glutamate photoswitches (MAG, Figure 1a),⁷ which were originally designed to enable optical control of the kainate-type ionotropic glutamate receptor GluK2. In MAG compounds, a lateral maleimide unit is used to covalently bind the PTL to a cysteine residue genetically engineered into the ligand binding domain of the receptor, while its photoswitchable azobenzene core enables light control of the separation distance between the glutamate moiety and the binding pocket. In particular, *trans-to-cis* isomerization of the system approaches the ligand toward the binding site and induces glutamate recognition, which leads to ion channel opening and, eventually, to the generation of a photoinduced neuronal excitatory potential (Figure 1b).⁷ Afterward, irradiation or thermal relaxation of the metastable *cis* isomer allows reversion of the process and recovery of the initial closed state of the channel.⁷ For the first MAG switch developed (Figure 1a), irradiation with UV-violet ($\lambda_{trans-cis} \approx 340\text{--}400\text{ nm}$) and blue-green light ($\lambda_{cis-trans} \approx 440\text{--}580\text{ nm}$) were required to photochemically operate the tagged GluK2 receptors,⁷ while spontaneous *cis-to-trans* isomerization

occurred in the dark on the minute time scale ($\tau_{cis} = 17.7\text{ min}$)^{7b} owing to the large thermal stability of the *cis* isomer resulting from its symmetric 4,4'-diamide substitution pattern.⁴ Although some efforts have been devoted to shift MAG photoactivity to longer wavelengths,^{8,9} the application of these compounds *in vivo* is severely compromised by the reduced penetration depth¹⁰ and low axial resolution associated with the conventional one-photon excitation of azobenzene-based switches with UV–visible light. To overcome these drawbacks and take full advantage of the light-regulated activity of MAG-labeled receptors, multiphoton excitation with near-infrared (NIR) light would be highly desirable, since it provides submicrometric resolution in three dimensions,¹¹ deep penetration into tissue,¹² and patterned illumination.^{13,14} Unfortunately, the two-photon absorption cross sections of common azobenzene chromophores are rather low,¹⁵ which makes multiphoton operation of MAG PTLs under NIR irradiation a challenge.

To tackle this problem, we have recently introduced two different molecular strategies, which allowed for the first time the two-photon optical control of light-gated ionotropic glutamate receptors based on cysteine-tagged GluK2 and MAG switches (LiGluR-MAG).¹⁶ On the one hand, the intrinsic two-photon activity of MAG compounds was shown to significantly increase upon push–pull substitution of their azobenzene core, as subsequently demonstrated by Isacoff and co-workers for analogous PTLs.¹⁷ However, a decrease of the thermal stability of the *cis* state of these systems was concomitantly observed,^{16,17} which imposes a severe constraint on their photochemical behavior: the ligand-bound state of the LiGluR-MAG tether dissociates as soon as illumination ceases, since the *cis* liganding configuration of the switch back-isomerizes in the dark on the ms scale. While this might be advantageous for certain applications requiring fast operation of ionotropic glutamate receptors,¹⁶ rapid thermal


relaxation of the switch prevents building up a large population of the open state of the ion channel and, therefore, the obtaining of large photoresponse amplitudes. Alternatively, multiphoton operation of MAG-type PTLs can be attained by incorporating a photosensitizer unit to the structure of the switch, which should absorb NIR light via a two-photon process and efficiently transfer its electronic excitation energy to the azobenzene chromophore (Figure 1b). We named this new family of compounds maleimide–azobenzene–glutamate–antenna switches (MAGA),¹⁶ whose multiphoton response could then be disentangled from the substitution pattern of the azoaromatic core. Actually, the one- and two-photon sensitized isomerization of a MAGA-type PTL was already demonstrated in our previous work using a naphthalene antenna (1 in Figure 1c); however, this compound also suffered from low *cis* thermal stability owing to the push–pull substitution of the azobenzene core selected ($\tau_{cis} = 265$ ms at 298 K) and, consequently, displayed reduced multiphoton activity when conjugated to LiGluR.¹⁶ In view of this, we pursue herein the preparation of a new MAGA photoswitch featuring both sensitized isomerization using a two-photon absorber and long *cis* state lifetime in the dark. As such, this compound could be exploited to optically trigger the LiGluR receptor via multiphoton absorption of NIR light in neuroscience applications demanding large biological photoinduced signals (i.e., accumulation of the open state of the channel upon repeated excitation).

Figure 1c shows the structure of 2, the novel MAGA-type PTL targeted in this work. This compound was derived on account of two main design concepts. First, the use of a symmetrically substituted azobenzene chromophore displaying two similar *N*-amido substituents at positions 4 and 4', which should ensure a long thermal lifetime of its *cis* state at room temperature according to the previous results obtained for MAG⁷ and other related azoaromatic derivatives.¹⁸ Second, the selection of a proper multiphoton antenna which should (i) display large two-photon absorption under NIR irradiation and (ii) emit in the same spectral region where the azobenzene core absorbs, thus enabling efficient sensitization of the switch via resonance energy transfer (RET). Since the 4,4'-diamidoazobenzene core in 2 must present a large hypsochromic absorption shift with respect to the azoaromatic group of 1 (~70 nm),^{7,16,18} the naphthalene sensitizer of the latter compound did not fulfill the second of the requirements of our present design and could not be used as antenna for the development of switch 2. Instead, a monoalkylated pyrene derivative was chosen as the sensitizer. Although these type of dyes present moderate two-photon absorption cross sections,¹⁹ they are often used as two-photon fluorescent probes for labeling cell membranes in confocal fluorescence microscopy studies¹⁸ and can be detected even down to the single-molecule level under two-photon excitation.²⁰ More importantly, their emission spectra ($\lambda_{fl,max} \approx 390$ nm¹⁹) largely overlaps the absorption spectra of the $\pi \rightarrow \pi^*$ transition of *trans*-4,4'-diamidoazobenzenes ($\lambda_{abs,max} \approx 370$ nm¹⁸), a critical requirement for photosensitization of their *trans*-to-*cis* isomerization.

RESULTS AND DISCUSSION

Synthesis of Photoswitchable Ligand 2. A convergent synthetic strategy was devised for the preparation of 2, which relies on the use of five different building blocks (Scheme 1): the *N,N*-orthogonally diprotected L-lysine 3; *trans*-4,4'-


Scheme 1. Retrosynthetic Analysis Proposed for the Synthesis of New MAG Compound 2



diaminoazobenzene (4) as the symmetrically substituted azoaromatic core; the linear protected derivative of glutamate 5; 2-(pyren-1-yl)acetic acid (6) as a two-photon sensitizer; and maleimide derivative 7. Two key features differentiate our approach from the synthetic procedures previously reported for the synthesis of analogous MAG photoswitches.^{7–9} On the one hand, 3 is used as a central scaffold to which the distinct functional fragments of the target compound would be sequentially tethered, thus aiming to develop a modular route toward MAGA PTLs (or any other type of multifunctional ligand derived from the original maleimide–azobenzene–glutamate scheme). On the other hand, 5 was developed as an alternative glutamate precursor to typical cyclic pyroglutamate groups, providing an optimized strategy to achieve the final photoswitchable ligand in terms of both deprotection reaction conditions and intermediate manipulation.

Since all the other building blocks of 2 were commercially available or synthetically accessible via prior reported procedures, our initial task focused on the preparation of glutamate derivative 5. Although this compound has been recently used by us for the synthesis of 1,¹⁶ we describe herein the obtaining of 5 for the first time. The synthetic sequence commenced with the protection of the carboxylic acid and the nitrogen atom of commercial L-pyroglutamic acid as the corresponding *tert*-butyl ester and *tert*-butyl carbamate, respectively, to obtain known pyroglutamate 8²¹ in a 53% overall yield (Scheme 2). Next, this product was treated with LiHMDS in THF at -78 °C to form the corresponding enolate, which was allowed to react with allyl bromide at low temperature to afford a 3:1 diastereoisomeric mixture of already described allylated derivatives *trans* 9a and *cis* 9b.²² The major *trans* diastereoisomer 9a was isolated after purification by column chromatography in a 36% yield and used in the following steps of the synthesis. The relative configuration assignment of these two isomers was confirmed by comparison with the described NMR data.²² Thus, in the ¹H NMR spectrum of 9a, H-2 resonates at an upper field


Scheme 2. Synthesis of Key Intermediate 9a



compared to that of **9b** and shows constant couplings with H-3 protons being quite different (9.6 and 1.6 Hz), whereas the values of these constants are more similar in the *cis* isomer (9.3 and 5.7 Hz).

Subsequent lithium hydroxide ring opening of **9a** proceeded smoothly providing acid **10** in 93% yield (Scheme 3), which

Scheme 3. Synthesis of Glutamate Derivative 5



was transformed into the corresponding di-*tert*-butyl ester derivative **11** in 56% yield by treatment with *tert*-butyl 2,2,2-trichloroacetimidate in the presence of catalytic BF₃. The synthesis was continued with a cross-metathesis reaction of **11** with crotonaldehyde using the Hoveyda–Grubs second generation catalyst to afford aldehyde **12** in 89% yield, which was oxidized to the corresponding acid **13** in 77% yield following a Pinnick protocol with NaClO₂ in the presence of 2-methyl-2-butene as the scavenger of the hypochlorous acid generated. Direct metathesis reaction of intermediate **11** with acrylic acid afforded derivative **13** in lower yield (52%) than that obtained with the two consecutive transformations described above (69% overall yield for the two steps). Finally, standard hydrogenation using a palladium catalyst rendered almost quantitatively the target glutamate derivative **5**.

With glutamate derivative **5** in hand, we proceeded to prepare the final MAGA photoswitch **2** (Scheme 4) following

the modular synthetic strategy outlined above and previously implemented in our group to obtain **1**.¹⁶ Accordingly, the *N,N*-orthogonally diprotected L-lysine **3** was tethered to the symmetric azobenzene core **4** using the carbodiimide coupling reagent EDCI along with HOBT and DIPEA as base, which led to a chromatographically easily separable mixture of starting material and compound **14** in 85% yield (over consumed starting material). Removal of the *tert*-butyl carbamate protection of this compound by treatment with 37% HCl in MeOH for 1 h furnished amine **15** in 98% yield. Longer reaction times led to partial hydrolysis of the amide moiety and should be strictly avoided. Next, the glutamate derivative **5** was introduced into this intermediate using again EDCI under standard coupling conditions, affording derivative **16** in 88% yield. The incorporation of the pyrene sensitizer was then achieved by treatment of **16** with 20% piperidine solution in DMF to render the corresponding reactive amine **17** and successive EDCI-promoted amide formation with **6**. In this manner, **18** was obtained in 64% overall yield over the two steps. The subsequent reaction between this intermediate and the freshly made acid chloride of maleimide derivative **7**²³ afforded compound **19** in 74% yield, which features all the functional fragments of the final MAGA switch.

The preparation and manipulation of such an advanced intermediate was highly favored by the use of the new linear glutamate derivative **5**. In our hands, **5** turned to be a better glutamate precursor than the cyclic pyroglutamate group normally employed in the synthesis of MAG compounds,^{7–9} which we found to be troublesome due to (i) the moderate stability to the basic conditions required for the ring opening reaction of pyroglutamate of the maleimide fragment, which should therefore be introduced in a later stage of the route, and (ii) the low solubility of the dicarboxylic acid generated from the ring opening process of the pyroglutamate precursor,^{7–9} which compromised its manipulation and the efficiency of posterior reactions (e.g., incorporation of the maleimide group). These inconveniences are overcome by the use of the already opened derivative of glutamate **5** whose carboxylic acid groups are protected as the corresponding *tert*-butyl esters. Finally, removal of these moieties and of the additional *tert*-butyl carbamate protecting group of intermediate **19** took place in a single final step by treatment with a 2:1 mixture of trifluoroacetic acid and CH₂Cl₂, thus providing the monotrifluoroacetate salt of target ligand **2** in 79% yield. As expected, this compound was selectively isolated as its more stable *trans* isomer (*trans*-**2**), according to spectroscopic data.

One-Photon Optical Characterization of Photo-switchable Ligand 2. With the aim to assess the validity of our molecular design, the optical properties of compound **2** were investigated to demonstrate both sensitization of the *trans*–*cis* isomerization of the ligand upon photoexcitation of the pyrene antenna and the long thermal stability of the *cis* state of the switch at room temperature. For the sake of simplicity, this study was conducted under one-photon excitation with UV–vis light. However, based on our previous work¹⁶ and the well-known nonlinear optical properties of pyrene derivatives,^{19,20} the results obtained herein could be extrapolated to the photochemical behavior expected for **2** upon two-photon absorption of NIR light.

Figure 2a shows the absorption spectrum of *trans*-**2** in DMSO, which is compared to those recorded for its photoactive building blocks: pyrene derivative **6** and *trans*-

Scheme 4. Synthesis of the New MAGA Compound 2





Figure 2. (a) Normalized absorption spectra of *trans*-2, 6 and *trans*-20 in DMSO ($c = 5 \times 10^{-6}$ M). (b) Fluorescence emission spectra of *trans*-2 ($\times 10$, $\lambda_{\text{exc}} = 320$ nm) and 6 ($\lambda_{\text{exc}} = 305$ nm) in DMSO ($c = 1 \times 10^{-6}$ M). Both emission spectra were normalized relative to the absorption at the excitation wavelength.

4,4'-bis(acetylamino)azobenzene (*trans*-20), a model azoaromatic dye obtained by bis-acylation of 4,4'-diaminoazobenzene using standard amide formation conditions according to literature precedents.²⁴ Clearly, the narrow pyrene vibronic absorption peaks at $\lambda_{\text{abs}} \approx 315, 330$, and

345 nm and the broad *trans*-azoaromatic $\pi \rightarrow \pi^*$ ($\lambda_{\text{abs,max}} = 380$ nm) and $n \rightarrow \pi^*$ ($\lambda_{\text{abs}} > 450$ nm) absorption bands are both observed in the spectrum of *trans*-2. This demonstrates not only the successful introduction of the two chromophoric units but also that they are not electronically coupled in the ground state of the ligand. In contrast, non-negligible interactions should take place in the electronic excited state, as revealed by the fluorescence emission spectra measured for *trans*-2 and 6 (Figure 2b). While strong emission was detected for the separated pyrene derivative in DMSO with a rather high fluorescence quantum yield ($\Phi_{\text{fl,6}} = 0.30$), an ~ 43 -fold decrease was observed after incorporation into the ligand ($\Phi_{\text{fl,trans-2}} = 0.007$). A small fraction of this reduction must be attributed to the competitive absorption of the pyrene and azobenzene moieties of *trans*-2 at the excitation wavelength of the fluorescence quantum yield measurements ($\lambda_{\text{exc}} = 320$ nm). Actually, according to the extinction coefficients measured for the reference compounds 6 and *trans*-20 ($\epsilon_6^{320\text{ nm}}/\epsilon_{\text{trans-20}}^{320\text{ nm}} \approx 2$), about one-third of the excitation photons in these experiments were absorbed by the nonemissive azoaromatic core of *trans*-2, thus accounting for an apparent ~ 1.5 -fold decrease in Φ_{fl} for the pyrene antenna. However, the major part of the fluorescence quantum yield reduction (~ 30 -fold) can be ascribed to an intrinsic decrement of the emission efficiency of this fluorophore upon tethering to the azobenzene group in *trans*-2, thus uncovering additional nonradiative relaxation

pathways of its electronic excited state. As such, this result is indicative of the occurrence of the RET processes required for sensitization of the photoisomerization of the azobenzene moiety.

Despite this, we first investigated *trans*-to-*cis* isomerization of **2** upon direct excitation of the azoaromatic moiety. Figure 3 depicts the changes registered in the absorption spectra of




Figure 3. Variation of the absorption spectra of *trans*-**2** ($c = 5 \times 10^{-6}$ M) and *trans*-**20** ($c = 2.7 \times 10^{-5}$ M) in DMSO upon irradiation at $\lambda_{\text{exc}} = 365$ nm. The arrows in the figures indicate the spectral changes observed, while the dashed lines correspond to the absorption spectra of the resulting photostationary state mixtures.

trans-**2** and reference compound *trans*-**20** in DMSO when irradiated at $\lambda_{\text{exc}} = 365$ nm, an excitation wavelength where the pyrene moiety of the ligand shows minimal absorption ($\epsilon_{\text{trans-20}}^{365 \text{ nm}}/\epsilon_6^{365 \text{ nm}} = 13$; see Figure 2a). In both cases, a noteworthy decrease of the broad $\pi \rightarrow \pi^*$ azobenzene band around 380 nm and a very slight increase of the $n \rightarrow \pi^*$ azobenzene band at $\lambda_{\text{abs}} > 450$ nm were observed. These are clear signatures of the photoinduced conversion of *trans*-**2** and *trans*-**20** into their corresponding *cis* isomers,^{4,7,16} which should lead to the formation of an equilibrium photostationary state (PSS). According to the ¹H NMR data measured for DMSO-*d*₆ solutions of these compounds under equivalent irradiation conditions, the *cis* composition of the PSS mixtures generated was found to be 65% and 63% for **2** and **20**, respectively. Since nearly identical values were also determined for the *trans*-to-*cis* photoisomerization quantum yields of these azobenzene derivatives ($\Phi_{\text{trans-cis,2}} = \Phi_{\text{trans-cis,20}} = 0.28$), we could unambiguously conclude that the introduction of a pyrene group into the structure of ligand **2** did not modify the intrinsic photochemical behavior of its azoaromatic moiety when subjected to direct light excitation. A similar conclusion was drawn when analyzing the photoinduced *cis*-to-*trans* isomerization of **2** and **20** upon selective excitation of their azobenzene core. Thus, when illuminating with green light ($\lambda_{\text{exc}} = 532$ nm) the *trans/cis* mixtures of these compounds obtained by irradiation at $\lambda_{\text{exc}} = 365$ nm, our absorption measurements demonstrated that both *cis*-**2** and *cis*-**20** back-isomerized to their *trans* isomers nearly quantitatively and at similar rates (Figure S3). In particular, PSS mixtures with 94% and 93% content in *trans*-**2** and *trans*-**20** were obtained under these experimental conditions, respectively, according to our spectral data.

Next, the photosensitized isomerization of *trans*-**2** upon irradiation of the pyrene antenna was explored. Unfortunately, selective excitation of this chromophore to prevent direct *trans*-to-*cis* isomerization of **2** could not be achieved because the absorption spectrum of the pyrene photosensitizer selected overlaps with that of the azobenzene core (see Figure 2a). To minimize this effect and evaluate the actual contribution of the sensitized pathway to the photoswitching of **2**, two main actions were taken. On the one hand,

photosensitization was investigated at $\lambda_{\text{exc}} = 347$ nm, for which the largest difference in absorption between the pyrene and azoaromatic fragments of *trans*-**2** was determined ($\epsilon_6^{347 \text{ nm}}/\epsilon_{\text{trans-20}}^{347 \text{ nm}} = 2$). On the other hand, photoisomerization was simultaneously studied at $\lambda_{\text{exc}} = 365$ nm, which should selectively promote direct light-induced *trans*-to-*cis* conversion, as discussed above. Figure 4 shows the relative photo-




Figure 4. Photogenerated ratios of *cis*-**2** and *cis*-**20** ($\%_{\text{cis-2}}/\%_{\text{cis-20}}$) measured upon continuous irradiation of DMSO solutions of *trans*-**2** and *trans*-**20** ($c = 5 \times 10^{-6}$ M) at $\lambda_{\text{exc}} = 347$ and 365 nm.

isomerization efficiencies of *trans*-**2** and antenna-less compound *trans*-**20** in DMSO solution at $\lambda_{\text{exc}} = 347$ and 365 nm, which are plotted as the photogenerated ratios of *cis*-**2** and *cis*-**20** determined at different irradiation times from absorption measurements. Noticeably, a straight line around $\%_{\text{cis-2}}/\%_{\text{cis-20}} = 1$ was obtained at $\lambda_{\text{exc}} = 365$ nm, which indicates that direct *trans*-to-*cis* photoisomerization proceeds with equivalent rates and efficiencies for both azoaromatic compounds. This is indeed the result expected according to our previous experiments, where nearly identical PSS compositions and $\Phi_{\text{trans-cis}}$ values were measured upon selective excitation of the *trans*-azobenzene moiety of **2** and **20**.

However, a more complex time profile was obtained at $\lambda_{\text{exc}} = 347$ nm. At short irradiation times ($t < 50$ min), we found $\%_{\text{cis-2}}/\%_{\text{cis-20}}$ to be larger than 1, which indicates that *trans*-to-*cis* photoisomerization takes place faster for ligand **2**. This pinpoints the occurrence of an additional light-induced isomerization mechanism in this compound with respect to reference *trans*-**20**, which we ascribe to photosensitization evolving through pyrene absorption and subsequent resonance energy transfer toward the azobenzene core. Such a RET process would therefore account for the fluorescence quenching observed for the pyrene unit in the ligand (see Figure 2b), and actually, we estimated it to take place with a high critical Förster radius ($R_0 = 3.42$ nm) and efficiency ($E_{\text{RET}} = 99.8\%$) according to the optical properties of **6** and *trans*-**20** and the separation distances between the antenna and azoaromatic groups in *trans*-**2** (Figure S1). Nevertheless, it must be noted that energy transfer from pyrene to the *cis*-azobenzene moiety of **2** is also to be expected given the non-negligible spectral overlap between these groups, and indeed, only slightly lower R_0 and E_{RET} values were determined for this process with respect to those for *trans*-**2**: $R_0 = 2.47$ nm and $E_{\text{RET}} = 96.2\%$ (Figure S2). This suggests that the *cis*-to-*trans* isomerization of this compound can also be photosensitized by the antenna once the concentration of *cis*-**2** significantly increases in the sample. As such, the PSS generated at $\lambda_{\text{exc}} = 347$ nm would depend on the intricate balance between direct and sensitized *trans*-to-*cis* and *cis*-to-*trans* photoisomerization rates, which explains the different equilibrium mixtures obtained under long irradiation times at this wavelength with respect to $\lambda_{\text{exc}} = 365$ nm. Despite this, it

is clear that addition of the pyrene group to the ligand enhances the photoconversion efficiency of *trans*-**2** (up to 60% according to the data at short irradiation times), thus demonstrating that photosensitization with a two-photon absorber can also be exploited to trigger MAGA-type PTLs bearing a symmetrically substituted azobenzene core.

Once proven that target photoswitch **2** can operate under photosensitized isomerization, the thermal stability of the resulting *cis* isomer was analyzed. With this aim, the spontaneous *cis*-to-*trans* isomerization of this compound and reference **20** was studied in the dark and at room temperature by absorption spectroscopy. Spectral changes were observed in this way that are consistent with *cis*-**2** and *cis*-**20** back-isomerization, namely, the recovery of the intense *trans* absorption around 380 nm and a slight decrease of the $n \rightarrow \pi^*$ band at $\lambda_{\text{abs}} > 450$ nm (Figure S4). By plotting the time dependence of these data, the thermal *cis*-to-*trans* conversion rates could be retrieved from monoexponential fits,¹⁷ as shown in Figure 5. In particular, the following values were




Figure 5. Variation of the absorption at 382 nm of the photostationary state mixtures of **2** and **20** in the dark at 25 °C in DMSO, which reports on the recovery of the concentration of *trans*-**2** and *trans*-**20** upon thermal *cis*-to-*trans* isomerization. Points correspond to the experimental data, while dashed lines were obtained from monoexponential fits.

obtained: $k_{\text{cis} \rightarrow \text{trans}} = 3.2 \times 10^{-5}$ and $4.1 \times 10^{-5} \text{ s}^{-1}$ for **2** and **20**, respectively, which correspond to thermal lifetimes of their *cis* isomers at room temperature of 8.7 and 6.8 h. Clearly, very similar *cis*-to-*trans* isomerization kinetics were found for both compounds in the dark. This indicates that, under equivalent experimental conditions, the thermal stability of *cis*-azobenzenes is mainly governed by the electronic effects imparted by the substituents introduced, as already reported.⁴ As such, the equivalent 4,4'-diamide substituent pattern of **2** and **20** accounts for their comparable and long τ_{cis} values regardless of the different nature of their actual substituents. Despite this, a minor though non-negligible size effect could be inferred from our measurements, with the bulkier amide groups introduced in **2** slightly retarding its thermal *cis*-to-*trans* back-isomerization process and leading to a slightly larger *cis* state lifetime. On the other hand, the large thermal stability encountered for *cis*-**2** and *cis*-**20** is in agreement with the behavior previously reported for *cis*-MAG ($\tau_{\text{cis}} = 17.7$ min in aqueous buffer),^{7b} which also bears a 4,4'-diamido-azobenzene core. The differences in τ_{cis} observed between these compounds must be attributed to the distinct media where the measurements were conducted, since it is known that the *cis* lifetime of azobenzene derivatives largely decreases in polar solvents.^{4a,25} In fact, when monitoring the thermal *cis*-to-*trans* isomerization kinetics of *cis*-**2** in a 1:1 DMSO/water mixture (Figure S5), we determined a much shorter τ_{cis} value ($\tau_{\text{cis}} = 2.0$ h) than in pure DMSO and closer to that found for *cis*-MAG in aqueous buffer. Finally, the large thermal *cis* state

stability obtained herein for **2** is in striking contrast with the short *cis* lifetime measured for the previous MAGA ligand **1** ($\tau_{\text{cis}} = 67$ ms in PBS/DMSO 4:1 and 265 ms when tethered to LiGluR¹⁶), which arose from its push-pull substitution pattern. This unequivocally demonstrates that our photosensitized approach toward two-photon responsive azobenzene-based PTLs also enables the use of switches displaying long-lived *cis* isomers and, consequently, can be applied to the generation of large light-induced biological responses upon continuous irradiation.

Actually, a preliminary assessment of the two-photon activity of **2** can be made on the basis of the one-photon experiments conducted herein, the two-photon absorption cross sections reported in the literature for pyrene²⁶ and azobenzene¹⁷ chromophores, and our previous work on fast relaxing MAGA switch **1**.¹⁶ Since the same excited state of the pyrene antenna would be built, the efficiency of the RET process toward the azobenzene core of **2** should be the same under one- or two-photon excitation. As such, the photosensitized two-photon response of this switch would mainly depend on the following parameters: (a) the selectivity of the two-photon excitation of the pyrene antenna with NIR light; (b) the lifetime of the *cis* isomer of the switch created under such conditions. As discussed above, the one-photon excitation of the pyrene moiety of **2** is poorly selective and the maximum photosensitization effect was found at $\lambda_{\text{exc}} = 347$ nm, where a 2:1 pyrene/azobenzene absorption ratio was estimated. Noticeably, we expect this absorption difference to increase under NIR irradiation due to the larger two-photon activity of the pyrene antenna, thus favoring the sensitized pathway for photoisomerization. In fact, 5-fold higher two-photon absorption cross sections at 820 nm have been reported for monosubstituted pyrene derivatives ($\sigma_2 = 55$ GM; 1 GM = $10^{-50} \text{ cm}^4 \text{ s } \gamma^{-1}$) with respect to 4,4'-diamidoazobenzenes ($\sigma_2 = 10$ GM).¹⁷ As a result, sensitization via the pyrene antenna should be the major pathway for the two-photon isomerization of **2**. It must be noted that the intrinsic efficiency of this process should be lower than for our previous MAGA switch **1** bearing a higher absorbing naphthalene sensitizer ($\sigma_2 \approx 200$ GM at 780 nm).^{16,27} Despite this, an overall large two-photon response would be expected for **2** on account of the longer lifetime of its *cis* isomer. Actually, the fast thermal back-isomerization of **1** resulted in an ~ 10 -fold lower biological signal for this compound with respect to MAG under continuous one-photon excitation,¹⁶ which indicated that a less *cis*-enriched photostationary state could be prepared for this MAGA switch. Taking into account that **2** also presents a long-lived *cis* state as MAG, a similar factor can be assumed between the one-photon responses of the two MAGA-type compounds. This should counterbalance the ~ 4 -fold lower two-photon absorptivity of the pyrene sensitizer with respect to the naphthalene antenna of **1** and eventually lead to larger biological signals for the switch developed herein under NIR irradiation.

CONCLUSIONS

In summary, we report herein the synthesis and photochemical characterization of a new azobenzene-based photo-switchable ligand designed to enable optical control of ionotropic glutamate receptors in neurons via two-photon excitation of a pyrene sensitizing unit with NIR light. Aiming to develop an efficient and versatile synthetic route for this

and other related ligands, a modular strategy is described which benefits from the use of a new linear full protected glutamate derivative stable in basic media. In this way, target compound **2** could be prepared in 7 steps with a 27% overall yield starting from a commercial *N,N*-orthogonally diprotected L-lysine derivative as a central scaffold, to which its different functional fragments were sequentially introduced: the azoaromatic switch, the glutamate moiety, the pyrene sensitizer, and the anchoring maleimide group. As originally planned, **2** presents one-photon photoisomerization either upon selective irradiation of its azobenzene core or via photoexcitation of the pyrene antenna. In addition, by using a symmetrically substituted azoaromatic group, the thermal lifetime of its *cis* isomer could be enhanced in a factor $>10^5$ with respect to the analogous two-photon responsive ligands previously reported, which rely on the intrinsic nonlinear optical properties of push–pull azobenzenes. This is an essential requirement for enabling accumulation of the protein-bound *cis* state of the switch upon repeated illumination of tagged glutamate receptors, which makes compound **2** a promising candidate for evoking large photoinduced biological responses under multiphoton operation of these systems with NIR light.

EXPERIMENTAL SECTION

General Methods. Commercially available reagents were used as received. The solvents were dried by distillation over the appropriate drying agents. All reactions were performed avoiding moisture by standard procedures and under a nitrogen atmosphere. Flash column chromatography was performed using silica gel (230–400 mesh). ^1H NMR and ^{13}C NMR spectra were recorded at 250 and 62.5 MHz or 360 and 90 MHz or 500 and 125 MHz. NMR signals were assigned with the help of DEPT, COSY, HMBC, HMQC, and NOESY experiments. Proton chemical shifts are reported in ppm (δ) (CDCl_3 , δ 7.26; DMSO- d_6 , δ 2.50; or methanol- d_4 , δ 3.31). Carbon chemical shifts are reported in ppm (δ) (CDCl_3 , δ 77.2; DMSO- d_6 , δ 39.5 or methanol- d_4 , δ 49.0). Melting points were determined by a hot stage apparatus and are uncorrected. High resolution mass spectra (HRMS) were recorded by a micrOTOFQ spectrometer using ESIMS (QTOF). Optical rotations were measured at 22 ± 2 °C.

Photochemical Characterization. All spectroscopic and photochemical experiments were carried out in HPLC or spectroscopy quality solvents and in Ar-degassed samples. Steady-state UV–vis absorption measurements were recorded on a UV–visible (190–1100 nm) spectrophotometer. Fluorescence emission spectra were measured in a fluorescence spectrometer using a high energy pulsed Xenon source for excitation. All the emission spectra registered were corrected by the wavelength dependence of the spectral response of the detection system. Fluorescence quantum yields were determined using the standard method²⁸ and relative to pyrene in ethanol ($\Phi_{\text{pyrene}} = 0.32$).²⁹ The same excitation wavelength was used for both the sample of interest and the standard ($\lambda_{\text{exc}} = 305$ (**6**) or 320 nm (*trans*-**2**)), and solutions with absorbance values below 0.05 were measured to prevent inner-filter effects. The fluorescence quantum yield of the target compound (Φ_{sample}) was determined by applying the following equation:²⁸

$$\Phi_{\text{sample}} = \Phi_{\text{ref}} \frac{F_{\text{sample}}}{F_{\text{ref}}} \frac{\text{Abs}_{\text{ref}}}{\text{Abs}_{\text{sample}}} \frac{n_{\text{sample}}^2}{n_{\text{ref}}^2} \quad (1)$$

In this equation, Φ_{ref} is the fluorescence quantum yield of the standard, F_{sample} and F_{ref} are the integrated emission intensities measured for the sample and the standard, $\text{Abs}_{\text{sample}}$ and Abs_{ref} are the optical densities of the sample and the standard at the excitation wavelength, and n_{sample} and n_{ref} are the refractive indices of the solvents used. Photoisomerization of **2** and **20** was investigated by (i) steady-state UV–vis absorption spectroscopy for *trans*-to-*cis* and

cis-to-*trans* photoisomerization and slow *cis*-to-*trans* thermal back-isomerization processes and (ii) ^1H NMR for the elucidation of the *trans*-to-*cis* photostationary state mixtures. Photoisomerization quantum yields were determined using the methodology reported in ref **30** and azobenzene as reference compound ($\Phi_{\text{trans-cis}} = 0.15$ in acetonitrile³¹). Different excitation sources were used in the photochemical experiments depending on the spectral requirements: a UV lamp equipped with two 4 W tubes emitting at 365 nm, a Xe lamp coupled to a spectrograph, the third harmonic of the Nd:YAG laser ($\lambda_{\text{exc}} = 355$ nm), or a cw laser diode at $\lambda_{\text{exc}} = 532$ nm (Z-Laser). All absorption, fluorescence, and photochemical measurements of **2**, **6**, and pyrene were carried at low enough concentrations as to prevent aggregation processes driven by π -stacking of pyrene units.

(2R)-2-((2S)-3-(tert-Butoxy)-2-[(tert-butoxycarbonyl)amino]-3-oxopropyl)-4-pentenoic Acid, 10. To an ice-cooled solution of **9a** (1.80 g, 5.53 mmol) in a mixture 5:2 of THF and water (75 mL), 1 M LiOH (8.0 mL, 8.00 mmol) was added. After stirring for 1 h at this temperature, the mixture was acidified to pH 2 with 1 M HCl and extracted with EtOAc (3 \times 50 mL). The combined organic extracts were dried over anhydrous MgSO_4 and the solvent was removed under vacuum. The residue was purified by column chromatography (hexanes/EtOAc, 3:1) to furnish **10** (1.78 g, 5.18 mmol, 93% yield) as a yellowish oil: $[\alpha]_{\text{D}}^{20} -0.8$ (*c* 1.47, CH_2Cl_2); ^1H NMR (360 MHz, CDCl_3) δ 5.74 (m, 1H, H-4), 5.39 (br d, $J_{\text{NH},2} = 8.5$ Hz, 1H, NH), 5.08 (m, 2H, H-5), 4.22 (m, 1H, H-2'), 2.49 (m, 2H, H-2, H-3), 2.31–2.04 (m, 2H, H-3, H-1'), 1.66 (m, 1H, H-1'), 1.46 (s, 9H, $\text{C}(\text{CH}_3)_3$), 1.45 (s, 9H, $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (90 MHz, CDCl_3) δ 178.6 (C-1), 171.3 (C-3'), 156.3 (CO, carbamate), 134.8 (C-4), 117.8 (C-5), 82.7 ($\text{C}(\text{CH}_3)_3$), 80.7 ($\text{C}(\text{CH}_3)_3$), 52.7 (C-2'), 41.8 (C-2), 36.4 (C-3), 35.2 (C-1'), 28.4 ($\text{C}(\text{CH}_3)_3$), 28.1 ($\text{C}(\text{CH}_3)_3$); IR (ATR) 2978, 1707, 1367, 1247, 1150 cm^{-1} ; HRMS (ESI+) calcd for $[\text{C}_{17}\text{H}_{29}\text{NO}_6+\text{Na}]$: 366.1887; found: 366.1893.

Di(tert-butyl) (2R,4S)-2-Allyl-4-[(tert-butoxycarbonyl)aminopentanedioate, 11. To a solution of compound **10** (3.50 g, 7.28 mmol) in CH_2Cl_2 (90 mL), *tert*-butyl 2,2,2-trichloroacetimidate (2.6 mL, 14.6 mmol) and $\text{BF}_3 \cdot \text{OEt}_2$ (370 μL , 2.96 mmol) were added. Then, a saturated aqueous solution of NaHCO_3 (20 mL) was added and the product was extracted with CH_2Cl_2 (3 \times 50 mL). The combined organic extracts were dried over anhydrous MgSO_4 and the solvent was removed under vacuum. The resulting yellow oil was purified by column chromatography (hexanes/EtOAc, 3:1) to give **11** (1.64 g, 4.10 mmol, 56%) as a yellow solid: mp 86–89 °C (from hexanes/EtOAc); $[\alpha]_{\text{D}}^{20} +10.7$ (*c* 0.59, CH_2Cl_2); ^1H NMR (250 MHz, CDCl_3) δ 5.70 (ddt, $J_{2',3'} = 17.0$ Hz, $J_{2',1'} = 10.1$ Hz, $J_{2',1'} = 6.8$ Hz, 1H, H-2'), 5.09 (m, 3H, 2 \times H-3', NH), 4.22 (m, 1H, H-4), 2.49–2.26 (m, 2H, H-1', H-2), 2.26–2.03 (m, 2H, H-1', H-3), 1.65 (m, 1H, H-3), 1.44 (m, 27H, 3 \times $\text{C}(\text{CH}_3)_3$); ^{13}C NMR (62.5 MHz, CDCl_3) δ 173.9/171.7 (C-1/C-5), 155.3 (CO, carbamate), 135.0 (C-2'), 117.2 (C-3'), 81.8 ($\text{C}(\text{CH}_3)_3$), 80.7 ($\text{C}(\text{CH}_3)_3$), 79.5 ($\text{C}(\text{CH}_3)_3$), 52.7 (C-4), 42.3 (C-2), 36.8 (C-1'), 34.1 (C-3), 28.3 ($\text{C}(\text{CH}_3)_3$), 28.1 ($\text{C}(\text{CH}_3)_3$), 28.0 ($\text{C}(\text{CH}_3)_3$); IR (ATR) 3382, 2979, 1742, 1703, 1526, 1366, 1229, 1150 cm^{-1} ; HRMS (ESI+) calcd for $[\text{C}_{21}\text{H}_{37}\text{NO}_6+\text{Na}]$: 422.2513; found: 422.2520.

Di(tert-butyl) (2S,4R)-2-[(tert-Butoxycarbonyl)amino]-4-[(2E)-4-oxo-2-butenyl]pentanedioate, 12. To a boiling solution of (4R)-**11** (2.30 g, 5.75 mmol) and crotonaldehyde (2.4 mL, 28.8 mmol) in dry CH_2Cl_2 (36 mL), the Hoveyda-Grubbs II catalyst (56 mg, 89 μmol) was added in three portions. The mixture was heated to reflux for 2.5 h, when TLC analysis (hexanes/EtOAc, 2:1) showed no starting material. The reaction mixture was filtered off through a pad of silica gel and concentrated. The crude was purified by column chromatography (hexanes/Et $_2$ O, from 3:1 to 1:1) to provide **12** (2.19 g, 5.12 mmol, 89% yield) as a yellow oil: $[\alpha]_{\text{D}}^{20} +88.4$ (*c* 0.48, CH_2Cl_2); ^1H NMR (250 MHz, CDCl_3) δ 9.46 (d, $J_{\text{CHO},3'} = 7.8$ Hz, 1H, CHO), 6.73 (dt, $J_{2',3'} = 15.7$ Hz, $J_{2',1'} = 6.7$ Hz, 1H, H-2'), 6.09 (dd, $J_{3',2'} = 15.7$ Hz, $J_{3',\text{CHO}} = 7.8$ Hz, 1H, H-3'), 5.03 (d, $J_{\text{NH},2} = 7.9$ Hz, 1H, NH), 4.20 (m, 1H, H-2), 2.68–2.39 (m, 3H, H-4, 2 \times H-

1ⁱ), 2.15 (ddd, $J_{\text{gem}} = 12.7$ Hz, $J_{3,2} = 8.6$ Hz, $J_{3,4} = 4.8$ Hz, 1H, H-3), 1.64 (ddd, $J_{\text{gem}} = 12.7$ Hz, $J_{3,2} = 8.6$ Hz, $J_{3,4} = 3.7$ Hz, 1H, H-3), 1.44 (m, 27H, 3 × C(CH₃)₃); ¹³C NMR (90 MHz, CDCl₃) δ 193.6 (CHO), 173.1/171.4 (C-1/C-5), 155.3 (CO, carbamate), 154.4 (C-2ⁱ), 134.7 (C-3ⁱ), 82.4 (C(CH₃)₃), 81.7 (C(CH₃)₃), 79.9 (C(CH₃)₃), 52.5 (C-2), 41.6 (C-4), 35.4 (C-1ⁱ), 34.8 (C-3), 28.4 (C(CH₃)₃), 28.2 (C(CH₃)₃), 28.1 (C(CH₃)₃); IR (ATR) 2977, 1699, 1392, 1367, 1251, 1150 cm⁻¹; HRMS (ESI+) calcd for [C₂₂H₃₇NO₇+Na]: 450.2462; found: 450.2460.

(2E,5R,7S)-8-tert-Butoxy-5-(tert-butoxycarbonyl)-7-[(tert-butoxycarbonyl)amino]-8-oxo-2-octenoic Acid, 13. To an ice-cooled solution of the aldehyde **12** (2.50 g, 5.85 mmol) in a 5:1 mixture of *t*-BuOH and water (240 mL), 2-methyl-2-butene (6.2 mL, 58.5 mmol), NaH₂PO₄·2H₂O (4.60 g, 29.2 mmol), and NaClO₂ (3.17 g, 35.1 mmol) were successively added. The resulting mixture was stirred at 0 °C for 1 h and warmed up to rt. The reaction mixture was stirred overnight. Then, water (30 mL) and a saturated aqueous solution of Na₂CO₃ were added until pH 8. The mixture was extracted with EtOAc (3 × 50 mL), dried over anhydrous MgSO₄, and concentrated under vacuum. Purification by column chromatography (hexanes/EtOAc/CH₃COOH, 4:1:0.1) gave **13** (2.00 g, 4.51 mmol, 77% yield) as a yellow oil: $[\alpha]_{\text{D}}^{20} +59.8$ (c 1.16, CH₂Cl₂); ¹H NMR (360 MHz, MeOH-*d*₄) δ 6.86 (dt, $J_{3,2} = 15.1$ Hz, $J_{3,4} = 7.3$ Hz, 1H, H-3), 5.84 (d, $J_{2,3} = 15.1$ Hz, 1H, H-2), 4.06 (dd, $J_{7,6} = 10.6$ Hz, $J_{7,8} = 3.8$ Hz, 1H, H-7), 2.56 (m, 1H, H-5), 2.41 (m, 2H, H-4), 2.08 (m, 1H, H-6), 1.65 (m, 1H, H-6), 1.45 (m, 27H, 3 × C(CH₃)₃); ¹³C NMR (62.5 MHz, MeOH-*d*₄) δ 174.5/172.8/169.0 (C-1/C-8/CO, ester), 157.3 (CO, carbamate), 146.7 (C-3), 124.4 (C-2), 82.4 (C(CH₃)₃), 82.1 (C(CH₃)₃), 80.2 (C(CH₃)₃), 53.6 (C-7), 42.9 (C-5), 35.9 (C-4), 34.4 (C-6), 28.8 (C(CH₃)₃), 28.4 (C(CH₃)₃), 28.3 (C(CH₃)₃); IR (ATR) 2977, 2932, 1703, 1366, 1247, 1147 cm⁻¹; HRMS (ESI+) calcd for [C₂₂H₃₇NO₈+Na]: 466.2411; found: 466.2417.

(5R,7S)-8-tert-Butoxy-5-(tert-butoxycarbonyl)-7-[(tert-butoxycarbonyl)amino]-8-oxooctanoic Acid, 5. To a solution of olefine **13** (1.50 g, 3.38 mmol) in EtOAc (60 mL), 10% Pd/C (0.15 g) was added. The resulting suspension was stirred in a H₂ atmosphere for 16 h. Then, it was filtered through Celite, and the solvent was evaporated under vacuum. The resulting oil was purified by column chromatography (hexanes/EtOAc/CH₃COOH, 5:4:1) to afford a colorless oil identified as **5** (1.50 g, 3.36 mmol, quantitative yield): $[\alpha]_{\text{D}}^{20} +66$ (c 0.30, CH₂Cl₂); ¹H NMR (360 MHz, MeOH-*d*₄) δ 6.83 (d, $J_{\text{NH},7} = 8.4$ Hz, 1H, NH), 3.98 (m, 1H, H-7), 2.44 (m, 1H, H-5), 2.29 (td, $J_{2,3} = 6.8$ Hz, $J_{2,4} = 3.7$ Hz, 2H, H-2), 2.03 (m, 1H, H-6), 1.60 (m, 5H, 2 × H-3, 2 × H-4, H-6), 1.46 (m, 27H, 3 × C(CH₃)₃); ¹³C NMR (90 MHz, MeOH-*d*₄) δ 176.8/175.9/173.3 (C-1/C-8/CO, ester), 157.8 (CO, carbamate), 82.5 (C(CH₃)₃), 82.0 (C(CH₃)₃), 80.3 (C(CH₃)₃), 54.1 (C-7), 43.9 (C-5), 35.0 (C-6), 34.5 (C-2), 33.4 (C-4), 28.8 (C(CH₃)₃), 28.4 (C(CH₃)₃), 28.3 (C(CH₃)₃), 23.6 (C-1); IR (ATR) 2976, 1709, 1392, 1366, 1248, 1146 cm⁻¹; HRMS (ESI+) calcd for [C₂₂H₃₉NO₈+Na]: 468.2568; found: 468.2570.

9H-Fluoren-9-ylmethyl N-[(5S)-6-{4-[(E)-2-(4-Aminophenyl)-1-diazenyl]anilino}-5-[(tert-butoxycarbonyl)amino]-6-oxohexyl]carbamate, 14. To a stirred solution of 4-[(E)-2-(4-aminophenyl)-1-diazenyl]aniline (**4**) (1.00 g, 4.71 mmol) in dry THF (100 mL) under a N₂ atmosphere, a solution of **3** (2.43 g, 5.19 mmol), EDCI (1.17 g, 6.10 mmol), HOBt (955 mg, 7.07 mmol), and DIPEA (3.3 mL, 18.8 mmol) in dry THF (100 mL) was added. The reaction mixture was stirred overnight at rt. Then, the mixture was diluted with EtOAc (200 mL) and washed with water (2 × 100 mL). The organic layer was dried over anhydrous MgSO₄ and concentrated under vacuum. The residue was purified by column chromatography (from CH₂Cl₂ to CH₂Cl₂/EtOAc, 4:1) to deliver amide **14** (1.62 g, 2.45 mmol, 85% yield considering starting material consumed) as an orange solid and starting material (388 mg, 1.83 mmol): mp 94–97 °C (from CH₂Cl₂/EtOAc); $[\alpha]_{\text{D}}^{20} -9.70$ (c 1.01, CHCl₃); ¹H NMR (250 MHz, CDCl₃) δ 8.73 (br s, 1H, C-1ⁱⁱⁱNH), 7.77 (m, 6H, H-4, H-5, H-3ⁱⁱⁱ, H-5ⁱⁱⁱ, H-2^{iv}, H-6^{iv}), 7.65 (d, $J_{2,3} = J_{6,5} = 8.0$ Hz, 2H, H-2ⁱⁱⁱ, H-6ⁱⁱⁱ), 7.58 (d, $J_{1,2} = J_{8,7} = 7.2$ Hz, 2H, H-

1, H-8), 7.38 (t, $J_{3,2} = J_{3,4} = J_{6,7} = J_{6,5} = 7.2$ Hz, 2H, H-3, H-6), 7.29 (td, $J_{2,3} = J_{2,1} = J_{7,8} = J_{7,6} = 7.2$ Hz, $J_{2,4} = J_{7,5} = 1.2$ Hz, 2H, H-2, H-7), 6.71 (d, $J_{3,2} = J_{5,6} = 8.8$ Hz, 2H, H-3^{iv}, H-5^{iv}), 5.34 (d, $J_{\text{C-5}^{\text{iii}}\text{NH}_5^{\text{iv}}} = 7.5$ Hz, 1H, C-5ⁱⁱⁱNH), 4.95 (t, $J_{\text{C-1}^{\text{iii}}\text{NH}_1^{\text{iv}}} = 6.0$ Hz, 1H, C-1ⁱⁱⁱNH), 4.41 (d, $J_{1,9} = 6.9$ Hz, 2H, H-1ⁱ), 4.20 (m, 2H, H-5ⁱⁱ, H-9), 4.06 (br s, 2H, NH₂), 3.19 (m, 2H, H-1ⁱⁱ), 1.95 (m, 1H, H-4ⁱⁱ), 1.72 (m, 1H, H-4ⁱⁱ), 1.62–1.36 (m, 13H, 2 × H-2ⁱⁱ, 2 × H-3ⁱⁱ, C(CH₃)₃); ¹³C NMR (100.6 MHz, CDCl₃) δ 170.8 (C-6ⁱⁱ), 156.9 (CO, carbamate), 156.5 (CO, carbamate), 149.5 (C-4ⁱⁱⁱ, C-4^{iv}), 145.7 (C-1^{iv}), 144.0 (C-9a, C-8a), 141.4 (C-4a, C-4b), 139.4 (C-1ⁱⁱⁱ), 127.8 (C-3, C-6), 127.2 (C-2, C-7), 125.1 (C-1, C-8, C-2^{iv}, C-6^{iv}), 123.4 (C-3ⁱⁱⁱ, C-5ⁱⁱⁱ), 120.1 (C-4, C-6, C-2ⁱⁱⁱ, C-6ⁱⁱⁱ), 114.8 (C-3^{iv}, C-5^{iv}), 80.7 (C(CH₃)₃), 66.7 (C-1ⁱ), 55.2 (C-5ⁱⁱ), 47.4 (C-9), 40.3 (C-1ⁱⁱ), 31.4 (C-4ⁱⁱ), 29.5 (C-2ⁱⁱ), 28.4 (C(CH₃)₃), 22.7 (C-3ⁱⁱ); IR (ATR) 3323, 2926, 2856, 1687, 1665, 1596, 1504, 1245 cm⁻¹; HRMS (ESI+) calcd for [C₃₈H₄₂N₆O₅+Na]: 685.3109; found: 685.3112.

9H-Fluoren-9-ylmethyl N-[(5S)-5-Amino-6-{4-[(E)-2-(4-aminophenyl)-1-diazenyl]anilino}-6-oxohexyl]carbamate, 15. To a solution of intermediate **14** (1.07 g, 1.62 mmol) in MeOH (100 mL), 37% HCl (21 mL, 0.22 mol) was added. The reaction mixture was stirred for 1 h at room temperature until TLC (EtOAc) showed no presence of starting material. Then, the mixture was concentrated under vacuum, diluted with EtOAc (20 mL), and neutralized with a saturated aqueous solution of NaHCO₃. Purification by column chromatography (from EtOAc to EtOAc/MeOH, 95:5) gave **15** (890 mg, 1.58 mmol, 98% yield) as an orange solid: mp 78–82 °C (from EtOAc); $[\alpha]_{\text{D}}^{20} -25.6$ (c 1.19, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 9.69 (br s, 1H, C-1ⁱⁱⁱNH), 7.84 (d, $J_{3,2} = J_{3,4} = J_{6,7} = J_{6,5} = 7.4$ Hz, 2H, H-3, H-6), 7.31 (td, $J_{2,3} = J_{2,1} = J_{7,8} = J_{7,6} = 7.4$ Hz, $J_{2,4} = J_{7,5} = 0.9$ Hz, 2H, H-2, H-7), 6.73 (d, $J_{3,2} = J_{5,6} = 8.7$ Hz, 2H, H-3^{iv}, H-5^{iv}), 4.90 (t, $J_{\text{C-1}^{\text{iii}}\text{NH}_1^{\text{iv}}} = 5.4$ Hz, 1H, C-1ⁱⁱⁱNH), 4.40 (d, $J_{1,9} = 6.9$ Hz, 2H, H-1ⁱ), 4.21 (t, $J_{9,1} = 6.5$ Hz, 1H, H-9), 3.49 (m, 1H, H-5ⁱⁱ), 3.21 (m, 2H, H-1ⁱⁱ), 1.96 (m, 1H, H-4ⁱⁱ), 1.68–1.29 (m, 5H, 2 × H-2ⁱⁱ, 2 × H-3ⁱⁱ, H-4ⁱⁱ); ¹³C NMR (100.6 MHz, CDCl₃) δ 173.5 (C-6ⁱⁱ), 156.7 (CO, carbamate), 149.5/149.4 (C-4ⁱⁱⁱ/C-4^{iv}), 145.7 (C-1^{iv}), 144.1 (C-9a, C-8a), 141.4 (C-4a, C-4b), 139.4 (C-1ⁱⁱⁱ), 127.8 (C-3, C-6), 127.2 (C-2, C-7), 125.1 (C-1, C-8, C-2^{iv}, C-6^{iv}), 123.5 (C-3ⁱⁱⁱ, C-5ⁱⁱⁱ), 120.1 (C-4, C-6), 119.6 (C-2ⁱⁱⁱ, C-6ⁱⁱⁱ), 114.8 (C-3^{iv}, C-5^{iv}), 66.7 (C-1ⁱ), 55.5 (C-5ⁱⁱ), 47.4 (C-9), 40.7 (C-1ⁱⁱ), 34.5 (C-4ⁱⁱ), 29.8 (C-2ⁱⁱ), 23.0 (C-3ⁱⁱ); IR (ATR) 3335, 2923, 1687, 1596, 1508, 1249, 1138 cm⁻¹; HRMS (ESI+) calcd for [C₃₃H₃₄N₆O₃+H]: 563.2765; found: 563.2769.

Di(tert-butyl) (2R,4S)-2-{4-[(1S)-1-{4-[(E)-2-(4-Aminophenyl)-1-diazenyl]anilino}carbonyl-5-[(9H-fluoren-9-ylmethoxy)carbonyl]amino]pentyl}amino-4-oxobutyl}-4-[(tert-butoxycarbonyl)amino]pentanedioate, 16. To a stirred solution of **15** (232 mg, 0.34 mmol) in dry THF (18 mL) under a N₂ atmosphere, a solution of **5** (169 mg, 0.38 mmol), EDCI (86 mg, 0.45 mmol), HOBt (70 mg, 0.52 mmol), and DIPEA (240 μL, 1.40 mmol) in dry THF (20 mL) was added. The reaction mixture was stirred overnight at rt. Then, the mixture was diluted with EtOAc (40 mL) and washed with water (2 × 40 mL). The organic layer was dried over anhydrous MgSO₄ and concentrated under vacuum. The residue was purified by column chromatography (hexanes/EtOAc, 1:9) to afford amide **16** (335 mg, 0.30 mmol, 88% yield) as an orange solid: mp 98–105 °C (from hexanes/EtOAc); $[\alpha]_{\text{D}}^{20} -22.5$ (c 1.18, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 8.91 (br s, 1H, C-1ⁱⁱⁱNH), 7.79 (m, 6H, H-4^{iv}, H-5^{iv}, H-3^v, H-5^v, H-2^{vi}, H-6^{vi}), 7.65 (d, $J_{2,3} = J_{6,5} = 8.7$ Hz, 2H, H-2^v, H-6^v), 7.57 (d, $J_{1,2} = J_{8,7} = 7.4$ Hz, 2H, H-1^{iv}, H-8^{iv}), 7.38 (t, $J_{3,2} = J_{3,4} = J_{6,7} = J_{6,5} = 7.4$ Hz, 2H, H-3^{iv}, H-6^{iv}), 7.31 (t, $J_{2,3} = J_{2,1} = J_{7,8} = J_{7,6} = 7.4$ Hz, 2H, H-2^{iv}, H-7^{iv}), 6.73 (d, $J_{3,2} = J_{5,6} = 8.7$ Hz, 2H, H-3^{iv}, H-5^{iv}), 6.48 (s, 1H, C-1ⁱⁱⁱNH), 5.10 (t, $J_{\text{C-5}^{\text{iii}}\text{NH}_5^{\text{iv}}} = 5.4$ Hz, 1H, C-5ⁱⁱⁱNH), 4.56 (m, 1H, H-1ⁱⁱ), 4.38 (d, $J_{1,9} = 6.9$ Hz, 2H, H-1ⁱ), 4.20 (m, 2H, H-4, H-9^{iv}), 4.05 (br s, 2H, NH₂), 3.22 (m, 2H, H-5ⁱⁱ), 2.34 (m, 1H, H-2), 2.24 (m, 2H, H-3ⁱ), 2.13–1.94 (m, 3H, H-3, H-1ⁱ, H-2ⁱⁱ), 1.86–1.37 (m, 7H, H-3, H-1ⁱ, 2 × H-2ⁱⁱ, H-2ⁱⁱ, 2 × H-4ⁱⁱ), 1.44 (m, 29H, 3 × C(CH₃)₃, 2 × H-3ⁱⁱ); ¹³C NMR (100.6 MHz, CDCl₃) δ 174.5/173.8/171.9/170.1 (C-1/C-

5/C-4ⁱ/CO, amide), 157.0 (CO, carbamate), 155.5 (CO, carbamate), 149.5 (C-4^v/C-4^{vi}), 145.7 (C-1^{vi}), 144.1 (C-9a^{iv}, C-8a^{iv}), 141.4 (C-4a^{iv}, C-4b^{iv}), 139.5 (C-1^v), 127.8 (C-3^{iv}, C-6^{iv}), 127.2 (C-2^{iv}, C-7^{iv}), 125.2/125.1 (C-1^{iv}, C-8^{iv}/C-2^{vi}, C-6^{vi}), 123.4 (C-3^v, C-5^v), 120.1 (C-4^{iv}, C-5^{iv}/C-2^v, C-6^v), 114.8 (C-3^{iv}, C-5^{iv}), 82.1 (C(CH₃)₃), 81.1 (C(CH₃)₃), 79.8 (C(CH₃)₃), 66.8 (C-1ⁱⁱⁱ), 54.0 (C-1ⁱⁱ), 52.8 (C-4), 47.4 (C-9^{iv}), 42.4 (C-2), 40.5 (C-5ⁱⁱ), 36.1 (C-3ⁱ), 34.9 (C-3), 32.1 (C-1ⁱ), 30.7 (C-2ⁱⁱ), 29.8 (C-4ⁱⁱ), 28.5 (C(CH₃)₃), 28.2 (C(CH₃)₃), 28.1 (C(CH₃)₃), 23.2 (C-2ⁱ), 22.6 (C-3ⁱⁱ); IR (ATR) 3300, 2976, 2931, 1691, 1598, 1530, 1366, 1248, 1148 cm⁻¹; HRMS (ESI+) calcd for [C₅₅H₇₁N₇O₁₀+H]⁺: 990.5335; found: 990.5347.

Di(*tert*-butyl) (2*R*,4*S*)-2-(4-((1*S*)-5-Amino-1-((4-((*E*)-(4-aminophenyl)-1-diazenyl)anilino)carbonyl)pentyl)amino)-4-oxobutyl)-4-((*tert*-butoxycarbonyl)amino)pentanedioate, 17. A commercially available solution of 20% piperidine in DMF (18 mL) was added to compound 16 (775 mg, 0.78 mmol). After 1 h of stirring at rt, TLC analysis (EtOAc/MeOH, 95:5) showed no presence of starting material. The mixture was diluted with water (18 mL) and washed with EtOAc (3 × 30 mL). The combined organic extracts were dried over anhydrous MgSO₄ and concentrated under vacuum. The resulting solid was purified by column chromatography (from EtOAc to EtOAc/MeOH/NH₃, 9:2:1) to deliver 17 (459 mg, 0.60 mmol, 76% yield) as an orange solid: mp 89–96 °C (from EtOAc); [α]_D²⁰ –31.1 (c 0.88, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 9.56 (br s, 1H, C-1ⁱⁱⁱNH), 7.81 (m, 4H, H-3ⁱⁱⁱ, H-5ⁱⁱⁱ, H-2^{iv}, H-6^{iv}), 7.70 (d, J_{2ⁱⁱⁱ,3ⁱⁱⁱ} = J_{6ⁱⁱⁱ,5ⁱⁱⁱ} = 7.7 Hz, 2H, H-2ⁱⁱⁱ, H-6ⁱⁱⁱ), 6.74 (d, J_{3^{iv},2^{iv}} = J_{5^{iv},6^{iv}} = 8.4 Hz, 2H, H-3^{iv}, H-5^{iv}), 6.68 (m, 1H, C-1^{iv}NH), 5.21 (m, 1H, C-4NH), 4.63 (m, 1H, H-1ⁱⁱ), 4.19 (m, 1H, H-4), 4.06 (br s, 2H, C-4^vNH₂), 2.82 (m, 2H, H-5ⁱⁱ), 2.58–2.31 (m, 3H, H-2, C-5ⁱⁱNH₂), 2.27 (m, 2H, H-3ⁱ), 2.20–1.90 (m, 3H, H-3, H-1ⁱ, H-2ⁱⁱ), 1.86–1.56 (m, 7H, H-3, H-1ⁱ, 2 × H-2ⁱ, H-2ⁱⁱ, 2 × H-4ⁱⁱ), 1.44 (m, 29H, 3 × C(CH₃)₃, 2 × H-3ⁱⁱ); ¹³C NMR (100.6 MHz, CDCl₃) δ 174.5/173.5/171.8/170.3 (C-1/C-5/C-4ⁱ/CO, amide), 155.4 (CO, carbamate), 149.5 (C-4ⁱⁱⁱ, C-4^{iv}), 145.7 (C-1^{iv}), 139.7 (C-1ⁱⁱⁱ), 125.1 (C-2^{iv}, C-6^{iv}), 123.4 (C-3ⁱⁱⁱ, C-5ⁱⁱⁱ), 120.1 (C-2ⁱⁱⁱ, C-6ⁱⁱⁱ), 114.8 (C-3^{iv}, C-5^{iv}), 82.7 (C(CH₃)₃), 81.1 (C(CH₃)₃), 79.7 (C(CH₃)₃), 53.9 (C-1ⁱⁱ), 52.8 (C-4), 42.4 (C-2), 41.2 (C-5ⁱⁱ), 36.1 (C-3ⁱ), 34.8 (C-3), 32.1 (C-1ⁱ/C-2ⁱⁱ), 31.6 (C-4ⁱⁱ), 28.5 (C(CH₃)₃), 28.2 (C(CH₃)₃), 28.1 (C(CH₃)₃), 23.2 (C-2ⁱ), 22.7 (C-3ⁱⁱ); IR (ATR) 3343, 2977, 2933, 1701, 1598, 1506, 1368, 1249, 1150 cm⁻¹; HRMS (ESI+) calcd for [C₄₀H₆₁N₇O₈+H]⁺: 768.4654; found: 768.4650.

Di(*tert*-butyl) (2*R*,4*S*)-2-(4-((1*S*)-1-((4-((*E*)-(4-Aminophenyl)-1-diazenyl)anilino)carbonyl)-5-((2-pyrenylacetyl)amino)pentyl)amino)-4-oxobutyl)-4-((*tert*-butoxycarbonyl)amino)pentanedioate, 18. To a stirred solution of 17 (410 mg, 0.53 mmol) in dry THF (20 mL) under a N₂ atmosphere, a solution of 1-pyreneacetic acid (6) (167 mg, 0.64 mmol), EDCl (133 mg, 0.69 mmol), and DIPEA (465 μL, 2.67 mmol) in dry THF (30 mL) was added. The reaction mixture was stirred overnight at rt. Then, the mixture was diluted with EtOAc (50 mL) and washed with water (2 × 20 mL). The organic layer was dried over anhydrous MgSO₄ and concentrated under vacuum. The residue was purified by column chromatography (EtOAc) and triturated with diethyl ether (2 × 10 mL) to furnish an orange solid identified as 18 (458 mg, 0.45 mmol, 84% yield): mp 157–161 °C (from EtOAc); [α]_D²⁰ +5.30 (c 0.96, CHCl₃); ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.3 (s, 1H, C-1^{iv}NH), 8.37 (d, J = 9.3 Hz, 1H, H-pyr), 8.29–8.17 (m, 5H, C-1ⁱⁱⁱNH, 4 × H-pyr), 8.12 (s, 2H, H-pyr), 8.08 (m, 1H, C-5ⁱⁱNH), 8.05 (t, J = 7.6 Hz, 1H, H-pyr), 7.99 (d, J = 7.9 Hz, 1H, H-pyr), 7.76 (d, J_{2ⁱⁱⁱ,3ⁱⁱⁱ} = J_{5^{iv},6^{iv}} = 9.1 Hz, 2H, H-2^{iv}, H-6^{iv}), 7.71 (d, J_{3^{iv},2^{iv}} = J_{5^{iv},6^{iv}} = 9.1 Hz, 2H, H-3^{iv}, H-5^{iv}), 7.63 (d, J_{2^{iv},3^{iv}} = J_{6^{iv},5^{iv}} = 8.8 Hz, 2H, H-2^v, H-6^v), 7.10 (d, J_{C-4NH,4} = 8.2 Hz, 1H, C-4NH), 6.66 (d, J_{3^{iv},2^{iv}} = J_{5^{iv},6^{iv}} = 8.8 Hz, 2H, H-3^{iv}, H-5^{iv}), 6.02 (s, 2H, NH₂), 4.40 (m, 1H, H-1ⁱⁱ), 4.17 (s, 2H, H-2ⁱⁱ), 3.76 (m, 1H, H-4), 3.09 (dd, J_{5ⁱⁱ,4ⁱⁱ} = 12.6 Hz, J_{5ⁱⁱ,C-5ⁱⁱNH} = 6.7 Hz, 2H, H-5ⁱⁱ), 2.33 (m, 1H, H-2), 2.12 (m, 2H, H-3ⁱ), 1.85–1.51 (m, 6H, 2 × H-3, 2 × H-1ⁱ, 2 × H-2ⁱ), 1.52–1.25 (m, 33H, 3 × C(CH₃)₃, 2 × H-2ⁱ, 2 × H-3ⁱⁱ, 2 × H-4ⁱⁱ); ¹³C NMR (100.6 MHz, DMSO-*d*₆) δ 173.8/172.1/171.7/171.2 (C-1/C-5/C-4ⁱ/CO, amide), 169.9 (C-1ⁱⁱⁱ), 155.5 (CO, carbamate), 152.4 (C-4^v), 148.1 (C-4^{iv}), 142.8 (C-1^v), 140.2 (C-1^{iv}), 131.1 (C-pyr), 130.8 (C-pyr), 130.4 (C-

pyr), 129.7 (C-pyr), 129.0 (C-pyr), 128.6 (CH-pyr), 127.4 (CH-pyr), 127.2 (CH-pyr), 126.8 (CH-pyr), 126.1 (CH-pyr), 125.1 (C-2^v, C-6^v), 124.9 (CH-pyr), 124.8 (CH-pyr), 124.7 (CH-pyr), 124.1 (C-pyr), 124.0 (CH-pyr), 123.9 (C-pyr), 122.5 (C-3^{iv}, C-5^{iv}), 119.5 (C-2^{iv}, C-6^{iv}), 113.4 (C-3^v, C-5^v), 80.3 (C(CH₃)₃), 79.7 (C(CH₃)₃), 78.0 (C(CH₃)₃), 53.4 (C-1ⁱⁱ), 52.5 (C-4), 42.2 (C-2), 40.1 (C-1ⁱⁱⁱ), 38.6 (C-5ⁱⁱ), 34.8 (C-3ⁱ), 33.0 (C-3), 32.0 (C-1ⁱ), 31.6 (C-2ⁱⁱ), 28.8 (C-4ⁱⁱ), 28.2 (C(CH₃)₃), 27.9 (C(CH₃)₃), 27.6 (C(CH₃)₃), 23.0 (C-3ⁱⁱ, C-2ⁱ); IR (ATR) 3277, 2975, 2935, 1720, 1638, 1599, 1531, 1506, 1368, 1251, 1151 cm⁻¹; HRMS (ESI+) calcd for [C₅₈H₇₁N₇O₉+H]⁺: 1010.5386; found: 1010.5377.

Di(*tert*-butyl) (2*S*,4*R*)-2-((*tert*-Butoxycarbonyl)amino)-4-((4-((1*S*)-1-((4-((*E*)-4-((2-(2,5-dioxo-2,5-dihydro-1*H*-1-pyrrolyl)acetyl)amino)phenyl)-1-diazenyl)anilino)carbonyl)-5-((2-pyrenylacetyl)amino)pentyl)amino)-4-oxobutyl)-pentanedioate, 19. To a stirred solution of 7 (120 mg, 0.77 mmol) and oxalyl chloride 2.0 M in THF (402 μL, 0.80 mmol) in dry CH₂Cl₂ (12 mL) was added one drop of DMF. After stirring for 2 h at rt, the mixture was concentrated. The resulting acid chloride was taken up in dry THF (5 mL) and slowly added to an ice-cooled solution of 18 (200 mg, 0.20 mmol), DIPEA (207 μL, 1.19 mmol) in dry THF (25 mL). After stirring for 10 min at this temperature, the mixture was warmed up to rt and stirred for an additional 4 h. Then, it was diluted with EtOAc (30 mL) and washed with water (3 × 10 mL). The organic extract was dried over anhydrous MgSO₄, concentrated under vacuum, and purified by column chromatography (EtOAc/MeOH, 9:1) to provide 19 (168 mg, 0.15 mmol, 74% yield) as an orange solid: mp 187–198 °C (from EtOAc); [α]_D²⁰ +68.5 (c 0.70, DMSO); ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.7 (s, 1H, C-4^{iv}NH), 10.3 (s, 1H, C-1^{iv}NH), 8.37 (d, J = 9.3 Hz, 1H, H-pyr), 8.29–8.17 (m, 5H, C-1ⁱⁱⁱNH, 4 × H-pyr), 8.14 (s, 2H, H-pyr), 8.09 (m, 1H, C-5ⁱⁱNH), 8.07 (t, J = 7.8 Hz, 1H, H-pyr), 7.98 (d, J = 7.8 Hz, 1H, H-pyr), 7.84 (m, 6H, H-2^{iv}, H-6^{iv}, H-3^{iv}, H-5^{iv}, H-3^v, H-5^v), 7.75 (d, J_{2ⁱⁱⁱ,3ⁱⁱⁱ} = J_{6ⁱⁱⁱ,5ⁱⁱⁱ} = 8.9 Hz, 2H, H-2^v, H-6^v), 7.16 (s, 2H, H-3ⁱⁱⁱ, H-4ⁱⁱⁱ), 7.09 (d, J_{C-2NH,2} = 8.0 Hz, 1H, C-2NH), 4.38 (m, 1H, H-1ⁱⁱ), 4.33 (s, 2H, H-2ⁱⁱ), 4.16 (s, 2H, H-2ⁱⁱⁱ), 3.76 (m, 1H, H-2), 3.08 (dd, J_{5ⁱⁱ,4ⁱⁱ} = 12.6 Hz, J_{5ⁱⁱ,C-5ⁱⁱNH} = 6.7 Hz, 2H, H-5ⁱⁱ), 2.31 (m, 1H, H-4), 2.11 (m, 2H, H-3ⁱ), 1.85–1.51 (m, 6H, 2 × H-3, 2 × H-1ⁱ, 2 × H-2ⁱ), 1.51–1.24 (m, 33H, 3 × C(CH₃)₃, 2 × H-2ⁱ, 2 × H-3ⁱⁱ, 2 × H-4ⁱⁱ); ¹³C NMR (100.6 MHz, DMSO-*d*₆) δ 172.8/171.1/170.7/170.5/169.7 (C-1/C-5/C-4ⁱ/C-2^{vi}, C-5^{vii}/CO, amide), 168.9 (C-1ⁱⁱⁱ), 164.3 (C-1^{iv}), 154.5 (CO, carbamate), 146.9/146.7 (C-4^{iv}/C-1^v), 140.7/140.1 (C-1^{iv}/C-4^v), 134.0 (C-3ⁱⁱⁱ, C-4ⁱⁱⁱ), 130.1 (C-pyr), 129.8 (C-pyr), 129.4 (C-pyr), 128.7 (C-pyr), 128.0 (C-pyr), 127.6 (CH-pyr), 126.4 (CH-pyr), 126.2 (CH-pyr), 125.8 (CH-pyr), 125.1 (CH-pyr), 124.1 (CH-pyr), 123.9 (CH-pyr), 123.7 (CH-pyr), 123.1 (C-pyr), 123.0 (CH-pyr), 122.9 (C-pyr), 122.5/122.4 (C-2^v, C-6^v/C-3^{iv}, C-5^{iv}), 118.5 (C-2^{iv}, C-6^{iv}, C-3^v, C-5^v), 79.3 (C(CH₃)₃), 78.7 (C(CH₃)₃), 77.0 (C(CH₃)₃), 52.4 (C-1ⁱⁱ), 51.6 (C-2), 41.2 (C-4), 39.5 (C-2ⁱⁱⁱ), 37.6 (C-5ⁱⁱ), 33.8 (C-3ⁱ), 32.0 (C-3), 31.0 (C-1ⁱ), 30.6 (C-2ⁱⁱ), 27.8 (C-4ⁱⁱ), 27.2 (C(CH₃)₃), 26.9 (C(CH₃)₃), 26.6 (C(CH₃)₃), 22.0 (C-3ⁱⁱ, C-2ⁱ); IR (ATR) 3274, 2976, 2933, 1716, 1528, 1366, 1247, 1148 cm⁻¹; HRMS (ESI+) calcd for [C₆₄H₇₄N₈O₁₂+Na]⁺: 1169.5318; found: 1169.5350.

{(1*S*,3*R*)-1,3-Dicarboxy-7-(((1*S*)-1-((4-((*E*)-2-(4-((2-(2,5-dioxo-2,5-dihydro-1*H*-1-pyrrolyl)acetyl)amino)phenyl)-1-diazenyl)anilino)carbonyl)-5-((2-(2-pyrenyl)acetyl)amino)pentyl)amino)-7-oxoheptyl)ammonium 2,2,2-Trifluoroacetate, 2. To a stirred solution of compound 19 (40 mg, 45 μmol) in CH₂Cl₂ (7.0 mL), trifluoroacetic acid (3.5 mL, 45.4 mmol) was added. The mixture was stirred at room temperature until the starting material was consumed as judged by TLC analysis (EtOAc/MeOH, 9:1). Then, the mixture was concentrated under vacuum, and the resulting purple solid was triturated with diethyl ether (2 × 8 mL) to furnish 2 (29 mg, 28 μmol, 79%) as a brown powder: [α]_D²⁰ +50.9 (c 0.43, DMSO); ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.7 (s, 1H, C-4^{iv}NH), 10.4 (s, 1H, C-1^{iv}NH), 8.37 (d, J = 9.3 Hz, 2H, H-pyr, C-1ⁱⁱⁱNH), 8.30–8.17 (m, 5H, C-5ⁱⁱNH, 4 × H-pyr), 8.14 (s, 2H, H-pyr), 8.05 (t, J = 7.6 Hz, 1H, H-pyr), 7.98 (d, J = 7.6 Hz, 1H, H-pyr), 7.84 (m, 6H, H-2^{iv}, H-6^{iv}, H-3^{iv}, H-5^{iv}, H-3^v, H-5^v), 7.76 (d, J_{2ⁱⁱⁱ,3ⁱⁱⁱ} = J_{6ⁱⁱⁱ,5ⁱⁱⁱ} =

8.6 Hz, 2H, H-2^{iv}, H-6^{iv}), 7.16 (s, 2H, H-3^{vi}, H-4^{vi}), 4.36 (m, 1H, H-1ⁱ), 4.33 (s, 2H, H-2^v), 4.19 (s, 3H, 2 × H-2ⁱⁱ, H-1), 3.08 (m, 2H, H-5ⁱ), 2.68 (m, 1H, H-3), 2.12 (m, 3H, H-2, 2 × H-6), 1.80–1.26 (m, 11H, H-2, 2 × H-4, 2 × H-5, 2 × H-4ⁱ, 2 × H-3ⁱ, 2 × H-2ⁱ); ¹³C NMR (100.6 MHz, DMSO-*d*₆) δ 175.9/172.3/171.7/170.7/170.0 (C-2^{vi}/C-7/C-1ⁱⁱ/CO, acid/CO, acid/CO, amide), 165.4 (C-1^v), 147.9/147.6 (C-4ⁱⁱⁱ/C-1^{iv}), 141.8/141.1 (C-1ⁱⁱⁱ/C-4^{iv}), 135.0 (C-3^{vi}, C-4^{vi}), 131.1 (C-pyr), 130.8 (C-pyr), 130.4 (C-pyr), 129.7 (C-pyr), 129.0 (C-pyr), 128.6 (CH-pyr), 127.4 (CH-pyr), 127.2 (CH-pyr), 126.8 (CH-pyr), 126.1 (CH-pyr), 125.1 (CH-pyr), 124.9 (CH-pyr), 124.7 (CH-pyr), 124.1 (2 × C-pyr), 123.9 (C-pyr), 123.4 (C-2^{iv}, C-6^{iv}, C-3ⁱⁱⁱ, C-5ⁱⁱⁱ), 119.5 (C-2ⁱⁱⁱ, C-6ⁱⁱⁱ, C-3^{iv}, C-5^{iv}), 53.8 (C-1ⁱ), 51.6 (C-1), 40.4 (C-3), 39.5 (C-2ⁱⁱ), 38.5 (C-5ⁱ), 36.6 (C-6), 32.3 (C-2), 31.4 (C-4), 30.8 (C-2ⁱ), 28.8 (C-4ⁱ), 23.0/22.7 (C-3ⁱ, C-5); IR (ATR) 3277, 3044, 2938, 1712, 1644, 1595, 1532, 1199, 1150 cm⁻¹; HRMS (ESI+) calcd for [C₅₁H₅₀N₈O₁₀+H]⁺: 935.3723; found: 935.3714.

■ ASSOCIATED CONTENT

● Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.5b01402.

Additional photochemical data, ¹H and ¹³C NMR spectra of all new compounds and HSQC spectrum for compounds 10, 16, and 18 (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: pau@icrea.cat.

*E-mail: felix.busque@uab.es.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We acknowledge financial support from Ministerio de Economía y Competitividad (MINECO) (Projects CTQ2012-30853 and CTQ2013-41161-R), and M.G.-M. thanks the “Universitat Autònoma de Barcelona” for a predoctoral grant. We also acknowledge financial support from the RecerCaixa foundation (2010ACUP00378); the Marató de TV3 Foundation (Grant 111531); the European Union’s Seventh Framework Programme for research, technological development and demonstration under Grant Agreements 270483, 210355, and 335011; the Catalan government (2012FI_B 01122 and 2014SGR-1251); the Spanish Government (CTQ2013-43892R); the Ramón Areces foundation, and the ERANET SynBio program.

■ REFERENCES

- (1) (a) Fenno, L.; Yizhar, O.; Deisseroth, K. *Annu. Rev. Neurosci.* **2011**, *34*, 389–412. (b) Deisseroth, K. *Nat. Methods* **2011**, *8*, 26–29.
- (2) Fehrentz, T.; Schönberger, M.; Trauner, D. *Angew. Chem., Int. Ed.* **2011**, *50*, 12156–12182.
- (3) Kramer, R. H.; Mouro, A.; Adesnik, H. *Nat. Neurosci.* **2013**, *16*, 816–823.
- (4) (a) Beharry, A. A.; Woolley, G. A. *Chem. Soc. Rev.* **2011**, *40*, 4422–4437. (b) Szymanski, W.; Beierle, J. M.; Kistemaker, H. A. V.; Velema, W. A.; Feringa, B. L. *Chem. Rev.* **2013**, *113*, 6114–6178.
- (5) Bautista-Barrufet, A.; Izquierdo-Serra, M.; Gorostiza, P. Photoswitchable ion channels and receptors. In *Novel Approaches for Single Molecule Activation and Detection, Advances in Atom and Single Molecule Machines*; Benfenati, F., et al., Eds.; Springer-Verlag: Berlin, Heidelberg, 2014; DOI: 10.1007/978-3-662-43367-6_9.
- (6) (a) Traynelis, S. F.; Wollmuth, L. P.; McBain, C. J.; Menniti, F. S.; Vance, K. M.; Ogden, K. K.; Hansen, K. B.; Yuan, H.; Myers, S.

- J.; Dingledine, R. *Pharmacol. Rev.* **2010**, *62*, 405–496. (b) Niswender, C. M.; Conn, P. J. *Annu. Rev. Pharmacol. Toxicol.* **2010**, *50*, 295–322.
- (7) (a) Volgraf, M.; Gorostiza, P.; Numano, R.; Kramer, R. H.; Isacoff, E. Y.; Trauner, D. *Nat. Chem. Biol.* **2006**, *2*, 47–52. (b) Gorostiza, P.; Volgraf, M.; Numano, R.; Szobota, S.; Trauner, D.; Isacoff, E. Y. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 10865–10870.
- (8) Kienzler, M. A.; Reiner, A.; Trautman, E.; Yoo, S.; Trauner, D.; Isacoff, E. Y. *J. Am. Chem. Soc.* **2013**, *135*, 17683–17686.
- (9) Rullo, A.; Reiner, A.; Reiter, A.; Trauner, D.; Isacoff, E. Y.; Woolley, G. A. *Chem. Commun.* **2014**, *50*, 14613–14615.
- (10) Yaroslavsky, A. N.; Schulze, P. C.; Yaroslavsky, I. V.; Schober, R.; Ulrich, F.; Schwarzaier, H. J. *Phys. Med. Biol.* **2002**, *47*, 2059–2073.
- (11) Denk, W.; Strickler, J. H.; Webb, W. W. *Science* **1990**, *248*, 73–76.
- (12) Papagiakoumou, E.; Begue, A.; Leshem, B.; Schwartz, O.; Stell, B. M.; Bradley, J.; Oron, D.; Emiliani, V. *Nat. Photonics* **2013**, *7*, 274–278.
- (13) Oron, D.; Papagiakoumou, E.; Anselmi, F.; Emiliani, V. *Prog. Brain Res.* **2012**, *196*, 119–143.
- (14) Watson, B. O.; Nikolenko, V.; Yuste, R. *Front. Neural Circuits* **2009**, *3*, 6.
- (15) (a) De Boni, L.; Misoguti, L.; Zilio, S. C.; Mendonça, C. R. *ChemPhysChem* **2005**, *6*, 1121–1125. (b) Silva, D. L.; Krawczyk, P.; Bartkowiak, W.; Mendonça, C. R. *J. Chem. Phys.* **2009**, *131*, 244516.
- (16) Izquierdo-Serra, M.; Gascón-Moya, M.; Hirtz, J. J.; Pittolo, S.; Poskanzer, K. E.; Ferrer, E.; Alibés, R.; Busqué, F.; Yuste, R.; Hernando, J.; Gorostiza, P. *J. Am. Chem. Soc.* **2014**, *136*, 8693–8701.
- (17) Carroll, E. C.; Berlin, S.; Levitz, J.; Kienzler, M. A.; Yuan, Z.; Madsen, D.; Larsen, D. S.; Isacoff, E. Y. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112*, E776–E785.
- (18) Blevins, A. A.; Blanchard, G. J. *J. Phys. Chem. B* **2004**, *108*, 4962–4968.
- (19) Xu, C.; Williams, R. M.; Zipfel, W.; Webb, W. W. *Bioimaging* **1996**, *4*, 198–207.
- (20) Chiric, G.; Cannone, F.; Baldini, G.; Diaspro, A. *Biophys. J.* **2003**, *84*, 588–598.
- (21) Ezquerro, J.; Pedregal, C.; Rubio, A.; Yrretagoyena, B.; Escribano, A.; Sánchez-Ferrando, F. *Tetrahedron* **1993**, *49*, 8665–8678.
- (22) Steger, M.; Young, D. W. *Tetrahedron* **1999**, *55*, 7935–7956.
- (23) Pearson, R. J.; Kassianidis, E.; Slawin, A. M. Z.; Philp, D. *Org. Biomol. Chem.* **2004**, *2*, 3434–3441.
- (24) Sousa, A. C.; Martins, L. O.; Robalo, M. P. *Adv. Synth. Catal.* **2013**, *355*, 2908–2917.
- (25) Nishimura, N.; Sueyoshi, T.; Yamanaka, H.; Imai, E.; Yamamoto, S.; Hasegawa, S. *Bull. Chem. Soc. Jpn.* **1976**, *49*, 1381–1387.
- (26) Kim, H. M.; Lee, Y. O.; Lim, C. S.; Kim, J. S.; Cho, B. R. *J. Org. Chem.* **2008**, *73*, 5127–5130.
- (27) Kim, H. M.; Cho, B. R. *Acc. Chem. Res.* **2009**, *42*, 863–872.
- (28) Lakowicz, J. R. *Principles of Fluorescence Spectroscopy*; Springer: New York, USA, 2006.
- (29) Berlman, I. B. *Handbook of Fluorescence Spectra of Aromatic Molecules*; Academic Press: New York and London, 1971.
- (30) Lees, A. J. *Anal. Chem.* **1996**, *68*, 226–229.
- (31) Bandara, H. M. D.; Burdette, S. C. *Chem. Soc. Rev.* **2012**, *41*, 1809–1825.