


Incorporation of Trifluoromethylated Proline and Surrogates into Peptides: Application to the Synthesis of Fluorinated Analogues of the Neuroprotective Glycine-Proline-Glutamate (GPE) Tripeptide

Julien Simon, Julien Pytkowicz, Nathalie Lensen,* Grégory Chaume,* and Thierry Brigaud*

Laboratoire de Chimie Biologique (LCB), Université de Cergy-Pontoise, EA 4505, 5 Mail Gay-Lussac, 95000 Cergy-Pontoise, France

S Supporting Information

ABSTRACT: The incorporation into a peptide chain of highly hindered and weakly nucleophilic trifluoromethylated prolines, pseudoproline and oxazolidinones has been achieved. As an application, the synthesis of a new class of fluorinated analogues of the neuroprotective tripeptide glycine-proline-glutamate (GPE) is reported. These analogues have been elaborated from a panel of five-membered ring trifluoromethylated amino acids (Tfm-AA) through the coupling reaction with a glutamate residue at the C-terminus and a glycine at the N-terminus. Although the peptide coupling reaction at the C-terminal position of the fluorinated amino acid was conveniently performed under standard conditions, the very challenging coupling reaction at the highly deactivated N-terminal position proved to be much more problematic. A methodological study was needed to identify suitable reaction conditions for this difficult peptide coupling.



INTRODUCTION

The use of peptides as therapeutic agents has been quite limited due to several major drawbacks such as structural flexibility, rapid degradation by peptidases and low lipophilicity. However, because of recent advances in peptides chemistry and biology, peptide-based drug discovery constitutes now a serious alternative for addressing new therapeutic challenges.¹ The introduction of fluorine atoms into biomolecules such as peptides is known to deeply influence their chemical and biological properties,^{2–5} and consequently, represents an option to develop the therapeutic potential of peptides. The synthesis of fluorinated amino acids has gained a considerable interest in peptide and protein chemistry^{6–8} as they are known to increase the chemical stability, the metabolic resistances^{9,10} and to provide a better affinity of peptides for lipid membranes. Moreover, their incorporation into peptides can induce stabilization of particular conformations and better autoassembly.^{11–15} Fluorinated peptides can also be used as efficient probes for ¹⁹F NMR studies.^{16,17} Trifluoromethylated amino acids (Tfm-AA) represent a special class of highly constrained nonproteogenic amino acids. While the coupling reactions of the Tfm-AA at their C-termini can be efficiently achieved using standard protocols, their incorporation into peptides at the N-terminal position still remains challenging due to the stereoelectronic effects imparted by the CF₃ group which strongly decrease the nitrogen nucleophilicity.¹⁸ Our group has developed the stereoselective synthesis of a panel of acyclic and cyclic Tfm-AA in enantiopure form^{19–26} and is now mainly focused on the development of efficient methodologies for their incorporation into peptides.^{27–30} Although α -CF₃-alanine has been incorporated into peptides, the peptide coupling at the N-

terminal position of α -CF₃-proline has not been reported so far. We report herein our investigations about the incorporation of α -CF₃-proline and various cyclic surrogates into short peptide sequences. The usefulness of this methodology for the synthesis of peptides of biological interest is illustrated by the synthesis of new CF₃-tripeptide analogues of the neuroprotective glycine-proline-glutamate peptide (GPE) (Figure 1). It is assumed that the GPE is the result of the proteolytic cleavage of the insulin-like growth factor (IGF-1).³¹ Even if the GPE neither binds to IGF-1 receptors nor has any neurotrophic effect, it displays remarkable CNS activities.^{32,33} In different animal models, the GPE shows neuroprotective effects on neurodegenerative processes, such as Alzheimer's, Parkinson's and Huntington's diseases. There is also evidence that the GPE exhibits neuromodulatory activities.^{34–36} Due to its structural simplicity, the GPE emerged as a lead for the development of potent neuroprotective agents for the treatment of various CNS injuries.³⁷ Numerous GPE analogues have been reported in the literature but most of them suffer of a lack of chemical stability and short half-time bioavailability.^{38–45} It has been shown that proline residue is crucial due to its unique conformational constraint.⁴⁶ The tripeptide NNZ-2566, containing a 2-methylproline residue, shows higher activity compared to GPE and other analogues and is currently in phase II for several neurological indications.⁴⁷ Replacement of the proline residue by a CF₃-containing surrogate is expected to enhance the chemical stability of the GPE tripeptide together with its bioavailability profile.

Received: April 1, 2016

Published: June 13, 2016




Figure 1. Chemical structure of GPE, NNZ-2566 and CF₃-GPE analogues.





Figure 2. Chemical structure of Tfm-AAs as proline surrogates.

Scheme 1. Synthesis of Dipeptides 6



Three distinctive trifluoromethylated cyclic surrogates were considered for replacing the proline residue in the GPE sequence varying the position and the configuration of the CF₃ group along the 5-membered ring: α -CF₃-proline **1**, CF₃-pseudoproline **2** and **3** derived from serine and cysteine respectively and CF₃-oxazolidines **4** and **5** derived from ethanolamine and (*R*)-phenylglycinol (Figure 2). Except for the oxazolidine **4**, the synthesis of enantiopure **1**,^{20,22} **2**,²³ **3**²³ and **5**^{20,21} have been previously described by our group.


RESULTS AND DISCUSSION

CF₃-Proline Containing GPE Analogues. The synthesis of CF₃-GPE analogues was first investigated using the (*S*)- and (*R*)- α -CF₃-proline **1**. Due to the very short GPE sequence, the synthesis was achieved in solution phase using the standard peptide elongation methodology. The coupling reaction between the glutamic acid residue (Glu) and the α -CF₃-proline **1** was performed following our reported procedure in order to avoid the diketopiperazine formation.²⁷ Because of the strong deactivation of the α -CF₃-proline **1** amino group induced by the electron-withdrawing effect of the neighboring CF₃ group,¹⁷ its *N*-protection was not required to perform the coupling

reaction in contrast with the nonfluorinated series. The four enantiopure dipeptides **6** [(*S,S*)-**6**, (*R,S*)-**6**, (*R,R*)-**6**, (*S,R*)-**6**] were obtained by coupling (*R*)- and (*S*)- α -CF₃-proline **1** with *L*- or *D*-glutamic acids (Scheme 1). Therefore, the addition of 1 equiv of α -CF₃-proline **1** to 2 equiv amount of glutamic acid dibenzyl ester in the presence of coupling reagents afforded the corresponding dipeptides **6** without any trace of diketopiperazine. The expected dipeptides were obtained in 42 to 89% yield depending on the *D*- or *L*-series of the glutamic acid (Scheme 1).

The next step involved the coupling reaction of the glycine residue (Gly) at the *N*-terminal position of the dipeptides **6**. Despite the small steric hindrance of the Gly residue, this reaction remained highly challenging because of the very low nucleophilicity of the dipeptide amino group and the steric bulkiness of the α -CF₃-group. Only very few examples are found in the literature for the *N*-terminal coupling of acyclic α -CF₃-amino acids.^{9,48–50} All these examples required specific activation methods, such as mixed anhydrides or acyl halides, to promote the peptide coupling in good yield. To our knowledge, the *N*-terminal coupling reaction has never been reported so far for cyclic α -CF₃-amino acids such as α -CF₃-proline. First, we


Table 1. Optimization of the Synthesis of the Tripeptide (S,S)-7 Using Amino Acid Chloride Activation



entry	base (equiv)	temperature (°C)	time (h)	tripeptide (S,S)-7 yield (%)
1	DIEA (1)	r.t.	48	0 ^a
2	DIEA (2)	60 ^b	24	15 ^c
3	–	60 ^b	24	88 ^c
4	–	microwave	0.5	54 ^c

^aStarting material was recovered. ^bReaction performed in a sealed tube. ^cObtained as a single diastereomer.

Scheme 2. Synthesis of Tripeptides 7



checked the use of a mixed anhydride obtained from isobutylchloroformate and Cbz-glycine. However, no coupling product was obtained from the dipeptide (S,S)-6 whatever the solvent used (ethyl acetate or DMF) and only starting material

was recovered. The use of acyl halide activation was then investigated. Amino acid fluorides are less moisture sensitive than amino acid chlorides and are recognized as excellent coupling reagents for both solution and solid-phase peptide


syntheses.⁵¹ They are particularly efficient for the coupling of sterically hindered amino acids but require the treatment with a silylating agent such as *N,O*-bis(trimethylsilyl)acetamide (BSA).⁵² A coupling test between the preactivated dipeptide (*S,S*)-**6** and the Fmoc-glycine fluoride, prepared according to Olah's method,⁵³ was attempted without success and only starting material was recovered. We postulated that the reactivity of the acyl fluoride was not enough to overcome the stereoelectronic deactivation of the fluorinated dipeptide **6** and we decided to try the more reactive amino acid chlorides. Because of the hydrolysis risk, epimerization and other side reactions, the acyl chlorides have been considered for a long time as overactivated species.⁵⁴ Thanks to the significant contribution of Carpino and his group, the use of the Fmoc-amino acid chlorides has turned out to be one of the most efficient coupling strategies.⁵⁵ However, particular attention is required to ensure the coupling. The Fmoc-glycyl chloride was prepared by treatment of the Fmoc-glycine with thionyl chloride in DCM under ultrasonication and obtained in pure form by crystallization from pentane.⁵⁶ Because of the release of HCl during the amide bond formation, the coupling reaction of the Fmoc-glycyl chloride with the dipeptide (*S,S*)-**6** was first performed in the presence of DIEA. As summarized in Table 1, no reaction occurred at room temperature and only the starting material was recovered even after 48 h (Table 1, entry 1). The increase of the temperature at 60 °C in a sealed tube led to the formation of the tripeptide (*S,S*)-**7** in a low yield (Table 1, entry 2). We postulated that the neutralization of the HCl released from the reaction would prevent the acidic activation of the amino acid chloride. As the amino group of the fluorinated dipeptide **6** is strongly deactivated, we anticipated that the HCl release would not be a problem and the reaction was performed without base. Indeed, the coupling reaction of the Fmoc-glycyl chloride with the dipeptide (*S,S*)-**6** at 60 °C in the absence of DIEA gave the corresponding tripeptide (*S,S*)-**7** in a very good yield (88%) without epimerization (Table 1, entry 3). The substitution of the conventional heating method by microwave irradiation, reported to accelerate the peptide syntheses, allowed the coupling in about 30 min but the yield decreased to 54% (Table 1, entry 4).

The optimized coupling conditions (Table 1, entry 3) were then successfully applied to the dipeptides (*R,S*)-**6**, (*S,R*)-**6** and (*R,R*)-**6** to afford the corresponding tripeptides **7** in 52–67% yield without epimerization (Scheme 2).

The access to the α -CF₃-proline containing GPE analogues required the full removal of the protecting groups. The sequential cleavages were first considered in classical manner as usually reported in peptide chemistry when using orthogonal protecting groups. Unexpectedly the standard Fmoc deprotection by treatment with piperidine in DMF led to degradation. The use of DBU in DCM gave the same result. Since both Fmoc and Bn protecting groups are sensitive to hydrogenolysis, their removal in a single step was envisioned. The hydrogenolysis of (*S,S*)-**7** and (*R,S*)-**7** tripeptides was carried out under hydrogen atmosphere (1 bar) in methanol in the presence of an excess amount of Pd/C catalyst (140%mol of Pd) to afford the CF₃-GPE analogues (*S,S*)-**8** and (*R,S*)-**8** in 55% and 54% yield respectively (Scheme 3). Surprisingly the use of higher hydrogen pressure (3–5 bar) led only to degradation.

CF₃-Pseudoproline Containing GPE Analogues. We then focused our attention on the CF₃-pseudoproline template. It is anticipated that this 5-CF₃-proline surrogate should

Scheme 3. Synthesis of CF₃-Proline Containing GPE Analogues (*S,S*)-**8** and (*R,S*)-**8**




constitute a valuable tool to control the *cis*–*trans* isomerization of the glycyl-pseudoprolyl bond.^{23,28,30} The synthesis was first attempted following the same peptide elongation sequence as used for the α -CF₃-proline. The pseudoproline (*S,S*)-**9** was prepared according to our reported procedure by condensation of serine methyl ester with fluoral.²³ After saponification of the ester function of the pseudoproline **9**, its coupling reaction with L-glutamic acid dibenzyl ester was carried out at the C-terminal position according to the standard procedure using EDCI and HOBT (Scheme 4). The corresponding dipeptide (*S,S,S*)-**10** was obtained in 72% yield as a diastereomeric mixture.

At this stage, the absolute configuration of the minor diastereomer could not be unambiguously assigned. This synthetic pathway being stereochemically irrelevant, we decided to adopt a reverse strategy for the peptide elongation and to start with the *N*-terminal coupling of pseudoprolines **9**. We have recently reported the coupling reactions of a mixture of *cis* and *trans* pseudoprolines **9** with various Fmoc protected amino acid chlorides in base free conditions.³⁰ We demonstrated that the reaction involved a dynamic kinetic resolution (DKR) process. Indeed only the corresponding dipeptides bearing a *cis* oxazolidine were obtained regardless the configuration at the C^δ position of the starting pseudoproline **9**. We postulated that the *trans* (*S,S*)-**9** and the *cis* (*R,S*)-**9** pseudoprolines interconvert rapidly throughout a ring opening equilibrium promoted by the acidic conditions (Scheme 5). Due to steric effects, the *N*-acylation of the *trans* pseudoproline (*S,S*)-**9** is strongly disfavored and only the *cis* (*R,S*)-**9** pseudoproline reacts to afford exclusively the dipeptide bearing the *cis* (*R,S*)-oxazolidine.^{57,58} It is important to note that no epimerization of the C^δ occurs once the pseudoprolines are *N*-acylated.


Applied to Fmoc-glycine chloride, the reaction of a diastereomeric mixture of pseudoprolines (*S,S*)-**9** and (*R,S*)-**9** gave the dipeptide (*R,S*)-**11** in 97% yield as a single diastereomer (Scheme 6). Compared to the conditions used with the CF₃-proline template (Table 1), the reaction was carried out in milder conditions at room temperature using only a slight excess amount (1.1 equiv) of acyl chloride.

These conditions were applied to the thiazolidines **12** prepared by condensation of cysteine methyl ester with fluoral.²³ Starting from a diastereomeric mixture of the *trans* (*S,R*)-**12** and *cis* (*R,R*)-**12** thiazolidines, the coupling reaction led to the expected dipeptide (*R,R*)-**13** but with a low conversion. The monitoring of the reaction progress by TLC


Scheme 4. C-Terminal Coupling Reaction of the Pseudoproline (S,S)-9



Scheme 5. Dynamic Kinetic Resolution Process during N-Coupling of Pseudoprolines 9



Scheme 6. N-Terminal Coupling Reaction of Pseudoprolines 9




and ¹⁹F NMR analyses shown a complete disappearance of the *cis* thiazolidine (R,R)-12 without any reaction of the *trans* (S,R)-12 diastereomer. The coupling reactions with Fmoc-Gly-Cl were then performed separately on the *trans* (S,R)-12 and *cis* (R,R)-12 thiazolidines (Scheme 7). The thiazolidine (R,R)-12 afforded the expected dipeptide (R,R)-13 in 89% yield as a single diastereomer while no reaction occurred with the (S,R)-12 thiazolidine (Scheme 7).




As observed with the oxazolidines 9, the N-coupling reaction is strongly dependent on steric effects and only the *cis* thiazolidine (R,S)-12 reacts to afford the corresponding (R,S)-13 dipeptide. In agreement with the literature in the nonfluorinated series,^{59,60} we assumed that, unlike the oxazolidines 9, the trifluoromethylated thiazolidines (R,R)-12 and (S,R)-12 are stable toward acidic media and cannot interconvert throughout a ring opening equilibrium. In a previous work,²³ we reported that a Lewis acid treatment (BF₃·

OEt₂) can mediate the ring opening and promote the epimerization of the *trans* (S,S)-9 oxazolidine into the *cis* (R,S)-9 (Scheme 8). The higher thermodynamic stability of the *trans* (S,S)-9 oxazolidine compared to the *cis* (R,S)-9 one induces a slow equilibrium shift toward the *trans* (S,S)-9. In contrast, no epimerization occurred when the *trans* (S,R)-12 thiazolidine was treated with BF₃·OEt₂, Ti(OiPr)₄ or Ti(OiPr)₃Cl as Lewis acids. The reaction with TiCl₄ or SnCl₄ gave only degradation products.

In order to synthesize oxazolidine-type CF₃-pseudoproline containing GPE analogues, the saponification of the methyl ester dipeptide (R,S)-11 was carried out under mild conditions to avoid early Fmoc deprotection.⁶¹ The treatment using NaOH in *i*PrOH/H₂O (7:3) solution of CaCl₂ (0.84 M) gave the corresponding acid which was directly engaged in the coupling reaction with respectively L- and D-glutamic acid dibenzyl ester (Scheme 9). Surprisingly, the reaction under the standard conditions (EDCl/HOBt) which were effective for the CF₃-proline template (Scheme 1) failed and the starting material was recovered. The use of BOP-Cl allowed the synthesis of the expected tripeptide (R,S,S)-14 in 68% yield without epimerization (Scheme 9). The removal of all the protecting groups was performed following the same procedure used for the CF₃-proline template. In a single step, the hydrogenolysis of the tripeptide (R,S,S)-14 afforded the corresponding CF₃-GPE analogues (R,S,S)-15 in good yield (72%).

Scheme 7. N-Terminal Coupling Reaction of Pseudoprolines 12




Scheme 8. Lewis Acid-Mediated Epimerization of Pseudoprolines **9** and **12**Scheme 9. Synthesis of CF₃-Pseudoproline Containing GPE Analogues (*R,S,S*)-**14** and (*R,S,S*)-**15**Scheme 10. *N*-Terminal Coupling Reaction of Oxazolidines **16** and **17**

These optimized conditions were applied for the coupling of the thiazolidine containing dipeptide (*R,R*)-**13** with *L*-glutamic acid dibenzyl ester. Unfortunately, the reaction led only to the recovery of the starting material. Other attempts using EDCI/HOBt coupling reagents or the mixed anhydride activation by treatment of (*R,R*)-**13** with isobutylchloroformate in the presence of NMM failed.

2-CF₃-Oxazolidine Containing GPE Analogues. The last investigated GPE analogue series ambitioned to incorporate the trifluoropyruvate oxazolidine template in place of the proline. According to our previously reported procedure,²¹ the CF₃-oxazolidine **16** was obtained in a high yield (95%) as a 75:25 diastereomeric mixture by the condensation of the ethyl trifluoropyruvate with *N*-Boc-(*R*)-phenylglycinol in the presence of PPTS (Scheme 10). We already reported that this oxazolidine proved to be a convenient building block for the synthesis of various enantiopure α-CF₃-amino acids^{20–22,24} and amino alcohols.⁶² In other respect, we consider herein this kind of oxazolidines as original stable proline surrogates. Indeed, the conjugated electron withdrawing effect of both the trifluoromethyl and the carboxylic group prevents the oxazolidine ring opening. Likewise, the unsubstituted racemic oxazolidine **17** was obtained in 44% yield through the condensation reaction of ethyl trifluoropyruvate with *N*-Boc-ethanolamine (Scheme 10).

As both diastereomers of **16** and the racemic oxazolidine **17** can be considered as 2-CF₃-proline surrogates (2-CF₃-ΨPro), we decided to investigate their incorporation into the GPE peptide sequence. Following the synthetic strategy successfully employed in the CF₃-pseudoproline series based on the *N*- to *C*-termini peptide elongation (vide supra), the coupling reaction between a diastereomeric mixture of oxazolidine **16** and the glycine residue was attempted (Scheme 10). However, the expected dipeptide was not obtained whatever was the glycine activation (acyl chloride or acyl bromide) or the protocol used (room temperature, conventional heating or microwave activation, zinc activation, stoichiometry and solvent nature). In order to decrease the steric impact of the oxazolidine, we decided to perform the coupling reaction starting from the unsubstituted racemic oxazolidine **17** (Scheme 10). Therefore, **17** was coupled with Fmoc-glycine chloride to afford the racemic dipeptide **18** in a good yield (75%).

The rest of the synthesis to get the GPE analogues was achieved following the protocol described for the CF₃-pseudoproline template. A saponification reaction followed by the coupling reaction of the dipeptide **18** with the dibenzylated glutamic acid gave the tripeptide **19** as a 57:43 diastereomeric mixture in 52% yield (Scheme 11). A pure fraction of each

Scheme 11. Synthesis of the 2-CF₃-Oxazolidine Containing GPE Analogue 20

diastereomer was obtained by flash chromatography separation. However, we were unable to assign the absolute configuration of the oxazolidine core of each diastereomer. Finally, the isolated diastereomer of the tripeptide **19**_{min} was submitted to hydrogenolysis to give the CF₃-GPE analogue **20** in 49% yield.

CONCLUSIONS

We have developed convenient methods for the incorporation of various CF₃-proline surrogates (α -CF₃-prolines, CF₃-pseudoproline and 2-CF₃-oxazolidines) within peptide sequences. As a result of this study, we demonstrated that the choice of the peptide elongation route (C- to N-terminus or N- to C-terminus) was crucial for the success of the peptide synthesis and also to guarantee its configuration integrity. For the CF₃-proline template, the standard peptide elongation route (C- to N-terminus) was applied, while the reverse strategy was required for the CF₃-pseudoproline and CF₃-oxazolidine templates. Noteworthy, the Fmoc-benzyl strategy allowed the final full removal of the protecting groups in single step by hydrogenolysis. The methods described in this paper have been successfully applied to the solution phase synthesis of several novel trifluoromethylated analogues of the neuroprotective GPE peptide. Evaluation of their neuroprotective effects on different types of neurons from diverse induced injuries activity are in progress and will be reported in due time. Moreover, for future developments, the fluorinated Fmoc-protected peptide intermediates elaborated in this work will constitute suitable peptide blocks for the solid phase synthesis of longer peptides incorporating CF₃-proline or its surrogates.

EXPERIMENTAL SECTION

General Methods. Unless otherwise mentioned, all the reagents were purchased from commercial source. All glassware was dried in an oven at 150 °C prior to use. All solvents were purified and dried by standard techniques and distilled prior to use. Dichloromethane was distilled over calcium hydride under argon. THF was distilled over sodium benzophenone ketyl under argon. All organic extracts were dried over MgSO₄, unless otherwise noted. Silica gel (230–400 mesh) was used for flash column chromatography, eluting (unless otherwise stated) with cyclohexane/ethyl acetate. Silica TLC plates were visualized under UV light, by a 10% solution of phosphomolybdic acid in ethanol followed by heating. Infrared spectra (IR) were obtained by Fourier transformation, and wave numbers are given in cm⁻¹. ¹H NMR, ¹³C NMR, and ¹⁹F NMR spectra were recorded in CDCl₃ (unless otherwise stated). ¹H NMR (400.00 MHz), ¹³C NMR (100.50 MHz) and ¹⁹F NMR (376.20 MHz) were measured on a

spectrometer operating at a ¹H frequency of 400 MHz. Chemical shifts of ¹H NMR are expressed in parts per million downfield from tetramethylsilane ($\delta = 0$) in CDCl₃. Chemical shifts of ¹³C NMR are expressed in parts per million downfield from CDCl₃ as internal standard ($\delta = 77.0$). Chemical shifts of ¹⁹F NMR are expressed in parts per million downfield from C₆F₆ as an internal standard ($\delta = -164.9$). Coupling constants are reported in Hertz. Assignments were obtained from the analysis of 2D ¹H–¹³C heteronuclear correlation (HETCOR) spectroscopy. Correlation spectroscopy (COSY) was used to correlate chemical shifts of protons coupled to one another. Melting points were uncorrected. High-resolution mass spectra were obtained using electrospray ionization (ESI) in positive ion mode and a TOF mass analyzer or using direct inlet probe (DI-HRMS).

CF₃-Proline Containing GPE Analogues. General Procedure for the Synthesis of Dipeptides (S,S)-6, (S,R)-6, (R,S)-6, (R,R)-6. Triethylamine (4.1 equiv), HOBT (1.5 equiv), EDCI (1.5 equiv), and finally α -Tfm-proline **1** (1 equiv) were successively added at 0 °C to a stirred solution of glutamic dibenzylester tosylate salt (2 equiv) in DMF (0.25 M/ α -Tfm amino acid). DMF was added and the resulting mixture was stirred at 0 °C for 20 min and then at room temperature for 24 h. The mixture was diluted with DCM and water. The aqueous layer was extracted with DCM (3 \times) and the combined organic layers were washed with water, dried with MgSO₄, filtered, and concentrated under reduced pressure. The crude residue was purified by flash chromatography to afford the corresponding dipeptides in 42–89% yields.

H-(S)- α -Tfm-Pro-L-Glu(OBn)-OBn (S,S)-6. The dipeptide (S,S)-6 was prepared by the General Procedure, with L-glutamic dibenzylester tosylate salt (961 mg, 1.92 mmol, 2 equiv), triethylamine (540 μ L, 3.94 mmol, 4.1 equiv), HOBT (195 mg, 1.44 mmol, 1.5 equiv), EDCI (275 mg, 1.44 mmol, 1.5 equiv), and (S)- α -Tfm proline **1** (176 mg, 0.96 mmol) in DMF (4 mL). Purification on silica gel (cyclohexane/AcOEt, 90:10) gave pure (S,S)-6 (388 mg, 82%) as a colorless oil: $[\alpha]_D^{25} -19.6$ (*c* 0.9, CHCl₃); IR (neat) 3021, 1740, 1685 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.70 (m, 1 H, H _{γ} Pro-Ha), 1.85 (m, 1 H, H _{γ} Pro-Hb), 2.07 (m, 1 H, H _{β} Glu-Ha), 2.15–2.46 (m, 6 H, NH Pro, H _{β} Glu-Hb, H _{γ} Glu and H _{β} Pro), 3.01–3.11 (m, 2 H, H _{δ} Pro-Ha), 4.62 (m, 1 H, Ha Glu), 5.09 (s, 2 H, Bz CH₂), 5.14 (d, *J* = 12.4 Hz, 1 H, Bn CH₂-Ha), 5.18 (d, *J* = 12.4 Hz, 1 H, Bn CH₂-Hb), 7.30–7.38 (m, 10 H, Bn arom.), 8.11 (d, *J* = 6.9 Hz, 1 H, NH Glu); ¹³C NMR (100.5 MHz, CDCl₃) δ 25.4 (CH₂, C _{γ} Pro), 26.9 (CH₂, C _{β} Glu), 30.1 (CH₂, C _{γ} Glu), 32.3 (CH₂, C _{β} Pro), 47.5 (CH₂, C _{δ} Pro), 51.9 (CH, C α Glu), 66.6 (CH₂, Bn CH₂), 67.4 (CH₂, Bn CH₂), 70.7 (q, *J* = 26.8 Hz, CH, C α Pro), 125.9 (q, *J* = 284.7 Hz, CF₃), 128.2, 128.3, 128.5, 128.6, 128.6, 135.0, 135.6, 169.5, 171.1, 172.3; ¹⁹F NMR (376.2 MHz, CDCl₃) δ -77.8 (s, CF₃); DI-HRMS calcd. for C₂₅H₂₇F₃N₂O₅ [M]⁺ 492.1872, found 492.1873.

H-(R)- α -Tfm-Pro-D-Glu(OBn)-OBn (R,R)-6. The dipeptide (R,R)-6 was prepared by the General Procedure, from D-glutamic dibenzylester

tosylate salt (332 mg, 0.67 mmol, 2 equiv), triethylamine (181 μ L, 1.36 mmol, 4.1 equiv), HOBt (68 mg, 0.50 mmol, 1.5 equiv), EDCI (96 mg, 0.50 mmol, 1.5 equiv), and (R)- α -Tfm proline **1** (61 mg, 0.33 mmol) in DMF (1.5 mL). Purification on silica gel (cyclohexane/AcOEt, 90:10) gave pure (R,R)-**6** (110 mg, 67%) as a colorless oil: $[\alpha]_D^{25} + 22.9$ (c 1.0, CHCl₃); The spectral data of (R,R)-**6** were identical to those of (S,S)-**6**; DI-HRMS calcd. for C₂₅H₂₇F₃N₂O₅ [M]⁺ 492.1872, found 492.1873.

H-(R)- α -Tfm-Pro-L-Glu(OBn)-OBn (R,S)-**6**. The dipeptide (R,S)-**6** was prepared by the General Procedure, with L-glutamic dibenzylester tosylate salt (764 mg, 1.53 mmol, 2 equiv), triethylamine (420 μ L, 3.14 mmol, 4.1 equiv), HOBt (155 mg, 1.15 mmol, 1.5 equiv), EDCI (219 mg, 1.15 mmol, 1.5 equiv), and (R)- α -Tfm proline **1** (140 mg, 0.76 mmol) in DMF (3 mL). Purification on silica gel (cyclohexane/AcOEt, 90:10) gave pure (R,S)-**6** (334 mg, 89%) as a colorless oil: $[\alpha]_D^{25} + 18.1$ (c 0.9, CHCl₃); IR (neat) 3332, 3019, 2925, 1736, 1685 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.69 (m, 1 H, H_γ Pro-Ha), 1.79 (m, 1 H, H_γ Pro-Hb), 2.04 (m, 1 H, H_β Glu-Ha), 2.11–2.48 (m, 6 H, NH Pro, H_β Glu-Hb, H_γ Glu and H_β Pro), 3.00–3.12 (m, 2 H, H_δ Pro), 4.60 (m, 1 H, H_α Glu), 5.09 (s, 2 H, Bn CH₂), 5.12 (d, J = 12.1 Hz, 1 H, Bn CH₂-Ha), 5.18 (d, J = 12.1 Hz, 1 H, Bn CH₂-Hb), 7.28–7.37 (m, 10 H, Bn arom.), 8.29 (d, J = 8.5 Hz, 1 H, NH Glu); ¹³C NMR (100.5 MHz, CDCl₃) δ 25.1 (CH₂, C_γ Pro), 26.9 (CH₂, C_β Glu), 30.2 (CH₂, C_γ Glu), 32.0 (CH₂, C_β Pro), 47.5 (CH₂, C_δ Pro), 51.8 (CH, C_α Glu), 66.5 (CH₂, Bn CH₂), 67.4 (CH₂, Bn CH₂), 71.0 (q, J = 26.8 Hz, CH, C_α Pro), 125.8 (q, J = 283.7 Hz, CF₃), 127.9, 127.9, 128.0, 128.2, 128.3, 135.1, 135.7, 169.4, 171.1, 172.4; ¹⁹F NMR (376.2 MHz, CDCl₃) δ -77.8 (s, CF₃); DI-HRMS calcd. for C₂₅H₂₇F₃N₂O₅ [M]⁺ 492.1872, found 492.1874.

H-(S)- α -Tfm-Pro-D-Glu(OBn)-OBn (S,R)-**6**. The dipeptide (S,R)-**6** was prepared by the General Procedure, with D-glutamic dibenzylester tosylate salt (605 mg, 1.12 mmol, 2 equiv), triethylamine (326 μ L, 2.46 mmol, 4.1 equiv), HOBt (122 mg, 0.90 mmol, 1.5 equiv), EDCI (173 mg, 0.90 mmol, 1.5 equiv), and (S)- α -Tfm proline **1** (111 mg, 0.60 mmol) in DMF (3 mL). Purification on silica gel (cyclohexane/AcOEt, 90:10) gave pure (S,R)-**6** (124 mg, 42%) as a colorless oil: $[\alpha]_D^{25} -16.6$ (c 1.0, CHCl₃); The spectral data of (S,R)-**6** were identical to those of (R,S)-**6**; DI-HRMS calcd. for C₂₅H₂₇F₃N₂O₅ [M]⁺ 492.1872, found 492.1874.

General Procedure for the Synthesis of Dipeptides (S,S)-7, (S,R)-7, (R,S)-7, (R,R)-7. To a solution of dipeptide **6** (1 equiv) in DCM under argon was added Fmoc-Gly-Cl (2 equiv) prepared according to our reported procedure.³⁰ The mixture was refluxed for 24 h and then quenched by saturated NaHCO₃ solution. The aqueous phase was extracted by DCM (3 \times) and the combined organic extracts were dried over MgSO₄ and concentrated under reduced pressure. The crude residue was purified by flash chromatography to afford the corresponding tripeptides in 52–88% yield.

Fmoc-Gly-(S)- α -Tfm-Pro-L-Glu(OBn)-OBn (S,S)-**7**. The tripeptide (S,S)-**7** was prepared following the General Procedure, using dipeptide (S,S)-**6** (552 mg, 1.12 mmol, 1 equiv) and Fmoc-Gly-Cl (708 mg, 2.24 mmol, 2 equiv) in DCM (4 mL). Purification by flash chromatography (70:30 cyclohexane/ethyl acetate) gave pure (S,S)-**7** (757 mg, 88%) as a white solid: mp 110–112 °C; $[\alpha]_D^{25} -53.7$ (c 0.23, CHCl₃); IR (neat) 3300, 1728, 1681 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.95–2.15 (m, 3 H, H_γ Pro and H_β Glu-Ha), 2.29 (m, 1 H, H_β Glu-Hb), 2.30–2.59 (m, 4 H, H_γ Glu and H_β Pro), 3.60 (q, J = 8.0 Hz 1 H, H_δ Pro-Ha), 3.71 (q, J = 8.0 Hz, 1 H, H_δ Pro-Hb), 4.03–4.09 (m, 2 H, H_α Gly), 4.20 (t, J = 7.2 Hz, 1 H, Fmoc CH), 4.30–4.40 (m, 2 H, Fmoc CH₂), 4.66 (m, 1 H, H_α Glu), 5.09 (s, 2 H, Bn CH₂), 5.14 (d, J = 12.1 Hz, 1 H, Bn CH₂-Ha), 5.16 (d, J = 12.1 Hz, 1 H, Bn CH₂-Hb), 5.73 (s, 1 H, NH Gly), 7.21 (d, J = 6.9 Hz, 1 H, NH Glu), 7.27–7.35 (m, 12 H, Fmoc arom. and Bn arom.) 7.41 (t, J = 7.3 Hz, 2 H, Fmoc arom.), 7.60 (d, J = 7.3 Hz, 2 H, Fmoc arom.), 7.76 (d, J = 7.3 Hz, 2 H, Fmoc arom.); ¹³C NMR (100.5 MHz, CDCl₃) δ 23.3 (CH₂, C_γ Pro), 26.4 (CH₂, C_β Glu), 29.9 (CH₂, C_γ Glu), 34.1 (CH₂, C_β Pro), 44.1 (CH₂, C_α Gly), 47.0 (CH, Fmoc CH), 48.5 (CH₂, C_δ Pro), 52.7 (CH, C_α Glu), 66.5 (CH₂, Bn CH₂), 67.1 (CH₂, Fmoc CH₂), 67.4 (CH₂, Bn CH₂), 72.0 (q, J = 28.8 Hz, CH, C_α Pro), 119.9, 124.9 (q, J = 286.6 Hz, CF₃), 125.1, 127.0, 127.6, 128.1, 128.2, 128.3, 128.5,

128.6, 135.0, 135.6, 141.2, 143.8, 156.1, 165.9, 167.9, 170.9, 172.9; ¹⁹F NMR (376.2 MHz, CDCl₃) δ -71.3 (s, CF₃); HRMS (ESI-TOF) calcd. for C₄₂H₄₁F₃N₃O₈ [M + H]⁺ 772.2846, found 772.2881.

Fmoc-Gly-(S)- α -Tfm-Pro-D-Glu(OBn)-OBn (S,R)-**7**. To a solution of dipeptide (S,R)-**6** (67 mg, 1.12 mmol, 1 equiv) in DCM (1 mL) under argon was added Fmoc-Gly-Cl (86 mg, 2.24 mmol, 2 equiv). Purification by flash chromatography (70:30 cyclohexane/ethyl acetate) gave pure (S,R)-**7** (55 mg, 52%) as a white solid: mp 110–112 °C; $[\alpha]_D^{25} -21.4$ (c 0.44, CHCl₃); IR (neat) 1728, 1681, 1519, 1450 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.92–2.15 (m, 3 H, H_γ Pro and H_β Glu-Ha), 2.20 (m, 1 H, H_β Glu-Hb), 2.34–2.52 (m, 4 H, H_γ Glu and H_β Pro), 3.55 (q, J = 8.0 Hz 1 H, H_δ Pro-Ha), 3.67 (q, J = 8.0 Hz, 1 H, H_δ Pro-Hb), 3.97 (dd, J = 17.4, 4.0 Hz, 1 H, H_α Gly-Ha), 4.11–4.21 (m, 2 H, H_α Gly-Hb and Fmoc CH), 4.28 (dd, J = 10.3, 6.0 Hz, 1 H, Fmoc CH₂-Ha), 4.37 (dd, J = 10.3, 7.3 Hz, 1 H, Fmoc CH₂-Hb), 4.63 (m, 1 H, H_α Glu), 5.01 (d, J = 12.1 Hz, 1 H, Bn CH₂-Ha), 5.09 (d, J = 12.1 Hz, 1 H, Bn CH₂-Hb), 5.10 (d, J = 12.1 Hz, 1 H, Bn CH₂-Ha), 5.19 (d, J = 12.1 Hz, 1 H, Bn CH₂-Hb), 5.71 (m, 1 H, NH-Gly), 7.27–7.35 (m, 13 H, NH Glu, Fmoc arom. and Bn arom.), 7.40 (t, J = 7.6 Hz, 2 H, Fmoc arom.), 7.57 (d, J = 7.6 Hz, 2 H, Fmoc arom.), 7.76 (d, J = 7.6 Hz, 2 H, Fmoc arom.); ¹³C NMR (100.5 MHz, CDCl₃) δ 22.9 (CH₂, C_γ Pro), 25.7 (CH₂, C_β Glu), 30.0 (CH₂, C_γ Glu), 34.4 (CH₂, C_β Pro), 44.2 (CH₂, C_α Gly), 46.9 (CH, Fmoc CH), 48.5 (CH₂, C_δ Pro), 52.3 (CH, C_α Glu), 66.8 (CH₂, Bn CH₂), 67.1 (CH₂, Fmoc CH₂), 67.3 (CH₂, Bn CH₂), 72.0 (q, J = 27.8 Hz, CH, C_α Pro), 119.9, 124.5 (q, J = 286.6 Hz, CF₃), 125.1, 127.0, 127.7, 127.9, 128.3, 128.4, 128.5, 128.6, 135.0, 135.3, 141.2, 143.8, 156.1, 166.2, 167.9, 171.0, 174.2; ¹⁹F NMR (376.2 MHz, CDCl₃) δ -71.1 (s, CF₃); HRMS (ESI-TOF) calcd. for C₄₂H₄₁F₃N₃O₈ [M + H]⁺ 772.2846, found 772.2864.

Fmoc-Gly-(R)- α -Tfm-Pro-L-Glu(OBn)-OBn (R,S)-**7**. The tripeptide (R,S)-**7** was prepared following the General Procedure, using dipeptide (R,S)-**6** (190 mg, 0.39 mmol, 1 equiv) and Fmoc-Gly-Cl (243 mg, 0.77 mmol, 2 equiv) in DCM (2 mL). Purification by flash chromatography (70:30 cyclohexane/ethyl acetate) gave pure (R,S)-**7** (159 mg, 53%) as a white solid: mp 110–112 °C; $[\alpha]_D^{25} + 23.1$ (c 0.45, CHCl₃); The spectral data of (R,S)-**7** were identical to those of (S,R)-**7**; HRMS (ESI-TOF) calcd. for C₄₂H₄₁F₃N₃O₈ [M + H]⁺ 772.2846, found 772.2864.

Fmoc-Gly-(R)- α -Tfm-Pro-D-Glu(OBn)-OBn (R,R)-**7**. The tripeptide (R,R)-**7** was prepared following the General Procedure, using dipeptide (R,R)-**6** (62 mg, 0.13 mmol, 1 equiv) in DCM (1 mL) under argon was added Fmoc-Gly-Cl (80 mg, 0.26 mmol, 2 equiv). Purification by flash chromatography (70:30 cyclohexane/ethyl acetate) gave pure (R,R)-**7** (70 mg, 72%) as a white solid: mp 110–112 °C; $[\alpha]_D^{25} + 52.2$ (c 0.25, CHCl₃); The spectral data of (R,R)-**7** were identical to those of (S,S)-**7**; HRMS (ESI-TOF) calcd. for C₄₂H₄₁F₃N₃O₈ [M + H]⁺ 772.2846, found 772.2864.

H-Gly-(S)- α -Tfm-Pro-L-Glu(OH)-OH (S,S)-**8**. A solution of tripeptide (S,S)-**7** (440 mg, 0.57 mmol, 1 equiv) in MeOH (2.5 mL) was hydrogenated over 20% Pd/C (440 mg) at room temperature for 24 h under hydrogen atmosphere (1 bar). The reaction mixture was filtered and evaporated. The crude was precipitated in pentane and filtrated to afford pure (S,S)-**8** (116 mg, 55%) as a white solid: $[\alpha]_D^{25} -55.5$ (c 1.3, H₂O); IR (neat) 3600–2750, 1667, 1538 cm⁻¹; ¹H NMR (400 MHz, D₂O) δ 1.90 (m, 1 H, H_β Glu-Ha), 1.98–2.18 (m, 3 H, H_γ Pro-H and H_β Glu-Hb), 2.28–2.38 (m, 3 H, H_β Pro-Ha and H_γ Glu-H), 2.55 (m, 1 H, H_β Pro-Hb), 3.65 (q, J = 8.0 Hz 1 H, H_δ Pro-Ha), 3.79–3.88 (m, 1 H, H_δ Pro-Hb), 3.98 (d, J = 16.7 Hz, 1 H, H_α Gly-Ha), 4.09 (d, J = 16.7 Hz, 1 H, H_α Gly-Hb), 4.21 (m, 1 H, H_α Glu-H); ¹³C NMR (100.5 MHz, D₂O) δ 22.7 (CH₂, C_γ Pro), 27.0 (CH₂, C_β Glu), 30.9 (CH₂, C_γ Glu), 34.5 (CH₂, C_β Pro), 41.3 (CH₂, C_α Gly), 48.7 (CH₂, C_δ Pro), 54.4 (CH, C_α Glu), 71.8 (q, J = 28.8 Hz, CH, C_α Pro), 124.5 (q, J = 285.6 Hz, CF₃), 166.0, 167.8, 176.8, 178.6; ¹⁹F NMR (376.2 MHz, D₂O) δ -71.7 (s, CF₃); HRMS (ESI-TOF) calcd. for C₁₃H₁₉F₃N₃O₆ [M + H]⁺ 370.1226, found 370.1221.

H-Gly-(R)- α -Tfm-Pro-L-Glu(OH)-OH (R,S)-**8**. A solution of tripeptide (R,S)-**7** (159 mg, 0.21 mmol, 1 equiv) in MeOH (1.0 mL) was hydrogenated over 20% Pd/C (159 mg) at room temperature for 24 h under hydrogen atmosphere (1 bar). The reaction mixture was filtered

and evaporated. The crude was precipitated in pentane and filtrated to afford pure (*R,S*)-**8** (41 mg, 54%) as a white solid: $[\alpha]_D^{25}$ -49.3 (*c* 0.85, H₂O); IR (neat) 3600–2750, 1667, 1538 cm⁻¹; ¹H NMR (400 MHz, D₂O) δ 1.90 (m, 1 H, H _{β} Glu-Ha), 2.00–2.18 (m, 3 H, H _{γ} Pro-H and H _{β} Glu-Hb), 2.28–2.48 (m, 3 H, H _{β} Pro-Ha and H _{γ} Glu-H), 2.58 (m, 1 H, H _{β} Pro-Hb), 3.69 (m, 1 H, H _{δ} Pro-Ha), 3.87 (m, 1 H, H _{δ} Pro-Hb), 4.01 (d, *J* = 16.5 Hz, 1 H, H _{α} Gly-Ha), 4.15 (d, *J* = 16.5 Hz, 1 H, H _{α} Gly-Hb), 4.20 (m, 1 H, H _{α} Glu-H); ¹³C NMR (100.5 MHz, D₂O) δ 22.7 (CH₂, C _{γ} Pro), 26.1 (CH₂, C _{β} Glu), 31.7 (CH₂, C _{γ} Glu), 34.5 (CH₂, C _{β} Pro), 41.5 (CH₂, C _{α} Gly), 48.9 (CH₂, C _{δ} Pro), 54.9 (CH, C _{α} Glu), 71.8 (q, *J* = 27.8 Hz, CH, C _{α} Pro), 124.5 (q, *J* = 285.6 Hz, CF₃), 166.3, 168.0, 177.0, 179.6; ¹⁹F NMR (376.2 MHz, D₂O) δ -71.5 (s, CF₃); HRMS (ESI-TOF) calcd. for C₁₃H₁₉F₃N₃O₆ [M + H]⁺ 370.1226, found 370.1222.

CF₃-Pseudoproline Containing GPE Analogues. *H*-(*S*)-Ser($\Psi^{CF_3,H}$ Pro)-L-Glu(OBn)-OBn (*S,S,S*)-**10**. To a solution of oxazolidine (*S,S*)-**9** (2 g, 10.0 mmol) in THF (40 mL) was slowly added at 0 °C 1 M aqueous solution of LiOH (11 mL, 11 mmol, 1.1 equiv). The solution mixture was vigorously stirred at 0 °C until the disappearance of the starting material (usually 1 h). Subsequently, Et₂O was added, and the reaction mixture was extracted with water (3 \times). The aqueous layers were combined, and water was removed under reduced pressure to give the corresponding lithium carboxylate, which was directly used without further purification. A fraction of the crude lithium carboxylate (210 mg, 1.1 mmol) was diluted in DMF (6 mL), and L-glutamic acid dibenzylester tosylate salt (1.1 g, 2.2 mmol, 2.0 equiv), NEt₃ (630 μ L, 4.5 mmol, 4.1 equiv), HOBt (223 mg, 1.65 mmol, 1.5 equiv), and EDCI (316 mg, 1.65 mmol, 1.5 equiv) were successively added at room temperature. The reaction mixture was stirred overnight at room temperature and then diluted with DCM and water. The layers were separated, and the aqueous layer was extracted with DCM (3 \times). The combined organic layers were dried over MgSO₄, filtered, and concentrated under reduced pressure. Purification by flash chromatography (70:30 cyclohexane/ethyl acetate) gave the dipeptide (*S,S,S*)-**10** (392 mg, 72%) as an inseparable 75:25 diastereomeric mixture: ¹H NMR (400 MHz, CDCl₃) (*major*) δ 2.08–2.12 (m, 1 H, H _{β} Glu-Ha), 2.20–2.31 (m, 1 H, H _{β} Glu-Hb), 2.32–2.50 (m, 2 H, H _{γ} Glu), 2.85 (m, 1 H, NH Ψ Pro), 3.81 (t, *J* = 7.0 Hz, 1 H, H _{β} Ψ Pro-Ha), 3.93 (t, *J* = 7.0 Hz, 1 H, H _{α} Ψ Pro), 4.17 (t, *J* = 7.8 Hz, 1 H, H _{β} Ψ Pro-Hb), 4.62–4.69 (m, 1 H, H _{α} Glu), 5.01 (q, *J* = 5.5 Hz, 1 H, H _{δ} Ψ Pro), 5.10 (s, 2 H, Bn CH₂-H), 5.14 (d, *J* = 12.1 Hz, 1 H, Bn CH₂-Ha), 5.18 (d, *J* = 12.1 Hz, 1 H, Bn CH₂-Hb), 7.10 (d, *J* = 7.1 Hz, 1 H, NH-Glu), 7.26–7.40 (m, 10 H, Bn arom.); (*minor*) δ 2.08–2.12 (m, 1 H, H _{β} Glu-Ha), 2.20–2.31 (m, 1 H, H _{β} Glu-Hb), 2.32–2.50 (m, 2 H, H _{γ} Glu), 2.85 (m, 1 H, NH Ψ Pro), 3.86–3.93 (m, 2 H, H _{β} Ψ Pro-Ha and H _{α} Ψ Pro), 4.11 (t, *J* = 7.1 Hz, 1 H, H _{β} Ψ Pro-Hb), 4.62–4.69 (m, 1 H, H _{α} Glu), 4.90 (q, *J* = 5.1 Hz, 1 H, H _{δ} Ψ Pro), 5.10 (s, 2 H, Bn CH₂-H), 5.14 (d, *J* = 12.1 Hz, 1 H, Bn CH₂-Ha), 5.18 (d, *J* = 12.1 Hz, 1 H, Bn CH₂-Hb), 7.10 (d, *J* = 7.1 Hz, 1 H, NH-Glu), 7.26–7.40 (m, 10 H, Bn arom.); ¹³C NMR (100.5 MHz, CDCl₃) (*major*) δ 26.7 (CH₂, C _{β} Glu), 30.2 (CH₂, C _{γ} Glu), 52.0 (CH, C _{α} Glu), 59.3 (CH, C _{α} Ψ Pro), 66.7 (CH₂, Bn CH₂), 67.6 (CH₂, Bn CH₂), 70.2 (CH₂, C _{β} Ψ Pro), 87.9 (q, *J* = 33.6 Hz, CH, C _{δ} Ψ Pro), 123.1 (q, *J* = 283.7 Hz, CF₃), 128.2, 128.3, 128.4, 128.5, 128.6, 128.7, 134.9, 135.5, 170.0, 171.1, 172.7; (*minor*) δ 26.8 (CH₂, C _{β} Glu), 30.2 (CH₂, C _{γ} Glu), 51.8 (CH, C _{α} Glu), 59.3 (CH, C _{α} Ψ Pro), 66.6 (CH₂, Bn CH₂), 67.6 (CH₂, Bn CH₂), 70.0 (CH₂, C _{β} Ψ Pro), 87.9 (q, *J* = 33.6 Hz, CH, C _{δ} Ψ Pro), 123.0 (q, *J* = 282.8 Hz, CF₃), 128.3, 128.3, 128.4, 128.5, 128.6, 128.7, 134.9, 135.6, 170.2, 171.0, 172.5; ¹⁹F NMR (376.2 MHz, CDCl₃) (*major*) δ -84.6 (d, *J* = 5.5 Hz, CF₃); (*minor*) δ -84.4 (d, *J* = 5.1 Hz, CF₃); HRMS (ESI-TOF) calcd. for C₂₄H₂₆F₃N₂O₆ [M + H]⁺ 495.1743, found 495.1732.

The Fmoc-Gly-(*R*)-Ser($\Psi^{CF_3,H}$ Pro)-OMe (*R,S*)-**11** was prepared according to our previously reported procedure.³⁰

Fmoc-Gly-(*R*)-Cys($\Psi^{CF_3,H}$ Pro)-OMe (*R,R*)-13**.** To a solution of (*R,R*)-**12** (3.0 g, 13.9 mmol, 1 equiv) in DCM (150 mL) was added Fmoc-Gly-Cl (4.820 g, 15.3 mmol, 1.1 equiv). The mixture is stirred at room temperature for 24 h then quenched with saturated NaHCO₃ solution (150 mL). The aqueous solution was extracted with DCM (3 \times 100 mL). The combined organic extracts were washed with brine

(150 mL) and were dried over MgSO₄, filtered and concentrated under reduced pressure. The crude mixture is purified by flash chromatography (80:20 cyclohexane/ethyl acetate) to afford pure dipeptide (*R,R*)-**13** (6.08 g, 89%) as a 57/43 inseparable mixture of rotational isomers in CDCl₃ at 298 K. Yellow solid: mp 72–74 °C; $[\alpha]_D^{25}$ -95.9 (*c* 1.07, CHCl₃); IR (neat) 3342, 1716, 1682, 1519, 1118 cm⁻¹; ¹H NMR (400 MHz, CDCl₃, 323 K) δ 3.33–3.65 (m, 2 H, H _{β} Ψ Pro), 3.79 (s, 3 H, OMe), 4.12 (m, 1 H, H _{α} Gly-Ha), 4.22 (t, *J* = 6.9 Hz, 1 H, Fmoc CH), 4.35–4.47 (m, 3 H, H _{α} Gly-Hb and Fmoc CH₂), 5.03 (m, 1 H, H _{α} Ψ Pro), 5.69 (m, 1 H, NH, Gly), 6.02 (m, 1 H, H _{δ} Ψ Pro), 7.31 (t, *J* = 7.3 Hz, 2 H, Fmoc arom.), 7.40 (t, *J* = 7.3 Hz, 2 H, Fmoc arom.), 7.59 (d, *J* = 7.3 Hz, 2 H, Fmoc arom.), 7.76 (d, *J* = 7.3 Hz, 2 H, Fmoc arom.); ¹³C NMR (100.5 MHz, CDCl₃, 323 K) δ 31.9 (CH₂, C _{β} Ψ Pro), 43.2 (CH₂, C _{α} Gly), 47.2 (CH, Fmoc CH), 53.0 (CH₃, OMe), 62.2 (q, *J* = 30.7 Hz, C _{δ} Ψ Pro), 62.8 (CH, C _{α} Ψ Pro), 67.4 (CH₂, Fmoc CH₂), 119.9, 124.2 (q, *J* = 281.8 Hz, CF₃), 125.0, 127.1, 127.7, 141.3, 143.8, 156.3, 168.4, 168.9; ¹⁹F NMR (376.2 MHz, CDCl₃, 298 K) δ (*majo* rotamer) δ -77.1 (s, CF₃), (*mino* rotamer) δ -77.3 (s, CF₃); HRMS (ESI-TOF) calcd. for C₂₃H₂₁F₃N₂O₅Na [M + Na]⁺ 517.1021, found 517.1028.

Fmoc-Gly-(*R*)-Ser($\Psi^{CF_3,H}$ Pro)-L-Glu(OBn)-OBn (*R,S,S*)-14**.** The saponification of the dipeptide methyl ester (*R,S*)-**11** (15.0 g, 31.4 mmol) was performed following a described method by addition of NaOH (1.5 g, 37.6 mmol, 1.2 equiv) to a 0.8 M CaCl₂ solution in iPrOH:H₂O 7:3 (715 mL).⁶¹ The reaction mixture was stirred for 12 h at room temperature, quenched with 1 M HCl, concentrated under reduced pressure and diluted with H₂O (200 mL). The aqueous solution was extracted with ethyl acetate (3 \times). The combined organic layers were washed with H₂O, dried over MgSO₄, filtered and concentrated under reduced pressure. The crude acid was used in the next step without further purification. To a solution of the crude acid (6.0 g, 12.9 mmol, 1 equiv) in DCM (690 mL) were added L-glutamic dibenzylester tosylate salt (8.06 g, 16.1 mmol, 1.25 equiv) and Et₃N (5.5 mL, 39.7 mmol, 3.2 equiv). After 20 min at room temperature BOP-Cl (4.26 g, 16.8 mmol, 1.3 equiv) was added and the mixture was stirred at room temperature for 24 h then quenched with water. The aqueous solution was extracted with DCM (3 \times 200 mL). The organic layer was washed with brine (100 mL). The combined organic extracts were dried over MgSO₄, filtered and concentrated under reduced pressure. The crude mixture is purified by flash chromatography (80:20 cyclohexane/ethyl acetate) to give pure (*R,S,S*)-**14** (6.82 g, 68%) as a 57/43 inseparable mixture of rotational isomers in CDCl₃ at 298 K. Yellow oil: $[\alpha]_D^{25}$ -27.2 (*c* 0.9, CHCl₃); IR (neat) 3310, 2928, 1731, 1677, 1529, 1150, 737 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ (mixture of rotational isomers) 2.00–2.12 (m, 1 H, H _{β} Glu-Ha), 2.21–2.32 (m, 1 H, H _{β} Glu-Hb), 2.34–2.54 (m, 2 H, H _{γ} Glu), 3.82–4.02 (m, 2 H, H _{α} Gly), 4.17–4.26 (m, 2 H, Fmoc CH and H _{β} Ψ Pro-Ha), 4.34–4.44 (m, 3 H, Fmoc CH₂ and H _{β} Ψ Pro-Hb), 4.67–4.77 (m, 1 H, H _{α} Glu), 5.02–5.20 (m, 5 H, H _{α} Ψ Pro and Bn CH₂), 5.69–6.09 (m, 2 H, H _{δ} Pro and NH-Gly), 7.12–7.21 (m, 1 H, NH Glu), 7.27–7.37 (m, 12 H, Bn arom. and Fmoc arom.), 7.41 (t, *J* = 7.3 Hz, 2 H, Fmoc arom.), 7.62 (d, *J* = 7.3 Hz, 2 H, Fmoc arom.), 7.77 (d, *J* = 7.3 Hz, 2 H, Fmoc arom.); ¹³C NMR (100.5 MHz, CDCl₃) δ (mixture of rotational isomers) 26.8 (CH₂, C _{β} Glu), 30.0 (CH₂, C _{γ} Glu), 43.3 and 44.0 (CH₂, C _{α} Gly), 46.8 (CH, Fmoc CH), 51.6 and 51.9 (CH, C _{α} Glu), 58.4 (CH, C _{α} Ψ Pro), 66.4 (CH₂, Bn CH₂), 67.0 (CH₂, Fmoc CH₂), 67.2 (CH₂, Bn CH₂), 67.3 (Fmoc CH₂), 67.4 (CH₂, C _{β} Ψ Pro), 84.7 (CH, C _{δ} Pro), 119.8, 122.5 (q, *J* = 287.5 Hz, CF₃), 124.9, 126.9, 127.6, 128.0, 128.1, 128.3, 128.4, 134.9, 135.4, 141.0, 143.5, 143.6, 156.5, 168.0, 169.2, 170.9, 171.3, 172.4, 172.5; ¹⁹F NMR (376.2 MHz, CDCl₃) δ (mixture of rotational isomers) -81.8 (s, CF₃) and -81.3 (s, CF₃); HRMS (ESI-TOF) calcd. for C₄₁H₃₉F₃N₃O₉ [M + H]⁺ 774.2638, found 774.2653.

H-Gly-(*R*)-Ser($\Psi^{CF_3,H}$ Pro)-L-Glu(OH)-OH (*R,S,S*)-15**.** A solution of tripeptide (*R,S,S*)-**14** (4.0 g, 5.16 mmol, 1 equiv) in MeOH (40 mL) was hydrogenated over 20% Pd/C (4.0 g) at room temperature for 24 h under 1 bar pressure of hydrogen. The reaction mixture was filtered and concentrated under reduced pressure. Water and Et₂O were added, aqueous phase was washed with Et₂O (3 \times 50 mL). The aqueous phase was concentrated under reduced pressure to give pure

deprotected peptide (*R,S,S*)-**15** (1.75 g, 72%) as a single rotational isomer in D₂O at 353 K. Yellow solid: $[\alpha]_D^{25} -31.1$ (*c* 1.2, H₂O); IR (neat) 3600–2750, 1667, 1538 cm⁻¹; ¹H NMR (400 MHz, D₂O, 353 K) δ 2.38–2.52 (m, 1 H, H _{β} Glu-Ha), 2.56–2.70 (m, 1 H, H _{β} Glu-Hb), 2.82–2.91 (m, 2 H, H _{γ} Glu), 4.57–4.62 (m, 2 H, H _{α} Gly), 4.80–4.92 (m, 2 H, H _{α} Glu, H _{β} Ψ Pro-Ha), 5.10–5.20 (m, 1 H, H _{β} Ψ Pro-Hb), 5.40–5.50 (m, 1 H, H _{α} Ψ Pro), 6.45–6.52 (m, 1 H, H _{δ} Ψ Pro); ¹³C NMR (100.5 MHz, D₂O, 353 K) δ 27.7 (CH₂, C _{β} Glu), 31.7 (CH₂, C _{γ} Glu), 41.6 (CH₂, C _{α} Gly), 55.3 (CH, C _{α} Glu), 59.4 (CH, C _{α} Ψ Pro), 71.1 (CH₂, C _{β} Ψ Pro), 85.5 (q, *J* = 34.5 Hz, C _{δ} Ψ Pro), 123.0 (q, *J* = 285.6 Hz, CF₃), 168.3, 169.1, 177.2, 178.8; ¹⁹F NMR (376.2 MHz, D₂O, 353 K) δ -80.4 (s, CF₃); HRMS (ESI-TOF) calcd. for C₁₂H₁₇F₃N₃O₇ [M + H]⁺ 372.1019, found 372.1019.

2-CF₃-Oxazolidine Containing GPE Analogues. The compound **16** was prepared as a mixture of diastereomer (75:25) according to our previously reported procedure.²⁰

2-CF₃- Ψ Pro-OEt-17. To a solution of *N*-Boc-ethanolamine (5.200 g, 33 mmol, 1 equiv) in toluene (100 mL) was added PPTS (829 mg, 3.3 mmol, 0.1 equiv) and ethyl trifluoropyruvate (5.610 g, 102 mmol, 1.2 equiv). After stirring the mixture for 1 h at room temperature, the reaction was heated to reflux with a Dean–Stark apparatus for 24 h then cooled to 0 °C, filtrated and concentrated under reduced pressure. Purification by flash chromatography (90:10 cyclohexane/ethyl acetate) gave a racemic mixture of **17** (5.320 g, 44%) as a yellow oil: IR (neat) 3342, 2987, 2904, 1742, 1448, 1372, 1322, 1279, 1232, 1177, 1016, 987, 943, 858, 817, 747, 681 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.35 (t, 3H, *J* = 7.1 Hz, H_{Et}), 3.16 (q, 1H, *J* = 9.9 Hz, H _{δ} -Ha), 3.32 (bs, 1H, NH), 3.38–3.45 (m, 1H, H _{δ} -Hb), 3.77 (q, 1H, *J* = 7.9 Hz, H _{γ} -Ha), 4.15 (ddd, 1H, *J* = 7.9 Hz, 6.9 Hz, 2.9 Hz, H _{γ} -Hb), 4.27–4.42 (m, 2H, H_{Et}); ¹³C NMR (100.5 MHz, CDCl₃) δ 13.6 (C_{Et}), 46.6 (C _{δ}), 63.6 (C_{Et}), 68.5 (C _{γ}), 93.3 (q, *J* = 31.6 Hz, C _{α}), 122.7 (q, *J* = 286.6 Hz, CF₃), 166.2 (CO); ¹⁹F NMR (376.2 MHz, CDCl₃) δ -83.1 (s, CF₃). HRMS (ESI-TOF) calcd. C₇H₁₁F₃NO₃ [M + H]⁺ 214.0691, found 214.0695.

Fmoc-Gly-2-CF₃- Ψ Pro-OEt (18). To a solution of **17** (220 mg, 1.03 mmol, 1 equiv) in DCM (3 mL) was added Fmoc-Gly-Cl (490 mg, 1.55 mmol, 1.5 equiv). The mixture is stirred at room temperature for 24 h then quenched with saturated NaHCO₃ solution (5 mL). The aqueous solution was extracted with DCM (3 × 10 mL). The organic layer was washed with a saturated NaCl solution (10 mL). The combined organic extracts were dried over MgSO₄, filtered and concentrated under reduced pressure. The crude mixture is purified by flash chromatography (80:20 cyclohexane/ethyl acetate) to give the racemic dipeptide **18** (378 mg, 75%) as a white solid: mp 54–56 °C; IR (neat) 3330, 2922, 1758, 1715, 1682, 1203, 1170, 1033, 759, 738 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.30 (t, *J* = 7.3 Hz, 3H, Et), 3.7 (dd, *J* = 8.2, 7.8 Hz, 1H, H _{δ} Ψ Pro-Ha), 3.92 (m, 1H, H _{δ} Ψ Pro-Hb), 4.02 (dd, *J* = 17.4, 4.6 Hz, 1H, H _{α} Gly-Ha), 4.11 (dd, *J* = 17.4, 6.9 Hz, 1H, H _{α} Gly-Hb), 4.18 (dd, *J* = 7.3, 7.3 Hz, 1H, CHFmoc), 4.29 (q, *J* = 7.3 Hz, 2H, H_{Et}), 4.35–4.39 (m, *J* = 7.3 Hz, 4H, H _{γ} Ψ Pro and CH₂Fmoc), 5.72 (bs, 1H, NH), 7.28 (t, *J* = 7.3, 2H, H_{Ar}Fmoc), 7.38 (t, *J* = 7.3 Hz, 2H, H_{Ar}Fmoc), 7.57 (d, *J* = 7.3 Hz, 2H, H_{Ar}Fmoc), 7.74 (d, *J* = 7.3 Hz, 2H, H_{Ar}Fmoc); ¹³C NMR (100.5 MHz, CDCl₃) δ 13.9 (C_{Et}), 44.1 (C _{α} Gly), 45.2 (C _{δ} Ψ Pro), 47.1 (CHFmoc), 63.1 (C_{Et}), 67.4 (C _{γ} Ψ Pro), 69.0 (CH₂Fmoc), 89.5 (q, *J* = 32.6 Hz, C _{α} Ψ Pro), 120.1 (CFmoc), 122.4 (q, *J* = 288.5 Hz, CF₃), 125.2, 127.2, 127.9, 141.4, 143.8 (CFmoc), 156.3, 163.5, 167.1 (CO); ¹⁹F NMR (376.2 MHz, CDCl₃) δ -79.7 (s, CF₃); DI-HRMS calcd. for C₂₄H₂₃F₃N₃O₆ [M]⁺ 492.1508, found 492.1491.

Fmoc-Gly-2-CF₃- Ψ Pro-L-Glu(OBn)-OBn (19). To a solution of calcium chloride (0.84 M) in isopropanol/water (7:3) (11 mL) were added the compound **18** (237 mg, 0.48 mmol, 1 equiv) and NaOH (23 mg, 0.58 mmol, 1.2 equiv).⁶¹ The mixture was stirred at room temperature for 24 h then quenched by HCl 1 M. The aqueous solution was extracted with ethyl acetate (3 × 10 mL). The organic layer was washed with a saturated NaCl solution (10 mL). The combined organic extracts were dried over MgSO₄, filtered and concentrated under reduced pressure to give the intermediate acid which was used directly in the next step without further purification. To a solution of the crude carboxylic acid intermediate (173 mg, 0.37

mmol, 1 equiv) in DCM (20 mL) were added L-Glu(OBn)-OBn (240 mg, 0.48 mmol, 1.30 equiv) and Et₃N (120 mg, 1.19 mmol, 3.2 equiv). After 20 min at room temperature BOP-Cl (122 mg, 254.57 mmol, 1.3 equiv) was added and the mixture was stirred at room temperature for 24 h then quenched with water. The aqueous solution was extracted with DCM (3 × 10 mL). The organic layer was washed with a saturated NaCl solution (10 mL). The combined organic extracts were dried over MgSO₄, filtered and concentrated under reduced pressure. The crude mixture was purified by flash chromatography (50:50 cyclohexane/ethyl acetate) to give access to the separate tripeptide diastereomers **19**_{maj} (85 mg, 30%) and **19**_{min} (64 mg, 22%).

The diastereomer **19**_{maj} was obtained as colorless foam: $[\alpha]_D^{20} +15.0$ (*c* 0.2, CHCl₃); IR (neat) 3375, 2924, 1694, 1524, 1451, 1164, 1125, 1003, 740, 696 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 2.01–2.10 (m, 1H, H _{β} Glu-Ha), 2.19–2.29 (m, 1H, H _{β} Glu-Hb), 2.36–2.43 (m, 2H, H _{γ} Glu), 3.71 (dd, *J* = 8.2, 7.3 Hz, 1H, H _{δ} Ψ Pro-Ha), 3.86 (dd, *J* = 8.0, 6.2 Hz, 1H, H _{δ} Ψ Pro-Hb), 4.02–4.07 (m, 2H, H _{α} Gly), 4.18 (dd, *J* = 7.3, 6.9 Hz, 1H, CHFmoc), 4.29–4.40 (m, 4H, CH₂Fmoc and H _{γ} Ψ Pro), 4.63–4.70 (m, 1H, H _{α} Glu), 5.03 (d, *J* = 14.6 Hz, 1H, HBn), 5.06 (d, *J* = 14.6 Hz, 1H, HBn), 5.12 (d, *J* = 12.4 Hz, 1H, HBn), 5.14 (s, 1H, NH), 5.19 (d, *J* = 12.4 Hz, 1H, HBn), 5.65 (s, 1H, NH), 7.28–7.33 (m, 12 H, H_{Ar}Fmoc and H_{Ar}), 7.40 (t, *J* = 7.3 Hz, 2 H, H_{Ar}Fmoc), 7.51 (d, *J* = 7.3 Hz, 1H, NH), 7.59 (d, *J* = 7.8 Hz, 2 H, H_{Ar}Fmoc), 7.75 (d, *J* = 7.8 Hz, 2 H, H_{Ar}Fmoc); ¹³C NMR (100.5 MHz, CDCl₃) δ 26.6 (C _{β} Glu), 30.0 (C _{γ} Glu), 44.3 (C _{α} Gly), 45.5 (C _{δ} Ψ Pro), 47.1 (CHFmoc), 52.4 (C _{α} Glu), 66.8, 67.4, 67.7, 67.9 (CBn CH₂Fmoc and C _{γ} Ψ Pro), 90.9 (q, *J* = 30.7 Hz, C _{α} Ψ Pro), 120.1 (CFmoc), 122.6 (q, *J* = 289.5 Hz, CF₃), 125.2, 127.2, 127.8, 128.2, 128.4, 128.7, 128.7 (CFmoc), 135.5, 141.2, 143.7 (CFmoc), 156.2, 162.6, 167.0, 170.7, 173.2 (CO); ¹⁹F NMR (376.2 MHz, CDCl₃) δ -77.5 (s, CF₃). HRMS (ESI-TOF) calcd. for C₄₁H₃₉F₃N₃O₉ [M + H]⁺ 774.2638, found 774.2643.

The diastereomer **19**_{min} is obtained as colorless foam: $[\alpha]_D^{20} -15.5$ (*c* 0.5, CHCl₃); IR (neat) 3333, 2916, 1694, 1246, 1207, 1164, 1125, 1103, 1081, 1002, 758, 740, 697 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 2.00–2.17 (m, 1H, H _{β} Glu-Ha), 2.20–2.36 (m, 1H, H _{β} Glu-Hb), 2.39–2.53 (m, 2H, H _{γ} Glu), 3.73 (dd, *J* = 15.1, 7.3 Hz, 1H, H _{δ} Ψ Pro-Ha), 3.94 (dd, *J* = 15.1, 6.9 Hz, 1H, H _{δ} Ψ Pro-Hb), 4.05–4.09 (m, 2H, H _{α} Gly), 4.21 (dd, *J* = 7.8, 6.9 Hz, 1H, CHFmoc), 4.30–4.40 (m, 4H, CH₂Fmoc and H _{γ} Ψ Pro), 4.60–4.70 (m, 1H, H _{α} Glu), 5.08 (AB, 2H, HBn), 5.14 (AB, 2H, HBn), 5.75 (s, 1H, NH), 7.25–7.43 (m, 15 H, H_{Ar}Fmoc, H_{Ar} and NH), 7.60 (d, *J* = 7.3 Hz, 2 H, H_{Ar}Fmoc), 7.76 (d, *J* = 7.3 Hz, 2H, H_{Ar}Fmoc); ¹³C NMR (100.5 MHz, CDCl₃) δ 26.4 (C _{β} Glu), 29.9 (C _{γ} Glu), 44.1 (C _{α} Gly), 45.5 (C _{δ} Ψ Pro), 47.0 (CHFmoc), 52.3 (C _{α} Glu), 66.5, 67.2, 67.4, 67.8 (CBn CH₂Fmoc and C _{γ} Ψ Pro), 90.7 (q, *J* = 32.6 Hz, C _{α} Ψ Pro), 119.9 (CFmoc), 122.4 (q, *J* = 291.7 Hz, CF₃), 125.0, 127.0, 127.6, 128.1, 128.2, 128.4, 128.5 (CFmoc), 134.9, 135.6, 141.1, 143.7 (CFmoc), 156.2, 162.7, 167.2, 170.8, 172.8 (CO); ¹⁹F NMR (376.2 MHz, CDCl₃) δ -77.5 (s, CF₃). HRMS (ESI-TOF) calcd. for C₄₁H₃₉F₃N₃O₉ [M + H]⁺ 774.2638, found 774.2660.

H-Gly-2CF₃- Ψ Pro-L-Glu(OH)-OH (20). The diastereomer **19**_{min} (86 mg, 0.11 mmol, 1 equiv) in MeOH (2 mL) was hydrogenated over 10% Pd/C (86 mg) at room temperature for 24 h under 1 bar pressure of hydrogen. The reaction mixture was filtered, concentrated under reduced pressure and purified by reverse phase semipreparative HPLC to give pure deprotected tripeptide **23** as a mixture of two conformers (59/41 determined by ¹⁹F NMR) (20 mg, 49%). White solid: ¹H NMR (400 MHz, D₂O, 353 K) δ 1.99 (ddd, *J* = 14.2, 7.1, 6.9 Hz, 1H, H _{β} Glu-Ha), 2.22 (ddd, *J* = 14.2, 8.2, 6.9 Hz, 1H, H _{β} Glu-Hb), 2.42 (t, *J* = 6.9 Hz, 2H, H _{γ} Glu), 3.91 (dt, *J* = 8.2, 7.1 Hz, 1H, H _{α} Gly-Ha), 4.02–4.10 (m, 3 H, H _{α} Gly-Hb, H _{δ} Ψ Pro), 4.42–4.52 (m, 3 H, H _{α} Glu, H _{γ} Ψ Pro); ¹³C NMR (100.5 MHz, D₂O, 353 K) δ 27.3 (CH₂, C _{β} Glu minor), 27.8 (CH₂, C _{β} Glu major), 31.1 (CH₂, C _{γ} Glu minor), 31.5 (CH₂, C _{γ} Glu major), 42.4 (CH₂, C _{α} Gly major), 45.1 (CH₂, C _{α} Gly minor), 46.6 (CH₂, C _{δ} Ψ Pro major), 46.9 (CH₂, C _{δ} Ψ Pro minor), 54.4 (CH, C _{α} Glu), 67.8 (CH₂, C _{γ} Ψ Pro minor), 69.5 (CH₂, C _{γ} Ψ Pro major), 91.8 (C _{α} Ψ Pro), 127.3 (q, *J* = 302.9 Hz, CF₃), 162.5 (CO Gly minor), 164.9 (CO Gly major), 165.9 (CO Ψ Pro), 175.9 (C _{δ} Glu), 177.0 (CO Glu); ¹⁹F NMR (376.2 MHz, D₂O, 353 K) δ -79.9 (s, CF₃

major), -78.8 (s, CF_3 , minor); HRMS (ESI-TOF) calcd. for $\text{C}_{12}\text{H}_{17}\text{F}_3\text{N}_3\text{O}_7$ $[\text{M} + \text{H}]^+$ 372.1019, found 372.1023.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.6b00704.

NMR spectra for all new compounds. (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: nathalie.lensen@u-cergy.fr.

*E-mail: gregory.chaume@u-cergy.fr.

*E-mail: thierry.brigaud@u-cergy.fr.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the Agence Nationale de la Recherche for financial support (No. ANR-09-JCJC-0060). We also thank the French Fluorine Network. This work has benefited from the facilities and expertise of the Small Molecule Mass Spectrometry platform of IMAGIF (Centre de Recherche de Gif, www.imagif.cnrs.fr).

■ REFERENCES

- (1) Albericio, F.; Kruger, H. G. *Future Med. Chem.* **2012**, *4*, 1527–1531.
- (2) *Bioorganic and Medicinal Chemistry of Fluorine*; Begue, J.-P., Bonnet-Delpon, D., Eds.; Wiley: Hoboken, NJ, 2008.
- (3) Pursler, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320–330.
- (4) *Fluorine in Medicinal Chemistry and Chemical Biology*; Ojima, I., Ed.; Wiley-Blackwell: Hoboken, NJ, 2009.
- (5) *Fluorine in Pharmaceutical and Medicinal Chemistry: From Biophysical Aspects to Clinical Applications*; Gouverneur, V., Müller, K., Eds.; Imperial College Press: London, 2012.
- (6) Smits, R.; Cadicamo, C. D.; Burger, K.; Kokscho, B. *Chem. Soc. Rev.* **2008**, *37*, 1727–1739.
- (7) Qiu, X.-L.; Qing, F.-L. *Eur. J. Org. Chem.* **2011**, *2011*, 3261–3278.
- (8) Aceña, J. L.; Sorochinsky, A. E.; Soloshonok, V. A. *Synthesis* **2012**, *44*, 1591–1602.
- (9) Kokscho, B.; Sewald, N.; Hofmann, H.-J.; Burger, K.; Jakubke, H.-D. *J. Pept. Sci.* **1997**, *3*, 157–167.
- (10) Asante, V.; Mortier, J.; Wolber, G.; Kokscho, B. *Amino Acids* **2014**, *46*, 2733–2744.
- (11) Yoder, N. C.; Kumar, K. *Chem. Soc. Rev.* **2002**, *31*, 335–341.
- (12) Jaeckel, C.; Kokscho, B. *Eur. J. Org. Chem.* **2005**, *2005*, 4483–4503.
- (13) Meng, H.; Kumar, K. *J. Am. Chem. Soc.* **2007**, *129*, 15615–15622.
- (14) Salwiczek, M.; Nyakatura, E. K.; Gerling, U. I. M.; Ye, S.; Kokscho, B. *Chem. Soc. Rev.* **2012**, *41*, 2135–2171.
- (15) Gerling, U. I. M.; Salwiczek, M.; Cadicamo, C. D.; Erdbrink, H.; Czekelius, C.; Grage, S. L.; Wadhvani, P.; Ulrich, A. S.; Behrends, M.; Haufe, G.; Kokscho, B. *Chem. Sci.* **2014**, *5*, 819–830.
- (16) Kubyshkin, V. S.; Komarov, I. V.; Afonin, S.; Mykhailiuk, P. K.; Grage, S. L.; Ulrich, A. S. Trifluoromethyl-substituted α -amino acids as solid-state ^{19}F NMR labels for structural studies of membrane-bound peptides. In *Fluorine in Pharmaceutical and Medicinal Chemistry: From Biophysical Aspects to Clinical Applications*; Gouverneur, V., Müller, K., Eds.; Imperial College Press: London, 2012; pp 91–138.
- (17) Kubyshkin, V.; Afonin, S.; Kara, S.; Budisa, N.; Mykhailiuk, P. K.; Ulrich, A. S. *Org. Biomol. Chem.* **2015**, *13*, 3171–3181 and references cited therein.

- (18) Schlosser, M. *Angew. Chem., Int. Ed.* **1998**, *37*, 1496–1513.
- (19) Huguenot, F.; Brigaud, T. *J. Org. Chem.* **2006**, *71*, 7075–7078.
- (20) Chaume, G.; Van Severen, M.-C.; Marinkovic, S.; Brigaud, T. *Org. Lett.* **2006**, *8*, 6123–6126.
- (21) Chaume, G.; Van Severen, M.-C.; Ricard, L.; Brigaud, T. *J. Fluorine Chem.* **2008**, *129*, 1104–1109.
- (22) Caupène, C.; Chaume, G.; Ricard, L.; Brigaud, T. *Org. Lett.* **2009**, *11*, 209–212.
- (23) Chaume, G.; Barbeau, O.; Lesot, P.; Brigaud, T. *J. Org. Chem.* **2010**, *75*, 4135–4145.
- (24) Simon, J.; Nguyen, T. T.; Chelain, E.; Lensen, N.; Pytkowicz, J.; Chaume, G.; Brigaud, T. *Tetrahedron: Asymmetry* **2011**, *22*, 309–314.
- (25) Lensen, N.; Marais, J.; Brigaud, T. *Org. Lett.* **2015**, *17*, 342–345.
- (26) Lubin, H.; Pytkowicz, J.; Chaume, G.; Sizon-Thomé, G.; Brigaud, T. *J. Org. Chem.* **2015**, *80*, 2700–2708.
- (27) Chaume, G.; Lensen, N.; Caupène, C.; Brigaud, T. *Eur. J. Org. Chem.* **2009**, *2009*, 5717–5724.
- (28) Chaume, G.; Feytens, D.; Chassaing, G.; Lavielle, S.; Brigaud, T.; Miclet, E. *New J. Chem.* **2013**, *37*, 1336–1342.
- (29) Jlalila, I.; Lensen, N.; Chaume, G.; Dzhabazova, E.; Astasidi, L.; Hadjiolova, R.; Bocheva, A.; Brigaud, T. *Eur. J. Med. Chem.* **2013**, *62*, 122–129.
- (30) Chaume, G.; Simon, J.; Caupène, C.; Lensen, N.; Miclet, E.; Brigaud, T. *J. Org. Chem.* **2013**, *78*, 10144–10153.
- (31) Sara, V. R.; Carlsson-Skwirut, C.; Bergman, T.; Jörnvall, H.; Roberts, P. J.; Crawford, M.; Nilsson-Hakansson, L.; Civalero, I.; Nordberg, A. *Biochem. Biophys. Res. Commun.* **1989**, *165*, 766–771.
- (32) Bourguignon, J. P.; Gerard, A.; Alvarez Gonzalez, M. L.; Franchimont, P. *Neuroendocrinology* **1993**, *58*, 525–530.
- (33) Sara, V. R.; Carlsson-Skwirut, C.; Drakenberg, K.; Giacobini, M. B.; Hakanson, L.; Mirmiran, M.; Nordberg, A.; Olson, L.; Reinecke, M.; Stahlbom, P. A.; Sandberg Nordqvist, A. C. *Ann. N. Y. Acad. Sci.* **1993**, *692*, 183–191.
- (34) Guan, J. *Frontiers in CNS Drug Discovery* **2010**, *1*, 51–75.
- (35) Guan, J.; Mathai, S.; Liang, H.-P.; Gunn, A. J. *Recent Pat. CNS Drug Discovery* **2013**, *8*, 142–160.
- (36) Guan, J.; Harris, P.; Brimble, M.; Lei, Y.; Lu, J.; Yang, Y.; Gunn, A. J. *Expert Opin. Ther. Targets* **2015**, *19*, 785–793.
- (37) Cacciatore, L.; Cornacchia, C.; Baldassarre, L.; Fornasari, E.; Mollica, A.; Stefanucci, A.; Pinnen, F. *Mini-Rev. Med. Chem.* **2012**, *12*, 13–23 and references cited therein.
- (38) Trotter, N. S.; Brimble, M. A.; Callis, D. J.; Sieg, F. *Bioorg. Med. Chem.* **2005**, *13*, 501–517.
- (39) Brimble, M. A.; Trotter, N. S.; Harris, P. W. R.; Sieg, F. *Bioorg. Med. Chem.* **2005**, *13*, 519–532.
- (40) Lai, M. Y. H.; Brimble, M. A.; Callis, D. J.; Harris, P. W. R.; Levi, M. S.; Sieg, F. *Bioorg. Med. Chem.* **2005**, *13*, 533–548.
- (41) Harris, P. W. R.; Brimble, M. A.; Muir, V. J.; Lai, M. Y. H.; Trotter, N. S.; Callis, D. J. *Tetrahedron* **2005**, *61*, 10018–10035.
- (42) Harris, P. W. R.; Brimble, M. A. *Org. Biomol. Chem.* **2006**, *4*, 2696–2709.
- (43) De Diego, S. A. A.; Munoz, P.; Gonzalez-Muniz, R.; Herranz, R.; Martin-Martinez, M.; Cenarruzabeitia, E.; Frechilla, D.; Del Rio, J.; Jimeno, M. L.; Garcia-Lopez, T. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 2279–2283.
- (44) De Diego, S. A. A.; Gutierrez-Rodriguez, M.; Perez de Vega, M. J.; Casabona, D.; Catiuela, C.; Gonzalez-Muniz, R.; Herranz, R.; Cenarruzabeitia, E.; Frechilla, D.; Del Rio, J.; Jimeno, M. L.; Garcia-Lopez, T. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 1392–1396.
- (45) De Diego, S. A. A.; Gutierrez-Rodriguez, M.; Perez de Vega, M. J.; Gonzalez-Muniz, R.; Herranz, R.; Martin-Martinez, M.; Cenarruzabeitia, E.; Frechilla, D.; Del Rio, J.; Jimeno, M. L.; Garcia-Lopez, T. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 3396–3400.
- (46) Yaron, A.; Naider, F. *Crit. Rev. Biochem. Mol. Biol.* **1993**, *28*, 31.
- (47) Brimble, M. A.; Harris, P. W. R. *Chem. N. Z.* **2011**, *75*, 133–136.
- (48) DalPozzo, A.; Bergonzi, R.; Ni, M. *Tetrahedron Lett.* **2001**, *42*, 3925–3927.

- (49) DalPozzo, A.; Ni, M.; Muzi, M.; Caporale, A.; de Castiglione, R.; Kaptein, B.; Broxterman, Q. B.; Formaggio, F. *J. Org. Chem.* **2002**, *67*, 6372–6375.
- (50) DalPozzo, A.; Ni, M.; Muzi, M.; de Castiglione, R.; Mondelli, R.; Mazzini, S.; Penco, S.; Pisano, C.; Castorina, M.; Giannini, G. *J. Med. Chem.* **2006**, *49*, 1808–1817.
- (51) Carpino, L. A.; Sadat-Aalae, D.; Chao, H. G.; DeSelms, R. H. *J. Am. Chem. Soc.* **1990**, *112*, 9651–9652.
- (52) Wenschuh, H.; Beyermann, M.; Winter, R.; Bienert, M.; Ionescu, D.; Carpino, L. A. *Tetrahedron Lett.* **1996**, *37*, 5483–5486.
- (53) Olah, G. A.; Nojima, M.; Kerekes, I. *Synthesis* **1973**, *1973*, 487.
- (54) Prabhu, G.; Basavaprabhu, Narendra, N.; Vishwanatha, T. M.; Sureshbabu, V. V. *Tetrahedron* **2015**, *71*, 2785–2832.
- (55) Carpino, L. A.; Cohen, B. J.; Stephens, K. E.; Sadat-Aalae, Y.; Tien, J. H.; Langridge, D. C. *J. Org. Chem.* **1986**, *51*, 3732–3734.
- (56) Kantharaju; Patil, B. S.; Sureshbabu, V. V. *Lett. Pept. Sci.* **2002**, *9*, 227–229.
- (57) Ando, W.; Igarashi, Y.; Huang, L. *Chem. Lett.* **1987**, *16*, 1361–1364.
- (58) Jeong, Y.-C.; Anwar, M.; Nguyen, T. M.; Tan, B. S. W.; Chai, C. L. L.; Moloney, M. G. *Org. Biomol. Chem.* **2011**, *9*, 6663–6669.
- (59) Fülöp, F.; Mattinen, J.; Pihlaja, K. *Tetrahedron* **1990**, *46*, 6545–6552.
- (60) Fülöp, F.; Pihlaja, K. *Tetrahedron* **1993**, *49*, 6701–6706.
- (61) Pascal, R.; Sola, R. *Tetrahedron Lett.* **1998**, *39*, 5031–5034.
- (62) Pytkowicz, J.; Stéphany, O.; Marinkovic, S.; Inagaki, S.; Brigaud, T. *Org. Biomol. Chem.* **2010**, *8*, 4540–4542.