

Synthesis of 6,12-Epiminodibenzo[*b,f*][1,5]diazocines via an Ytterbium Triflate-Catalyzed, AB₂ Three-Component Reaction

Isravel Muthukrishnan,[†] Muthu Karuppasamy,[†] Subbiah Nagarajan,[†] C. Uma Maheswari,[†] Vittorio Pace,[‡] J. Carlos Menéndez,[§] and Vellaisamy Sridharan^{*,†}


[†]Organic Synthesis Group, Department of Chemistry, School of Chemical and Biotechnology, SASTRA University, Thanjavur, Tamil Nadu 613401, India

[‡]Department of Pharmaceutical Chemistry, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria

[§]Departamento de Química Orgánica y Farmacéutica, Facultad de Farmacia, Universidad Complutense, 28040 Madrid, Spain

Supporting Information

ABSTRACT: An efficient and selective procedure for the synthesis of epiminodibenzo[*b,f*][1,5]diazocines involving a AB₂ three-component reaction is developed. Two equivalents of suitably substituted 2-aminoarylaldehydes reacted with arylamines in the presence of Yb(OTf)₃ to afford the desired products in high yields. The reaction is highly atom-economic and waste-free, in addition to allowing the generation of two heterocyclic rings and four C–N bonds in a single operation. Significantly, this approach is complementary to the existing literature procedures, affording arylamine-derived products that could not be accessed previously. A plausible mechanism is proposed involving an imine formation-intermolecular annulation-intramolecular iminium ion cyclization sequence.



INTRODUCTION

Tröger's base **1** and its analogs are recognized as a very important class of nitrogen heterocycles with potential applications in various areas of supramolecular chemistry.¹ The presence of two stereogenic nitrogen atoms and its unique V-shaped rigid structure with a hydrophobic cavity between the aryl rings explains its role in asymmetric catalysis, molecular recognition, self-assembly, sensors, optoelectronics, and enzyme inhibition.^{2,3} Consequently, considerable effort has been devoted to the synthesis, functionalization, and chiral resolution of Tröger's base-related compounds.⁴ Likewise, one of the most closely related analogs of Tröger's base, epiminodibenzo[*b,f*][1,5]diazocine **2**, was also found to possess similar applications owing to its structural resemblance (Figure 1).^{5–7} Consequently, development of efficient synthetic procedures for this class of compounds is an essential task. Very few methodologies to access epiminodibenzo[*b,f*][1,5]diazocines are known in the literature that include a porous grapheme oxide-catalyzed tandem reaction of 2-aminobenzylamine,⁵ a TMSOTf/Et₃N-catalyzed domino process starting




Figure 1. Structures of Tröger's base and 6,12-epiminodibenzo[*b,f*][1,5]diazocine.

from *N*-alkyl-aldimines,⁶ and a metal-free cyclization of 2-aminophenyl ketimines.⁷


Although epiminodibenzo[*b,f*][1,5]diazocines may seem, in principle, easy to access from 2 equiv of 2-aminoarylaldehydes and a primary amine, this approach presents a major limitation associated with the low stability of these aldehydes, where the presence of two highly reactive adjacent functionalities often affords trimerization products rather than the desired compounds, especially in the presence of nucleophiles.⁸ For this reason, functionalized 2-aminoarylaldehydes including iminophosphoranes⁹ and *N*-Ts/Ms derivatives¹⁰ have been employed as precursors for the synthesis of epiminodibenzo[*b,f*][1,5]diazocines combined with primary amines, albeit in low yields. Recently, Yu, Wang, and co-workers reported a simple cascade reaction between 2-aminoarylaldehydes **3** and primary amines in the presence of scandium pentafluorobenzoate.¹¹ This reaction afforded the expected epiminodibenzo[*b,f*][1,5]diazocine derivatives **4** in good yields, but worked exclusively for benzylamines and other alkylamines. On the other hand, 2-aminoarylaldehydes reacted with arylamines furnishing the ring-fused quinazoline derivatives **5** (Scheme 1, eq 1). In this context, we envisioned to develop a simple procedure for the synthesis of epiminodibenzo[*b,f*][1,5]diazocines **7** starting from 2-aminoarylaldehydes **6** and arylamines with an aim to provide a complementary approach to the existing Lu and Wang's protocol (Scheme 1, eq 2).

Received: July 23, 2016

Published: September 28, 2016


Scheme 1. Reactivity of 2-Aminoarylaldehydes with Primary Amines

Yu, Wang and co-workers (2013)




RESULTS AND DISCUSSION

Recently, we have reported the synthesis of 5,6-dihydrodibenzo[*b,h*][1,6]naphthyridines via copper(II) bromide-catalyzed intramolecular [4 + 2] hetero-Diels–Alder reactions from 2-(*N*-propargylamino-*N*-tosyl)benzaldehydes and arylamines.¹² During this investigation, we were drawn to study whether the reaction between 2-(*N*-propargylamino)benzaldehyde **6a** and *p*-toluidine would afford the completely aromatized dibenzo[*b,h*][1,6]naphthyridine **8**. However, under our experimental conditions, in the presence of 10 mol % of CuBr₂ in toluene at 25 °C, we observed only the imine intermediate **A**. On the other hand, at 80 °C, 15% of dibenzo[*b,h*][1,6]naphthyridine **8** was isolated together with unreacted aldehyde and imine **A** (Table 1, entries 1 and 2). Surprisingly, switching the reaction solvent to acetonitrile, under similar experimental conditions, the reaction furnished 58% of epiminodibenzo[*b,f*][1,5]diazocines **7a** via an AB₂ three-component reaction, together with the imine intermediate **A** (Table 1, entry 3). Among the other tested copper catalysts, CuCl₂ furnished comparable yield, nonetheless, in the presence of CuCl, and complete decomposition of starting material **6a** was observed (entries 4 and 5). In an effort to improve the reaction yield, subsequently, we moved to screen some Lewis acids. As shown in entries 6–9, with the exception of InCl₃, other tested catalysts cerium(IV) ammonium nitrate (CAN), Sc(OTf)₃, and Yb(OTf)₃ furnished the product **7a** in good yields. The reaction was completed in 8 h in the presence

Table 1. Optimization of the AB₂ Three-Component Synthesis of Epiminodibenzo[*b,f*][1,5]diazocines^a

entry	compd	catalyst	mol %	solvent	temperature (°C)	reaction time (h)	yield of 7a/7b ^b (%)
1	7a	CuBr ₂	10	toluene	25	24	— ^c
2	7a	CuBr ₂	10	toluene	80	24	— ^d
3	7a	CuBr ₂	10	MeCN	80 ^c	12	58 ^e
4	7a	CuCl ₂	10	MeCN	80 ^c	16	55 ^e
5	7a	CuCl	10	MeCN	80 ^c	24	— ^f
6	7a	InCl ₃	10	MeCN	80 ^c	24	21 ^e
7	7a	CAN	10	MeCN	80 ^c	8	63
8	7a	Sc(OTf) ₃	10	MeCN	80 ^c	16	69 ^e
9	7a	Yb(OTf) ₃	10	MeCN	80 ^c	8	75
10 ^g	7a	Yb(OTf) ₃	5	MeCN	80	8	74
11	7a	Yb(OTf) ₃	5	DCM	45	12	65
12	7a	Yb(OTf) ₃	5	DCE	80	24	57
13	7a	Yb(OTf) ₃	5	THF	70	12	55
14	7a	Yb(OTf) ₃	5	toluene	80	24	51 ^e
15	7a	Yb(OTf) ₃	5	EtOH	80	24	25 ^e
16	7a	Yb(OTf) ₃	5	water	80	24	— ^h
17	7a	Yb(OTf) ₃	5	PEG-200	80	24	40 ^e
18	7a	Yb(OTf) ₃	5	glycerol	80	24	28 ^e
19 ^g	7b	Yb(OTf) ₃	5	MeCN	80	8	95

^aAll reactions were carried out with **6a/6b** (1.0 mmol) and *p*-toluidine (0.5 mmol) with catalyst in 2 mL of solvent. ^bIsolated yield. ^cOnly imine **A** was observed at 25 °C. ^d15% of compound **8** was isolated together with unreacted **6a** and intermediate **A**. ^eUnreacted imine **A** was noticed in the crude ¹H NMR spectra. ^fDecomposition of compound **6a** was observed. ^gOptimized reaction conditions. ^hOnly imine **A** was formed.


Scheme 2. Synthesis of Epiminodibenzo[*b,f*][1,5]diazocines **7**^a

^aReaction conditions: aldehydes **6** (1.0 mmol), arylamines (0.5 mmol), Yb(OTf)₃ (5 mol %), MeCN (2 mL), 80 °C, 6–12 h.

of 10 mol % of Yb(OTf)₃ at 80 °C, and a maximum yield of 75% was observed (entry 9). Decreasing the catalyst loading to 5 mol % did not affect the rate of the reaction significantly and delivered 74% of the product under similar reaction time (entry 10). It should be highlighted here that all the tested catalysts, including the copper salts, afforded only the imine intermediate **A** at 25 °C, and in no case, the [4 + 2] cycloaddition product **8**

was isolated. Further screening of solvents including DCM, DCE, THF, toluene, and ethanol revealed that DCM gave comparable yield, and in the cases of toluene and ethanol, the reaction was not completed even after 24 h (entries 11–15). Inferior outcomes were observed when the reaction was carried out in green solvents such as water, PEG-200, and glycerol (entries 16–18). Remarkably, the bromo derivative **7b** was

Scheme 3. Proposed Mechanism



obtained in 95% yield under optimized reaction conditions (entry 19).


With the optimized reaction conditions in hand, we next turned our attention to investigate the scope and limitations of the AB₂ three-component synthesis of epiminodibenzo[*b,f*]-[1,5]diazocines **7**. Unlike Yu and Wang's methodology,¹¹ our conditions worked very well for all the tested arylamines to afford the target compounds in good to excellent yields (Scheme 2). Initially, we examined the effect of the substituents on the aldehyde component **6**, and it revealed that the presence of electron-withdrawing substituents including bromine and chlorine afforded the products in excellent yields compared to the unsubstituted aldehyde, presumably due to the stability and controlled reactivity of aldehydes **6** (e.g., **7b**, **7g–7n**). These electron-withdrawing groups decrease the nucleophilicity of the nitrogen atom, thus confirming the smooth reactivity of compounds **6**. The nitrogen atom also bears propargyl and benzyl substituents. We then investigated the nature of the substituents on the arylamines. All moderately electron-releasing groups including *p*-methyl (**7a** and **7b**), *m*-methyl (**7j**, **7k**, and **7o**) and 3,5-dimethyl (**7i** and **7p**) and strongly electron-releasing *p*- and *m*-methoxy (**7d**, **7l** and **7q**) arylamines furnished the products in high yields. Similarly, electron-deficient arylamines including fluoro- (**7f**), chloro- (**7h** and **7r**), bromo- (**7e** and **7m**) substituted analogs also reacted smoothly to yield the corresponding epiminodibenzo[*b,f*][1,5]diazocines **7** in comparable yields. Notably, *m*-nitroaniline, bearing a strong electron-withdrawing group, also yielded the product **7n** in 87% yield albeit in a longer reaction time. In order to show the generality of the protocol, we then investigated the reactivity of benzylamine and butylamine under the optimized reaction conditions. Interestingly, benzylamine tolerated the reaction conditions to afford the product **7s** in 81% yield. However, the reaction was sluggish and needed longer reaction time (20 h) than arylamines. On the other hand, the reaction failed to yield the desired product in the case of butylamine presumably due to the poor stability of the intermediate imine. Thus, this methodology allows general and direct access to arylamine-derived epiminodibenzo[*b,f*][1,5]diazocines **7**, which could not be obtained by the existing procedure. Furthermore, this AB₂ three-component approach generates two heterocyclic rings and four new C–N bonds in a single operation. In addition, the reaction is highly atom economic, and no waste is produced except two molecules of water. The

reaction between 2-aminobenzaldehyde and *p*-toluidine under our optimized conditions afforded a mixture of epiminodibenzo[*b,f*][1,5]diazocine **7u**^{9a} and quinazoline derivative **5a**.¹¹ Thus, the presence of *N*-alkyl group on the 2-aminoarylaldehyde is essential to yield the desired products in a controlled manner.

As shown in Scheme 3, eq 1, this reaction is proposed to proceed via imine intermediate **A** generated from aldehyde **6** and arylamines, as the imine formation was observed immediately after the addition arylamines to the aldehydes and confirmed by crude ¹H NMR analysis. Subsequent intermolecular self-cyclization of two molecules of imine **A** would generate the aminal **B**, which could eliminate a molecule of arylamine followed by intramolecular 6-endo-trig cyclization to afford compound **7** through the intermediacy of iminium cation **C**. This mechanism is supported by the observation that, in some occasions, the initially formed imine **A** was converted into the product **7** in the absence of aldehyde **6**. Alternatively, as shown in Scheme 3, eq 2, Lewis acid-catalyzed reaction between imine **A** and a second equivalent of aldehyde **6** would afford 1,2,3,4-tetrahydroquinazolin-4-ol intermediate **D**, which subsequently generates the iminium cation **C** triggered by the Lewis acid. A final intramolecular cyclization of intermediate **C** furnishes the product **7**.


We next sought to showcase the usefulness of the terminal alkyne group in many of our compounds to introduce an additional diversity point. 1,2,3-Triazoles are important class of nitrogen heterocycles known to exhibit broad-spectrum pharmacological activities exemplified by anti-influenza A agents,¹³ σ_2 receptor ligands,¹⁴ Nrf2–Keap1 protein–protein interaction inhibitors,¹⁵ and many others. The Cu(I)-catalyzed 1,3-dipolar reaction between alkyne and azide, popularly known as the “click reaction” is a well-explored protocol to access 1,2,3-triazoles.¹⁶ Consequently, we envisioned to utilize the terminal alkynes of the synthesized epiminodibenzo[*b,f*][1,5]diazocines **7** to introduce the triazole moiety by the reaction with benzyl azide. As expected, bis-triazole tethered epiminodibenzo[*b,f*][1,5]diazocines **9** were obtained in excellent yields under mild experimental conditions in the presence of CuI (Scheme 4). Owing to the biological interest of both 1,2,3-triazole and epiminodibenzo[*b,f*][1,5]diazocines, this approach could be used to generate a library of compounds bearing both heterocycles.

Scheme 4. Synthesis of Bis-triazole Tethered Epiminodibenzo[*b,f*][1,5]diazocines **8**



Finally, we derivatized the terminal alkynes under Sonogashira reaction conditions as shown in Scheme 5. Under standard conditions, phenyl group was successfully introduced to epiminodibenzo[*b,f*][1,5]diazocines **7a** and **7b** to obtain compounds **7v** and **7w** in high yields.

Scheme 5. Sonogashira Reaction of Epiminodibenzo[*b,f*][1,5]diazocines **7a** and **7b**



In conclusion, we have demonstrated the synthesis of Tröger's base analogs epiminodibenzo[*b,f*][1,5]diazocines from readily available starting materials involving a Yb(OTf)₃-catalyzed AB₂ three-component reaction in high yields. The reaction generates two heterocyclic rings and four C–N bonds in a single operation in addition to its high atom economy. This methodology is complementary to the existing procedure since this reaction is completely selective for arylamines, whereas the literature method works only for benzylamines and alkyl amines. We have also proposed a plausible mechanism to explain the product formation involving imine formation-intermolecular annulation-intramolecular iminium ion cyclization steps.

EXPERIMENTAL SECTION

General Information. Commercially available reagents were used without further purification, and the reaction solvents (THF, MeCN, DCM etc.) were purified using standard procedures. The reactions were monitored by thin-layer chromatography visualized by UV detection or using *p*-anisaldehyde or 2,4-DNP stains or molecular iodine. Flash column chromatography was performed with silica gel (230–400 mesh). Melting points were recorded on a melting point apparatus in capillaries and are uncorrected. ¹H and ¹³C NMR spectra were recorded in CDCl₃ at room temperature in a spectrometer operating at 300 MHz for ¹H and 75 MHz for ¹³C. Chemical shifts (δ) are expressed in ppm using TMS as internal standard, and coupling constants (*J*) are given in hertz. Infrared (IR) spectra were obtained in a FTIR spectrometer with a diamond ATR accessory for solid and liquid samples, requiring no sample preparation, and the major frequencies were reported in cm⁻¹. Elemental analyses were determined by using a CHNS combustion microanalyzer.

General Procedure for the Synthesis of 2-(Prop-2-yn-1-ylamino)arylaldehyde **6a–d.** To a stirred solution of suitably

substituted 2-aminobenzyl alcohol (30 mmol) in acetonitrile (80 mL) were added K₂CO₃ (45 mmol) and propargyl bromide (40 mmol) and stirring was continued for 6–8 h at 75 °C. The reaction mixture was diluted with water, extracted with ethyl acetate (3 × 30 mL), and washed with brine solution. The organic layer was dried over anhydrous Na₂SO₄ and then concentrated under reduced pressure. The residue was purified by silica gel column chromatography using petroleum ether-ethyl acetate mixture as eluent (70:30 v/v). Subsequently, a solution of 2-(prop-2-ynylamino)benzylamine obtained in the previous step in CH₂Cl₂ (150 mL) was added MnO₂ (10 equiv), and the mixture was stirred at room temperature for 12–16 h. The reaction mixture was filtered and washed with CH₂Cl₂ (3 × 50 mL). The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by silica gel column chromatography using petroleum ether-ethyl acetate mixture as eluent (96:4 v/v) to yield the products **6a–c**. Compound **6d** was prepared by using the literature procedure.¹⁷

2-(Prop-2-ynylamino)benzaldehyde (6a**).** Yellow viscous liquid; yield: 2.29 g, 48%; IR (neat) 3331.9, 3287.5, 3047.3, 2830.8, 2111.5, 1655.3, 1575.3, 1515.7, 1335.5, 1199.3 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 2.44 (t, *J* = 2.4 Hz, 1H), 4.06 (dd, *J* = 6.0, 2.4 Hz, 2H), 6.77–6.83 (m, 2H), 7.47 (td, *J* = 7.8, 1.5 Hz, 1H), 7.52 (dd, *J* = 7.8, 1.5 Hz, 1H), 8.44 (br s, 1H), 9.85 (s, 1H). ¹³C NMR (75 MHz, CDCl₃): δ 32.1, 71.6, 79.8, 111.2, 116.1, 119.1, 135.8, 136.6, 149.4, 194.2. Anal. calcd for C₁₀H₉NO: C, 75.45; H, 5.70; N, 8.80. Found: C, 75.09; H, 5.80; N, 8.87.

5-Bromo-2-(prop-2-ynylamino)benzaldehyde (6b**).** Yellow solid; mp: 112–115 °C; yield: 3.64 g, 51%; IR (neat) 3314.2, 3244.8, 3050.0, 2849.6, 2109.3, 1654.6, 1562.5, 1504.4, 1303.5, 1174.7 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 2.19 (s, 1H), 3.96 (d, *J* = 5.7 Hz, 2H), 6.66 (d, *J* = 9.0 Hz, 1H), 7.45 (d, *J* = 9.0 Hz, 1H), 7.54 (s, 1H), 8.36 (br s, 1H), 9.70 (s, 1H). ¹³C NMR (75 MHz, CDCl₃): δ 32.2, 71.9, 79.2, 107.3, 113.4, 120.4, 138.3, 138.4, 148.3, 192.9. Anal. calcd for C₁₀H₈BrNO: C, 50.45; H, 3.39; N, 5.88. Found: C, 50.14; H, 3.31; N, 6.06.

5-Chloro-2-(prop-2-yn-1-ylamino)benzaldehyde (6c**).** Yellow solid; mp: 105–108 °C; yield: 3.25 g, 56%; IR (neat) 3318.2, 3267.3, 3027.0, 2820.0, 1577.8, 1555.7, 1340.3, 1161.1 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 2.26 (t, *J* = 2.4 Hz, 1H), 4.04 (dd, *J* = 6.0, 2.4 Hz, 2H), 6.78 (d, *J* = 9.0 Hz, 1H), 7.40 (dd, *J* = 9.0, 2.7 Hz, 1H), 7.49 (d, *J* = 2.7 Hz, 1H), 8.42 (br s, 1H), 9.78 (s, 1H). ¹³C NMR (75 MHz, CDCl₃): δ 32.2, 71.9, 79.3, 113.0, 119.7, 120.7, 135.2, 135.7, 147.9, 193.0. Anal. calcd for C₁₀H₈ClNO: C, 62.03; H, 4.16; N, 7.23. Found: C, 61.94; H, 4.21; N, 7.29.

General Procedure for the AB₂ Three-Component Synthesis of Epiminodiazocines **7.** To a stirred solution of *N*-substituted-2-aminoarylaldehyde **6** (1.0 mmol) and primary amines (0.5 mmol) in acetonitrile (2 mL) was added Yb(OTf)₃ (5 mol %), and stirring was continued until completion of the reaction (6–12 h) at 80 °C. After completion of the reaction, as indicated by TLC, the reaction mixture was diluted with CH₂Cl₂ and washed with water. The organic layer was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified through silica gel column chromatography using petroleum ether-ethyl acetate (97:5, v/v) as eluent.

5,11-Di(prop-2-yn-1-yl)-13-(*p*-tolyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (7a**).** Colorless solid; mp: 139–142 °C; yield: 144 mg, 74%; IR (neat) 3286.3, 3258.1, 2988.5, 2120.2, 1602.2, 1493.1, 1458.4, 1325.1, 1225.6, 1165.5, 1098.4 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 2.27 (s, 3H), 2.46 (t, *J* = 2.4 Hz, 2H), 4.07 (dd, *J* = 17.1, 2.4 Hz, 2H), 4.29 (dd, *J* = 17.1, 2.4 Hz, 2H), 5.97 (s, 2H), 6.77–6.85 (m, 4H), 7.01–7.09 (m, 4H), 7.15 (td, *J* = 7.5, 1.5 Hz, 2H), 7.37 (d, *J* = 7.5 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃): δ 20.6, 40.4, 69.7, 73.6, 79.9, 115.2, 118.3, 119.1, 123.3, 128.2, 128.9, 129.8, 130.6, 142.4, 144.7. Anal. calcd for C₂₇H₂₃N₃: C, 83.26; H, 5.95; N, 10.79. Found: C, 82.98; H, 6.06; N, 10.61.

2,8-Dibromo-5,11-di(prop-2-yn-1-yl)-13-(*p*-tolyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (7b**).** Pale brown solid; mp: 73–75 °C; yield: 262 mg, 96%; IR (neat) 3280.8, 3255.2, 2977.0, 2116.7, 1605.9, 1485.5, 1344.6, 1234.6, 1157.3, 1088.2 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 2.26 (s, 3H), 2.45 (t, *J* = 2.4 Hz, 2H),

4.03 (dd, $J = 17.4, 2.4$ Hz, 2H), 4.26 (dd, $J = 17.4, 2.4$ Hz, 2H), 5.83 (s, 2H), 6.65 (d, $J = 8.7$ Hz, 2H), 6.93 (d, $J = 8.7$ Hz, 2H), 7.06 (d, $J = 8.7$ Hz, 2H), 7.24 (dd, $J = 8.7, 2.4$ Hz, 2H), 7.45 (d, $J = 2.4$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 20.5, 40.5, 69.6, 74.1, 79.0, 111.4, 116.9, 118.5, 124.9, 129.9, 130.7, 131.4, 131.8, 141.1, 143.9. Anal. calcd for $\text{C}_{27}\text{H}_{21}\text{Br}_2\text{N}_3$: C, 59.25; H, 3.87; N, 7.68. Found: C, 58.97; H, 3.80; N, 7.74.

13-(4-Phenyl-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7c). Colorless solid; mp: 133–136 °C; yield: 135 mg, 72%; IR (neat) 3288.1, 3257.5, 3037.5, 2950.1, 2113.2, 1597.7, 1490.7, 1227.4, 1148.5, 1097.2 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.39 (t, $J = 2.1$ Hz, 2H), 3.98 (dd, $J = 17.4, 2.1$ Hz, 2H), 4.21 (dd, $J = 17.4, 2.1$ Hz, 2H), 5.95 (s, 2H), 6.69–6.80 (m, 4H), 6.82 (t, $J = 7.2$ Hz, 1H), 7.04–7.21 (m, 6H), 7.28 (d, $J = 7.5$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.4, 69.4, 73.8, 79.8, 115.3, 118.1, 119.2, 121.1, 123.3, 128.2, 129.0, 129.3, 142.4, 147.0. Anal. calcd for $\text{C}_{26}\text{H}_{21}\text{N}_3$: C, 83.17; H, 5.64; N, 11.19. Found: C, 82.80; H, 5.75; N, 11.02.

13-(4-Methoxyphenyl)-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7d). Colorless solid; mp: 128–132 °C; yield: 156 mg, 77%; IR (neat) 3280.5, 3245.0, 2932.2, 2118.6, 1601.2, 1506.1, 1453.8, 1356.4, 1236.4, 1094.5, 1026.1 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.44 (t, $J = 2.4$ Hz, 2H), 3.76 (s, 3H), 4.06 (dd, $J = 17.1, 2.4$ Hz, 2H), 4.28 (dd, $J = 17.1, 2.4$ Hz, 2H), 5.86 (s, 2H), 6.78–6.85 (m, 6H), 7.05 (d, $J = 9.0$ Hz, 2H), 7.16 (td, $J = 8.7, 1.5$ Hz, 2H), 7.36 (dd, $J = 7.5, 1.2$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.4, 55.5, 70.4, 73.5, 79.9, 114.5, 115.1, 119.7, 120.3, 123.3, 128.2, 128.9, 140.9, 142.4, 154.6. Anal. calcd for $\text{C}_{27}\text{H}_{23}\text{N}_3\text{O}$: C, 79.97; H, 5.72; N, 10.36. Found: C, 79.66; H, 5.78; N, 10.20.

13-(4-Bromophenyl)-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7e). Colorless solid; mp: 96–98 °C; yield: 182 mg, 80%; IR (neat) 3285.9, 3025.8, 2824.9, 2185.9, 1601.2, 1584.0, 1490.1, 1366.4, 1232.9, 1097.4 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.47 (t, $J = 2.7$ Hz, 2H), 4.04 (dd, $J = 17.1, 2.7$ Hz, 2H), 4.28 (dd, $J = 17.1, 2.7$ Hz, 2H), 5.97 (s, 2H), 6.77 (d, $J = 8.1$ Hz, 2H), 6.82 (td, $J = 7.5, 1.2$ Hz, 2H), 7.01 (d, $J = 9.0$ Hz, 2H), 7.15 (td, $J = 8.1, 1.5$ Hz, 2H), 7.33–7.36 (m, 4H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.3, 69.3, 74.0, 79.6, 113.5, 115.3, 119.4, 119.8, 122.9, 128.2, 129.2, 132.1, 142.2, 146.2. Anal. calcd for $\text{C}_{26}\text{H}_{20}\text{BrN}_3$: C, 68.73; H, 4.44; N, 9.25. Found: C, 68.51; H, 4.39; N, 9.02.

13-(4-Fluorophenyl)-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7f). Colorless solid; mp: 143–146 °C; yield: 140 mg, 71%; IR (neat) 3293.4, 3275.3, 3070.4, 2825.2, 2115.0, 1600.8, 1491.0, 1363.4, 1231.6, 1091.1 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.44 (t, $J = 2.4$ Hz, 2H), 4.04 (dd, $J = 17.1, 2.4$ Hz, 2H), 4.27 (dd, $J = 17.1, 2.4$ Hz, 2H), 5.89 (s, 2H), 6.78 (d, $J = 8.1$ Hz, 2H), 6.82 (td, $J = 7.2, 1.2$ Hz, 2H), 6.94 (t, $J = 9.0$ Hz, 2H), 7.02–7.09 (m, 2H), 7.15 (td, $J = 9.0, 1.8$ Hz, 2H), 7.34 (dd, $J = 7.5, 1.5$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.3, 70.1, 73.8, 79.7, 115.2, 115.8 (d, $J = 21.8$ Hz), 119.3, 120.1 (d, $J = 8.3$ Hz), 123.1, 128.2, 129.1, 142.3, 143.5 (d, $J = 2.3$ Hz), 157.9 (d, $J = 239.3$ Hz). Anal. calcd for $\text{C}_{26}\text{H}_{20}\text{FN}_3$: C, 79.37; H, 5.12; N, 10.68. Found: C, 79.01; H, 5.18; N, 10.72.

2,8-Dibromo-13-(4-phenyl)-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7g). Colorless solid; mp: 148–151 °C; yield: 224 mg, 84%; IR (neat) 3276.8, 3037.7, 2827.0, 2118.9, 1593.7, 1488.3, 1355.9, 1233.8, 1113.8 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.40 (t, $J = 2.4$ Hz, 2H), 3.97 (dd, $J = 17.4, 2.4$ Hz, 2H), 4.15 (dd, $J = 17.4, 2.4$ Hz, 2H), 5.83 (s, 2H), 6.60 (d, $J = 8.7$ Hz, 2H), 6.87 (t, $J = 7.5$ Hz, 1H), 6.97 (d, $J = 8.1$ Hz, 2H), 7.16–7.22 (m, 4H), 7.39 (d, $J = 2.4$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.6, 69.4, 74.3, 79.0, 111.6, 117.1, 118.3, 121.9, 125.0, 129.4, 130.8, 131.9, 141.2, 146.4. Anal. calcd for $\text{C}_{26}\text{H}_{19}\text{Br}_2\text{N}_3$: C, 58.56; H, 3.59; N, 7.88. Found: C, 58.29; H, 3.63; N, 7.86.

2,8-Dibromo-13-(4-chlorophenyl)-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7h). Pale yellow solid; mp: 150–153 °C; yield: 259 mg, 91%; IR (neat) 3289.4, 3264.8, 2923.6, 2837.4, 2153.6, 1593.6, 1479.2, 1346.2, 1232.8, 1058.2 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.48 (t, $J = 2.4$ Hz, 2H), 4.04 (dd, $J = 17.4, 2.4$ Hz, 2H), 4.22 (dd, $J = 17.4, 2.4$ Hz, 2H), 5.85

(s, 2H), 6.66 (d, $J = 8.7$ Hz, 2H), 6.98 (d, $J = 9.0$ Hz, 2H), 7.20–7.30 (m, 4H), 7.44 (d, $J = 2.1$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.5, 69.4, 74.5, 78.8, 111.7, 117.1, 119.7, 124.6, 126.9, 129.4, 130.8, 132.1, 141.0, 145.0. Anal. calcd for $\text{C}_{26}\text{H}_{18}\text{Br}_2\text{ClN}_3$: C, 55.01; H, 3.20; N, 7.40. Found: C, 54.80; H, 3.32; N, 7.18.

2,8-Dibromo-13-(3,5-dimethylphenyl)-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7i). Pale brown solid; mp: 191–194 °C; yield: 267 mg, 95%; IR (neat) 3296.6, 2915.8, 2843.4, 2121.5, 1588.9, 1484.1, 1367.1, 1268.4, 1155.9, 1058.2 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.26 (s, 6H), 2.48 (t, $J = 2.4$ Hz, 2H), 4.05 (dd, $J = 17.4, 2.4$ Hz, 2H), 4.23 (dd, $J = 17.4, 2.4$ Hz, 2H), 5.89 (s, 2H), 6.58–6.68 (m, 5H), 7.24 (dd, $J = 9.0, 2.1$ Hz, 2H), 7.44 (d, $J = 2.1$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 21.7, 40.5, 69.3, 74.2, 79.1, 111.4, 115.8, 117.1, 123.5, 125.1, 130.8, 131.9, 139.0, 141.2, 146.2. Anal. calcd for $\text{C}_{28}\text{H}_{23}\text{Br}_2\text{N}_3$: C, 59.91; H, 4.13; N, 7.49. Found: C, 59.66; H, 4.18; N, 7.53.

2,8-Dibromo-5,11-di(prop-2-yn-1-yl)-13-(m-tolyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7j). Colorless solid; mp: 163–167 °C; yield: 258 mg, 94%; IR (neat) 3283.9, 3235.4, 2966.2, 2845.9, 2114.9, 1601.3, 1489.2, 1353.9, 1255.2, 1190.8, 1091.8 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.30 (s, 3H), 2.47 (t, $J = 2.4$ Hz, 2H), 4.05 (dd, $J = 17.4, 2.4$ Hz, 2H), 4.23 (dd, $J = 17.4, 2.4$ Hz, 2H), 5.89 (s, 2H), 6.66 (d, $J = 8.7$ Hz, 2H), 6.75 (d, $J = 7.5$ Hz, 1H), 6.83 (dd, $J = 7.8, 1.8$ Hz, 1H), 6.90 (s, 1H), 7.14 (t, $J = 7.8$ Hz, 1H), 7.24 (dd, $J = 8.7, 2.1$ Hz, 2H), 7.45 (d, $J = 2.4$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 21.7, 40.5, 69.3, 74.2, 79.0, 111.5, 115.2, 117.1, 119.0, 122.6, 125.0, 129.2, 130.8, 131.9, 139.2, 141.2, 146.3. Anal. calcd for $\text{C}_{27}\text{H}_{21}\text{Br}_2\text{N}_3$: C, 59.25; H, 3.87; N, 7.68. Found: C, 58.88; H, 3.89; N, 7.61.

2,8-Dichloro-5,11-di(prop-2-yn-1-yl)-13-(m-tolyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7k). Colorless solid; mp: 159–161 °C; yield: 213 mg, 93%; IR (neat) 3302.0, 3039.7, 2919.5, 2829.6, 2132.2, 1599.3, 1492.8, 1368.8, 1255.0, 1180.2, 1034.2 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.31 (s, 3H), 2.48 (t, $J = 2.4$ Hz, 2H), 4.04 (dd, $J = 17.4, 2.4$ Hz, 2H), 4.23 (dd, $J = 17.4, 2.4$ Hz, 2H), 5.91 (s, 2H), 6.71 (d, $J = 8.7$ Hz, 2H), 6.75 (d, $J = 7.5$ Hz, 1H), 6.84 (dd, $J = 8.1, 2.1$ Hz, 1H), 6.91 (s, 1H), 7.12 (dd, $J = 9.0, 2.4$ Hz, 2H), 7.15 (t, $J = 7.8$ Hz, 1H), 7.32 (d, $J = 2.4$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 21.8, 40.7, 69.3, 74.3, 79.2, 115.2, 116.8, 118.9, 122.6, 124.3, 124.7, 127.9, 129.1, 129.3, 139.2, 140.8, 146.4. Anal. calcd for $\text{C}_{27}\text{H}_{21}\text{Cl}_2\text{N}_3$: C, 70.75; H, 4.62; N, 9.17. Found: C, 70.37; H, 4.58; N, 9.11.

2,8-Dichloro-13-(3-methoxyphenyl)-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7l). Pale brown solid; mp: 154–157 °C; yield: 188 mg, 79%; IR (neat) 3306.3, 3025.1, 2932.6, 2122.0, 1597.2, 1488.7, 1322.6, 1188.9, 1078.5 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.50 (t, $J = 2.4$ Hz, 2H), 3.78 (s, 3H), 4.04 (dd, $J = 17.4, 2.4$ Hz, 2H), 4.23 (dd, $J = 17.4, 2.4$ Hz, 2H), 5.93 (s, 2H), 6.49 (dd, $J = 8.1, 2.1$ Hz, 1H), 6.62–6.68 (m, 2H), 6.71 (d, $J = 8.7$ Hz, 2H), 7.11 (dd, $J = 8.7, 2.7$ Hz, 2H), 7.17 (t, $J = 8.1$ Hz, 1H), 7.32 (d, $J = 2.4$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.7, 55.3, 69.2, 74.3, 79.1, 104.5, 106.7, 110.6, 117.0, 124.5, 124.6, 127.9, 129.1, 130.1, 140.8, 147.8, 160.6. Anal. calcd for $\text{C}_{27}\text{H}_{21}\text{Cl}_2\text{N}_3\text{O}$: C, 68.36; H, 4.46; N, 8.86. Found: C, 68.02; H, 4.39; N, 8.81.

13-(4-Bromophenyl)-2,8-dichloro-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7m). Pale yellow solid; mp: 183–186 °C; yield: 249 mg, 95%; IR (neat) 3286.2, 3044.5, 2993.2, 1599.6, 1497.2, 1366.1, 1159.1, 1021.5 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.49 (t, $J = 2.4$ Hz, 2H), 4.02 (dd, $J = 17.4, 2.4$ Hz, 2H), 4.22 (dd, $J = 17.4, 2.4$ Hz, 2H), 5.87 (s, 2H), 6.71 (d, $J = 8.7$ Hz, 2H), 6.94 (d, $J = 9.0$ Hz, 2H), 7.12 (dd, $J = 8.7, 2.4$ Hz, 2H), 7.31 (d, $J = 2.4$ Hz, 2H), 7.36 (d, $J = 9.0$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.6, 69.2, 74.5, 78.9, 114.2, 116.8, 120.0, 124.2, 124.6, 127.9, 129.3, 132.3, 140.6, 145.5. Anal. calcd for $\text{C}_{26}\text{H}_{18}\text{BrCl}_2\text{N}_3$: C, 59.68; H, 3.47; N, 8.03. Found: C, 59.32; H, 3.37; N, 8.01.

2,8-Dichloro-13-(3-nitrophenyl)-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7n). Pale yellow solid; mp: 169–171 °C; yield: 213 mg, 87%; IR (neat) 3294.9, 3108.7, 2924.5, 2849.0, 2124.6, 1602.9, 1520.8, 1493.8, 1351.4

1246.2, 1118.0, 1006.5 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.66 (t, $J = 2.4$ Hz, 2H), 4.07 (dd, $J = 17.4$, 2.4 Hz, 2H), 4.30 (dd, $J = 17.4$, 2.4 Hz, 2H), 6.08 (s, 2H), 6.73 (d, $J = 8.7$ Hz, 2H), 7.14 (dd, $J = 8.7$, 2.4 Hz, 2H), 7.34 (d, $J = 2.4$ Hz, 2H), 7.36–7.50 (m, 2H), 7.76 (dt, $J = 7.5$, 1.5 Hz, 1H), 8.04 (t, $J = 1.8$ Hz, 1H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.4, 68.6, 75.1, 78.5, 112.3, 115.9, 117.0, 123.1, 123.9, 124.9, 127.9, 129.5, 130.2, 140.5, 147.3, 149.3. Anal. calcd for $\text{C}_{26}\text{H}_{18}\text{Cl}_2\text{N}_4\text{O}_2$: C, 63.82; H, 3.71; N, 11.45. Found: C, 63.62; H, 3.60; N, 11.32.

5,11-Dibenzyl-13-(*m*-tolyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (7o). Colorless solid; mp: 141–143 °C; yield: 178 mg, 72%; IR (neat) 3033.7, 2969.9, 2916.7, 1599.0, 1489.1, 1449.5, 1353.1, 1248.5, 1099.0 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.22 (s, 3H), 4.65 (s, 4H), 5.67 (s, 2H), 6.60 (d, $J = 8.4$ Hz, 2H), 6.65–6.75 (m, 5H), 7.01–7.11 (m, 5H), 7.25–7.30 (m, 10H); ^{13}C NMR (75 MHz, CDCl_3): δ 21.6, 55.2, 71.1, 114.7, 115.5, 117.8, 119.5, 122.1, 123.1, 127.1, 127.3, 127.7, 128.6, 128.7, 129.0, 138.6, 138.9, 143.7, 147.3. Anal. calcd for $\text{C}_{35}\text{H}_{31}\text{N}_3$: C, 85.16; H, 6.33; N, 8.51. Found: C, 85.20; H, 6.39; N, 8.43.

5,11-Dibenzyl-13-(3,5-dimethylphenyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (7p). Colorless solid; mp: 110–113 °C; yield: 188 mg, 74%; IR (neat) 3019.8, 2911.8, 2842.6, 1595.2, 1490.8, 1448.8, 1357.4, 1119.1, 1050.8 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.17 (s, 6H), 4.62 (d, $J = 16.2$ Hz, 2H), 4.69 (d, $J = 16.2$ Hz, 2H), 5.67 (s, 2H), 6.52 (s, 2H), 6.55 (s, 1H), 6.61 (d, $J = 8.1$ Hz, 2H), 6.69 (td, $J = 7.5$, 0.9 Hz, 2H), 7.03 (td, $J = 8.7$, 1.5 Hz, 2H), 7.10 (dd, $J = 7.5$, 1.5 Hz, 2H), 7.25–7.33 (m, 10H); ^{13}C NMR (75 MHz, CDCl_3): δ 21.6, 55.2, 70.9, 114.8, 116.2, 117.8, 122.9, 123.3, 127.2, 127.5, 127.7, 128.7, 128.8, 138.7, 138.9, 143.8, 147.2. Anal. calcd for $\text{C}_{36}\text{H}_{33}\text{N}_3$: C, 85.17; H, 6.55; N, 8.28. Found: C, 84.90; H, 6.39; N, 8.17.

5,11-Dibenzyl-13-(4-methoxyphenyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (7q). Colorless solid; mp: 168–172 °C; yield: 189 mg, 69%; IR (neat) 3055.4, 3021.9, 2947.5, 1599.8, 1492.2, 1352.3, 1242.3, 1095.9 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 3.75 (s, 3H), 4.62 (s, 4H), 5.51 (s, 2H), 6.61 (d, $J = 8.1$ Hz, 2H), 6.70 (td, $J = 7.5$, 1.2 Hz, 2H), 6.74 (d, $J = 9.0$ Hz, 2H), 6.84 (d, $J = 9.0$ Hz, 2H), 7.01–7.09 (m, 4H), 7.20–7.31 (m, 10H); ^{13}C NMR (75 MHz, CDCl_3): δ 55.4, 55.6, 72.2, 114.5, 114.7, 117.9, 121.2, 123.3, 127.2, 127.4, 127.7, 128.6, 128.8, 138.8, 141.4, 143.8, 155.0. Anal. calcd for $\text{C}_{35}\text{H}_{31}\text{N}_3\text{O}$: C, 82.48; H, 6.13; N, 8.25. Found: C, 82.11; H, 6.02; N, 8.14.

5,11-Dibenzyl-13-(3-chlorophenyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (7r). Colorless solid; mp: 129–131 °C; yield: 183 mg, 71%; IR (neat) 3043.1, 2944.5, 1599.2, 1467.8, 1379.3, 1158.1, 1034.9 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 4.64 (s, 4H), 5.63 (s, 2H), 6.65 (d, $J = 8.1$ Hz, 2H), 6.69–6.75 (m, 3H), 6.87–6.90 (m, 2H), 7.03–7.12 (m, 5H), 7.28–7.36 (m, 10H); ^{13}C NMR (75 MHz, CDCl_3): δ 55.2, 70.7, 114.9, 116.3, 118.1, 118.7, 121.2, 122.7, 127.3, 127.4, 127.7, 128.7, 128.9, 130.1, 134.8, 138.3, 143.5, 148.5. Anal. calcd for $\text{C}_{33}\text{H}_{28}\text{ClN}_3$: C, 79.44; H, 5.49; N, 8.17. Found: C, 79.17; H, 5.41; N, 8.01.

13-Benzyl-2,8-dibromo-5,11-di(prop-2-yn-1-yl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (7s). Colorless solid; mp: 78–80 °C; yield: 222 mg, 81%; IR (neat) 3256.5, 3028.7, 2857.0, 2135.5, 1603.8, 1488.3, 1365.3, 1213.8, 1123.4, 1032.2 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.31 (t, $J = 2.4$ Hz, 2H), 3.62 (d, $J = 12.9$ Hz, 1H), 3.70 (d, $J = 12.9$ Hz, 1H), 4.03 (dd, $J = 17.7$, 2.4 Hz, 2H), 4.17 (dd, $J = 17.7$, 2.4 Hz, 2H), 4.90 (s, 2H), 6.66 (d, $J = 8.7$ Hz, 2H), 7.25 (dd, $J = 8.7$, 2.1 Hz, 2H), 7.29–7.38 (m, 7H); ^{13}C NMR (75 MHz, CDCl_3): δ 40.0, 54.2, 70.7, 73.6, 79.0, 110.6, 115.8, 124.0, 127.7, 128.6, 129.3, 131.1, 131.7, 136.9, 140.5. Anal. calcd for $\text{C}_{27}\text{H}_{21}\text{Br}_2\text{N}_3$: C, 59.25; H, 3.87; N, 7.68. Found: C, 58.96; H, 3.81; N, 7.54.

9-Methylidibenzo[*b,h*][1,6]naphthyridine (8). Pale brown solid; mp: 172–175 °C; yield: 37 mg, 15%; IR (neat) 3057.8, 3006.8, 2919.7, 2857.8, 1600.9, 1491.6, 1382.5, 1294.2, 1036.2 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.62 (s, 3H), 7.75–7.88 (m, 4H), 8.18 (dd, $J = 7.8$, 1.2 Hz, 1H), 8.28 (d, $J = 8.7$ Hz, 1H), 8.75 (s, 1H), 9.31 (dd, $J = 7.8$, 1.5 Hz, 1H), 9.37 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3): δ 21.7, 119.5, 124.1, 125.3, 126.9, 127.0, 127.6, 129.3, 129.4, 130.4, 134.5,

135.9, 136.5, 145.8, 147.0, 148.9, 154.0. Anal. calcd for $\text{C}_{17}\text{H}_{12}\text{N}_2$: C, 83.58; H, 4.95; N, 11.47. Found: C, 83.19; H, 5.07; N, 11.31.

General Procedure for the Synthesis of Triazoles 9. To a solution of epiminodiazocine 7 (0.5 mmol) and benzyl azide (1.5 mmol) in acetonitrile (3 mL) was added CuI (10 mol %) under nitrogen atmosphere at 25 °C and stirred for 4 h. After completion of reaction, the mixture was diluted with CH_2Cl_2 and washed with water. The organic layer was separated, dried over Na_2SO_4 , and solvent removed under reduced pressure. The residue was purified through silica gel column chromatography using 40–60% ethyl acetate in petroleum ether as eluent.

5-((1-Benzyl-1H-1,2,3-triazol-4-yl)methyl)-11-((1-benzyl-1H-1,2,3-triazol-5-yl)methyl)-13-(*p*-tolyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (9a). Colorless solid; mp: 96–98 °C; yield: 295 mg, 90%; IR (neat) 3134.4, 3028.8, 2918.9, 1602.1, 1493.7, 1454.1, 1324.0, 1214.2, 1045.2 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.19 (s, 3H), 4.72 (d, $J = 16.8$ Hz, 2H), 4.82 (d, $J = 16.8$ Hz, 2H), 5.33 (d, $J = 15.0$ Hz, 2H), 5.39 (d, $J = 15.0$ Hz, 2H), 5.61 (s, 2H), 6.55–6.61 (m, 4H), 6.71 (d, $J = 8.7$ Hz, 2H), 6.77 (s, 2H), 6.85 (d, $J = 8.1$ Hz, 2H), 7.01 (td, $J = 7.8$, 1.5 Hz, 2H), 7.06 (d, $J = 7.2$ Hz, 2H), 7.10–7.14 (m, 4H), 7.30–7.36 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3): δ 20.6, 47.0, 54.0, 71.8, 113.8, 117.9, 119.0, 122.4, 122.8, 127.8, 128.6, 128.9, 129.7, 131.0, 134.8, 142.2, 144.7, 146.4. Anal. calcd for $\text{C}_{41}\text{H}_{37}\text{N}_6$: C, 75.09; H, 5.69; N, 19.22. Found: C, 74.83; H, 5.70; N, 18.91. One aromatic carbon is merged with others.

5-((1-Benzyl-1H-1,2,3-triazol-4-yl)methyl)-11-((1-benzyl-1H-1,2,3-triazol-5-yl)methyl)-2,8-dichloro-13-(*m*-tolyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (9b). Colorless solid; mp: 95–98 °C; yield: 323 mg, 89%; IR (neat) 3136.9, 3033.8, 2924.0, 2855.5, 1602.6, 1495.1, 1357.2, 1252.5, 1115.5, 1047.4 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.13 (s, 3H), 4.63 (d, $J = 16.2$ Hz, 2H), 4.80 (d, $J = 16.2$ Hz, 2H), 5.34 (d, $J = 15.0$ Hz, 2H), 5.43 (d, $J = 15.0$ Hz, 2H), 5.61 (s, 2H), 6.51–6.57 (m, 3H), 6.67–6.74 (m, 4H), 6.93 (t, $J = 7.8$ Hz, 1H), 7.00 (dd, $J = 8.7$, 2.4 Hz, 2H), 7.06 (d, $J = 2.4$ Hz, 2H), 7.14–7.17 (m, 4H), 7.32–7.40 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3): δ 21.6, 47.4, 54.1, 71.3, 115.8, 115.9, 119.9, 122.1, 123.0, 123.2, 124.3, 127.4, 127.9, 128.2, 128.7, 129.0, 129.1, 134.6, 139.2, 140.9, 145.6, 146.6. Anal. calcd for $\text{C}_{41}\text{H}_{35}\text{Cl}_2\text{N}_6$: C, 67.95; H, 4.87; N, 17.40. Found: C, 67.67; H, 4.98; N, 17.31.

5-((1-Benzyl-1H-1,2,3-triazol-4-yl)methyl)-11-((1-benzyl-1H-1,2,3-triazol-5-yl)methyl)-13-(4-bromophenyl)-2,8-dichloro-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (9c). Colorless solid; mp: 215–218 °C; yield: 372 mg, 94%; IR (neat) 3302.4, 3039.6, 2921.5, 2852.1, 1599.6, 1493.2, 1369.1, 1255.2, 1180.4, 1034.6 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 4.61 (d, $J = 16.2$ Hz, 2H), 4.77 (d, $J = 16.2$ Hz, 2H), 5.43 (s, 4H), 5.56 (s, 2H), 6.60 (d, $J = 8.4$ Hz, 2H), 6.64 (d, $J = 9.0$ Hz, 2H), 6.87 (s, 2H), 7.02 (dd, $J = 8.7$, 2.4 Hz, 2H), 7.07 (d, $J = 2.4$ Hz, 2H), 7.14 (d, $J = 9.0$ Hz, 2H), 7.18–7.21 (m, 4H), 7.35–7.37 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3): δ 46.8, 54.1, 70.3, 114.1, 116.0, 120.5, 122.2, 123.3, 124.0, 127.4, 127.9, 128.7, 129.1, 132.0, 134.5, 140.7, 145.5. Anal. calcd for $\text{C}_{40}\text{H}_{34}\text{BrCl}_2\text{N}_6$: C, 60.69; H, 4.33; N, 15.93. Found: C, 60.37; H, 4.20; N, 15.81. Two aromatic carbons are merged with others.

Sonogashira Coupling of Epiminodibenzo[*b,f*][1,5]diazocines 7. To a stirred solution of epiminodiazocine 7 (0.5 mmol) in acetonitrile (2 mL) under nitrogen atmosphere were added iodobenzene (1.25 mmol), triethylamine (1.5 mmol), palladium acetate (5 mol %), triphenylphosphine (10 mol %), and copper(I) iodide (10 mol %). Stirring was continued at 25 °C until the completion of the reaction (3 h), and then the reaction mixture was diluted with ethyl acetate and washed with water. The organic layer was separated, dried over Na_2SO_4 , and concentrated under reduced pressure. The residue was purified through silica gel column chromatography using petroleum ether-ethyl acetate (98:2, v/v) as eluent.

5,11-Bis(3-phenylprop-2-yn-1-yl)-13-(*p*-tolyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[*b,f*][1,5]diazocine (7v). Pale brown solid; mp: 132–134 °C; yield: 236 mg, 87%; IR (neat) 3031.3, 2923.9, 2245.0, 1603.5, 1513.4, 1491.5, 1359.8, 1237.6, 1097.5 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.23 (s, 3H), 4.26 (d, $J = 17.1$ Hz, 2H), 4.51 (d, $J = 17.1$ Hz, 2H), 6.10 (s, 2H), 6.78–6.83 (m, 4H), 7.02–7.17 (m,

6H), 7.31–7.36 (m, 6H), 7.40–7.48 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3): δ 20.6, 41.0, 69.7, 85.3, 85.6, 115.2, 118.3, 119.0, 122.8, 123.4, 128.2, 128.4, 128.5, 128.9, 129.8, 130.4, 131.8, 142.6, 144.8. Anal. calcd for $\text{C}_{39}\text{H}_{31}\text{N}_3$: C, 86.47; H, 5.77; N, 7.76. Found: C, 86.17; H, 5.69; N, 7.71.

2,8-Dibromo-5,11-bis(3-phenylprop-2-yn-1-yl)-13-(p-tolyl)-5,6,11,12-tetrahydro-6,12-epiminodibenzo[b,f][1,5]diazocine (7w). Pale brown solid; mp: 93–95 °C; yield: 294 mg, 84%; IR (neat) 3042.1, 2918.8, 2221.4, 1597.5, 1477.7, 1367.3, 1247.9, 1161.0, 1045.3 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3): δ 2.24 (s, 3H), 4.28 (d, $J = 17.4$ Hz, 2H), 4.44 (d, $J = 17.4$ Hz, 2H), 5.95 (s, 2H), 6.72 (d, $J = 8.7$ Hz, 2H), 6.99 (d, $J = 8.4$ Hz, 2H), 7.05 (d, $J = 8.4$ Hz, 2H), 7.25 (dd, $J = 8.7, 2.4$ Hz, 2H), 7.31–7.35 (m, 6H), 7.43–7.47 (m, 4H), 7.53 (d, $J = 2.4$ Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 19.4, 40.1, 68.7, 83.2, 84.8, 110.0, 115.6, 117.4, 121.3, 123.9, 127.2, 127.5, 128.8, 129.7, 130.2, 130.7, 140.1, 142.9. Anal. calcd for $\text{C}_{39}\text{H}_{29}\text{Br}_2\text{N}_3$: C, 66.97; H, 4.18; N, 6.01. Found: C, 66.62; H, 4.12; N, 5.88. One aromatic carbon is merged with others.

■ ASSOCIATED CONTENT

■ Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.6b01764.

Copies of ^1H and ^{13}C NMR spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: vsridharan@scbt.sastra.edu; vesridharan@gmail.com.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Financial support from the Department of Science and Technology, DST (nos. SB/FT/CS-006/2013 and INT/AUA/BMWF/P-26/2015), and the Council of Scientific and Industrial Research, CSIR (no. 02(0219)/14/EMR-II) is gratefully acknowledged. J.C.M. acknowledges financial support from MINECO, grant CTQ2012-33272-BQU. This article is dedicated to Professor S. Sivasubramanian on the occasion of his 73rd birthday.

■ REFERENCES

- (1) For a review on Tröger's bases, see: Dolenský, B.; Havlík, M.; Král, V. *Chem. Soc. Rev.* **2012**, *41*, 3839–3858.
- (2) For selected examples of applications of Tröger's bases, see: (a) Reddy, M. B.; Shailaja, M.; Manjula, A.; Premkumar, J. R.; Sastry, G. N.; Sirisha, K.; Sarma, A. V. S. *Org. Biomol. Chem.* **2015**, *13*, 1141–1149. (b) Yang, Z.-Z.; Zhang, H.; Yu, B.; Zhao, Y.; Ji, G.; Liu, Z. *Chem. Commun.* **2015**, *51*, 1271–1274. (c) Banerjee, S.; Bright, S. A.; Smith, J. A.; Burgeat, J.; Martinez-Calvo, M.; Williams, D. C.; Kelly, J. M.; Gunnlaugsson, T. *J. Org. Chem.* **2014**, *79*, 9272–9283. (d) Boyle, E. M.; Comby, S.; Molloy, J. K.; Gunnlaugsson, T. *J. Org. Chem.* **2013**, *78*, 8312–8319. (e) Baldeyrou, B.; Tardy, C.; Bailly, C.; Colson, P.; Houssier, C.; Charmantray, F.; Demeunynck, M. *Eur. J. Med. Chem.* **2002**, *37*, 315–322.
- (3) (a) For biological applications, see: Paul, A.; Maji, B.; Misra, S. K.; Jain, A. K.; Muniyappa, K.; Bhattacharya, S. *J. Med. Chem.* **2012**, *55*, 7460–7471. (b) Johnson, R. A.; Gorman, R. R.; Wnuk, R. J.; Crittenden, N. J.; Aiken, J. W. *J. Med. Chem.* **1993**, *36*, 3202–3206.
- (4) (a) Jarzebski, A.; Bannwarth, C.; Tenten, C.; Benkhäuser, C.; Schnakenburg, G.; Grimme, S.; Lützen, A. *Synthesis* **2015**, *47*, 3118–3132. (b) Dawaigher, S.; Månsson, K.; Ascic, E.; Artacho, J.; Mårtensson, R.; Loganathan, N.; Wendt, O. F.; Harmata, M.; Snieckus, V.; Wärnmark, K. *J. Org. Chem.* **2015**, *80*, 12006–12014. (c) Jensen, J.; Tejler, J.; Wärnmark, K. *J. Org. Chem.* **2002**, *67*, 6008–6014. (d) Sathyanarayana, A.; Suresh, P.; Prabhusankar, G. *J. Organomet.*

Chem. **2012**, *720*, 46–51. (e) Sharma, A.; Besnard, B.; Guénée, L.; Lacour, J. *Org. Biomol. Chem.* **2012**, *10*, 966–969. (f) Artacho, J.; Ascic, E.; Rantanen, T.; Wallentin, C.-J.; Dawaigher, S.; Bergquist, K.-E.; Harmata, M.; Snieckus, V.; Wärnmark, K. *Org. Lett.* **2012**, *14*, 4706–4709.

(5) Su, C.; Tandiana, R.; Balapanuru, J.; Tang, W.; Pareek, K.; Nai, C. T.; Hayashi, T.; Loh, K. P. *J. Am. Chem. Soc.* **2015**, *137*, 685–690.

(6) (a) Frank, K. E.; Aubé, J. *J. Org. Chem.* **2000**, *65*, 655–666. (b) Frank, K. E.; Aubé, J. *Tetrahedron Lett.* **1998**, *39*, 7239–7242.

(7) (a) Leganza, A.; Bezze, C.; Zonta, C.; Fabris, F.; De Lucchi, O.; Linden, A. *Eur. J. Org. Chem.* **2006**, *2006*, 2987–2990. (b) Pettersson, B.; Bergman, J.; Svensson, P. H. *Tetrahedron* **2013**, *69*, 2647–2654.

(8) (a) Sridharan, V.; Ribelles, P.; Ramos, M. T.; Menéndez, J. C. *J. Org. Chem.* **2009**, *74*, 5715–5718. (b) Albert, A.; Yamamoto, H. *J. Chem. Soc. B* **1966**, 956–963.

(9) (a) Molina, P.; Arques, A.; Tárraga, A.; del Rosario Obón, M.; Foces-Foces, C.; Jagerovic, N.; Elguero, J. *Tetrahedron* **1998**, *54*, 997–1004. (b) Molina, P.; Arques, A.; Cartagena, I.; Obón, R. *Tetrahedron Lett.* **1991**, *32*, 2521–2524.

(10) Ukhin, L. Y.; Kuźmina, L. G. *Russ. Chem. Bull.* **2004**, *53*, 2262–2268.

(11) Mao, D.; Tang, J.; Wang, W.; Wu, S.; Liu, X.; Yu, J.; Wang, L. *J. Org. Chem.* **2013**, *78*, 12848–12854.

(12) Muthukrishnan, I.; Vinoth, P.; Vivekanand, T.; Nagarajan, S.; Maheswari, C. U.; Menéndez, J. C.; Sridharan, V. *J. Org. Chem.* **2016**, *81*, 1116–1124.

(13) Cheng, H.; Wan, J.; Lin, M.-I.; Liu, Y.; Lu, X.; Liu, J.; Xu, Y.; Chen, J.; Tu, Z.; Cheng, Y.-Z. E.; Ding, K. *J. Med. Chem.* **2012**, *55*, 2144–2153.

(14) Bai, S.; Li, S.; Xu, J.; Peng, X.; Sai, K.; Chu, W.; Tu, Z.; Zeng, C.; Mach, R. H. *J. Med. Chem.* **2014**, *57*, 4239–4251.

(15) Bertrand, H. C.; Schaap, M.; Baird, L.; Georgakopoulos, N. D.; Fowkes, A.; Thiollier, C.; Kachi, H.; Dinkova-Kostova, A. T.; Wells, G. *J. Med. Chem.* **2015**, *58*, 7186–7194.

(16) For recent reviews, see: (a) Tiwari, V. K.; Mishra, B. B.; Mishra, K. B.; Mishra, N.; Singh, A. S.; Chen, X. *Chem. Rev.* **2016**, *116*, 3086–3240. (b) Kacprzak, K.; Skiera, I.; Piasecka, M.; Paryzek, Z. *Chem. Rev.* **2016**, *116*, 5689–5743.

(17) Levesque, P.; Fournier, P.-A. *J. Org. Chem.* **2010**, *75*, 7033–7036.